


| G | Study | Total | Correlation | COR | 95%-CI | Weight (fixed) | Weight (random) |
|---|--------------------------------------|----------------------|---|-------|---------------|-------------------|-----------------|
| | Tubic | 46 | | 0.19 | [-0.09; 0.47] | 44.1% | 46.2% |
| | Tubic | 62 | | -0.05 | [-0.30; 0.20] | 55.9% | 53.8% |
| | Fixed effect model | 108 | | 0.06 | [-0.13; 0.24] | 100.0% | |
| | Random effects model | | | 0.06 | [-0.17; 0.30] | | 100.0% |
| | Heterogeneity: $I^2 = 36\%$, τ | $\epsilon^2 = 0.010$ | | | | | |
| | | - | -0.4 0 0.2 0.4 0.6 0.8 1 | | | | |
| | | | Correlation tOC and BMI in HS | | | | |

| H | Study | Total | Correlation | COR | 95%-CI | Weight (fixed) | Weight (random) |
|---|--|----------------------------|--|-------------|--------------------------------|-------------------|--------------------|
| | Tubic Tubic | 46 62 | | | [0.12; 0.62] [-0.02; 0.46] | 47.3% 52.7% | 47.3% 52.7% |
| | Fixed effect model Random effects mode Heterogeneity: $I^2 = 0\%$, τ^2 | $p^2 = 0, p = 0.40$ -0.4 | 0 0.2 0.4 0.6 0.8 ion tOC and WC in HS | 0.29 | [0.12; 0.46] [0.12; 0.46] | 100.0% | 100.0% |

| Study | Total (| Correlation | COR | 95%-CI | Weight (fixed) | Weight (random) |
|--|----------------------|----------------------|------|--------------|-------------------|--------------------|
| Tubic | 46 | - • - | 0.47 | [0.24; 0.70] | 50.5% | 50.4% |
| Tubic | 62 | | 0.29 | [0.06; 0.52] | 49.5% | 49.6% |
| Fixed effect model | 108 | | 0.38 | [0.22; 0.54] | 100.0% | |
| Random effects model | | | 0.38 | [0.20; 0.56] | | 100.0% |
| Heterogeneity: $I^2 = 16\%$, τ^2 | p = 0.0026, p = 0.28 | | | | | |
| | -0.4 | 0 0.2 0.4 0.6 0.8 1 | 1 | | | |
| | Correlation | tOC and Height in HS | 3 | | | |



| K | Study | Total | Correlation | | COR | 95%-CI | Weight (fixed) | Weight (random) |
|---|--|---------|--------------------|------------|-------|----------------|-------------------|--------------------|
| | Faienza | 186 | : - | | -0.16 | [-0.31; -0.02] | 68.0% | 54.1% |
| | Guarneri | 62 | - | | -0.43 | [-0.63; -0.23] | 32.0% | 45.9% |
| | Fixed effect model | 248 | | | -0.25 | [-0.37; -0.13] | 100.0% | |
| | Random effects model Heterogeneity: $I^2 = 77\%$, τ^2 | | = 0.04 | | | [-0.55; -0.03] | | 100.0% |
| | 110.010 gonony. 1 = 71 70, 0 | -1 -0.8 | | 0.2 0.4 | | | | |
| | | Correla | tion tOC and Hb | A1c in T1I |) | | | |