

Supplementary Material

FBXO43 increases CCND1 stability to promote hepatocellular carcinoma cell proliferation and migration

Chun-Ming Li¹, Jie Zhang², Wu Wu¹, Zhu Zhu¹, Feng Li³, Di Wu², Xiao-Jun Wang^{2*}, Chuan-Ming Xie^{2*}, Jian-Ping Gong^{1*}

¹Department of Hepatobiliary Surgery, The Second Affiliated Hospital of Chongqing Medical University, Chongqing 400010, China

²Key Laboratory of Hepatobiliary and Pancreatic Surgery, Institute of Hepatobiliary Surgery, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing 400038, China

³Department of Hepatobiliary Surgery, The Third Affiliated Hospital of Chongqing Medical University, Chongqing 401120, China

* Correspondence:

Jian-Ping Gong, E-mail: gongjianping11@126.com

Chuan-Ming Xie, E-mail: cmxie@tmmu.edu.cn; chuanming506@126.com; Tel: +86 23 68765809, Fax: +86 23 68765809.

Xiao-Jun Wang, E-mail: yh830@163.com.

1 Supplementary Figures and Tables

1.1 Supplementary Fig. 1



Supplementary Fig. 1. FBXO43 regulates CCND1 without the involvement of lysosomes

The Western blots to detect the level of CCND1 in siCtrl- or siFBXO43-treated Huh7 cells with or without a lysosome inhibitor (50 μ M chloroquine).