

Table 1. Clinical characteristics of the whole cohort of MIS-C patients and of the groups of patients with ESS and with normal thyroid function tests (no ESS)

Variable	Overall - n.42(%)	ESS - n.36 (%)	No ESS - n.6 (%)	<i>p value</i>
Age (median)	9 years (0.7 – 17.7)	9 years (0.7 – 17.7)	8 years (2.2 – 13.9)	0.774
Sex no. (%)				
Male	29 (69)	26 (72)	3 (50)	0.353
Race no. (%)				
Caucasian	37 (88)	32 (89)	5 (83)	0.557
African	4 (9)	3 (8)	1 (16)	0.474
Other	1 (1)	1 (3)		> 0.999
Comorbidity no. (%)	4 (9)	2 (4)	2 (33)	0.091
Clinical features no. (%)				
Gastrointestinal	39 (92)	33 (92)	6 (100)	> 0.999
Abdominal pain	28 (67)	23 (64)	5 (83)	0.645
Vomit	19 (45)	18 (50)	1 (16)	0.197
Diarrhea	17 (40)	14 (39)	3 (50)	0.672
Cardiovascular	34 (81)	30 (83)	4 (66)	0.319
Hypotension	14 (33)	13 (36)	1 (16)	0.645
↑ Troponin	23 (55)	20 (55)	3 (50)	> 0.999
↑ BNP	25 (59)	23 (64)	2 (33)	0.202
Coronary involvement	7 (17)	7 (19)	0 (0)	0.567
EF ≤55	16 (38)	14 (39)	2 (33)	> 0.999
Skin	28 (67)	25 (69)	3 (50)	0.383
Renal	8 (19)	8 (22)	0 (0)	0.576
Central Nervous System	4 (9)	4 (11)	0 (0)	> 0.999
ICU admission no. (%)	6 (14)	6 (17)	0 (0)	0.569

p value is significant when <0.05