

Supplementary Material

Predicting responses of an Adélie penguin population to future sea ice change in the Ross Sea

Tosca Ballerini ^{1,2,3,*}, Giacomo Tavecchia ⁴, Francesco Pezzo ^{1,5}, Stéphanie Jenouvrier ^{6,7}, Silvia Olmastroni ^{1,8}

¹Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente, Università degli Studi di Siena, Italia

²Université de Toulon, CNRS/INSU, IRD, Mediterranean Institute of Oceanography (MIO), UM 110, 83957 La Garde, France

³ Aix Marseille Université, CNRS/INSU, IRD, Mediterranean Institute of Oceanography (MIO), UM 110, 13288 Marseille, France

⁴Institut Mediterrani d'Estudis Avançats (IMEDEA), CSIC-UIB, 07190 Esporles, Mallorca, Spain

⁵Museo di Storia Naturale della Maremma, Strada Corsini 5, 58100 Grosseto, Italy

⁶Biology Dept., MS-34, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

⁷Centre d'Etudes Biologiques de Chizé, Centre National de la Recherche Scientifique, UPR1934, F-79360 Villiers en Bois, France

⁸Museo Nazionale dell'Antartide, Via del Laterino 8, 53100 Siena, Italy

*** Correspondence:** Tosca Ballerini ² Université de Toulon, CNRS/INSU, IRD, Mediterranean Institute of Oceanography (MIO), UM 110, 83957 La Garde, France
toscaballerini@gmail.com, tosca.ballerini@mio.osupytheas.fr





Figure A1. Comparison of Ross Sea sea ice extent values derived from satellite observations (red line) and Ross Sea sea ice values derived from the AOGCMs.





Figure A2. Comparison of Ross Sea sea ice extent values derived from satellite observations (red line) and Ross Sea sea ice values derived from the AOGCMs.