

Table 1. Agronomic practices in field experiments to determine seed-to-seedling transmission of *Ascochyta pisi* in CDC Patrick pea at Outlook, Milden and Saskatoon, Canada, in 2012-2014.

Agronomic practices	2012		2013		2014	
	Outlook	Saskatoon	Milden	Saskatoon	Milden	Saskatoon
Pre-seeding herbicide (fall application)	69.19 mL ha ⁻¹ ¹ Pursuit (22.87% imazethapyr) 29.16 kg ha ⁻¹ Edge (5% ethalfluralin)	69.19 mL ha ⁻¹ ¹ Pursuit (22.87% imazethapyr) 29.16 kg ha ⁻¹ Edge (5% ethalfluralin)	29.16 kg ha ⁻¹ Edge (5% ethalfluralin)	29.16 kg ha ⁻¹ Edge (5% ethalfluralin)	29.16 kg ha ⁻¹ Edge (5% ethalfluralin)	69.19 mL ha ⁻¹ ¹ Pursuit (22.87% imazethapyr) 29.16 kg ha ⁻¹ Edge (5% ethalfluralin)
Seeding	25 May	30 May	7 May	11 May	10 May	12 May
Post-seeding herbicide	None	25 June 2.2. L ha ⁻¹ Basagran forte (480g L ⁻¹ ¹ Bentazon)	11 June 0.59 L ha ⁻¹ Axial (50 g L ⁻¹ Pinoxaden)	11 June 0.59 L ha ⁻¹ Axial (50 g L ⁻¹ Pinoxaden)	6 June 0.59 L ha ⁻¹ Axial (50 g L ⁻¹ ¹ Pinoxaden)	6 June 0.59 L ha ⁻¹ Axial (50 g L ⁻¹ Pinoxaden)
Emergence count	21 June	11 July	11 June 185 mL ha ⁻¹ ¹ Centurion (240g L ⁻¹ ¹ Clethodim)	11 June 185 mL ha ⁻¹ ¹ Centurion (240g L ⁻¹ ¹ Clethodim)	11 June 185 mL ha ⁻¹ ¹ Centurion (240g L ⁻¹ ¹ Clethodim)	11 June 185 mL ha ⁻¹ ¹ Centurion (240g L ⁻¹ ¹ Clethodim)
1st Disease rating	21 June	11 July	12 June	14 June	10 June	10 June
2nd Disease rating	26 July	11 July	12 July	10 July	25 June	26 June
3rd Disease rating	26 July	26 July	30 July	13 August	15 July	16 July
Desiccant application	9 August	13 August	14 August	21 August	30 July	26 July
Harvest date	15 August 2.5 L ha ⁻¹ Reglone (240 g L ⁻¹ Diquat)	17 August 2.5 L ha ⁻¹ Reglone (240 g L ⁻¹ Diquat)	23 August 2.5 L ha ⁻¹ Reglone (240 g L ⁻¹ Diquat)	16 August 2.5 L ha ⁻¹ Reglone (240 g L ⁻¹ Diquat)	18 August 2.5 L ha ⁻¹ Reglone (240 g L ⁻¹ Diquat)	18 August 2.5 L ha ⁻¹ Reglone (240 g L ⁻¹ Diquat)
Harvest date	17 August	29 August	30 August	29 August	20 August	25 August