

Supplementary Table 4

Clinical characteristics of patients with delayed graft function. N.S.: Non-significant. The variables that were selected for the multivariate analysis are marked in bold.

Condition.	Delayed graft function		Rest of patients		p
	N / mean	% / se	N / mean	% / se	
Sex (women)	72	(37.7%)	233	(42.4%)	N.S.
Age (years)	52.5	±0.9	48	±0.6	<0.001
Donor age (years)	50.5	±1.2	44	±0.8	<0.001
Time on dialysis (months)	36.2	±2.8	32.6	±1.9	0.004
Body mass index	26.2	±0.4	24.7	±0.2	<0.001
Diabetes mellitus	40	(20.9%)	83	(15.1%)	N.S.
Type 1 Diabetes	6	(3.1%)	25	(4.6%)	N.S.
Type 2 Diabetes	17	(8.9%)	29	(5.3%)	N.S.
Dyslipidemia	95	(49.7%)	263	(47.9%)	N.S.
Hypertension	153	(80.1%)	388	(70.7%)	0.015
Patients IgA aB2GP1 positive	103	(53.9%)	185	(33.7%)	<0.001
Causes CKD					
Chronic glomerulonephritis	52	(27.2%)	158	(28.8%)	N.S.
Interstitial kidney disease	21	(11%)	79	(14.4%)	N.S.
Nephroangiosclerosis	21	(11%)	39	(7.1%)	N.S.
Polycystic kidney disease	28	(14.7%)	90	(16.4%)	N.S.
Diabetes mellitus	17	(8.9%)	39	(7.1%)	N.S.
Unknown	26	(13.6%)	86	(15.7%)	N.S.
Other	26	(13.6%)	58	(10.6%)	N.S.
Transplant-associated factors					
Previous kidney transplant	36	(18.8%)	74	(13.5%)	N.S.
PRA at time of transplant >50%	4	(2.1%)	20	(3.6%)	N.S.
Historical PRA >50%	18	(9.4%)	55	(10%)	N.S.
Cold ischemia time (hours)	20.5	±0.4	19.3	±0.2	0.001