

Table S1. Minimum inhibitory concentration (MIC- mg/mL) of azoles drugs for *Aspergillus fumigatus* strains.

Strain/parameter	Voriconazole	Posaconazole	Itraconazole
<i>Aspergillus fumigatus</i> CEA10 ^S	0.50	1.00	0.50
<i>A. fumigatus</i> Af293 ^S	0.50	1.00	1.00
<i>A. fumigatus</i> AfS35 ^S	0.50	1.00	0.50
<i>A. fumigatus</i> CEA17 ^S	0.25	0.50	0.50
<i>A. fumigatus</i> CYP-15-75 ^{AR}	4.00	1.00	>8.00
<i>A. fumigatus</i> CYP-15-91 ^{AR}	4.00	1.00	>16.00
<i>A. fumigatus</i> CYP-15-93 ^{AR}	4.00	1.00	>8.00
<i>A. fumigatus</i> CYP-15-106 ^{AR}	4.00	0.50	>8.00
<i>A. fumigatus</i> CYP-15-108 ^{AR}	8.00	1.00	>16.00
<i>A. fumigatus</i> CYP-15-109 ^{AR}	>8.00	1.00	2.00
<i>A. fumigatus</i> CYP-15-115 ^{AR}	4.00	0.50	>16.00
<i>A. fumigatus</i> CYP-15-117 ^{AR}	0.25	0.50	>16.00

S= susceptible, *AR*=Azole-resistant

Table S2. Minimum inhibitory concentration (MIC- mg/mL) of amphotericin B, fluconazole and caspofungin for susceptible *Candida* spp.

Strains	Amphotericin B	Fluconazole ^a	Caspofungin
<i>Candida haemulonii</i> sensu stricto 768	0.12	0.12	0.12
<i>C. haemulonii</i> sensu stricto 3834A	0.50	0.50	0.06
<i>C. haemulonii</i> sensu stricto 6083	0.50	0.25	0.06
<i>C. haemulonii</i> sensu stricto 145/18	0.50	0.12	0.03
<i>C. albicans</i> ATCC 90025	0.06	2.00	0.12
<i>C. albicans</i> 16	0.06	0.12	0.12
<i>C. albicans</i> 83	0.06	0.25	0.12
<i>C. albicans</i> 106	0.06	0.12	0.06
<i>C. albicans</i> 123	0.06	0.12	0.06
<i>C. glabrata</i> ATCC 90030	0.25	0.25	0.06
<i>C. glabrata</i> 614	0.06	0.12	0.06
<i>C. glabrata</i> 636	0.06	0.50	0.12
<i>C. glabrata</i> 558	0.06	0.25	0.12
<i>C. parapsilosis</i> ATCC 22019	0.06	0.50	0.25
<i>C. krusei</i> ATCC 6558	0.06	4.00	0.25
<i>C. auris</i> CBS 10913	0.50	2.00	0.06

a- MIC endpoint considering 50% of growth inhibition