

Supplementary Table 1. Primers sequences for the specific genes used in this study.

Genes	Primer Sequence		Reference
<i>Rn18s</i>	Forward:	5'-TTCGTATTGCGCCGCTAGA-3'	dos Santos, A. G., Mendes, É. A., de Oliveira, R. P., Faria, A. M. C., de Sousa, A. O., Pirovani, C. P., et al. (2017). Trichoderma asperelloides Spores Downregulate dectin1/2 and TLR2 Receptors of Mice Macrophages and Decrease Candida parapsilosis Phagocytosis Independent of the M1/M2 Polarization. <i>Front. Microbiol.</i> 8, 1–13. doi:10.3389/fmicb.2017.01681.
	Reverse:	5'-CTTTCGCTCTGGTCCGTCTT-3'	
<i>Clec7a</i>	Forward:	5'-GAACCACAAGCCCACAGAAT-3'	
	Reverse:	5'-CATGGCCCTTCACTCTGATT-3'	
<i>Tlr2</i>	Forward:	5'- GTGGTACCTGAGAATGATGTGGG-3'	
	Reverse:	5'- GTTAATTAAGTCAGGAAGTGGGTG-3'	
<i>Tlr4</i>	Forward:	5'-CTGGGTGAGAAATGAGCTGG-3'	
	Reverse:	5'-GATACAATTCCACCTGCTGCC-3'	
<i>Rn18s</i> : 18S RNA ribosomal RNA; <i>Clec7a</i> : receptor dectin-1; <i>Tlr2</i> : toll-like receptor 2; <i>Tlr4</i> : toll-like receptor 4.			