APPENDIX Quezada et al, 2020

Supplementary Figures







Supplementary figure 2. MCU and TOM20 bands on PVDF membranes. In a) and c) the MCU and TOM20 band (n = 4) is observed. In b) and d) the membranes are observed with Ponceau red (n=4). The samples were taken from complete muscles of the fdb and soleus.



Supplementary figure 3. MICU1 and TOM20 bands on PVDF membranes. In a), d) and g) the MICU band is observed in the membranes (n = 5). In b), e) and h) the TOM20 band is observed in the membranes (n = 5). In c), f) and i) the membranes are observed with Ponceau red (n = 5). The samples were taken from the complete muscles of the *fdb* and soleus.



Supplementary figure 4. Pooled data indicates that low-frequency electrical stimulation (ES) transiently reduces mRNA levels of MCU complex in skeletal muscle fibers. Muscle fibers isolated from the *Flexor Digitorium Brevis* (*fdb*) muscle were stimulated with 270 pulses, 0.3 ms each (a-d) at 20 Hz. mRNA levels of MCU (n=22), MICU1 (n=15), MICU2 (n=16) and EMRE (n=17) decreased 1h post ES. The plates with isolated *fdb* fibers for each condition come from the same animal. For (a) and (b) using the Mann-Whitney test was applied. Values are presented as mean \pm S.E.M. (n = 4-6). 18s was used as a normalizer. *, p < 0.05; **, p < 0.01; ****, p < 0.001.