

	<i>In-hospital mortality, n=166 of 1169 patients</i>			<i>Three-month mortality (N=147, only survivors of the acute hospital stay were considered)</i>		
	<i>Multivariable logistic regression analyses</i>					
	OR	95%-CI	<i>P</i>	OR	95%-CI	<i>P</i>
Age (per year increasing)	1.02	1.01-1.05	0.016	1.04	1.02-1.07	<0.001
NIHSS at admission (per point increasing)	1.06	1.03-1.09	<0.001	1.11	1.07-1.15	<0.001
Preexisting functional impairment (pmRS>1 vs. ≤1)	0.79	0.48-1.32	0.368	2.53	1.51-4.23	<0.001
PC-AKI (vs. no PC-AKI)	3.63	1.35-9.74	0.011	2.38	0.63-8.99	0.201
Posterior circulation stroke (vs. anterior circulation stroke)	3.31	1.82-6.04	<0.001	0.36	0.13-1.03	0.056
Failed recanalization (TICI 0-2a vs. 2b-3)	1.73	1.02-2.94	0.042	3.12	1.79-5.46	<0.001
sICH vs. no sICH	4.79	2.20-10.42	<0.001	2.95	0.99-8.79	0.052

Supplementary Table 6: Multivariable logistic regression analysis for in-hospital mortality and three-month mortality in patients without baseline renal impairment (eGFR at admission <60mL/min/1.73 m²)

NIHSS, National Institutes of Health Stroke Scale; pmRS, premorbid modified Rankin Scale; PC-AKI, post-contrast-AKI; TICI, Thrombolysis In Cerebral Infarction; sICH, symptomatic intracerebral hemorrhage. P-values ≤0.5 are displayed in bold.