

	<i>In-hospital mortality, n=127 of 1053 patients, only anterior-circulation stroke</i>			<i>3-month mortality (N=138, only survivors of the acute hospital stay were considered), only anterior-circulation stroke</i>		
	<i>Multivariable logistic regression analyses</i>					
	OR	95%-CI	<i>P</i>	OR	95%-CI	<i>P</i>
Age (per year increasing)	1.04	1.02-1.06	<0.001	1.04	1.02-1.06	<0.001
NIHSS at admission (per point increasing)	1.05	1.03-1.08	<0.001	1.09	1.06-1.13	<0.001
Preexisting functional impairment (pmRS>1 vs. ≤1)	0.98	0.65-1.47	0.904	4.55	1.52-13.59	0.007
Baseline renal impairment (eGFR<60 vs. ≥60 at admission)	1.22	0.81-1.84	0.334	1.22	0.800-1.86	0.355
PC-AKI (vs. no PC-AKI)	2.26	0.82-6.20	0.114	2.51	1.67-3.78	<0.001
Failed recanalization (TICI 0-2a vs. 2b-3)	2.07	1.36-3.16	0.001	2.80	1.78-4.38	<0.001
sICH vs. no sICH	2.97	1.51-5.83	0.002	2.96	1.90-6.75	0.010

Supplementary Table 8: Multivariable logistic regression analysis for in-hospital mortality and 3-month mortality in patients with anterior circulation stroke

NIHSS, National Institutes of Health Stroke Scale; pmRS, premorbid modified Rankin Scale; eGFR, estimated glomerular filtration rate (mL/min/1.73 m²); PC-AKI, post-contrast-AKI; TICI, Thrombolysis In Cerebral Infarction; sICH, symptomatic intracerebral hemorrhage. P-values ≤0.5 are displayed in bold.