Figure S5: Pooled SMD about social functioning for patients with first-episode shizophrenia Std. Mean Difference Std. Mean Difference Experimental Control SD Total Mean SD Total Weight IV. Random, 95% CI Study or Subgroup Mean IV. Random, 95% CI 9.1.1 first-episode Liu L. et al 2018 19.36 1.72 45 21.57 2.02 20.5% -1.17 [-1.62, -0.72] Liu SS. et al 2017 19.32 1.76 58 21.53 2.01 20.6% -1.18 [-1.61, -0.74] -2.89 [-3.46, -2.32] Shen YR, et al 2016 4.32 1.06 81 7.47 1.14 20.0% Subtotal (95% CI) 184 115 61.1% -1.73 [-2.75, -0.71] Heterogeneity: $Tau^2 = 0.75$: $Chi^2 = 26.84$. df = 2 (P < 0.00001): $I^2 = 93\%$ Test for overall effect: Z = 3.33 (P = 0.0009) 9.1.2 Not sepecific Yang J. et al 2017 -21.44 4.57 41 -7.93 1 44 19.0% -3.95 [-4.71, -3.19] Ying C. et al 2020 -72.59 11.02 28 -57.7 17.17 19.9% -1.04 [-1.63, -0.45] Subtotal (95% CI) 69 38.9% -2.48 [-5.34, 0.37] Heterogeneity: $Tau^2 = 4.12$: $Chi^2 = 35.41$. df = 1 (P < 0.00001): $I^2 = 97\%$ Test for overall effect: Z = 1.71 (P = 0.09) Total (95% CI) 253 179 100.0% -2.02 [-3.00, -1.03] Heterogeneity: $Tau^2 = 1.18$: $Chi^2 = 66.56$. df = 4 (P < 0.00001): $I^2 = 94\%$ Test for overall effect: Z = 4.02 (P < 0.0001) Favours [experimental] Favours [control] Test for subgroup differences: $Chi^2 = 0.24$ df = 1 (P = 0.63) $I^2 = 0\%$