## Figure S7: Pooled SMD about family burden for patients with first-episode schizophrenia Experimental Control Std. Mean Difference Std. Mean Difference

	Expointional			00114101			Otal Moall Dillorolloo		otal mount pintoronico	
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Random, 95% CI	IV, Random, 95% CI	
3.2.1 First-episode										
Liu L. et al 2018	13.2	6.43	45	20.29	6.6	45	49.3%	-1.08 [-1.52, -0.64]	<del></del>	
Liu SS. et al 2017	14.37	3.68	58	20.12	6.11	40	50.7%	-1.18 [-1.62, -0.75]	<del></del>	
Subtotal (95% CI)			103			85	100.0%	-1.13 [-1.44, -0.82]	•	
Heterogeneity: Tau <sup>2</sup> =	0.00; Ch	ni² = 0.	11, df =	1 (P =	0.74);	$I^2 = 0\%$	, •			
Test for overall effect:	Z = 7.13	(P < 0	.00001	)						
3.2.2 Not specific										
Xiang JF. et al 2019	13.22	3.44	45	19.31	5.84	45	100.0%	-1.26 [-1.71, -0.81]	<b>—</b>	
Subtotal (95% CI)			45			45	100.0%	-1.26 [-1.71, -0.81]	•	
Heterogeneity: Not ap	plicable									
Test for overall effect:	Z = 5.44	(P < 0	.00001	)						
				•						
								-	-2 -1 0 1 2	
Test for subgroup diffe	rences.	Chi² =	0.21 d	f = 1 (P	= 0.65	5) I <sup>2</sup> = (	0%		Favours [experimental] Favours [control]	