Figure S9: Pooled SMD about quality of life for patients with first-episode schizophrenia

Study or Subgroup	Experimental Mean	Experimental SD	Experimental Total	Control Mean	Control SD	Control Total	Weight	Std. Mean Difference IV, Random, 95% CI
4.2.1 first-episode								
Liu L. et al 2018	-16.35	2.73	45	-12.55	2.21	45	47.9%	-1.52 [-1.99, -1.05]
Liu SS. et al 2017	-15.51	3.03	58	-12.54	2.58	40	52.1%	-1.03 [-1.46, -0.60]
Subtotal (95% CI)			**103**			**85**	**100.0%**	**-1.26 [-1.74, -0.79]**
Heterogeneity: Tau² = 0.06; Chi² = 2.23, df = 1 (P = 0.14); I² = 55%								
Test for overall effect: Z = 5.22 (P < 0.00001)								
4.2.2 not specific								
Ying C. et al 2020	38.95	16.95	28	42	22.33	23	100.0%	-0.15 [-0.71, 0.40]
Subtotal (95% CI)			**28**			**23**	**100.0%**	**-0.15 [-0.71, 0.40]**
Heterogeneity: Not applicable								
Test for overall effect: Z = 0.55 (P = 0.59)								



Test for subgroup differences: Chi² = 8.93, df = 1 (P = 0.003), I² = 88.8%