

Supplementary Material

Table S2. Neglect screening and assessment tools with documented protocols - analysis of neglect subtypes and ICF categories

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Behavioural Inattention Test (BIT)[1] modified scoring [2]	✓	X		✓		X	✓			X	✓	✓	
BIT-CS (conventional subtests)	✓	X		✓		X	X			X	✓		
BIT-bs (behavioural subtest)	✓	X		✓			✓			X	✓	✓	
sBIT (3-item version) [3]	✓	X		✓			X			X	✓		
mBIT for acute stroke (8 item version) [4]	✓			✓	✓		✓			X	✓	✓	
Chinese Behavioural Inattention Test–Hong Kong version (CBIT-HK) [5]	✓	X		✓		X	✓			X	✓	✓	
BIT Japanese version [6,7]	✓	X		✓		X	✓			X	✓	✓	
BIT Hebrew version [8,9]	✓	X		✓		X	✓			X	✓	✓	
Neglect Test (NET) (Fels & Geissner, 1996) (German BIT)[10, 11]	✓	X		✓		X	✓			X	✓	✓	
BIT-E (Spanish version) [12]	✓	X		✓		X	✓			X	✓	✓	
Sunnybrook Neglect Assessment Procedure (SNAP)[13]	✓	X		✓			X			X	✓		
Batterie d’Evaluation de la Négligence (BEN) [14,15] (GEREN neglect battery)	✓	X		✓		X	✓			X	✓	✓	
Rivermead Perceptual Assessment Battery [16]	X	X		✓			X			X	✓		
Northwick Park Examination of Cognition [17] Perception component	✓	X		✓			X			X	✓		
Strokevision app [18]	✓	X		✓			X			X	✓		
Vision Impairment Screening Assessment (VISA) tool[19]	✓	X		✓			X			X	✓		
Screening Instrument for neuropsychological impairment in stroke (SINS)[20]	✓	X	X	X		X	X			X	✓		
Battery for the evaluation of hemineglect [21]	✓			✓			✓			X	✓	✓	
OT-APST (Adult Perceptual Screening test) neglect subtest [22]	X	X		✓		X	✓			X	✓	✓	
OSOT (Ontario Society of Occupational Therapists) Evaluation of Perceptual Function [23]	X	X		✓		X	X			X	✓		
The Ontario Society of Occupational Therapy (OSOT) Perceptual Evaluation French version [24] - Visual scanning and visuo-spatial neglect subtest	X	X		✓		X	X			X	✓		
The Visuospatial Neglect Test Battery [25]	✓	X		✓		X	X			X	✓		
Circle Monitor CM Tasks [10]	✓	X		✓			X			X	✓		

✓ Tool differentiates the neglect subtype.

X Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(X) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Cancellation tasks													
Bell's test[26] (modified protocols [27,28])	✓			✓			X			X	✓		
Star cancellation test [1]	✓			✓			X			X	✓		
Star cancellation test - Russian version [29]	✓			✓			X			X	✓		
line cancellation test [1], Albert's Test[30,31] (40 lines)	✓			✓			X			X	✓		
Mesulam shape cancellation task [32]	✓			✓			X			X	✓		
Line cancellation task 30 x short lines [33,34]	✓			✓			X			X	✓		
Line cancellation (Left, middle and right of midline)[35]	✓			✓			X			X	✓		
Barrage test/ Line cancellation test) [21] (21 lines, A3 page)	✓			✓			X			X	✓		
Ellipses cancellation task [36]	✓			✓			X			X	✓		
BIT - short screening test [37] (2 items: Letter & star)	✓			✓			X			X	✓		
BIT cancellation tasks (letter, line, star)	✓			✓			X			X	✓		
Computerised cancellation tasks [38]	✓			✓		X	X			X	✓		
Balloons test [39]	✓			✓			X			X	✓		
Digitized shape cancellation task [40]	✓			✓			X			X	✓		
Behavioural task [34]	✓			✓			X			X	✓		
Head mounted display (HMD) - line & star cancellation[6]	✓	X		✓			X			X	✓		
VR and haptics cancellation task [41]	✓			✓			X			X	✓		
Touch screen cancellation tasks) (letter and shape) [42]	✓			✓			X			X	✓		
Letter cancellation task (SLCT/H cancellation Task) [43,44]	✓			✓			X			X	✓		
(Random) Letter cancellation task [32,45]	✓			✓			X			X	✓		
Double Letter Cancellation Test (DLCT) [44,46]	✓			✓			X			X	✓		
Letter cancellation test [47]	✓			✓			X			X	✓		
O/X cancellation task with graphics tablet [48]	✓			✓			X			X	✓		
Structured Chinese Word Cancellation Test (SCWCT) [49]	✓			✓			X			X	✓		
Random Chinese Word Cancellation Test (RCWCT) [49–51]	✓			✓			X			X	✓		
Custom cancellation test "T" [52]	✓			✓			X			X	✓		
Letter cancellation (BIT subtest)[1] (timed version [28])	✓			✓			X			X	✓		
O cancellation test (visible/invisible marks) [53]	✓			✓			X			X	✓		
Letter cancellation test [54]	✓			✓			X			X	✓		

✓ Tool differentiates the neglect subtype.

✗ Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(✗) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Egocentric and allocentric cancellation tasks													
Apples cancellation test [55]	✓	✓		✓			X			X	✓		
Birmingham Cognitive Screen (BCoS) [56]	✓	✓		✓			X			X	✓		
Hong Kong Birmingham Cognitive Screen (HKBCOS) [57]	✓	✓		✓			X			X	✓		
Cantonese Birmingham Cognitive Screen (Cantonese BCoS) [58]	✓	✓		✓			X			X	✓		
Oxford Cognitive Screen: Broken hearts cancellation task [59]	✓	✓		✓			X			X	✓		
Hong Kong version of Oxford Cognitive Screen: Broken hearts cancellation task (HK-OCS) [60]	✓	✓		✓			X			X	✓		
Russion Version of the Oxford Cognitive Screen (Rus-OCS) [61]	✓	✓		✓			X			X	✓		
3S Spreadsheet Test (3S Test) [62]	✓	✓		✓			X			X	✓		
Ota circle cancellation [63] (Gap detection task, Ota search task)	✓	✓		✓			X			X	✓		
modified Gap Detection Test [64] (Visual and Tactile)	✓	✓		✓			X	X		X	✓		
Defect detection task - circle and triangle [63]	✓	✓		✓			X			X	✓		
Apples cancellation test Italian standardization [55,65]	✓	✓		✓			X			X	✓		
Motor tasks													
Lateralised motor performance [66]	✓		✓								✓	✓	
Computerised line bisection Task - Fractionated where and aiming measures [66]	X	X		✓			✓			✓	✓	✓	
Closed circuit camera of line bisection and cancellation tasks [67]	✓			✓			✓			✓	✓	✓	
Paradigm for assessing directional motor bias [68]	✓			✓			✓			✓	✓	✓	
Motor and perceptual RT task [69]	✓			✓			✓			✓	✓	✓	
Landmark line bisection task (Landmark Test) [70]	✓			✓			✓			✓	✓	✓	
Modified landmark task (mLMT) [71]	X	X		✓			✓			✓	✓	✓	
Congruent and non-congruent bisection task [72]	X	X		✓			✓			✓	✓	✓	
Wooden handle [73]	✓			✓							✓	✓	
Exploratory-motor test [74]	✓			✓			X			X	✓		

✓ Tool differentiates the neglect subtype.

✗ Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(✗) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Line bisection tasks													
Schenkenberg Line bisection task (modified scoring [75])	X	X		✓			X			X	✓		
Line bisection - (2 x 10cm, 2 x 15cm, 2 x 25cm lines) [76]	X	X		✓			X			X	✓		
Line bisection test (BIT)[1] (timed version [28])	X	X		✓			X			X	✓		
Line bisection [77] (24 x 20cm)	X	X		✓			X			X	✓		
Line bisection test (3 x 16cm and 3 x 24cm) [78]	X	X		✓			X			X	✓		
Line bisection (2 x 5cm and 2 x 20cm) [14] (% deviation)	X	X		✓			X			X	✓		
Line bisection task (5 x 6cm, 5 x 7cm, 5 x 8cm [79])	X	X		✓			X			X	✓		
Line bisection task 3 x 20cm line [33]	X	X		✓			X			X	✓		
Solid line bisection task (10 x 24.2cm) [80,81]	X	X		✓			X			X	✓		
Line bisection 5 x 24cm [82,83]	X	X		✓			X			X	✓		
Line bisection (3 x 17cm lines) [84]	X	X		✓			X			X	✓		
Line bisection (3 x 22cm lines on A4 page) [85]	X	X		✓			X			X	✓		
Line bisection task (5 x 18cm) [86]	X	X		✓			X			X	✓		
Line bisection (27cm) (page to the L, R and midline) [87]	X	X		✓			X			X	✓		
Complex line bisection test [88,89]	X	X		✓			X			X	✓		
Line bisection [90] (36 lines 26cm/30cm (numbers at either end, presented left and right of midline)	X	X		✓			X			X	✓		
Stroke Impairment Assessment Set (SIAS) - visuospatial perception subtest [91,92] (Bisect 50cm tape x 2)	X	X		✓			X			X	✓		
Line bisection test (10 x 10.7cm) [54]	X	X		✓			X			X	✓		
Line bisection task (5 x 18cm) [36]	X	X		✓			X			X	✓		
Horizontal line bisection [93]	X	X		✓			X			X	✓		
Line bisection task (5 x 242mm) [94]	X	X		✓			X			X	✓		
Line bisection (6 lines on page offset from centre) [18]	X	X		✓			X			X	✓		
Line bisection 25 x 15 cm [95]	X	X		✓			X			X	✓		
Line bisection (10 x 24cm: 5 left of midline, 5 right of midline) [96]	X	X		✓			X			X	✓		
Bisection of a 20cm horizontal line [97]	X	X		✓			X			X	✓		

✓ Tool differentiates the neglect subtype.

✗ Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(✗) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Line bisection tasks (continued)													
Line bisection (8 lines 3 x 6.2cm, 3 x 10cm, 2 x 18cm, cumulative % deviation) [98]	X	X		✓			X			X		✓	
Character Line Bisection Task (CLBT: Letter-line and Star-line Bisection Tasks) [80]	X	X		✓			X			X		✓	
Computerized touch screen line bisection test (300 x (20cm, 25cm, 30cm lines) (in middle, L, R space with/o distractions) [99]	X	X		✓			X			X		✓	
Rebisection task [100]	X	X		✓			X			X		✓	
Line quadrisection task [101]	X	X		✓			X			X		✓	
Line extension task (5x6cm, 5x7cm, 5x8cm in centre of A4 page) [79]	X	X		✓			✓			✓		✓	
L figure line extension task [102]	X	X		✓			X			X		✓	
Bisection of Judd stimuli (10cm and 12cm rods with visual, haptic, and visuo-haptic presentation) [103]	X	X		✓		X	X	X		X		✓	
Extended Line Bisection task (LB) [104,105]	X	X		✓			X			X		✓	
Pointing/grasping objects													
Pointing/grasping centre of rod [106]	X	X		✓			X			X		✓	
Baking Tray Task (BTT) (Tham & Tegner) [3,107]	✓			✓			X			X		✓	
Baking Tray Task with E-TAN (E-BTT) [108]	✓			✓			X			X		✓	
modified version of Baking Tray Task [109]	✓			✓			X			X		✓	
Table Test (Kerkhoff) [52]	✓			✓			X			X		✓	
Visual search board (VSB; Kimura) [13]	✓			✓			X			X		✓	
Robotic object hitting task [110]	✓			✓			X			X		✓	
Two-part picture test [111]	✓			✓			X			X		✓	
Object-finding test [89]	✓			✓			X			X		✓	
The shape test [112]	✓			✓		X	X	X		X		✓	

✓ Tool differentiates the neglect subtype.

✗ Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(✗) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Copying tasks													
5-element complex drawing (a house and four trees) (Gainotti drawing, Landscape drawing) [113]	x	x		✓			x			x	✓		
Copying Task (a fence, a car, a house and a tree) [114]	x	x		✓			x			x	✓		
Ogden Scene (tree, fence, house, tree) [115](landscape drawing, scene/ figure copy) [14]	x	x		✓			x			x	✓		
Figure copy (BIT) (star, cube, flower, triangles) [1,2]	x	x		✓			x			x	✓		
Design copy test (house, flower, clock, diamond) [116]	x	x		✓			x			x	✓		
Copy a daisy/Daisy drawing [103]	x	x		✓			x			x	✓		
Two daisy figure /Complex flower copying task [117]	x	x		✓			x			x	✓		
Rey-Osterrieth complex figure test, copy [118]	x	x		✓			x			x	✓		
Drawing tasks													
Clock drawing task (19 pt scale) [119]	x	x		✓		x	x			x	✓		
Clock drawing test (Freedman 15 pt scale) [120]	x	x		✓		x	x			x	✓		
Clock drawing task (CLOX) (15pts) [121]	x	x		✓		x	x			x	✓		
Clock drawing (from memory) (12 pt scale) [103,122]	x	x		✓		x	x			x	✓		
Clock drawing test (Sunderland/Manos 10 pts) [123,124]	x	x		✓		x	x			x	✓		
Clock drawing test (7 pt scale) - Watson system [5]; Seven Minute Screen (Solomon & Pendlebury) [124]	x	x		✓		x	x			x	✓		
Clock drawing test (4 pt scale) (printed 8cm circle) [125]	x	x		✓		x	x			x	✓		
Clock drawing (4 pt scale) [126]	x	x		✓		x	x			x	✓		
OT-APST (Occupational Therapy Adult Perceptual Screening Test) - clock drawing subtest [22]	x	x		✓		x	x			x	✓		
Clock drawing test (2 pt scale) [14,127]	x	x		✓		x	x			x	✓		
Clock drawing task (Copy) (impaired/normal) [81]	x	x		✓		x	x			x	✓		
Number clockface (printed circle) (impaired/normal) [16]	x	x		✓		x	x			x	✓		
Five-Point test (Regard 1982) [128] (5 Dot test)	✓			✓			x			x	✓		
BIT representational drawing (clock, person, butterfly) [1,2]	x	x		✓		x	x			x	✓		
Draw-A-Man test [129]	x	x		✓		x	x			x	✓		
Writing task [14]	✓			✓		x	x			x	✓		✓
Arrow location in 3 conditions (blank, lines, numbers) [130]	✓			✓			x			x	✓		

✓ Tool differentiates the neglect subtype.

x Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(x) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Reading tasks													
Sentence reading [131]	✓	✗		✓			✓				✓	✓	
Reading task/ text reading [14]	✓	✗		✓			✓				✓	✓	
Sentence reading [123,132]	✓	✗		✓			✓				✓	✓	
Single word reading (Vallar 1996) [123]	✗	✗		✓			✓				✓		
Text reading [133]	✓	✗		✓			✓				✓	✓	
Indented paragraph reading test [134]	✓	✗		✓			✓				✓	✓	
Indented paragraph test modified [46,134,135]	✓			✓			✓				✓	✓	
Reading of compound-words [133]	✓	✓		✓			✓				✓		
Reading task (individual words & sentences) [35]	✓	✓		✓			✓				✓	✓	
Reading task [136]	✓	✗		✓			✓				✓	✓	
Munich reading tests [137]	✓			✓			✓				✓	✓	
Menu subtests of the BIT [1]	✗	✗		✓			✓				✓	✓	
Article reading subtests of the BIT [1]	✗	✗		✓			✓				✓	✓	
Neglect dyslexia test - Two-word phrases [138]	✗	✗		✓			✓				✓		
Neglect dyslexia test - Single words [138]	✗	✗		✓			✓				✓		
Word reading test (WR; range 0–35) [139]	✗	✗		✓			✓				✓	✓	
Experimental word reading test (EWR; range:0–35) [139]	✗	✗		✓			✓				✓		
Single word reading task [140]	✗	✗		✓			✓				✓		
Article reading (Finnish folk story) [89]	✓	✓		✓			✓				✓	✓	
Neglect dyslexia test (Korean) [141]	✗	✗		✓			✓				✓		
Conventional text reading task [140]	✗	✗		✓			✓				✓	✓	
Number reading task (Max 9pts) [121]	✗	✗		✓			✓				✓		
Other visual tasks													
Ravens Coloured Progressive Matrices (RCPM) (proportion of right and left responses) [46]	✓			✓			✓			(x)	✓		
Visual-counting test [74]	✓			✓			✓				✓		
Picture scanning subtest of BIT [1]	✓			✓			✓				✓		
Overlapping figures test [14,142]	✓	✓		✓			✓				✓		

✓ Tool differentiates the neglect subtype.

✗ Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(x) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Other visual tasks (continued)													
Object search task [143]	x	x		✓			✓			(x)	✓		
Independent segment estimation task [71]	x	x		✓		x	x				✓		
Chimeric object naming test [144,145]	x	x		✓			✓				✓		
Chimeric figures [146]	x	x		✓			✓				✓		
Chimeric faces [147]	x	x		✓			✓				✓		
Wundt–Jastrow area illusion test [148]	x	x		✓			✓				✓		
Eye tracking free exploration task [121]	✓			✓			✓				✓		
Eye tracking visual search task [121]	✓			✓			✓				✓		
Free visual exploration task [149]	✓			✓			✓				✓		
Gaze and head orientation at rest and during active visual exploration [150]	✓			✓			✓			x	✓		
Spontaneous head and eye gaze [151]	✓				✓		✓			x	✓		
Video-oculography with FVE paradigm [152]	✓			✓			✓				✓		
Eye tracking with IADL task [153]	✓				x	x	✓				✓		✓
Inhibition of return (IOR) - covert/overt attention [154]	✓			✓			✓				✓		
Orientation singletons [155]	✓			✓			✓				✓		
Representational tasks													
The “Bisiach-Ogden test” [156]	x	x		✓		x	x				✓		
Familiar Square Description Test [157,158]	✓						✓				✓		
3 Roman Squares [159]	✓						✓				✓		
Imaginal RT task (towns in France) [160]	✓						✓			x	✓		
Towns on coast of Italy [159]	✓						✓				✓		
Countries of Europe [159]	✓						✓				✓		
Description from memory of map of France [161]	✓						✓				✓		
map of France (and inverted map) [162]	✓						✓				✓		
O’Clock Test [158,163]	x	x		✓			✓				✓		
Map of France (eyes open and eyes closed) [161]	✓						✓				✓		
Tactuo-Motor search (TMS) task [89]	x	x		✓		x	x				✓		

✓ Tool differentiates the neglect subtype.

x Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(x) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Personal space tasks													
Fluff test [164]	✓		✓			X		X		X	✓		
modified version of the Fluff Test (6x Left, 4 Right) [28]	✓		✓			X		X		X	✓		
modified version of Fluff test - (12 x upper body) [165]	✓		✓			X		X		X	✓		
modified version of Fluff test - (5xL side, 2xR) [126]	✓		✓			X		X		X	✓		
Vest Test [166]	✓		✓			X		X		X	✓		
Personal neglect evaluation [89]	✓		✓			X	X			X	✓		
Extension of Personal neglect test [167]	✓		✓			X	X			X	✓		
Personal neglect test /Bisiach test (Touching hand) [168]	✓		✓			X	X			X	✓		
Far space tasks					✓		✓			X	✓	✓	✓
Dublin Extrapersonal neglect assessment (DENA) [169]	✓				✓								
Room description task [4]	✓				✓		✓					✓	
The Visual Scanning Test (VST) [170]	✓				✓		✓					✓	
Large screen tasks (Active space): Detection task and crash task (Single and dual task) [171]	✓				✓		✓			?		✓	
Slide (far space) test [89]	✓				✓			✓				✓	
laser line-bisection task [28]	X	X			✓		X			X		✓	
Far and near space tasks					✓	✓		X		X		✓	
Line bisection and Bells Test in near and far space [172]	✓				✓	✓							
Digitized object cancellation in peripersonal and extrapersonal space) [173]	✓				✓	✓		X		X		✓	
Digitized line bisection task in peripersonal and extrapersonal space) (3 x 22cm lines on page x 4) [173]	✓				✓	✓		X		X		✓	
Line bisection task (10 x lines at 50cm, 1.5m, 3m) [174]	✓				✓	✓		X		X		✓	
Letter cancellation task [173]	✓				✓	✓		X		X		✓	
Schenkenberg line bisection & star cancellation test in near and far extrapersonal space [174,175]	✓				✓	✓		X		X		✓	
Wundt-Jastro area illusion: peri-personal and extra-personal presentation [176]	X	X			✓	✓		✓				✓	
Doorway bisection task [174]	✓				✓	✓		X		X	✓	✓	✓

✓ Tool differentiates the neglect subtype.

✗ Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(✗) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Personal, far and near space tasks												
Halifax Visual Scanning Test (HVST)[177]	✓	✓	✓	✓		✓				✓		
Dynamic tasks – VR and computer-based												
Test Battery for Attentional Performance (TAP) - neglect subtest [178]	✓		✓			✓			?	✓		
Virtual Reality Lateralized Attention Test (VRLAT) [28, 179]	✓		X	X		X			X	✓	✓	
Virtual reality wheelchair navigation test [180]	✓		X	X		X			X	✓	✓	
VR-DiSTRO (VR-SCT, VR-LB, VR-BTT, VR-EXT)[181]	✓		✓			X			X	✓		
Virtual Environment for Spatial Neglect Assessment (VESNA) [182]	✓		X	X		X			X	✓		
Virtual reality assessment for visuospatial neglect [183]	✓		? ?			✓			X	✓	✓	
VR Assessment and Training system for Unilateral Neglect[184]	✓		? ?			✓			X	✓		
VR -three-dimensional virtual street crossing program[185,186]	✓		X X			✓				✓	✓	
Virtual Street Crossing system [187]	✓		X X			X			X	✓	✓	
Ecological VR-based Evaluation of Neglect Symptoms (EVENS) [188]	✓		X X			X			X	✓	✓	
VR-based goal-directed locomotion task [175]	✓			✓	X	X			X	✓	✓	
Visual Search Task - Cognitive rehabilitation platform Guttmann-NeuroPersonal Trainer (García-Molina et al., 2010; Solana et al., 2014; Solana et al., 2015).	✓		✓			X			X	✓		
VISSTA (Visual Spatial Search Task) [189]	✓		? ?			✓				✓		
Computerised Table Setting Test [94]	✓		✓			X			X	✓	✓	
MonAmour Test (Robot Diagnosis Test for Hemineglect) [178]	✓ ✓		✓			✓				✓		
Starry Night Test (SNT) [190]	✓		✓			✓				✓		
Adaptive visual conjunction search task [191,192]	✓		? ?			✓				✓		
Vienna Test System - peripheral perception subtest [193]	✓		✓			✓			X	✓		
TVA - whole or partial report task [194]	✓		✓			✓				✓		
Useful Field of View (UFOV) [195]	✓		✓			X			X	✓		
Simulated driving task [196,197]	✓			✓		X			X	✓	✓	

✓ Tool differentiates the neglect subtype.

✗ Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(✗) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Dual tasks													
E-prime single task & dual task [198,199]	✓			✓			✓				✓		
Computerized Visual RT Task: single (CVRT) and dual (CVRT-D) task [196]	✓			✓			✓				✓		
Multi-tasking paradigm (Single and Dual Tasks) [200]	✓			✓			✓		✓		✓		
Dual task test [180]	✓			✓			✓				✓		
Reaction time tasks													
Posner cuing task - Reaction time test / Valid versus invalid cue paradigm [120]	✓			✓			✓				✓		
Spatial cueing (Posner) task [201]	✓			✓			✓				✓		
Eye movement RT task [202]	✓			✓			✓				✓		
Spatial orientation task [203]	✓			✓			✓				✓		
Perceptual RT task/ Reaction time task [69,160]	✓			✓			✓				✓		
Auditory tasks											✓		
Dichotic listening task [204]	✓										✓		
Temporal Order Judgement (TOJ) task [205]	✓			✓			✓				(x)		
Lateralisation of dichotic sound [206]	✓										✓		
Auditory Spatial Discrimination Task [207]	✓										✓		
Psychoacoustic pattern discrimination test (PPDT) modified non-adaptive version [208]	✓										✓		
Visual and Auditory TOJ Task [209]	✓			✓			✓				✓		
Extinction Tasks													
National Institute of Health Stroke Scale (NIHSS) - Neglect /extinction item	x		x	x			x	x	x		✓		
Cog-4 (NIHSS Subscale of 4 Cognitive Items) [210]	x		x	x			x	x	x		✓		
Tactile extinction test DSS (8L, 8R, 8B) [33]	x		x					x			✓		
Visual extinction test DSS (8, 8, 8) [33]	x		x				x				✓		
Thermal pain DSS protocol [33]	x		x					x			✓		
DSS with electromagnetic solenoids (18x/12x + 6 catch trials) [211]	x		x					x			✓		
Confrontation method of extinction (Tactile, auditory, visual) (score 0-9) [212]	x		x	x	x		x	x	x		✓		
Computerised test of visual neglect and extinction [213]	x			x			x			x		✓	
Novel computerised test for visual extinction/neglect [214]	x				?		x					✓	

✓ Tool differentiates the neglect subtype.

x Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(x) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
ADL based tasks												
Catherine Bergego Scale (CBS) [215]	✓	x	x	x	x	x	x	x	x		✓	
Catherine Bergego Scale via KF-NAP [216,217]	✓	x	x	x	x	x	x	x	x		✓	
Subjective Neglect Questionnaire [218]	✓	x	x	x	x	x	x	x	x		✓	
ADL checklist [8]	✓	x	x	x		x	x	x	x		✓	✓
Árnadóttir OT-ADL Neurobehavioral Evaluation (A-ONE) [219]	✓	x	x		x	x		x	x		✓	
Catherine Bergego Scale Dutch version [220,221]	✓	x	x	x	x	x	x	x	x		✓	
Catherine Bergego Scale -extrapersonal items (CBS-E) [169]	✓			✓		x		x	x		✓	
OT checklist of everyday neglect behaviours [2]	✓	x	x	x		x	x	x	x		✓	✓
Modified checklist of everyday neglect behaviors [2,6]	✓	x	x	x	x	x	x	x	x		✓	✓
Novel ADL-based neglect test (Activities of Daily Living Scale) [11]	✓	x	x	x	x	x	x		x		✓	
Semi-structured scale for the functional evaluation of extrapersonal neglect [222]	✓			✓	✓			✓		x		✓
Semi-structured scale for the functional evaluation of personal neglect [223]	✓		✓			x	x		x		✓	
Tea Making (TM) Task [224]	✓			✓		x			x		✓	
Document Filing (DF) Task [224]	✓			✓		x			x		✓	
Comb and razor test [225]	✓		✓			x	x		x		✓	
Reformulated comb and razor test [225,226]	✓	x			x	x			x		✓	
Key removal, grocery naming, face washing, tray wiping [165]	✓	x	x			x			x		✓	
Visual Neglect Recovery Index (VNRI) [227]	✓		x	x		x			x		✓	✓
Naturalistic Action Test (NAT) [228] (Lateralised Attention Score) [229]	✓		x	x		x			x		✓	✓
Multi-Level Action Test (MLAT) (making toast, wrapping present, packing lunchbox) [230]	✓		✓			x			x		✓	

✓ Tool differentiates the neglect subtype.

x Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(x) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

Tool name (reference)	Egocentric	Allocentric	Personal	Peripersonal	Extrapersonal	Representation	Visual	Tactile	Auditory	Motor	Impairment	Activity Limitation	Participation Restriction
Mobility based tasks													
Mobility Assessment Course [85,231]	✓			X	X		✓			X		✓	
Doorway Accuracy Test (DAT) [232]	✓			X	X		X			X		✓	
Wheelchair assessment Course (WAC) [233]	✓			X	X		X			X		✓	
Cone Evasion Walk Test [234]	✓			X	X		✓			X		✓	
Moss–Magee Wheelchair Navigation Test (MMWNT) [45,180,235]	✓			X	X		X			X	✓	✓	
modified MossReal World Navigation (RW) test [28, 180]	✓			X	X		X			X	✓	✓	
Wheelchair obstacle course [177,236]	✓			X	X		X			X	✓	✓	
Wheelchair Collision Test (WCT) [237]	✓			X	X		X			X	✓	✓	
Wheelchair assessment Course (WAC) [233]	✓			X	X		X			X	✓	✓	

✓ Tool differentiates behaviours for the neglect subtype.

X Deficient performance may be attributable to the subtype, but the scoring does not allow for it to be differentiated from other subtypes.

? Unable to determine from documented protocols.

(x) Alternative protocols of the test involve this subtype (for example motor response if unable to verbalise response)

References:

- [1] Wilson B, Cockburn J, Halligan P. Development of a behavioral test of visuospatial neglect. *Arch Phys Med Rehabil* 1987;68:98–102.
- [2] Halligan PW, Cockburn J, Wilson BA. The behavioural assessment of visual neglect. *Neuropsychological Rehabilitation* 1991;1:5–32. <https://doi.org/10.1080/09602019108401377>.
- [3] Appelros P, Karlsson G, Thorwalls A, Tham K, Nydevik I. Unilateral neglect: further validation of the baking tray task. *Journal of Rehabilitation Medicine* 2004;36:258–61.
- [4] Stone SP, Wilson B, Wroot A, Halligan PW, Lange LS, Marshall JC, et al. The assessment of visuo-spatial neglect after acute stroke. *J Neurol Neurosurg Psychiatry* 1991;54:345–50.
- [5] Fong KNK, Chan MKL, Chan BYB, Ng PPK, Fung ML, Tsang MHM, et al. Reliability and validity of the Chinese behavioural inattention test-Hong Kong version (CBIT-HK) for patients with stroke and unilateral neglect. *Hong Kong Journal of Occupational Therapy* 2007;17:23–33. <https://doi.org/10.1016/S1569-1861%2807%2970004-9>.
- [6] Tanaka T, Sugihara S, Nara H, Ino S, Ifukube T. A preliminary study of clinical assessment of left unilateral spatial neglect using a head mounted display system (HMD) in rehabilitation engineering technology. *Journal of NeuroEngineering and Rehabilitation* 2005;2:31. <https://doi.org/10.1186/1743-0003-2-31>.
- [7] Isai S. Behavioural Inattention Test, Japanese edition. Tokyo, Japan: Shinkoh Igaku Shuppan, Co Ltd; 1999.
- [8] Hartman-Maeir A, Katz N. Validity of the Behavioral Inattention Test (BIT): Relationships with functional tasks. *American Journal of Occupational Therapy* 1995;49:507–16. <https://doi.org/10.5014/ajot.49.6.507>.
- [9] Katz N, Averbuch S, Itzkovich M. Hebrew translation of behavioural inattention test manual. Fareham, Hants, England: Thames Valley Test Co; 1991.
- [10] Ulm L, Wohlraup D, Meinzer M, Steinicke R, Schatz A, Denzler P, et al. A circle-monitor for computerised assessment of visual neglect in peripersonal space. *PloS One* 2013;8:e82892. <https://doi.org/10.1371/journal.pone.0082892>.
- [11] Eschenbeck P, Vossel S, Weiss PH, Saliger J, Karbe H, Fink GR. Testing for neglect in right-hemispheric stroke patients using a new assessment battery based upon standardized activities of daily living (ADL). *Neuropsychologia* 2010;48:3488–96. <https://doi.org/10.1016/j.neuropsychologia.2010.07.034>.
- [12] Sánchez-Cabeza A, Huertas-Hoyas E, Máximo-Bocanegra N, Martínez-Piédrola RM, Pérez-de-Heredia-Torres M, Alegre-Ayala J. Spanish transcultural adaptation and validity of the behavioral inattention test. *Occup Ther Int* 2017;2017. <https://doi.org/10.1155/2017/1423647>.
- [13] Leibovitch FS, Vasquez BP, Ebert PL, Beresford KL, Black SE. A short bedside battery for visuoconstructive hemispatial neglect: Sunnybrook Neglect Assessment Procedure (SNAP). *Journal of Clinical and Experimental Neuropsychology* 2012;34:359–68. <https://doi.org/10.1080/13803395.2011.645016>.

- [14] Azouvi P, Samuel C, Louis-Dreyfus A, Bernati T, Bartolomeo P, Beis JM, et al. Sensitivity of clinical and behavioural tests of spatial neglect after right hemisphere stroke. *Journal of Neurology Neurosurgery and Psychiatry* 2002;73:160–6. <https://doi.org/10.1136/jnnp.73.2.160>.
- [15] Azouvi P, Bartolomeo P, Beis J-M, Perennou D, Pradat-Diehl P, Rousseaux M. A battery of tests for the quantitative assessment of unilateral neglect. *Restorative Neurology and Neuroscience* 2006;24:273–85.
- [16] Friedman PJ, Leong L. The Rivermead Perceptual Assessment Battery in acute stroke. *The British Journal of Occupational Therapy* 1992;55:233–7. <https://doi.org/10.1177/030802269205500608>.
- [17] Williams PM, Johnson C, Swan S, Barber C, Murphy P, Devine J, et al. The northwick park examination of cognition: A brief cognitive assessment tool for use in acute stroke services. *International Journal of Therapy and Rehabilitation* 2016;23:314–22.
- [18] Quinn TJ, Livingstone I, Weir A, Shaw R, Breckenridge A, McAlpine C, et al. Accuracy and Feasibility of an Android-Based Digital Assessment Tool for Post Stroke Visual Disorders-The StrokeVision App. *Front Neurol* 2018;9:146. <https://doi.org/10.3389/fneur.2018.00146>.
- [19] Rowe F.J., Hepworth L.R., Hanna K.L., Howard C. Visual Impairment Screening Assessment (VISA) tool: Pilot validation. *BMJ Open* 2018;8:e020562. <https://doi.org/10.1136/bmjopen-2017-020562>.
- [20] Sodring KM, Laake K, Sveen U, Wyller TB, Bautz-Holter E. Validation of the screening instrument for neuropsychological impairment in stroke. *Physiother Res Int* 1998;3:15–26.
- [21] Pizzamiglio L, Judica A, Razzano C, Zoccolotti P. Toward a comprehensive diagnosis of visual-spatial disorders in unilateral brain damaged patients. *Evaluación Psicológica* 1989;5:199–218.
- [22] Cooke D.M., McKenna K., Fleming J., Darnell R. The reliability of the Occupational Therapy Adult Perceptual Screening Test (OT-APST). *Br J Occup Ther* 2005;68:509–17.
- [23] Boys M, Fisher P, Holzberg C, Reid DW. The OSOT Perceptual Evaluation: a research perspective. *The American Journal of Occupational Therapy* 1988;92–8.
- [24] Desrosiers J, Mercier L, Rochette A. Fidélité test-retest et interjugés de la version française du OSOT Perceptual Evaluation. *Can J Occup Ther* 1999;66:134–9. <https://doi.org/10.1177/000841749906600305>.
- [25] Vaes N, Lafosse C, Nys G, Schevernels H, Dereymaeker L, Oostra K, et al. Capturing peripersonal spatial neglect: An electronic method to quantify visuospatial processes. *Behavior Research Methods* 2015;47:27–44. <https://doi.org/10.3758/s13428-014-0448-0>.
- [26] Gauthier L, Dehaut F, Joanette Y. The bells test: a quantitative and qualitative test for visual neglect. *International Journal of Clinical Neuropsychology* 1989;11:49–54.
- [27] Basagni B, De Tanti A, Damora A, Abbruzzese L, Varalta V, Antonucci G, et al. The assessment of hemineglect syndrome with cancellation tasks: a comparison between the Bells test and the Apples test. *Neurol Sci* 2017;38:2171–6. <https://doi.org/10.1007/s10072-017-3139-7>.
- [28] Buxbaum LJ, Dawson AM, Linsley D. Reliability and validity of the Virtual Reality Lateralized Attention Test in assessing hemispatial neglect in right-hemisphere stroke. *Neuropsychology* 2012;26:430–41. <https://doi.org/10.1037/a0028674>.
- [29] Friedman PJ. The Star Cancellation Test in acute stroke. *Clinical Rehabilitation* 1992;6:23–30. <https://doi.org/10.1177/026921559200600104>.
- [30] Albert ML. A simple test of visual neglect. *Neurology* 1973;23:658–64. <https://doi.org/10.1212/WNL.23.6.658>.
- [31] Fullerton KJ, Mcsherry D, Stout RW. Albert's test: A neglected test of perceptual neglect. *The Lancet* 1986;327:430–2. [https://doi.org/10.1016/S0140-6736\(86\)92381-0](https://doi.org/10.1016/S0140-6736(86)92381-0).
- [32] Weintraub S, Mesulam MM. Principles of behavioral neurology. Mental state assessment of young and elderly adults in behavioral neurology, Philadelphia: PA: Davis FA; 1985, p. 71–123.
- [33] Liu CC, Veldhuijzen DS, Ohara S, Winberry J, Greenspan JD, Lenz FA. Spatial attention to thermal pain stimuli in subjects with visual spatial hemi-neglect: Extinction, mislocalization and misidentification of stimulus modality. *Pain* 2011;152:498–506. <https://doi.org/10.1016/j.pain.2010.10.017>.
- [34] Toba MN, Rabuffetti M, Duret C, Pradat-Diehl P, Gainotti G, Bartolomeo P. Component deficits of visual neglect: “Magnetic” attraction of attention vs. impaired spatial working memory. *Neuropsychologia* 2018;109:52–62. <https://doi.org/10.1016/j.neuropsychologia.2017.11.034>.
- [35] Khurshed S, Trupe LA, Newhart M, Davis C, Molitoris JJ, Medina J, et al. Reperfusion of specific cortical areas is associated with improvement in distinct forms of hemispatial neglect. *Cortex* 2012;48:530–9. <https://doi.org/10.1016/j.cortex.2011.01.003>.
- [36] Pia L, Ricci R, Gindri P, Vallar G. Drawing perseveration in neglect: Effects of target density. *Journal of Neuropsychology* 2013;7:45–57. <https://doi.org/10.1111/j.1748-6653.2012.02034.x>.
- [37] Halligan P, Wilson B, Cockburn J. A short screening test for visual neglect in stroke patients. *Int Disabil Stud* 1990;12:95–9.
- [38] Wansard M, Meulemans T, Gillet S, Segovia F, Bastin C, Toba MN, et al. Visual neglect: Is there a relationship between impaired spatial working memory and re-cancellation? *Experimental Brain Research* 2014;232:3333–43. <https://doi.org/10.1007/s00221-014-4028-4>.
- [39] Edgeworth Jennifer A, MacMillan TM, Robertson IH. The balloons test manual 1998.
- [40] Spreij LA, Ten Brink AF, Visser-Meily JMA, Nijboer TCW. Simulated driving: The added value of dynamic testing in the assessment of visuo-spatial neglect after stroke. *Journal of Neuropsychology* 2020;14:28–45. <https://doi.org/10.1111/jnp.12172>.

- [41] Broeren J, Samuelsson H, Stibrant-Sunnerhagen K, Blomstrand C, Rydmark M. Neglect assessment as an application of virtual reality. *Acta Neurologica Scandinavica* 2007;116:157–63. <https://doi.org/10.1111/j.1600-0404.2007.00821.x>.
- [42] Rabuffetti M, Ferrarin M, Spadone R, Pellegatta D, Gentileschi V, Vallar G, et al. Touch-screen system for assessing visuo-motor exploratory skills in neuropsychological disorders of spatial cognition. *Medical and Biological Engineering and Computing* 2002;40:675–86.
- [43] Vallar G, Guariglia C, Nico D, Bisiach E. Spatial hemineglect in back space. *Brain* 1995;118:467–72. <https://doi.org/10.1093/brain/118.2.467>.
- [44] Diller L, Weinberg J. Hemi-inattention in rehabilitation. The evolution of a rational remediation program. In: Weinstein EA, Friedland RP, editors. *Hemi-inattention and hemisphere specialization: Advances in neurology*, New York: Raven Press; 1977, p. 62–82.
- [45] Webster JS, Rapport LJ, Godlewski MC, Abadee PS. Effect of attentional bias to right space on wheelchair mobility. *Journal of Clinical and Experimental Neuropsychology* 1994;16:129–37. <https://doi.org/10.1080/01688639408402623>.
- [46] Bachman L, Fein G, Davenport L, Price L. The indented paragraph reading test in the assessment of left hemi-neglect. *Archives of Clinical Neuropsychology* 1993;8:485–96. <https://doi.org/10.1016/0887-6177%2893%2990049-7>.
- [47] Verhoeven CL, Post MW, Schiemanck SK, van Zandvoort MJ, Vrancken PH, van Heugten CM. Is Cognitive Functioning 1 Year Poststroke Related to Quality of Life Domain? *Journal of Stroke & Cerebrovascular Diseases* 2011;20:450–458 9p.
- [48] Donnelly N, Guest R, Fairhurst M, Potter J, Deighton A, Patel M. Developing algorithms to enhance the sensitivity of cancellation tests of visuospatial neglect. *Behavior Research Methods, Instruments, & Computers : A Journal of the Psychonomic Society, Inc* 1999;31:668–73.
- [49] Sea M-JC, Henderson A. The reliability and validity of visuospatial inattention tests with stroke patients. *Occupational Therapy International* 1994;1:36–48. <https://doi.org/10.1002/oti.6150010106>.
- [50] Chen Sea M, Cermack SA, Henderson A. Performance of normal Chinese adults and right CVA patients on the Random Chinese Word Cancellation Test. *Clinical Neuropsychologist* 1993;7:239–49. <https://doi.org/10.1080/13854049308401896>.
- [51] Chen-Sea M. Unilateral neglect and functional significance among patients with stroke. *Occupational Therapy Journal of Research* 2001;21:223–240 18p.
- [52] Goldenberg G, Münsinger U, Karnath HO. Severity of neglect predicts accuracy of imitation in patients with right hemisphere lesions. *Neuropsychologia* 2009;47:2948–52. <https://doi.org/10.1016/j.neuropsychologia.2009.06.024>.
- [53] Wojciulik E, Rorden C, Clarke K, Husain M, Driver J. Group study of an “undercover” test for visuospatial neglect: Invisible cancellation can reveal more neglect than standard cancellation. *Journal of Neurology, Neurosurgery and Psychiatry* 2004;75:1356–8. <https://doi.org/10.1136/jnnp.2003.021931>.
- [54] Nijboer TCW, Kollen BJ, Kwakkel G. Time course of visuospatial neglect early after stroke: A longitudinal cohort study. *Cortex* 2013;49:2021–7. <https://doi.org/10.1016/j.cortex.2012.11.006>.
- [55] Bickerton WL, Samson D, Williamson J, Humphreys GW. Separating forms of neglect using the Apples Test: Validation and functional prediction in chronic and acute stroke. *Neuropsychology* 2011;25:567–80. <https://doi.org/10.1037/a0023501>.
- [56] Bickerton W-L, Demeyere N, Francis D, Kumar V, Remoundou M, Balani A, et al. The BCoS cognitive profile screen: Utility and predictive value for stroke. *Neuropsychology* 2015;29:638–48. <https://doi.org/10.1037/neu0000160>.
- [57] Kong A.P.-H., Chan J., Lau J.K.-L., Bickerton W.-L., Weekes B., Humphreys G. Developing a Cantonese Version of Birmingham Cognitive Screen for Stroke Survivors in Hong Kong. *Commun Disord Q* 2018;39:387–401. <https://doi.org/10.1177/1525740117720382>.
- [58] Pan X, Chen H, Bickerton WL, Lau JKL, Kong APH, Rotshtein P, et al. Preliminary findings on the reliability and validity of the cantonese birmingham cognitive screen in patients with acute ischemic stroke. *Neuropsychiatric Disease and Treatment* 2015;11:2377–90. <https://doi.org/10.2147/NDT.S85698>.
- [59] Demeyere N, Riddoch MJ, Slavkova ED, Bickerton WL, Humphreys GW. The Oxford Cognitive Screen (OCS): validation of a stroke-specific short cognitive screening tool. *Psychol Assess* 2015;27:883–94. <https://doi.org/10.1037/pas0000082>.
- [60] Kong A.P.-H., Lam P.H.-P., Ho D.W.-L., Lau J.K., Humphreys G.W., Riddoch J., et al. The Hong Kong version of the Oxford Cognitive Screen (HK-OCS): validation study for Cantonese-speaking chronic stroke survivors. *Aging Neuropsychol Cogn* 2016;23:530–48. <https://doi.org/10.1080/13825585.2015.1127321>.
- [61] Shendyapina M, Kuzmina E, Kazymae S, Petrova A, Demeyere N, Weekes BS. The Russian version of the Oxford Cognitive Screen: Validation study on stroke survivors. *Neuropsychology* 2019;33:77–92. <https://doi.org/10.1037/neu0000491>.
- [62] Chen P, Caulfield MD, Hartman AJ, O'Rourke J, Toglia J. Assessing viewer-centered and stimulus-centered spatial bias: The 3s spreadsheet test version 1. *Appl Neuropsychol Adult* 2017;24:532–9. <https://doi.org/10.1080/23279095.2016.1220382>.
- [63] Ota H, Fujii T, Suzuki K, Fukatsu R, Yamadori A. Dissociation of body-centered and stimulus-centered representations in unilateral neglect. *Neurology* 2001;57:2064–9.
- [64] Marsh EB, Hillis AE. Dissociation between egocentric and allocentric visuospatial and tactile neglect in acute stroke. *Cortex* 2008;44:1215–20. <https://doi.org/10.1016/j.cortex.2006.02.002>.

- [65] Mancuso M, Rosadoni S, Capitani D, Bickerton WL, Humphreys GW, De Tanti A, et al. Italian standardization of the Apples Cancellation Test. *Neurological Sciences* 2015;36:1233–40. <https://doi.org/10.1007/s10072-015-2088-2>.
- [66] Goedert KM, Chen P, Botticello A, Masmela JR, Adler U, Barrett AM. Psychometric evaluation of neglect assessment reveals motor-exploratory predictor of functional disability in acute-stage spatial neglect. *Archives of Physical Medicine and Rehabilitation* 2012;93:137–42. <https://doi.org/10.1016/j.apmr.2011.06.036>.
- [67] Na DL, Adair JC, Williamson DJG, Schwartz RL, Haws B, Heilman KM. Dissociation of sensory-attentional from motor-intentional neglect. *Journal of Neurology Neurosurgery and Psychiatry* 1998;64:331–8.
- [68] Husain M, Mattingley JB, Rorden C, Kennard C, Driver J. Distinguishing sensory and motor biases in parietal and frontal neglect. *Brain* 2000;123:1643–59.
- [69] Bartolomeo P, D’Erme P, Perri R, Gainotti G. Perception and action in hemispatial neglect. *Neuropsychologia* 1998;36:227–37. [https://doi.org/10.1016/S0028-3932\(97\)00104-8](https://doi.org/10.1016/S0028-3932(97)00104-8).
- [70] Bisiach E, Ricci R, Lualdi M, Colombo MR. Perceptual and response bias in unilateral neglect: two modified versions of the Milner Landmark Task. *Brain & Cognition* 1998;37:369–386.
- [71] Riestra AR, Crucian GP, Burks DW, Womack KB, Heilman KM. Extinction, working memory, and line bisection in spatial neglect. *Neurology* 2001;57:147–9. <https://doi.org/10.1212/WNL.57.1.147>.
- [72] Bisiach E, Geminiani G, Berti A, Rusconi ML. Perceptual and premotor factors of unilateral neglect. *Neurology* 1990;40:1278. <https://doi.org/10.1212/WNL.40.8.1278>.
- [73] Heilman KM, Bowers D, Coslett HB, Whelan H, Watson RT. Directional hypokinesia: Prolonged reaction times for leftward movements in patients with right hemisphere lesions and neglect. *Neurology* 1985;35:855–9.
- [74] Maeshima S, Truman G, Smith DS, Dohi N, Nakai K, Itakura T, et al. Is unilateral spatial neglect a single phenomenon? A comparative study between exploratory-motor and visual-counting tests. *J Neurol* 1997;244:412–7. <https://doi.org/10.1007/s004150050114>.
- [75] Stone A., Cooke D., Morton D., Steele M. Reliability of revised scoring methods for the Schenkenberg Line Bisection Test with adults following stroke: Preliminary findings. *Br J Occup Ther* 2019;82:750–8. <https://doi.org/10.1177/0308022619866377>.
- [76] Fortis P, Maravita A, Gallucci M, Ronchi R, Grassi E, Senna I, et al. Rehabilitating patients with left spatial neglect by prism exposure during a visuomotor activity. *Neuropsychology* 2010;24:681–97. <https://doi.org/10.1037/a0019476>.
- [77] Bailey MJ, Riddoch MJ, Crome P. Evaluation of a test battery for hemineglect in elderly stroke patients for use by therapists in clinical practice. *NeuroRehabilitation* 2000;14:139–50.
- [78] Vallar G, Zilli T, Gandola M, Bottini G. Productive and defective impairments in the neglect syndrome: graphic perseveration, drawing productions and optic prism exposure. *Cortex* 2006;42:911–20.
- [79] Perri R, Bartolomeo P, Gainotti G. Lack of impairments on leftward and rightward line extension tasks in neglect patients. *International Journal of Neuroscience* 2000;103:101–13.
- [80] Lee BH, Kang SJ, Park JM, Son Y, Lee KH, Adair JC, et al. The Character-line Bisection Task: A new test for hemispatial neglect. *Neuropsychologia* 2004;42:1715–24. <https://doi.org/10.1016/j.neuropsychologia.2004.02.015>.
- [81] Ye L-L, Cao L, Xie H-X, Shan G-X, Zhang Y-M, Song W-Q. Visual-spatial neglect after right-hemisphere stroke: behavioral and electrophysiological evidence. *Chin Med J (Engl)* 2019;132:1063–70. <https://doi.org/10.1097/CM9.0000000000000218>.
- [82] Karnath HO, Perenin MT. Tactile exploration of peripersonal space in patients with neglect. *NeuroReport* 1998;9:2273–7. <https://doi.org/10.1097/00001756-199807130-00024>.
- [83] Niemeier M, Karnath HO. Exploratory saccades show no direction-specific deficit in neglect. *Neurology* 2000;54:515–8.
- [84] Machner B, Mah YH, Gorgoraptis N, Husain M. How reliable is repeated testing for hemispatial neglect? Implications for clinical follow-up and treatment trials. *Journal of Neurology, Neurosurgery and Psychiatry* 2012;83:1032–4. <https://doi.org/10.1136/jnnp-2012-303296>.
- [85] Ten Brink AF, Visser-Meily JM, Nijboer TC. Dynamic assessment of visual neglect: The mobility assessment course as a diagnostic tool. *Journal of Clinical and Experimental Neuropsychology* 2018;40:161–72. <https://doi.org/10.1080/13803395.2017.1324562>.
- [86] Bisiach E, Ricci R, Silani G, Cossa FM, Crespi MG. Hypernesia in unilateral neglect. *Cortex: A Journal Devoted to the Study of the Nervous System & Behavior* 1999;35:701–711.
- [87] Kleinman JT, Gottesman RF, Davis C, Newhart M, Heidler-Gary J, Hillis AE. Gender differences in unilateral spatial neglect within 24 hours of ischemic stroke. *Brain Cogn* 2008;68:49–52. <https://doi.org/10.1016/j.bandc.2008.02.122>.
- [88] Butter CM, Mark VW, Heilman KM. An experimental analysis of factors underlying neglect in line bisection. *Journal of Neurology* 1988;51:1581–3.
- [89] Lindell AB, Jalas MJ, Tenovuo O, Brunila T, Voeten MJ, Hamalainen H. Clinical assessment of hemispatial neglect: Evaluation of different measures and dimensions. *The Clinical Neuropsychologist* 2007;21:479–97. <https://doi.org/10.1080/13854040600630061>.
- [90] Speedie LJ, Wertman E, Verfaellie M, Butter C, Silberman N, Liechtenstein M, et al. Reading direction and spatial neglect. *Cortex* 2002;38:59–67. [https://doi.org/10.1016/S0010-9452\(08\)70638-5](https://doi.org/10.1016/S0010-9452(08)70638-5).
- [91] Liu M, Chino N, Tuji T, Masakado Y, Hase K, Kimura A. Psychometric properties of the Stroke Impairment Assessment Set (SIAS). *Neurorehabilitation & Neural Repair* 2002;16:339–351.

- [92] Tsuji T, Liu M, Sonoda S, Domen K, Chino N. The Stroke Impairment Assessment Set: its internal consistency and predictive validity. *Archives of Physical Medicine & Rehabilitation* 2000;81:863-868.
- [93] Heilman KM, Valenstein E. Mechanisms underlying hemispatial neglect. *Annals of Neurology* 1979;5:166–70. <https://doi.org/10.1002/ana.410050210>.
- [94] Chung SJ, Park E, Ye BS, Lee HS, Chang H-J, Song D, et al. The Computerized Table Setting Test for Detecting Unilateral Neglect. *PLoS ONE* 2016;11:e0147030. <https://doi.org/10.1371/journal.pone.0147030>.
- [95] Pitzalis S, Spinelli D, Vallar G, Di Russo F. Transcutaneous electrical nerve stimulation effects on neglect: A visual-evoked potential study. *Frontiers in Human Neuroscience* 2013. <https://doi.org/10.3389/fnhum.2013.00111>.
- [96] Sperber C, Karnath HO. Diagnostic validity of line bisection in the acute phase of stroke. *Neuropsychologia* 2016;82:200–4. <https://doi.org/10.1016/j.neuropsychologia.2016.01.026>.
- [97] Agrell BM, Dehlin OI, Dahlgren CJ. Neglect in elderly stroke patients: a comparison of five tests. *Psychiatry Clin Neurosci* 1997;51:295–300.
- [98] Bartolomeo P, Chokron S. Egocentric frame of reference: its role in spatial bias after right hemisphere lesions. *Neuropsychologia* 1999;37:881–94. [https://doi.org/10.1016/S0028-3932\(98\)00150-X](https://doi.org/10.1016/S0028-3932(98)00150-X).
- [99] Williamson JB, Lamb DG, Burtis DB, Haque S, Zilli EM, Kesayan T, et al. Right hemispatial ipsilesional neglect with chronic right hemisphere strokes. *Journal of Clinical and Experimental Neuropsychology* 2017;1–10. <https://doi.org/10.1080/13803395.2017.1347606>.
- [100] Wang Q, Sonoda S, Hanamura M, Okazaki H, Saitoh E. Line bisection and rebisection: the crossover effect of space location. *Neurorehabilitation & Neural Repair* 2005;19:84–92 9p.
- [101] Lee BH, Kwon SU, Kwon JC, Baek MJ, Lee KH, Kim GH, et al. Line quadrisection errors in patients with hemispatial neglect. *Neurocase* 2011;17:372–80. <https://doi.org/10.1080/13554794.2010.532136>.
- [102] Charras P, Lupiáñez J, Bartolomeo P. Assessing the weights of visual neglect: A new approach to dissociate defective symptoms from productive phenomena in length estimation. *Neuropsychologia* 2010;48:3371–5. <https://doi.org/10.1016/j.neuropsychologia.2010.07.006>.
- [103] Mancini F, Bricolo E, Mattioli FC, Vallar G. Visuo-haptic interactions in unilateral spatial neglect: The cross modal Judd illusion. *Frontiers in Psychology* 2011;2. <https://doi.org/10.3389/fpsyg.2011.00341>.
- [104] Pirondini E, Goldshuv-Ezra N, Zinger N, Britz J, Soroker N, Deouell LY, et al. Resting-state EEG topographies: Reliable and sensitive signatures of unilateral spatial neglect. *NeuroImage-Clin* 2020;26:102237. <https://doi.org/10.1016/j.nicl.2020.102237>.
- [105] Anderson B. A mathematical model of line bisection behaviour in neglect. *Brain* 1996;119 (Pt 3):841–50. <https://doi.org/10.1093/brain/119.3.841>.
- [106] Edwards MG, Humphreys GW. Pointing and grasping in unilateral visual neglect: Effect of on-line visual feedback in grasping. *Neuropsychologia* 1999;37:959–73. [https://doi.org/10.1016/S0028-3932\(98\)00132-8](https://doi.org/10.1016/S0028-3932(98)00132-8).
- [107] Bailey MJ, Riddoch M, Crome P. Test-retest stability of three tests for unilateral visual neglect in patients with stroke: Star Cancellation, Line Bisection, and the Baking Tray Task. *Neuropsychological Rehabilitation* 2004;14:403–19. <https://doi.org/10.1080/09602010343000282>.
- [108] Cerrato A, Pacella D, Palumbo F, Beauvais D, Ponticorvo M, Miglino O, et al. E-TAN, a technology-enhanced platform with tangible objects for the assessment of visual neglect: A multiple single-case study. *Neuropsychol Rehabil* 2020. <https://doi.org/10.1080/09602011.2020.1762671>.
- [109] Wenman R, Bowen A, Tallis RC, Gardener E, Cross S, Niven D. Use of a randomised single case experimental design to evaluate therapy for unilateral neglect. *Neuropsychol Rehabil* 2003;13:441–59. <https://doi.org/10.1080/09602010343000048>.
- [110] Tyryshkin K, Coderre AM, Glasgow JI, Herter TM, Bagg SD, Dukelow SP, et al. A robotic object hitting task to quantify sensorimotor impairments in participants with stroke. *J NeuroEng Rehabil* 2014;11:47. <https://doi.org/10.1186/1743-0003-11-47>.
- [111] Brunila T, Jalas M, Lindell JA, Tenovuo O, Hamalainen H. The two part picture in detection of visuospatial neglect. *Clin Neuropsychol* 2003;17:45–53.
- [112] Làdavas E, Umiltà C, Ziani P, Brogi A, Minarini M. The role of right side objects in left side neglect: A dissociation between perceptual and directional motor neglect. *Neuropsychologia* 1993;31:761–73. [https://doi.org/10.1016/0028-3932\(93\)90127-L](https://doi.org/10.1016/0028-3932(93)90127-L).
- [113] Gainotti G, Messerli P, Tissot R. Qualitative analysis of unilateral spatial neglect in relation to laterality of cerebral lesions. *J Neurol Neurosurg Psychiatry* 1972;35:545–50. <https://doi.org/10.1136/jnnp.35.4.545>.
- [114] Johannsen L, Karnath HO. How efficient is a simple copying task to diagnose spatial neglect in its chronic phase? *Journal of Clinical and Experimental Neuropsychology* 2004;26:251–6. <https://doi.org/10.1076/jcen.26.2.251.28085>.
- [115] Ogden JA. Contralesional neglect of constructed visual images in right and left brain-damaged patients. *Neuropsychologia* 1985;23:273–7. [https://doi.org/10.1016/0028-3932\(85\)90112-5](https://doi.org/10.1016/0028-3932(85)90112-5).
- [116] Warren M. Identification of visual scanning deficits in adults after cerebrovascular accident. *Journal of Occupational Therapy* 1990;44:391–9.
- [117] Marshall JC, Halligan PW. Visuo-spatial neglect: A new copying test to assess perceptual parsing. *Journal of Neurology* 1993;240:37–40. <https://doi.org/10.1007/BF00838444>.

- [118] Osterrieth PA. Le test de copie d'une figure complexe; contribution à l'étude de la perception et de la mémoire. [Test of copying a complex figure; contribution to the study of perception and memory.]. *Archives de Psychologie* 1944;30:206–356.
- [119] Chen P, Goedert KM. Clock Drawing in Spatial Neglect: A Comprehensive Analysis of Clock Perimeter, Placement, and Accuracy. *Journal of Neuropsychology* May 3;6:270–89. <https://doi.org/10.1111/j.1748-6653.2012.02028.x>.
- [120] Rengachary J, d'Avossa G, Sapir A, Shulman GL, Corbetta M. Is the Posner Reaction Time Test more accurate than clinical tests in detecting left neglect in acute and chronic stroke? *Archives of Physical Medicine & Rehabilitation* 2009;90:2081–2088 8p. <https://doi.org/10.1016/j.apmr.2009.07.014>.
- [121] Delazer M, Sojer M, Ellmerer P, Boehme C, Benke T. Eye-Tracking Provides a Sensitive Measure of Exploration Deficits After Acute Right MCA Stroke. *Front Neurol* 2018;9:359. <https://doi.org/10.3389/fneur.2018.00359>.
- [122] Sambo CF, Vallar G, Fortis P, Ronchi R, Posteraro L, Forster B, et al. Visual and spatial modulation of tactile extinction: Behavioural and electrophysiological evidence. *Frontiers in Human Neuroscience* 2012;6:217. <https://doi.org/10.3389/fnhum.2012.00217>.
- [123] Veronelli L, Vallar G, Marinelli CV, Primativo S, Arduino LS. Line and word bisection in right-brain-damaged patients with left spatial neglect. *Experimental Brain Research* 2014;232:133–46. <https://doi.org/10.1007/s00221-013-3726-7>.
- [124] Nokleby K, Boland E, Bergersen H, Schanke A-K, Farner L, Wagle J, et al. Screening for cognitive deficits after stroke: A comparison of three screening tools. *Clinical Rehabilitation* 2008;22:1095–104. <https://doi.org/10.1177/0269215508094711>.
- [125] Ishiai S, Ishiai S, Sugishita M, Sugishita M, Ichikawa T, Ichikawa T, et al. Clock-drawing test and unilateral spatial neglect. *Neurology* 1993;43:106–10. https://doi.org/10.1212/WNL.43.1_Part_1.106.
- [126] Klinke ME, Hjaltason H, Tryggvadóttir GB, Jónsdóttir H. Hemispatial neglect following right hemisphere stroke: clinical course and sensitivity of diagnostic tasks. *Top Stroke Rehabil* 2018;25:120–30. <https://doi.org/10.1080/10749357.2017.1394632>.
- [127] Rousseaux M, Beis JM, Pradat-Diehl P, Martin Y, Bartolomeo P, Bernati T, et al. A battery for assessing spatial neglect: Norms and effects of age, educational level, sex, hand, and laterality. *Revue Neurologique* 2001;157:1385–400.
- [128] Kaufmann B, Frey J, Pflugshaupt T, Wyss P, Paladini R, Vanbellingen T, et al. The spatial distribution of perseverations in neglect patients during a nonverbal fluency task depends on the integrity of the right putamen. *Neuropsychologia* 2018;115:42–50. <https://doi.org/10.1016/j.neuropsychologia.2018.01.025>.
- [129] Chen-Sea MJ. Validating the Draw-A-Man Test as a personal neglect test. *Journal of Occupational Therapy* 2000;54:391–7.
- [130] Bartolomeo P, Urbanski M, Chokron S, Chainay H, Moroni C, Siéroff E, et al. Neglected attention in apparent spatial compression. *Neuropsychologia* 2004;42:49–61. [https://doi.org/10.1016/S0028-3932\(03\)00146-5](https://doi.org/10.1016/S0028-3932(03)00146-5).
- [131] Pizzamiglio L, Antonucci G, Judica A, Montenero P, Razzano C, Zoccolotti P. Cognitive rehabilitation of the hemineglect disorder in chronic patients with unilateral right brain damage. *J Clin Exp Neuropsychol* 1992;14:901–23.
- [132] Zoccolotti P, Antonucci G, Judica A, Montenero P, Pizzamiglio L, Razzano C. Incidence and Evolution of the Hemineglect Disorder in Chronic Patients with Unilateral Right Brain Damage. *Null* 1989;47:209–16. <https://doi.org/10.3109/00207458908987435>.
- [133] Verdon V, Schwartz S, Lovblad K-O, Hauert C-A, Vuilleumier P. Neuroanatomy of hemispatial neglect and its functional components: A study using voxel-based lesion-symptom mapping. *Brain* 2010;133:880–94. <https://doi.org/10.1093/brain/awp305>.
- [134] Caplan B. Assessment of unilateral neglect: A new reading test. *Journal of Clinical and Experimental Neuropsychology* 1987;9:359–64. <https://doi.org/10.1080/01688638708405056>.
- [135] Towle D, Lincoln NB. Use of the Indented Paragraph test with right hemisphere-damaged stroke patients. *Br J Clin Psychol* 1991;30:37–45.
- [136] Emerson RL, García-Molina A, López Carballo J, García Fernández J, Aparicio-López C, Novo J, et al. Visual search in unilateral spatial neglect: The effects of distractors on a dynamic visual search task. *Appl Neuropsychol Adult* 2019;26:401–10. <https://doi.org/10.1080/23279095.2018.1434522>.
- [137] Cazzoli D., Muri R.M., Schumacher R., Von Arx S., Chaves S., Gutbrod K., et al. Theta burst stimulation reduces disability during the activities of daily living in spatial neglect. *Brain* 2012;135:3426–39. <https://doi.org/10.1093/brain/aws182>.
- [138] Galletta EE, Campanelli L, Maul KK, Barrett AM. Assessment of neglect dyslexia with functional reading materials. *Top* 2014;21:75–86. <https://doi.org/10.1310/tsr2101-75>.
- [139] Ronchi R, Algeri L, Chiapella L, Gallucci M, Spada MS, Vallar G. Left neglect dyslexia: Perseveration and reading error types. *Neuropsychologia* 2016;89:453–64. <https://doi.org/10.1016/j.neuropsychologia.2016.07.023>.
- [140] Reinhart S, Schunck A, Schaadt AK, Adams M, Simon A, Kerkhoff G. Assessing neglect dyslexia with compound words. *Neuropsychology* 2016;30:869–73. <https://doi.org/10.1037/neu0000307>.
- [141] Lee BH, Suh MK, Kim EJ, Seo SW, Choi KM, Kim GM, et al. Neglect dyslexia: Frequency, association with other hemispatial neglects, and lesion localization. *Neuropsychologia* 2009;47:704–10. <https://doi.org/10.1016/j.neuropsychologia.2008.11.027>.

- [142] Gainotti G, D'Erme P, Bartolomeo P. Early orientation of attention toward the half space ipsilateral to the lesion in patients with unilateral brain damage. *J Neurol Neurosurg Psychiatry* 1991;54:1082–9. <https://doi.org/10.1136/jnnp.54.12.1082>.
- [143] Toglia J, Cermak SA. Dynamic Assessment and Prediction of Learning Potential in Clients With Unilateral Neglect. *Am J Occup Ther* 2009;63:569–79. <https://doi.org/10.5014/ajot.63.5.569>.
- [144] Buxbaum LJ, Coslett HB. Neglect of chimeric figures: Two halves are better than a whole. *Neuropsychologia* 1994;32:275–88. [https://doi.org/10.1016/0028-3932\(94\)90131-7](https://doi.org/10.1016/0028-3932(94)90131-7).
- [145] Sarri M, Greenwood R, Kalra L, Papps B, Husain M, Driver J. Prism adaptation aftereffects in stroke patients with spatial neglect: Pathological effects on subjective straight ahead but not visual open-loop pointing. *Neuropsychologia* 2008;46:1069–80. <https://doi.org/10.1016/j.neuropsychologia.2007.11.005>.
- [146] di Pellegrino G, Frassinetti F, Basso G. Coordinate Frames for Naming Misoriented Chimerics: A Case Study of Visuo-Spatial Neglect. *Cortex* 1995;31:767–77. [https://doi.org/10.1016/S0010-9452\(13\)80027-5](https://doi.org/10.1016/S0010-9452(13)80027-5).
- [147] Mattingley JB, Bradshaw JL, Phillips JG, Bradshaw JA. Reversed perceptual asymmetry for faces in left unilateral neglect. *Brain and Cognition* 1993;23:145–65. <https://doi.org/10.1006/brcg.1993.1052>.
- [148] Massironi M, Antonucci G, Pizzamiglio L, Vitale MV, Zoccolotti P. The Wundt-Jastrow illusion in the study of spatial hemi-inattention. *Neuropsychologia* 1988;26:161–6.
- [149] Paladini RE, Wyss P, Kaufmann BC, Urwyler P, Nef T, Cazzoli D, et al. Re-fixation and perseveration patterns in neglect patients during free visual exploration. *Journal of Neuroscience* 2019;49:1244–53. <https://doi.org/10.1111/ejn.14309>.
- [150] Berger MF, Johannsen L, Karnath HO. Time Course of Eye and Head Deviation in Spatial Neglect. *Neuropsychology* 2008;22:697–702. <https://doi.org/10.1037/a0013351>.
- [151] Karnath HO, Niemeier M, Dichgans J. Space exploration in neglect. *Brain* 1998;121 (Pt 12):2357–67. <https://doi.org/10.1093/brain/121.12.2357>.
- [152] Kaufmann BC, Cazzoli D, Pflugshaupt T, Bohlhalter S, Vanbellingen T, Müri RM, et al. Eyetracking during free visual exploration detects neglect more reliably than paper-pencil tests. *Cortex* 2020;129:223–35. <https://doi.org/10.1016/j.cortex.2020.04.021>.
- [153] Kortman B, Nicholls K. Assessing for Unilateral Spatial Neglect Using Eye-Tracking Glasses: A Feasibility Study. *Occup Ther Health Care* 2016;30:344–55. <https://doi.org/10.1080/07380577.2016.1208858>.
- [154] Bourgeois A, Chica AB, Migliaccio R, de Schotten MT, Bartolomeo P. Cortical control of inhibition of return: Evidence from patients with inferior parietal damage and visual neglect. *Neuropsychologia* 2012;50:800–9. <https://doi.org/10.1016/j.neuropsychologia.2012.01.014>.
- [155] Dombrowe I, Donk M, Wright H, Olivers CNL, Humphreys GW. The contribution of stimulus-driven and goal-driven mechanisms to feature-based selection in patients with spatial attention deficits. *Cognitive Neuropsychology* 2012;29:249–74. <https://doi.org/10.1080/02643294.2012.712509>.
- [156] Pitteri M, Chen P, Passarini L, Albanese S, Meneghelli F, Barrett AM. Conventional and functional assessment of spatial neglect: Clinical practice suggestions. *Neuropsychology* 2018;32:835–42. <https://doi.org/10.1037/neu0000469>.
- [157] Bisiach E, Luzzatti C. Unilateral Neglect of Representational Space. *Cortex* 1978;14:129–33. [https://doi.org/10.1016/S0010-9452\(78\)80016-1](https://doi.org/10.1016/S0010-9452(78)80016-1).
- [158] Guariglia C, Palermo L, Piccardi L, Iaria G, Incoccia C. Neglecting the left side of a city square but not the left side of its clock: prevalence and characteristics of representational neglect. *PLoS One* 2013;8:e67390. <https://doi.org/10.1371/journal.pone.0067390>.
- [159] Bartolomeo P, D'erme P, Gainotti G. The relationship between visuospatial and representational neglect. *Neurology* 1994;44:1710–4.
- [160] Bourlon C, Pradat-Diehl P, Duret C, Azouvi P, Bartolomeo P. Seeing and imagining the “same” objects in unilateral neglect. *Neuropsychologia* 2008;46:2602–6. <https://doi.org/10.1016/j.neuropsychologia.2008.04.002>.
- [161] Rode G, Revol P, Rossetti Y, Boisson D, Bartolomeo P. Looking while imagining: The influence of visual input on representational neglect. *Neurology* 2007;68:432–7. <https://doi.org/10.1212/01.wnl.0000252936.54063.b0>.
- [162] Rode G, Cotton F, Revol P, Jacquin-Courtois S, Rossetti Y, Bartolomeo P. Representation and disconnection in imaginal neglect. *Neuropsychologia* 2010;48:2903–11. <https://doi.org/10.1016/j.neuropsychologia.2010.05.032>.
- [163] Grossi D, Angelini R, Pecchinenda A, Pizzamiglio L. Left imaginal neglect in heminattention: experimental study with the O'clock Test. *Behavioural Neurology* 1993;6:155–8. <https://doi.org/10.3233/BEN-1993-6307>.
- [164] Cocchini G, Beschin N, Jehkonen M. The Fluff Test: A simple task to assess body representation neglect. *Neuropsychological Rehabilitation* 2001;11:17–31. <https://doi.org/10.1080/09602010042000132>.
- [165] Bowen A, Gardener E, Cross S, Tyrrell P, Graham J. Developing functional outcome measures for unilateral neglect: A pilot study. *Neuropsychological Rehabilitation* 2005;15:97–113. <https://doi.org/10.1080/09602010443000010>.
- [166] Glockner D, Bittl P, Kerkhoff G. Construction and psychometric properties of a novel test for body representational neglect (Vest Test). *Restorative Neurology and Neuroscience* 2006;24:303–17.

- [167] Bolognini N, Casanova D, Maravita A, Vallar G. Bisecting real and fake body parts: Effects of prism adaptation after right brain damage. *Frontiers in Human Neuroscience* 2012. <https://doi.org/10.3389/fnhum.2012.00154>.
- [168] Bisiach E, Perani D, Vallar G, Berti A. Unilateral neglect: Personal and extra-personal. *Neuropsychologia* 1986;24:759–67. [https://doi.org/10.1016/0028-3932\(86\)90075-8](https://doi.org/10.1016/0028-3932(86)90075-8).
- [169] Cunningham LJ, O'Rourke K, Finlay C, Gallagher M. A preliminary investigation into the psychometric properties of the Dublin Extrapersonal Neglect Assessment (DENA): A novel screening tool for extrapersonal neglect. *Neuropsychological Rehabilitation* 2017;27:349–68. <https://doi.org/10.1080/09602011.2015.1084334>.
- [170] Borsotti M, Mosca IE, Di Lauro F, Pancani S, Bracali C, Dore T, et al. The Visual Scanning Test: a newly developed neuropsychological tool to assess and target rehabilitation of extrapersonal visual unilateral spatial neglect. *Neurol Sci* 2020;41:1145–52. <https://doi.org/10.1007/s10072-019-04218-2>.
- [171] Villarreal S, Linnavuo M, Sepponen R, Vuori O, Jokinen H, Hietanen M. Dual-Task in Large Perceptual Space Reveals Subclinical Hemispatial Neglect. *Journal of the International Neuropsychological Society* 2020;26:993–1005. <https://doi.org/10.1017/S1355617720000508>.
- [172] Aimola L, Schindler I, Simone AM, Venneri A. Near and far space neglect: Task sensitivity and anatomical substrates. *Neuropsychologia* 2012;50:1115–23. <https://doi.org/10.1016/j.neuropsychologia.2012.01.022>.
- [173] Van Der Stoep N, Visser-Meily JMA, Kappelle LJ, De Kort PLM, Huisman KD, Eijssackers ALH, et al. Exploring near and far regions of space: Distance-specific visuospatial neglect after stroke. *Journal of Clinical and Experimental Neuropsychology* 2013;35:799–811. <https://doi.org/10.1080/13803395.2013.824555>.
- [174] Berti A, Smania N, Rabuffetti M, Ferrarin M, Spinazzola L, D'Amico A, et al. Coding of far and near space during walking in neglect patients. *Neuropsychology* 2002;16:390–9. <https://doi.org/10.1037/0894-4105.16.3.390>.
- [175] Ogourtsova T, Archambault PS, Lamontagne A. Post-stroke visual neglect affects goal-directed locomotion in different perceptuo-cognitive conditions and on a wide visual spectrum. *Restorative Neurology and Neuroscience* 2018;36:313–31. <https://doi.org/10.3233/RNN-170766>.
- [176] Pizzamiglio L, Cappa S, Vallar G, Zoccolotti P, Bottini G, Ciurli P, et al. Visual Neglect for Far and Near Extra-Personal Space in Humans. *Cortex* 1989;25:471–7. [https://doi.org/10.1016/S0010-9452\(89\)80060-7](https://doi.org/10.1016/S0010-9452(89)80060-7).
- [177] Whitehouse CE, Green J, Giles SM, Rahman R, Coolican J, Eskes GA. Development of the Halifax Visual Scanning Test: A new measure of visual-spatial neglect for personal, peripersonal, and extrapersonal space. *Journal of the International Neuropsychological Society* 2019;25:490–500. <https://doi.org/10.1017/S135561771900002X>.
- [178] Montedoro V, Alsamour M, Dehem S, Lejeune T, Dehez B, Edwards MG. Robot Diagnosis Test for Egocentric and Allocentric Hemineglect. *Archives of Clinical Neuropsychology : The Official Journal of the National Academy of Neuropsychologists* 2019;34:481–94. <https://doi.org/10.1093/arclin/acy062>.
- [179] Dawson AM, Buxbaum LJ, Rizzo AA. The virtual reality lateralized attention test: Sensitivity and validity of a new clinical tool for assessing hemispatial neglect. *Virtual Rehabil., IWVR*, Vancouver, BC: 2008, p. 77–82. <https://doi.org/10.1109/ICVR.2008.4625140>.
- [180] Buxbaum LJ, Palermo MA, Mastrogiovanni D, Read MS, Rosenberg-Pitonyak E, Rizzo AA, et al. Assessment of spatial attention and neglect with a virtual wheelchair navigation task. *Journal of Clinical and Experimental Neuropsychology* 2008;30:650–60. <https://doi.org/10.1080/13803390701625821>.
- [181] Fordell H, Bodin K, Bucht G, Malm J. A virtual reality test battery for assessment and screening of spatial neglect. *Acta Neurologica Scandinavica* 2011;123:167–74. <https://doi.org/10.1111/j.1600-0404.2010.01390.x>.
- [182] Dvorkin AY, Bogey RA, Harvey RL, Patton JL. Mapping the neglected space: gradients of detection revealed by virtual reality. *Neurorehabil Neural Repair* 2012;26:120–31. <https://doi.org/10.1177/1545968311410068>.
- [183] Peskine A, Rosso C, Box N, Galland A, Caron E, Rautureau G, et al. Virtual reality assessment for visuospatial neglect: importance of a dynamic task. *Journal of Neurology, Neurosurgery and Psychiatry* 2011;82:1407. <https://doi.org/10.1136/jnnp.2010.217513>.
- [184] Kim K, Kim J, Ku J, Kim DY, Chang WH, Shin DI, et al. A Virtual Reality Assessment and Training System for Unilateral Neglect. *CyberPsychology & Behavior* 2004;7:742–9. <https://doi.org/10.1089/cpb.2004.7.742>.
- [185] Kim J, Kim K, Kim DY, Chang WH, Park C-I, Ohn SH, et al. Virtual environment training system for rehabilitation of stroke patients with unilateral neglect: Crossing the virtual street. *CyberPsychology & Behavior* 2007;10:7–15. <https://doi.org/10.1089/cpb.2006.9998>.
- [186] Kim DY, Ku J, Chang WH, Park TH, Lim JY, Han K, et al. Assessment of post-stroke extrapersonal neglect using a three-dimensional immersive virtual street crossing program. *Acta Neurologica Scandinavica* 2010;121:171–7. <https://doi.org/10.1111/j.1600-0404.2009.01194.x>.
- [187] Navarro MD, Llorens R, Noe E, Ferri J, Alcaniz M. Validation of a low-cost virtual reality system for training street-crossing. A comparative study in healthy, neglected and non-neglected stroke individuals. *Neuropsychol* 2013;23:597–618. <https://doi.org/10.1080/09602011.2013.806269>.
- [188] Ogourtsova T, Archambault P, Sangani S, Lamontagne A. Ecological Virtual Reality Evaluation of Neglect Symptoms (EVENS): Effects of Virtual Scene Complexity in the Assessment of Poststroke Unilateral Spatial Neglect. *Neurorehabilitation and Neural Repair* 2018;32:46–61. <https://doi.org/10.1177/1545968317751677>.
- [189] Erez AB-H, Katz N, Ring H, Soroker N. Assessment of spatial neglect using computerised feature and conjunction visual search tasks. *Neuropsychological Rehabilitation* 2009;19:677–95. <https://doi.org/10.1080/09602010802711160>.

- [190] Deouell LY, Sacher Y, Soroker N. Assessment of spatial attention after brain damage with a dynamic reaction time test. *Journal of the International Neuropsychological Society* 2005;11:697–707. <https://doi.org/10.1017/S1355617705050824>.
- [191] Schendel K, Dronkers NF, Turken AU. Not Just Language: Persisting Lateralized Visuospatial Impairment after Left Hemisphere Stroke. *J Int Neuropsychol Soc* 8;22:695–704. <https://doi.org/10.1017/S1355617716000515>.
- [192] List A, Brooks JL, Esterman M, Flevaris AV, Landau AN, Bowman G, et al. Visual hemispatial neglect, re-assessed. *Journal of the International Neuropsychological Society* 2008;14:243–56. <https://doi.org/10.1017/S1355617708080284>.
- [193] Nyffeler T, Cazzoli D, Hess CW, Muri RM. One session of repeated parietal theta burst stimulation trains induces long-lasting improvement of visual neglect. *Stroke* 2009;40:2791–6. <https://doi.org/10.1161/STROKEAHA.109.552323>.
- [194] Bublak P, Finke K, Krummenacher J, Preger R, Kyllingsbaek S, Muller HJ, et al. Usability of a theory of visual attention (TVA) for parameter-based measurement of attention II: Evidence from two patients with frontal or parietal damage. *Journal of the International Neuropsychological Society* 2005;11:843–54. <https://doi.org/10.1017/S1355617705050988>.
- [195] Mazer BL, Sofer S, Korner-Bitensky N, Gelinas N. Use of the UFOV to evaluate and retrain visual attention skills in clients with stroke: A pilot study. *American Journal of Occupational Therapy* 2001;55:552–7. <https://doi.org/10.5014/ajot.55.5.552>.
- [196] van Kessel ME, van Nes IJW, Geurts ACH, Brouwer WH, Fasotti L. Visuospatial asymmetry in dual-task performance after subacute stroke. *J Neuropsychol* 2013;7:72–90. <https://doi.org/10.1111/j.1748-6653.2012.02036.x>.
- [197] van Kessel ME, van Nes IJW, Brouwer WH, Geurts ACH, Fasotti L. Visuospatial asymmetry and non-spatial attention in subacute stroke patients with and without neglect. *Cortex* 2010;46:602–12. <https://doi.org/10.1016/j.cortex.2009.06.004>.
- [198] Bonato M, Priftis K, Marenzi R, Umiltà C, Zorzi M. Increased attentional demands impair contralesional space awareness following stroke. *Neuropsychologia* 2010;48:3934–40. <https://doi.org/10.1016/j.neuropsychologia.2010.08.022>.
- [199] Bonato M. Unveiling residual, spontaneous recovery from subtle hemispatial neglect three years after stroke. *Front Human Neurosci* 2015;9:1–9. <https://doi.org/10.3389/fnhum.2015.00413>.
- [200] Blini E, Romeo Z, Spironelli C, Pitteri M, Meneghelli F, Bonato M, et al. Multi-tasking uncovers right spatial neglect and extinction in chronic left-hemisphere stroke patients. *Neuropsychologia* 2016;92:147–57. <https://doi.org/10.1016/j.neuropsychologia.2016.02.028>.
- [201] Sacher Y, Serfaty C, Deouell L, Sapir A, Henik A, Soroker N. Role of disengagement failure and attentional gradient in unilateral spatial neglect -- a longitudinal study. *Disability & Rehabilitation* 2004;26:746–755 10p.
- [202] D'Erme P, Robertson I, Bartolomeo P, Daniele A, Gainotti G. Early rightwards orienting of attention on simple reaction time performance in patients with left-sided neglect. *Neuropsychologia* 1992;30:989–1000. [https://doi.org/10.1016/0028-3932\(92\)90050-V](https://doi.org/10.1016/0028-3932(92)90050-V).
- [203] Carter AR, McAvoy MP, Siegel JS, Hong X, Astafiev SV, Rengachary J, et al. Differential white matter involvement associated with distinct visuospatial deficits after right hemisphere stroke. *Cortex* 2017;1:81–97. <https://doi.org/10.1016/j.cortex.2016.12.009>.
- [204] Jacquin-Courtois S, Rode G, Pavani F, O'Shea J, Giard MH, Boisson D, et al. Effect of prism adaptation on left dichotic listening deficit in neglect patients: glasses to hear better? *Brain* 2010;133:895–908. <https://doi.org/10.1093/brain/awp327>.
- [205] Roberts KL, Lau JKL, Chechlacz M, Humphreys GW. Spatial and temporal attention deficits following brain injury: A neuroanatomical decomposition of the temporal order judgement task. *Cognitive Neuropsychology* 2012;29:300–24. <https://doi.org/10.1080/02643294.2012.722548>.
- [206] Zimmer U, Lewald J, Karnath HO. Disturbed sound lateralization in patients with spatial neglect. *Journal of Cognitive Neuroscience* 2003;15:694–703. <https://doi.org/10.1162/089892903322307410>.
- [207] Pavani F, Làdavas E, Driver J. Selective deficit of auditory localisation in patients with visuospatial neglect. *Neuropsychologia* 2002;40:291–301. [https://doi.org/10.1016/S0028-3932\(01\)00091-4](https://doi.org/10.1016/S0028-3932(01)00091-4).
- [208] Prilop L, Gutschalk A. Auditory-cortex lesions impair contralateral tone-pattern detection under informational masking. *Cortex* 2017;95:1–14. <https://doi.org/10.1016/j.cortex.2017.07.007>.
- [209] Sinnett S, Juncadella M, Rafal R, Azañón E, Soto-Faraco S. A dissociation between visual and auditory hemi-inattention: Evidence from temporal order judgements. *Neuropsychologia* 2007;45:552–60. <https://doi.org/10.1016/j.neuropsychologia.2006.03.006>.
- [210] Cumming TB, Blomstrand C, Bernhardt J, Linden T. The NIH stroke scale can establish cognitive function after stroke. *Cerebrovasc Dis* 2010;30:7–14. <https://doi.org/10.1159/000313438>.
- [211] Maravita A, McNeil J, Malhotra P, Greenwood R, Husain M, Driver J. Prism adaptation can improve contralesional tactile perception in neglect. *Neurology* 2003;60:1829–31. <https://doi.org/10.1212/WNL.60.11.1829>.

- [212] Kinsella G, Packer S, Ng K, Olver J, Stark R. Continuing issues in the assessment of neglect. *Neuropsychological Rehabilitation* 1995;5:239–58. <https://doi.org/10.1080/09602019508401469>.
- [213] Anton HA, Hershler C, Lloyd P, Murray D. Visual neglect and extinction: a new test. *Arch Phys Med Rehabil* 1988;69:1013–6.
- [214] Vossel S, Eschenbeck P, Weiss PH, Weidner R, Saliger J, Karbe H, et al. Visual extinction in relation to visuospatial neglect after right-hemispheric stroke: quantitative assessment and statistical lesion-symptom mapping. *Journal of Neurology, Neurosurgery & Psychiatry* 2011;82:862–868 7p. <https://doi.org/10.1136/jnnp.2010.224261>.
- [215] Azouvi P. Functional consequences and awareness of unilateral neglect: Study of an evaluation scale. *Neuropsychological Rehabilitation* 1996;6:133–50. <https://doi.org/10.1080/713755501>.
- [216] Chen P, Hreha K, Kong Y, Barrett AM. Impact of spatial neglect on stroke rehabilitation: evidence from the setting of an inpatient rehabilitation facility. *Archives of Physical Medicine and Rehabilitation* 2015;96:1458–66. <https://doi.org/10.1016/j.apmr.2015.03.019>.
- [217] Chen P, Hreha K, Fortis P, Goedert K, Barrett A. Functional assessment of spatial neglect: A review of the catherine bergego scale and an introduction of the kessler foundation neglect assessment process. *Topics in Stroke Rehabilitation* 2012;19:423–35. <https://doi.org/10.1310/tsr1905-423>.
- [218] Towle D, Lincoln NB. Development of a questionnaire for detecting everyday problems in stroke patients with unilateral visual neglect. *Clinical Rehabilitation* 1991;5:135–40. <https://doi.org/10.1177/026921559100500208>.
- [219] Gardarsdóttir S, Kaplan S. Validity of the Arnadóttir OT-ADL Neurobehavioral Evaluation (A-ONE): performance in activities of daily living and neurobehavioral impairments of persons with left and right hemisphere damage. *Am J Occup Ther* 2002;56:499–508. <https://doi.org/10.5014/ajot.56.5.499>.
- [220] Azouvi P, Olivier S, de Montety G, Samuel C, Louis-Dreyfus A, Tesio L. Behavioral assessment of unilateral neglect: study of the psychometric properties of the Catherine Bergego Scale. *Arch Phys Med Rehabil* 2003;84:51–7.
- [221] Ten Brink AF, Verwer JH, Biesbroek J, Visser-Meily JM, Nijboer TC. Differences between left- and right-sided neglect revisited: A large cohort study across multiple domains. *Journal of Clinical and Experimental Neuropsychology* 2017;39:707–23. <https://doi.org/10.1080/13803395.2016.1262333>.
- [222] Zoccolotti P, Judica A. Functional evaluation of hemineglect by means of a semistructured scale: Personal extrapersonal differentiation. *Null* 1991;1:33–44. <https://doi.org/10.1080/09602019108401378>.
- [223] Zoccolotti P, Antonucci G, Judica A. Psychometric characteristics of two semi-structured scales for the functional evaluation of hemi-inattention in extrapersonal and personal space. *Null* 1992;2:179–91. <https://doi.org/10.1080/09602019208401407>.
- [224] Bienkiewicz MMN, Brandi ML, Hughes C, Voitl A, Hermsdorfer J. The complexity of the relationship between neuropsychological deficits and impairment in everyday tasks after stroke. *Brain and Behavior* 2015;5. <https://doi.org/10.1002/brb3.371>.
- [225] Beschin N, Robertson IH. Personal versus extrapersonal neglect: A group study of their dissociation using a reliable clinical test. *Cortex* 1997;33:379–84.
- [226] McIntosh RD, Brodie EE, Beschin N, Robertson IH. Improving the clinical diagnosis of personal neglect: a reformulated comb and razor test. *Cortex* 2000;36:289–92.
- [227] Levy D, Blizzard R, Halligan P, Stone S. Fluctuations in visual neglect after stroke? *European Neurology* 1995;35:341–3. <https://doi.org/10.1159/000117157>.
- [228] Schwartz M, Segal M, Veramonti T, Ferraro M, Buxbaum L. The Naturalistic Action Test: a standardised assessment for everyday action impairment. *Neuropsychological Rehabilitation* 2002;12:311–39.
- [229] Grattan ES, Woodbury ML. Do Neglect Assessments Detect Neglect Differently? *Am J Occup Ther* 2017;71:7103190050p1. <https://doi.org/10.5014/ajot.2017.025015>.
- [230] Schwartz MF, Buxbaum LJ, Montgomery MW, Fitzpatrick-DeSalme E, Hart T, Ferraro M, et al. Naturalistic action production following right hemisphere stroke. *Neuropsychologia* 1999;37:51–66. <https://doi.org/10.1016/S0028-3932%2898%2900066-9>.
- [231] Verlander D, Hayes A, McInnes JK, Liddle RJ, Liddle GW, Clarke GE, et al. Assessment of clients with visual spatial disorders: a pilot study. *Null* 2000;2:129–42. <https://doi.org/10.1076/vimr.2.3.129.4422>.
- [232] Punt TD, Kitadono K, Hulleman J, Humphreys GW, Riddoch MJ. Modulating wheelchair navigation in patients with spatial neglect. *Neuropsychological Rehabilitation* 2011;21:367–82. <https://doi.org/10.1080/09602011.2011.559132>.
- [233] Punt TD, Kitadono K, Hulleman J, Humphreys GW, Riddoch MJ. From both sides now: Crossover effects influence navigation in patients with unilateral neglect. *Journal of Neurology, Neurosurgery and Psychiatry* 2008;79:464–6. <https://doi.org/10.1136/jnnp.2007.129205>.
- [234] Sjoholm H, Hagg S, Nyberg L, Rolander B, Kammerlind A-S. The Cone Evasion Walk test: Reliability and validity in acute stroke. *Physiother Res Int* 2019;24:e1744. <https://doi.org/10.1002/pri.1744>.
- [235] Webster JS, Roades LA, Morrill B, Rapport LJ, Abadee PS, Sowa MV, et al. Rightward orienting bias, wheelchair maneuvering, and fall risk. *Archives of Physical Medicine and Rehabilitation* 1995;76:924–8. [https://doi.org/10.1016/S0003-9993\(95\)80068-9](https://doi.org/10.1016/S0003-9993(95)80068-9).
- [236] Webster JS, Cottam G, Gouver WD, Blanton P, Beissel GF, Wofford J. Wheelchair obstacle course performance in right cerebral vascular accident victims. *J Clin Exp Neuropsychol* 1989;11:295–310.

- [237] Qiang W, Sonoda S, Suzuki M, Okamoto S, Saitoh E. Reliability and validity of a wheelchair collision test for screening behavioral assessment of unilateral neglect after stroke. American Journal of Physical Medicine and Rehabilitation 2005;84:161–6. <https://doi.org/10.1097/01.PHM.0000154902.79990.12>.