

**List of references used to prepare distribution maps of
Paralicella tenuipes and *P. caperesca***

- Bühning S.I., and Christiansen B. (2001). Lipids in selected abyssal benthopelagic animals: links to the epipelagic zone? *Progr. Oceanogr.* 50, 369–382. doi: 10.1016/s0079-6611(01)00061-1
- Chevreaux, E. (1908). Diagnoses d'amphipodes nouveaux provenant des campagnes de la "Princesse-Alice" dans l' Atlantique nord. *Bulletin de l'Institut Océanographique de Monaco* 117, 1–13.
- Cousins N.J., Horton T., Wigham B.D., and Bagley P.M. (2013). Abyssal scavenging demersal fauna at two areas of contrasting productivity on the Subantarctic Crozet Plateau, southern Indian Ocean. *Afr. J. Mar. Sci.* 35, 299–306. doi: 10.2989/1814232X.2013.802747
- Duffy G.A., Horton T., and Billett D.S.M. (2012). Deep-sea scavenging amphipod assemblages from the submarine canyons of the Western Iberian Peninsula. *Biogeosciences* 9, 4861–4869. doi: 10.5194/bg-9-4861-2012
- Duffy, G. A., Gutteridge, Z. R., Thurston, M. H., and Horton, T. (2016). A comparative analysis of canyon and non-canyon populations of the deep-sea scavenging amphipod *Paralicella caperesca*. *J. Mar. Biol. Assoc. UK* 96(8), 1687–1699. doi: 10.1017/S0025315415002064
- France, S. C. (1994). Genetic population structure and gene flow among deep-sea amphipods, *Abyssorchomene* spp., from six California Continental Borderland basins. *Mar. Biol.* 118(1), 67–77. doi: 10.1007/BF00699220
- Fujii T., Kilgallen N.M., Rowden A.A., and Jamieson A.J. (2013). Deep-sea amphipod community structure across abyssal to hadal depths in the Peru-Chile and Kermadec trenches. *Mar. Ecol. Prog. Ser.* 492, 125–138. doi: 10.3354/meps10489
- Hasegawa M., Kurohiji Y., Takayanagi S., Sawadaishi S., and Yao M. (1986). Collection of fish and Amphipoda from abyssal sea-floor at 30 8N – 147 8E using traps tied to 10000 m wire of research vessel. *Bulletin of Tokai Regional Fisheries Research Laboratory* 119, 65–75
- Hendrycks E.A., and Conlan K.E. (2003). New and unusual abyssal gammaridean Amphipoda from the north-east Pacific. *J. Nat. Hist.* 37, 2303–2368. doi: 10.1080/00222930210138926
- Hessler, R. R., Ingram, C. L., Yayanos, A. A., and Burnett, B. R. (1978). Scavenging amphipods from the floor of the Philippine Trench. *Deep-Sea Res.* 25, 1029–1047. doi: [10.1016/0146-6291\(78\)90585-4](https://doi.org/10.1016/0146-6291(78)90585-4)
- Horton, T., and Thurston, M. (2009). *Hirondellea sindhusagar* (Crustacea: Amphipoda: Lysianassoidea), a new deep-water scavenger species from the Indian Ocean, with a key to the genus *Hirondellea*. *Zootaxa* 2096(1), 433–441. doi: 10.11646/zootaxa.2096.1.26
- Horton T., Thurston M.H., and Duffy G.A. (2013). Community composition of scavenging amphipods at bathyal depths on the Mid-Atlantic Ridge. *Deep-Sea Res. Part II Top. Stud. Oceanogr.* 98, 352–359. doi: 10.1016/j.dsr2.2013.01.032

- Horton, T., Thurston, M. H., Vlierboom, R., Gutteridge, Z., Pebody, C. A., Gates, A. R., et al. (2020). Are abyssal scavenging amphipod assemblages linked to climate cycles? *Progr Oceanogr* 184, 102318. doi: [10.1016/j.pocean.2020.102318](https://doi.org/10.1016/j.pocean.2020.102318)
- Iguchi, A., Nishijima, M., Yoshioka, Y., Miyagi, A., Miwa, R., Tanaka, Y., et al. (2020). Deep-sea amphipods around cobalt-rich ferromanganese crusts: Taxonomic diversity and selection of candidate species for connectivity analysis. *PLoS ONE* 15(2), e0228483. doi: [10.1371/journal.pone.0228483](https://doi.org/10.1371/journal.pone.0228483)
- Ingram, C.L., Hessler, R.R. (1983). Distribution and behavior of scavenging amphipods from the central North Pacific. *Deep-Sea Res.* 30(7A), 683–706. doi: [10.1016/0198-0149\(83\)90017-1](https://doi.org/10.1016/0198-0149(83)90017-1)
- Jazdzewska, A. M., and Mamos, T. (2019). High species richness of Northwest Pacific deep-sea amphipods revealed through DNA barcoding. *Prog. Oceanogr.* 178, 102184. doi: [10.1016/j.pocean.2019.102184](https://doi.org/10.1016/j.pocean.2019.102184)
- Jones E.G., Collins M.A., Bagley P.M., Addison S., and Priede I.G. (1998). The fate of cetacean carcasses in the deep sea: observations on consumption rates and succession of scavenging species in the abyssal north-east Atlantic Ocean. *Philos. T. Roy. Soc. B* 265, 1119–1127. doi: [10.1098/rspb.1998.0407](https://doi.org/10.1098/rspb.1998.0407)
- Kaufmann R.S. (1994). Structure and function of chemoreceptors in scavenging lysianassoid amphipods. *J. Crust. Biol.* 14, 54–71. doi: [10.2307/1549055](https://doi.org/10.2307/1549055)
- Kaufmann, R. S., Wakefield, W. W., and Genin, A. (1989). Distribution of epibenthic megafauna and lebensspuren on two central North Pacific seamounts. *Deep-Sea Res. Part I Oceanogr. Res. Pap.* 36(12), 1863–1896. doi: [10.1016/0198-0149\(89\)90116-7](https://doi.org/10.1016/0198-0149(89)90116-7)
- Lacey, N. C., Rowden, A. A., Clark, M. R., Kilgallen, N. M., Linley, T., Mayor, D. J. et al. (2016). Community structure and diversity of scavenging amphipods from bathyal to hadal depths in three South Pacific Trenches. *Deep-Sea Res. Part I Oceanogr. Res. Pap.* 111, 121–137. doi: [10.1016/j.dsr.2016.02.014](https://doi.org/10.1016/j.dsr.2016.02.014)
- Lampitt R.S., Merrett N.R., and Thurston M.H. (1983). Inter-relations of necrophagous amphipods, a fish predator, and tidal currents in the deep sea. *Mar. Biol.* 74, 73–78. doi: [10.1007/BF00394277](https://doi.org/10.1007/BF00394277)
- Macdonald A. G. and Gilchrist I. (1980). Effects of hydraulic decompression and compression on deep-sea amphipods. *Comp. Biochem. Phys. A* 67, 149–153. doi: [10.1016/0300-9629\(80\)90420-X](https://doi.org/10.1016/0300-9629(80)90420-X).
- Patel, T., Robert, H., d'Udekem d'Acoz, C., Martens, K., De Mesel, I., Degraer, S., et al. (2020). Biogeography and community structure of abyssal scavenging Amphipoda (Crustacea) in the Pacific Ocean. *Biogeosciences* 17, 2731–2744. doi: [10.5194/bg-17-2731-2020](https://doi.org/10.5194/bg-17-2731-2020)
- Ritchie, H., Jamieson, A. J., and Piertney, S. B. (2015). Phylogenetic relationships among hadal amphipods of the Superfamily Lysianassoidea: Implications for taxonomy and biogeography. *Deep-Sea Res. Part I Oceanogr. Res. Pap.* 105, 119–131. doi: [10.1016/j.dsr.2015.08.014](https://doi.org/10.1016/j.dsr.2015.08.014)

- Shulenberger, E., and Barnard, J. L. (1976). Amphipods from an abyssal trap set in the North Pacific Gyre. *Crustaceana* 31(3), 241–258. doi: 10.1163/156854076X00035
- Smith K.L., and Baldwin R. J. (1982). Scavenging deep-sea amphipods – effects of food odor on oxygen-consumption and a proposed metabolic strategy. *Mar. Biol.* 68, 287–298. doi: 10.1007/BF00409595
- Thurston, M.H. (1979). Scavenging abyssal amphipods from the North-East Atlantic Ocean. *Mar. Biol.* 51, 55–68. doi: 10.1007/BF00389031.
- Thurston, M. H. (1990). Abyssal necrophagous amphipods (Crustacea: Amphipoda) in the northeast and tropical Atlantic Ocean. *Progr. Oceanogr.* 24(1-4), 257–274. doi: 10.1016/0079-6611(90)90036-2
- Treude, T., Janßen, F., Queisser, W., and Witte, U. (2002). Metabolism and decompression tolerance of scavenging lysianassooid deep-sea amphipods. *Deep-Sea Res. Part I Oceanogr. Res. Pap.* 49(7), 1281–1289. doi: 10.1016/S0967-0637(02)00023-7
- Weston, J. N., Peart, R. A., and Jamieson, A. J. (2020). Amphipods from the Wallaby-Zenith Fracture Zone, Indian Ocean: new genus and two new species identified by integrative taxonomy. *Syst. Biodivers.* 18(1), 57–78. doi: 10.1080/14772000.2020.1729891
- Weston, J. N. J., Peart, R. A., Stewart, H. A., Ritchie, H., Piertney, S. B., Linley, T. D., et al. (2021). Scavenging amphipods from the Wallaby-Zenith Fracture Zone: Extending the hadal paradigm beyond subduction trenches. *Mar. Biol.* 168, 1. doi: 10.1007/s00227-020-03798-4
- Wilson, J. P., Schnabel, K. E., Rowden, A. A., Peart, R. A., Kitazato, H., and Ryan, K. G. (2018). Bait-attending amphipods of the Tonga Trench and depth-stratified population structure in the scavenging amphipod *Hirondellea dubia* Dahl, 1959. *PeerJ*, 6, e5994.
- Wilson R.R., Smith K.L., and Rosenblatt R.H. (1985). Megafauna associated with bathyal seamounts in the central North Pacific-Ocean. *Deep-Sea Res. Part I Oceanogr. Res. Pap.* 32, 1243–1254. doi: 10.1016/0198-0149(85)90007-X
- Yayanos, A. A. (1981). Reversible inactivation of deep-sea amphipods (*Paralicella capresca*) by a decompression from 601 bars to atmospheric pressure. *Comp. Biochem. Phys. A* 69(3), 563–565. doi: 10.1016/0300-9629(81)93020-6