

*Supporting Information for*

# Achieving NIR Light-Mediated Tumor-Specific Fenton Reaction-Assisted Oncotherapy by Using Magnetic Nanoclusters

*Shaoyou Qin,<sup>†</sup> Jinru Xue,<sup>‡</sup> Erna Jia,<sup>†</sup> Na Ren,<sup>‡</sup> Yongqiang Dong,<sup>§</sup> and Changyu Zhou <sup>\*,†</sup>*

<sup>†</sup> Department of Gastroenterology and Hepatology, China-Japan Union Hospital of Jilin University, Jilin University, Changchun 130033, P.R. China

<sup>‡</sup> Department of Thoracic Surgery, China-Japan Union Hospital of Jilin University, Jilin University, Changchun 130033, P.R. China

<sup>§</sup> Department of Thyroid Surgery, First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, P.R. China.

E-mail: cyzhou@jlu.edu.cn

**Table S1.** Mean value of hemolysis percentages after co-incubation with MNCs.

concentrations ( $\mu\text{g/mL}$ )	average value of hemolysis (%)
0.9% NaCl	0
D.I. water	100
3.125	0.198
6.25	0.198
12.5	0.396
25	0.495
50	0.594
100	0.693
200	0.792
400	1.188