

Supplementary Table 2. Retrograde-labeled cell counts and percent of total cells from tracer injections into ML (FR) and CPB (BDA)(Case 2). Sorted by cortical area in supragranular (S) and infragranular (I) layers). The results are graphically summarized in the charts below and in [Supplementary Figure 2](#).

	A1				R				RT				CM				MM			
	FR		BDA		FR		BDA		FR		BDA		FR		BDA		FR		BDA	
	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I
Count	273	54	1	0	5	4	2	1	0	0	0	0	45	15	15	1	69	18	85	10
%	7	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	2	0

	RM				RTM				ProA				CL				ML			
	FR		BDA		FR		BDA		FR		BDA		FR		BDA		FR		BDA	
	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I
Count	70	46	14	1	13	7	0	12	13	6	18	1	87	63	83	15	363	141	160	8
%	2	1	0	0	0	0	0	0	0	0	0	0	2	2	2	0	10	4	4	0

	AL				RTL				CPB				RPB				TPO			
	FR		BDA		FR		BDA		FR		BDA		FR		BDA		FR		BDA	
	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I	S	I
Count	40	28	50	13	5	17	19	5	169	136	429	167	39	53	156	99	19	49	264	265
%	1	1	1	0	0	0	1	0	5	4	12	5	1	1	4	3	1	1	7	7

