

Supplemental Material

Detecting Small Vessel Pathology in Cocaine Use Disorder

Marco Öchsner, Elijah Mak, Karen D Ersche[✉]

[✉] ke220@cam.ac.uk

Table S1: Demographics, personality traits and clinical data for participants who received a FLAIR, and those who did not [means and standard deviation, (Std.) in parentheses]

Demographics	Radiographer's choices			
	no FLAIR mean (\pm SD)	with FLAIR mean (\pm SD)	t statistic	p-value
Sample(n)	88	36	-	-
Age (years)	38.61 (\pm 9.7)	40.37 (\pm 9.8)	-0.92	0.364
Gender (% male)	95.5%	97.2%	Fisher's	1.000
Formal education (years)	12.86 (\pm 3.0)	12.22 (\pm 2.7)	1.16	0.249
Body Mass Index (BMI)	23.99 (\pm 3.5)	24.83 (\pm 3.4)	-1.25	0.214
Systolic blood pressure (mmHg)	128.61 (\pm 14.5)	128.42 (\pm 14.7)	0.07	0.946
Smoking status (% smokers)	46.6%	61.1%	Fisher's	0.168
Impulsivity (BIS11 total score)	65.89 (\pm 12.8)	70.89 (\pm 13.4)	-1.91	0.060
	CUD no FLAIR mean (\pm SD)	CUD with FLAIR mean (\pm SD)	t statistic	p-value
Sample (n)	44	20	-	-
Duration of cocaine use (years)	15.1 (\pm 5.9)	19.6 (\pm 7.5)	-2.38	0.024
Age at onset (years)	22.0 (\pm 6.0)	20.2 (\pm 4.8)	1.28	0.208
Compulsivity of drug use (OCDUS total)	23.9 (\pm 7.8)	25.7 (\pm 11.2)	-0.64	0.528