

Influenza A virus-driven airway inflammation may be dissociated from limb muscle atrophy in cigarette smoke-exposed mice

Kevin Mou¹, Stanley MH. Chan¹, Kurt Brassington¹, Aleksandar Dobric¹, Simone N. De Luca¹, Huei Jiunn Seow¹, Stavros Selemidis¹, Steven Bozinovski¹, Ross Vlahos^{1*}

¹School of Health and Biomedical Sciences, RMIT University, BUNDOORA, VIC 3083, Australia.

*** Correspondence:**

Professor Ross Vlahos, PhD

School of Health and Biomedical Sciences



RMIT University, PO Box 71, Bundoora, VIC 3083 Australia

Tel: +61 3 9925 7362

Email: ross.vlahos@rmit.edu.au

Keywords: viral exacerbation, COPD, muscle weakness, fiber type transformation, lung-to-muscle axis, myogenic disruption, conditioned medium.

Supplementary Material

1 SUPPLEMENTARY FIGURES**(A)****(B)**

Supplementary Figure 1. Representative images of cell morphology/viability following (A) direct exposure of Poly I:C (TLR3 agonist), cigarette smoke extract (CSE) or their combination in BEAS-2B cells, or (B) C2C12 myotubes exposed to conditioned media derived from the respective conditions in BEAS-2B.