



Fatty acid-derived biofuels and chemicals production in *Saccharomyces cerevisiae*

Yongjin J. Zhou, Nicolaas A. Buijs, Verena Siewers and Jens Nielsen*

Department of Chemical and Biological Engineering, Chalmers University of Technology, Gothenburg, Sweden

Edited by:

Patrik R. Jones, Imperial College London, UK

Reviewed by:

M. Kalim Akhtar, University College London, UK

Jun Ishii, Kobe University, Japan

Rebecca Lennen, Novo Nordisk

Foundation Center for

Biosustainability, Denmark

*Correspondence:

Jens Nielsen, Department of Chemical and Biological Engineering, Chalmers University of Technology, Gothenburg SE-412 96, Sweden
e-mail: nielsenj@chalmers.se

Volatile energy costs and environmental concerns have spurred interest in the development of alternative, renewable, sustainable, and cost-effective energy resources. Environment-friendly processes involving microbes can be used to synthesize advanced biofuels. These fuels have the potential to replace fossil fuels in supporting high-power demanding machinery such as aircrafts and trucks. From an engineering perspective, the pathway for fatty acid biosynthesis is an attractive route for the production of advanced fuels such as fatty acid ethyl esters, fatty alcohols, and alkanes. The robustness and excellent accessibility to molecular genetics make the yeast *Saccharomyces cerevisiae* a suitable host for the purpose of bio-manufacturing. Recent advances in metabolic engineering, as well as systems and synthetic biology, have now provided the opportunity to engineer yeast metabolism for the production of fatty acid-derived fuels and chemicals.

Keywords: fatty acid metabolism, fatty acid ethyl ester, fatty alcohol, alkanes/alkene, yeast

INTRODUCTION

Volatile energy costs and environmental concerns have motivated the development of sustainable, renewable, and cost-effective alternative energy sources that have reduced pollution emissions or carbon footprints (Fortman et al., 2008). Biofuels are such green alternatives to petroleum-based fuels, given the capacity of photosynthetic organisms to recycle CO₂ after biofuel combustion, thereby leading to near-zero net greenhouse gas emissions. Microbial synthesis is an attractive approach for biofuel production due to the large flexibility for pathway engineering and low environmental footprint. Among the metabolic pathways, fatty acid biosynthesis has attracted significant attention for production of highly reduced biofuels and chemicals with high energy densities (Lennen and Pfleger, 2013).


The yeast *Saccharomyces cerevisiae* is a well-studied model microorganism and is particularly suited for commercial scale processes due to its robustness and tolerance toward industrial conditions, the capability of high-density fermentations, and insusceptibility toward phage contamination (Nielsen et al., 2013; Mattanovich et al., 2014). In fact, *S. cerevisiae* has been successfully applied for the industrial production of bioethanol (Mussatto et al., 2010). Additional studies have shown that it can be engineered for the production of a variety of fuel molecules including isobutanol, butanol, and farnesene (Buijs et al., 2013). Recently, it has received increased attention as a host for the synthesis of fatty acid-derived biofuels and chemicals (Li et al., 2014; Rungtaphan and Keasling, 2014). This review will summarize recent progress in the engineering of *S. cerevisiae* for the production of fatty acid-derived biofuels and chemicals, in addition to analyzing the current challenges in increasing their productivity for commercial deployment.

FATTY ACID PRODUCTION IN *S. CEREVISIAE*

Free fatty acids (FFAs) can be used for the industrial manufacturing of detergents, soaps, lubricants, cosmetics, and

pharmaceutical ingredients (Tee et al., 2014). In addition, FFAs can also serve as precursors for the production of alkanes by catalytic decarboxylation (Lennen et al., 2010) or fatty acid methyl esters (FAMES) through esterification (Christie and Han, 2010).

The biosynthesis of fatty acids in *S. cerevisiae* differs from that in bacteria such as *Escherichia coli* (Figure 1). In bacteria, fatty acid synthesis is carried out by a type II fatty acid synthase (FAS) that consists of discrete, monofunctional enzymes (Figure 1B); while in *S. cerevisiae*, the *de novo* synthesis of fatty acids can take place in at least two subcellular compartments: cytoplasm (type I FAS) and mitochondria (type II FAS). Mitochondrial FAS II has been implicated as the sole mitochondrial source of octanoic acid, which is a precursor of the lipoic acid (LA) cofactor that is required for maintaining the function of several mitochondrial enzyme complexes such as pyruvate dehydrogenase (Hiltunen et al., 2009). However, most functional and storage lipids are synthesized by cytosol type I FAS (Koch et al., 2014), which is a large, multifunctional dimeric complex that is responsible for fatty acid synthesis from malonyl-CoA and acetyl-CoA (Figure 1A). This distinction is important as it has implications for further metabolic engineering of fatty acid metabolism. Considering its predominant role for fatty acids synthesis, we will focus on the FAS I system. This process starts with loading of acetyl-CoA to the acyl carrier protein (ACP) by the ACP acyltransferase (AT). Then consecutive catalytic steps of β -ketoacyl-ACP synthesis, β -ketoacyl-ACP reduction, β -hydroxyacyl-ACP dehydration, and enoyl-ACP reduction extend the chain length in a repetitive manner by using malonyl-CoA as building blocks. The malonyl-CoA is synthesized from acetyl-CoA by incorporation of CO₂, which is catalyzed by acetyl-CoA carboxylase (Acc1). The chain extension usually stops at palmitoyl-ACP after seven cycles, which is mainly determined by the ketoacyl synthase domain (Sangwallek et al., 2013). Finally, acyl-ACP and malonyl-CoA are transformed by malonyl transacylase (MPT) to form acyl-CoA and the activated malonyl-ACP, which is necessary for initiating the next acyl-CoA synthesis.




Acyl-CoA can be transformed into lipids or FFAs catalyzed by AT or thioesterase, respectively.

Disruption of two main fatty acyl-CoA synthetase-encoding genes *FAA1* and *FAA4* led to the production of 900 $\mu\text{mol/L}$ (≈ 240 mg/L) FFAs, of which 220 $\mu\text{mol/L}$ (≈ 60 mg/L) FFAs were secreted. The accumulation of FFAs was most likely the result of interrupted lipid remodeling processes (Scharnewski et al., 2008). Besides *FAA1* and *FAA4* disruption, overexpression of the FAS subunits (Fas1 and Fas2), acetyl-CoA carboxylase (Acc1), and *E. coli* acyl-ACP thioesterase (TesA) resulted in a higher FFAs production of 400 mg/L. Blockage of β -oxidation was also beneficial for FFA accumulation though it was found to be much less effective than elimination of fatty acyl-CoA synthetase (Runghpan and Keasling, 2014). Similarly, after disruption of the β -oxidation pathway, (i) elimination of the acyl-CoA synthetases, (ii) overexpression of different thioesterases, and (iii) enhancement of acetyl-CoA/malonyl-CoA, the engineered strain produced more than 140 mg/L FFAs (Li et al., 2014). Additionally, acetyl-CoA carboxylase (Acc1) was identified as a critical bottleneck for fatty acid synthesis in *S. cerevisiae* with a cell-free system (Li et al., 2014).

Fatty acid metabolism has also been widely tailored for producing fatty acid ethyl esters (FAEEs), fatty alcohols, and alkanes/alkenes with different chain length in prokaryotes, especially in *E. coli*, which has been reviewed elsewhere (Lennen and Pfeleger, 2013; Janssen and Steinbuechel, 2014). **Figure 2** illustrates how the different fatty acid-derived products can be obtained in yeast and is described in greater detail within the subsequent sections.

TRIACYLGLYCEROL PRODUCTION

Besides FFAs, microbial lipids have also been attracting great attention as alternative feedstocks to vegetable oils and animal fats for production of FAMES that are used as biodiesel (Hu et al., 2011). Triacylglycerols (TAGs), which can accumulate to very high levels in eukaryotic cells, play an essential role in energy storage (Sorger and Daum, 2003; Klug and Daum, 2014). Wild-type *S. cerevisiae* strains typically do not accumulate storage lipids to more than 10% dry cell weight (DCW), whereas some oleaginous yeasts and fungi can accumulate TAGs to much higher levels (Li et al., 2007). Overexpression of several genes encoding FAS (*FAS1* and *FAS2*), acetyl-CoA carboxylase (*ACC1*), and diacylglycerol AT (*DGA1*) made *S. cerevisiae* BY4742 accumulate total



TAGs to 17% DCW (Runguphan and Keasling, 2014). Mimicking the metabolism of oleaginous microorganisms, by introduction of ATP-citrate lyase (ACL) and disruption of isocitrate dehydrogenase genes *IDH1* and *IDH2*, increased the total fatty acid content by 21% (Tang et al., 2013). From transposon mutagenesis analysis, it was found that disruption of the transcription factor Snf2 resulted in a 70% increase of TAGs (Kamisaka et al., 2006). Furthermore, when the fatty acyl-CoA synthetase-encoding gene *FAA3* and the diacylglycerol AT-encoding gene *DGA1* were overexpressed, total lipid accumulation increased to 30% DCW, of which TAGs were the most abundant (Kamisaka et al., 2007). Overexpression of the active diacylglycerol AT variant *Dga1ΔN*,

lacking the N-terminal 29 amino acids in the *dga1Δ* and *snf2Δ* mutant, further increased lipid production up to 45% (Kamisaka et al., 2013). This is comparable to levels found in oleaginous yeast and is the highest lipid content reported for *S. cerevisiae* so far. It is worthy to mention that the disruption of *SNF2* (Kamisaka et al., 2007) is more efficient in boosting TAG accumulation than overexpression of fatty acid biosynthesis-related genes (Runguphan and Keasling, 2014), which indicates that TAG and fatty acid biosynthesis are under transcriptional control. The high TAG accumulation provides a feasible alternative route for FAME-based biodiesel production by direct methanolysis of oleaginous microbial biomass (Liu and Zhao, 2007).

FATTY ACID ETHYL ESTER PRODUCTION

Fatty acid ethyl esters are potentially attractive diesel fuel replacements due to their high energy density and low host toxicity (Zhang et al., 2012). FAEs can be synthesized by condensation of acyl-CoAs and ethanol by using a wax ester synthase/acyl-CoA:diacylglycerol AT (WS/DGAT). An early attempt at FAE biosynthesis in *S. cerevisiae* using a WS/DGAT from *Acinetobacter calcoaceticus* ADP1 resulted in a very poor yield (Kalscheuer et al., 2004). Recently, five different wax ester synthases were investigated for their ability to perform FAE biosynthesis (Shi et al., 2012). It was found that the wax ester synthase from *Marinobacter hydrocarbonoclasticus* had the best performance toward short-chain alcohols *in vitro*. This enzyme also led to production of the highest FAE titer of 6.3 mg/L when expressed in yeast. In order to overcome the instability and metabolic burden of the plasmid-based enzyme expression, the wax ester synthase gene was integrated, as multiple copies, into the yeast chromosome, and the FAE production increased sixfold up to 34 mg/L (Shi et al., 2014b).

Through overexpression of *ACC1* encoding acetyl coenzyme A carboxylase (ACCase), FAE production was further enhanced by 30%, resulting in FAE production of 8.2 mg/L (Shi et al., 2012). Acetyl-CoA carboxylation is a flux controlling step in fatty acid biosynthesis and overexpression of *ACC1* has been shown to increase fatty acid production by 58% (Runguphan and Keasling, 2014). However, *Acc1* activity is strictly regulated by phosphorylation under the control of Snf1. Thus, abolishing phosphorylation regulation of *Acc1* by introducing double mutants of S1157A and S659A, increased ACCase activity 3.1-fold in wild-type strain, which was an even higher improvement than *snf1Δ* mutation (Shi et al., 2014a). Overexpression of *ACC1*^{S1157A,S659A} increased FAEs threefold to 15.8 mg/L, while overexpression of wild-type *ACC1* just led to an increase of 20% (Shi et al., 2014a). This study should be helpful for increasing the production of other malonyl-CoA- and fatty acid-derived chemicals.

In order to enhance the fatty acyl-CoA level for FAE production, fatty acyl-CoAs consuming pathways, including β -oxidation, steryl esters (SEs), and triacylglycerols (TAG) biosynthesis were blocked. Compared to the wild-type, there was approximately a threefold increase in FAE production of 17.2 mg/L in the mutant strain. Additionally, the FFAs increased fivefold, which may be helpful for producing other fatty acid-derived molecules (Valle-Rodriguez et al., 2014). Another report showed that enhancing the fatty acid biosynthesis pathway and blocking β -oxidation in

S. cerevisiae BY4742 increased FAEE production of 4.5-fold (Runguphan and Keasling, 2014). However, the FAEE titer of 5.4 mg/L was less than the strain (17.2 mg/L) in which all the fatty acyl-CoAs consuming pathways were blocked (Valle-Rodriguez et al., 2014), indicating that a high level of fatty acyl-CoAs is essential for efficient FAEE production. Since fatty acyl-CoA biosynthesis requires large amounts of NADPH and acetyl-CoA, the ethanol degradation pathway was therefore up-regulated and a phosphoketolase pathway was introduced to increase acetyl-CoA and NADPH supply for FAEE production (de Jong et al., 2014). By overexpression of the ethanol degradation pathway enzymes alcohol dehydrogenase *Adh2*, acetaldehyde dehydrogenase *Ald6*, and the *Salmonella enterica* acetyl-CoA synthetase variant *AcsSE* (L641P) (Starai et al., 2005), the FAEE yield increased threefold. Introduction of the phosphoketolase pathway yielded a 1.7-fold improvement in FAEE production to 5.1 mg/gDCW in shake flasks (de Jong et al., 2014).

To date, the highest reported FAEE production in *S. cerevisiae* is 0.52 g/L by using glycerol as a carbon source after enhancing ethanol biosynthesis, down-regulating glycerol export, and also adding exogenous fatty acids (Yu et al., 2012). Given that the addition of exogenous fatty acids was essential for high FAEE production, the enhancement of fatty acid biosynthesis would be necessary for the increased *de novo* synthesis of FAEE from sugar-based carbon sources.

FATTY ALCOHOL PRODUCTION

Fatty alcohols are widely used as detergents, skin care products, cosmetics, and medicines and are also considered as potential biofuels (Liu et al., 2014a). Fatty alcohol biosynthesis can proceed via the reduction of a fatty aldehyde intermediate, by an aldehyde reductase. The fatty aldehyde can be generated from (i) fatty acyl-ACP via a fatty acyl-ACP reductase (Schirmer et al., 2010), (ii) acyl-CoA via an acyl-CoA reductase (Reiser and Somerville, 1997), or (iii) fatty acid via a carboxylic acid reductase (Akhtar et al., 2013). Fatty alcohols can also be synthesized directly from fatty acyl-CoA in a four-electron reduction manner catalyzed by a bi-functional fatty acyl-CoA reductase (Willis et al., 2011). Introduction of an NADPH-dependent bi-functional fatty acyl-CoA reductase from *Mus musculus* in wild-type *S. cerevisiae* BY4742 led to fatty alcohol production of 47.4 mg/L. After enhancing the fatty acid biosynthesis pathway and introducing a malic enzyme from *Mucor circinelloides* for improved NADPH supply, the fatty alcohol titer increased to 98 mg/L (Runguphan and Keasling, 2014), which showed the potential of *S. cerevisiae* as a platform for fatty alcohol production. Though this titer is not comparable with that reached in *E. coli* of more than 1.5 g/L in fed-batch cultivation (Youngquist et al., 2013), further increase of fatty acid biosynthesis and engineering of more efficient fatty acyl-CoA/fatty acid reduction pathways should be helpful for fatty alcohol overproduction (Akhtar et al., 2013).

ALKANE/ALKENE PRODUCTION

Fatty acid intermediates are ideal precursors for alkane production. Though alkane synthesis in microbes was discovered decades ago, the detailed biochemical pathway for converting fatty acids or their intermediates to alkanes was elucidated only recently (Schirmer et al., 2010). The pathway involves the activities of

fatty acyl-ACP reductase and aldehyde-deformylating oxygenase that catalyze the reduction of fatty acyl-ACP to the aldehyde, followed by its conversion to the alkane. Then, two other aldehyde decarboxylases that convert fatty aldehydes into their corresponding $n - 1$ alkanes were identified in *Drosophila melanogaster* (Qiu et al., 2012) and *Arabidopsis thaliana* (Bernard et al., 2012). Several other modification enzymes involved in alkane/alkene biosynthesis have also been identified and characterized including Claisen condensation enzymes (Beller et al., 2010; Sukovich et al., 2010; Frias et al., 2011), a novel cytochrome P450 fatty acid decarboxylase (Rude et al., 2011) and a PKS synthase (Mendez-Perez et al., 2011). Most of these enzymes were heterologously expressed and characterized in *E. coli* for alkane/alkene production (Schirmer et al., 2010; Lennen and Pflieger, 2013; Wang and Lu, 2013; Liu et al., 2014b), and the highest alkane titer reached 580 mg/L by using *Clostridium acetobutylicum* fatty acyl-CoA reductase and *A. thaliana* fatty aldehyde decarboxylase (Choi and Lee, 2013). However, the engineering of alkane production in *S. cerevisiae* is lagging behind. To date, there is only one report concerning alkane production in *S. cerevisiae*, which was achieved via reconstruction of the *A. thaliana* very long-chain alkane biosynthesis pathway (Bernard et al., 2012). Introduction of the aldehyde decarboxylase *CER1* and the cognate acyl-CoA reductase *CER3* in *S. cerevisiae* INVSur4#, a very long-chain fatty acid producer, enabled synthesis of 19 μ g/g DCW very long-chain alkanes, primarily 29 carbons in length. Furthermore, expression of *Arabidopsis* cytochrome *b5* isoforms (CYTB5s) and long-chain acyl-CoA synthetase 1 (*LACS1*) for enhancing *CER1*–*CER3* activity and precursor supply, respectively, increased alkane production up to 86 μ g/g DCW (Bernard et al., 2012). There were no medium-chain alkanes (<20 carbons in length) observed, even though it was recently shown that *CER1* is active toward short-chain fatty aldehydes in *E. coli* (Choi and Lee, 2013). The difficulty of alkane production in *S. cerevisiae* may be attributed to the complexity of yeast metabolism, difficulty in expressing complex bacterial enzymes, formation of by-products, and/or alkane toxicity.

In order to overcome the toxicity of alkanes, especially short-chain alkanes, to *S. cerevisiae*, heterologous efflux pumps *ABC2* and *ABC3* from *Yarrowia lipolytica*, an alkane-assimilating yeast, were shown to significantly increase the tolerance against decane and undecane in *S. cerevisiae* through maintaining lower intracellular alkane levels (Chen et al., 2013). In addition, the endogenous efflux pumps *Snq2* and *Pdr5* were identified to be involved in alkane export and tolerance by transcriptional analysis of *S. cerevisiae* cells that were exposed to decane and undecane (Ling et al., 2013). These efflux pumps serve as valuable tools for improving the cellular tolerance of yeast toward the production of alkanes.

CONCLUDING REMARKS

There is an urgent need to develop sustainable economic approaches for the production of fuels and other chemicals, traditionally derived from petroleum, using renewable feedstocks. With regard to engineering, the fatty acid biosynthetic pathway is such an attractive target for the production of a wide range of chemicals and transportation fuels. In recent years, several modifying enzymes have been identified for transforming fatty acids and

their intermediates to alkanes and alcohols with different chain lengths. So far, the potential of these enzymes for the production of fatty acid-derived biofuels and chemicals has been evaluated extensively in *E. coli* (Lennen and Pfeleger, 2013; Janssen and Steinbuchel, 2014). Developments are lagging behind in *S. cerevisiae*, even though this cell factory has been widely used for producing numerous types of chemicals (Hong and Nielsen, 2012; Zhou et al., 2012; Paddon et al., 2013). This is probably due to the more complicated cellular metabolism in addition to its complex regulatory mechanisms and cellular compartmentalization. Nonetheless, given its robustness and the ease with which it can be genetically manipulated, *S. cerevisiae* certainly merits further attention with regard to the production of advanced fuels, as reflected in several recent studies (Li et al., 2014; Runguphan and Keasling, 2014; Shi et al., 2014b).

ACKNOWLEDGMENTS

The authors acknowledge funding from Vetenskapsrådet and FORMAS.

REFERENCES

- Akhtar, M. K., Turner, N. J., and Jones, P. R. (2013). Carboxylic acid reductase is a versatile enzyme for the conversion of fatty acids into fuels and chemical commodities. *Proc. Natl. Acad. Sci. U.S.A.* 110, 87–92. doi:10.1073/pnas.1216516110
- Beller, H. R., Goh, E. B., and Keasling, J. D. (2010). Genes involved in long-chain alkene biosynthesis in *Micrococcus luteus*. *Appl. Environ. Microbiol.* 76, 1212–1223. doi:10.1128/AEM.02312-09
- Bernard, A., Domergue, F., Pascal, S., Jetter, R., Renne, C., Faure, J. D., et al. (2012). Reconstitution of plant alkane biosynthesis in yeast demonstrates that *Arabidopsis* ECERIFERUM1 and ECERIFERUM3 are core components of a very-long-chain alkane synthesis complex. *Plant Cell* 24, 3106–3118. doi:10.1105/tpc.112.099796
- Buijs, N. A., Siewers, V., and Nielsen, J. (2013). Advanced biofuel production by the yeast *Saccharomyces cerevisiae*. *Curr. Opin. Chem. Biol.* 17, 480–488. doi:10.1016/j.cbpa.2013.03.036
- Chen, B., Ling, H., and Chang, M. W. (2013). Transporter engineering for improved tolerance against alkane biofuels in *Saccharomyces cerevisiae*. *Biotechnol. Biofuels* 6, 21. doi:10.1186/1754-6834-6-21
- Choi, Y. J., and Lee, S. Y. (2013). Microbial production of short-chain alkanes. *Nature* 502, 571–574. doi:10.1038/nature12536
- Christie, W. W., and Han, X. (2010). *Lipid Analysis: Isolation, Separation, Identification and Lipidomic Analysis (Fourth Edition)*. Bridgewater: PJ Barnes & Associates.
- de Jong, B. W., Shi, S., Siewers, V., and Nielsen, J. (2014). Improved production of fatty acid ethyl esters in *Saccharomyces cerevisiae* through up-regulation of the ethanol degradation pathway and expression of the heterologous phosphoketolase pathway. *Microb. Cell Fact.* 13, 39. doi:10.1186/1475-2859-13-39
- Fortman, J. L., Chhabra, S., Mukhopadhyay, A., Chou, H., Lee, T. S., Steen, E., et al. (2008). Biofuel alternatives to ethanol: pumping the microbial well. *Trends Biotechnol.* 26, 375–381. doi:10.1016/j.tibtech.2008.03.008
- Frias, J. A., Richman, J. E., Erickson, J. S., and Wackett, L. P. (2011). Purification and characterization of OleA from *Xanthomonas campestris* and demonstration of a non-decarboxylative Claisen condensation reaction. *J. Biol. Chem.* 286, 10930–10938. doi:10.1074/jbc.M110.216127
- Hiltunen, J. K., Schonauer, M. S., Autio, K. J., Mittelmeier, T. M., Kastaniotis, A. J., and Dieckmann, C. L. (2009). Mitochondrial fatty acid synthesis type II: more than just fatty acids. *J. Biol. Chem.* 284, 9011–9015. doi:10.1074/jbc.R800068200
- Hong, K. K., and Nielsen, J. (2012). Metabolic engineering of *Saccharomyces cerevisiae*: a key cell factory platform for future biorefineries. *Cell. Mol. Life Sci.* 69, 2671–2690. doi:10.1007/s00018-012-0945-1
- Hu, C., Wu, S., Wang, Q., Jin, G., Shen, H., and Zhao, Z. K. (2011). Simultaneous utilization of glucose and xylose for lipid production by *Trichosporon cutaneum*. *Biotechnol. Biofuels* 4, 25. doi:10.1186/1754-6834-4-25
- Janssen, H. J., and Steinbuchel, A. (2014). Fatty acid synthesis in *Escherichia coli* and its applications towards the production of fatty acid based biofuels. *Biotechnol. Biofuels* 7, 7. doi:10.1186/1754-6834-7-7
- Kalscheuer, R., Luftmann, H., and Steinbuchel, A. (2004). Synthesis of novel lipids in *Saccharomyces cerevisiae* by heterologous expression of an unspecific bacterial acyltransferase. *Appl. Environ. Microbiol.* 70, 7119–7125. doi:10.1128/AEM.70.12.7119-7125.2004
- Kamisaka, Y., Kimura, K., Uemura, H., and Yamaoka, M. (2013). Overexpression of the active diacylglycerol acyltransferase variant transforms *Saccharomyces cerevisiae* into an oleaginous yeast. *Appl. Microbiol. Biotechnol.* 97, 7345–7355. doi:10.1007/s00253-013-4915-9
- Kamisaka, Y., Tomita, N., Kimura, K., Kodaki, T., and Hosaka, K. (2006). Identification of genes affecting lipid content using transposon mutagenesis in *Saccharomyces cerevisiae*. *Biosci. Biotechnol. Biochem.* 70, 646–653. doi:10.1271/bbb.70.646
- Kamisaka, Y., Tomita, N., Kimura, K., Kainou, K., and Uemura, H. (2007). DGA1 (diacylglycerol acyltransferase gene) overexpression and leucine biosynthesis significantly increase lipid accumulation in the Deltasnf2 disruptant of *Saccharomyces cerevisiae*. *Biochem. J.* 408, 61–68. doi:10.1042/BJ20070449
- Klug, L., and Daum, G. (2014). Yeast lipid metabolism at a glance. *FEMS Yeast Res.* 14, 369–388. doi:10.1111/1567-1364.12141
- Koch, B., Schmidt, C., and Daum, G. (2014). Storage lipids of yeasts: a survey of non-polar lipid metabolism in *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Yarrowia lipolytica*. *FEMS Microbiol. Rev.* doi:10.1111/1574-6976.12069
- Lennen, R. M., Braden, D. J., West, R. A., Dumesic, J. A., and Pfeleger, B. F. (2010). A process for microbial hydrocarbon synthesis: overproduction of fatty acids in *Escherichia coli* and catalytic conversion to alkanes. *Biotechnol. Bioeng.* 106, 193–202. doi:10.1002/bit.22660
- Lennen, R. M., and Pfeleger, B. F. (2013). Microbial production of fatty acid-derived fuels and chemicals. *Curr. Opin. Biotechnol.* 24, 1044–1053. doi:10.1016/j.copbio.2013.02.028
- Li, X., Guo, D., Cheng, Y., Zhu, F., Deng, Z., and Liu, T. (2014). Overproduction of fatty acids in engineered *Saccharomyces cerevisiae*. *Biotechnol. Bioeng.* 111, 1841–1852. doi:10.1002/bit.25239
- Li, Y. H., Zhao, Z. B., and Bai, F. W. (2007). High-density cultivation of oleaginous yeast *Rhodospiridium toruloides* Y4 in fed-batch culture. *Enzyme Microb. Technol.* 41, 312–317. doi:10.1016/j.enzmictec.2007.02.008
- Ling, H., Chen, B., Kang, A., Lee, J. M., and Chang, M. W. (2013). Transcriptome response to alkane biofuels in *Saccharomyces cerevisiae*: identification of efflux pumps involved in alkane tolerance. *Biotechnol. Biofuels* 6, 95. doi:10.1186/1754-6834-6-95
- Liu, B., and Zhao, Z. (2007). Biodiesel production by direct methanolysis of oleaginous microbial biomass. *J. Chem. Technol. Biotechnol.* 82, 775–780. doi:10.1016/j.jbiortech.2014.01.111
- Liu, R., Zhu, F., Lu, L., Fu, A., Lu, J., Deng, Z., et al. (2014a). Metabolic engineering of fatty acyl-ACP reductase-dependent pathway to improve fatty alcohol production in *Escherichia coli*. *Metab. Eng.* 22, 10–21. doi:10.1016/j.ymben.2013.12.004
- Liu, Y., Wang, C., Yan, J., Zhang, W., Guan, W., Lu, X., et al. (2014b). Hydrogen peroxide-independent production of alpha-alkenes by OleTJE P450 fatty acid decarboxylase. *Biotechnol. Biofuels* 7, 28. doi:10.1186/1754-6834-7-28
- Mattanovich, D., Sauer, M., and Gasser, B. (2014). Yeast biotechnology: teaching the old dog new tricks. *Microb. Cell Fact.* 13, 34. doi:10.1186/1475-2859-13-34
- Mendez-Perez, D., Begemann, M. B., and Pfeleger, B. F. (2011). Modular synthase-encoding gene involved in alpha-olefin biosynthesis in *Synechococcus* sp. strain PCC 7002. *Appl. Environ. Microbiol.* 77, 4264–4267. doi:10.1128/AEM.00467-11
- Mussatto, S. I., Dragone, G., Guimaraes, P. M., Silva, J. P., Carneiro, L. M., Roberto, I. C., et al. (2010). Technological trends, global market, and challenges of bio-ethanol production. *Biotechnol. Adv.* 28, 817–830. doi:10.1016/j.biotechadv.2010.07.001
- Nielsen, J., Larsson, C., van Maris, A., and Pronk, J. (2013). Metabolic engineering of yeast for production of fuels and chemicals. *Curr. Opin. Biotechnol.* 24, 398–404. doi:10.1016/j.copbio.2013.03.023
- Paddon, C. J., Westfall, P. J., Pitera, D. J., Benjamin, K., Fisher, K., McPhee, D., et al. (2013). High-level semi-synthetic production of the potent antimalarial artemisinin. *Nature* 496, 528–532. doi:10.1038/nature12051
- Qiu, Y., Tittiger, C., Wicker-Thomas, C., Le, G. offG., Young, S., Wajnberg, E., et al. (2012). An insect-specific P450 oxidative decarboxylase for cuticular hydrocarbon biosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* 109, 14858–14863. doi:10.1073/pnas.1208650109
- Reiser, S., and Somerville, C. (1997). Isolation of mutants of *Acinetobacter calcoaceticus* deficient in wax ester synthesis and complementation of one mutation with a gene encoding a fatty acyl coenzyme A reductase. *J. Bacteriol.* 179, 2969–2975.

- Rude, M. A., Baron, T. S., Brubaker, S., Alibhai, M., Del Cardayre S. B., and Schirmer, A. (2011). Terminal olefin (1-alkene) biosynthesis by a novel p450 fatty acid decarboxylase from *Jeotgalicoccus* species. *Appl. Environ. Microbiol.* 77, 1718–1727. doi:10.1128/AEM.02580-10
- Runguphan, W., and Keasling, J. D. (2014). Metabolic engineering of *Saccharomyces cerevisiae* for production of fatty acid-derived biofuels and chemicals. *Metab. Eng.* 21, 103–113. doi:10.1016/j.ymben.2013.07.003
- Sangwalak, J., Kaneko, Y., Sugiyama, M., Ono, H., Bamba, T., Fukusaki, E., et al. (2013). Ketoacyl synthase domain is a major determinant for fatty acyl chain length in *Saccharomyces cerevisiae*. *Arch. Microbiol.* 195, 843–852. doi:10.1007/s00203-013-0933-3
- Scharnewski, M., Pongdontri, P., Mora, G., Hoppert, M., and Fulda, M. (2008). Mutants of *Saccharomyces cerevisiae* deficient in acyl-CoA synthetases secrete fatty acids due to interrupted fatty acid recycling. *FEBS J.* 275, 2765–2778. doi:10.1111/j.1742-4658.2008.06417.x
- Schirmer, A., Rude, M. A., Li, X., Popova, E., and del Cardayre, S. B. (2010). Microbial biosynthesis of alkanes. *Science* 329, 559–562. doi:10.1126/science.1187936
- Shi, S., Chen, Y., Siewers, V., and Nielsen, J. (2014a). Improving production of malonyl coenzyme a-derived metabolites by abolishing Snf1-dependent regulation of Acc1. *MBio* 5, e01130–14. doi:10.1128/mBio.01130-14
- Shi, S., Valle-Rodriguez, J. O., Siewers, V., and Nielsen, J. (2014b). Engineering of chromosomal wax ester synthase integrated *Saccharomyces cerevisiae* mutants for improved biosynthesis of fatty acid ethyl esters. *Biotechnol. Bioeng.* 111, 1740–1747. doi:10.1002/bit.25234
- Shi, S., Valle-Rodriguez, J. O., Khoomrung, S., Siewers, V., and Nielsen, J. (2012). Functional expression and characterization of five wax ester synthases in *Saccharomyces cerevisiae* and their utility for biodiesel production. *Biotechnol. Biofuels* 5, 7. doi:10.1186/1754-6834-5-7
- Sorger, D., and Daum, G. (2003). Triacylglycerol biosynthesis in yeast. *Appl. Microbiol. Biotechnol.* 61, 289–299. doi:10.1007/s00253-002-1212-4
- Starai, V. J., Gardner, J. G., and Escalante-Semerena, J. C. (2005). Residue Leu-641 of acetyl-CoA synthetase is critical for the acetylation of residue Lys-609 by the protein acetyltransferase enzyme of *Salmonella enterica*. *J. Biol. Chem.* 280, 26200–26205. doi:10.1074/jbc.M504863200
- Sukovich, D. J., Seffernick, J. L., Richman, J. E., Hunt, K. A., Gralnick, J. A., and Wackett, L. P. (2010). Structure, function, and insights into the biosynthesis of a head-to-head hydrocarbon in *Shewanella oneidensis* strain MR-1. *Appl. Environ. Microbiol.* 76, 3842–3849. doi:10.1128/AEM.00433-10
- Tang, X., Feng, H., and Chen, W. N. (2013). Metabolic engineering for enhanced fatty acids synthesis in *Saccharomyces cerevisiae*. *Metab. Eng.* 16, 95–102. doi:10.1016/j.ymben.2013.01.003
- Tee, T. W., Chowdhury, A., Maranas, C. D., and Shanks, J. V. (2014). Systems metabolic engineering design: fatty acid production as an emerging case study. *Biotechnol. Bioeng.* 111, 849–857. doi:10.1002/bit.25205
- Valle-Rodriguez, J. O., Shi, S. B., Siewers, V., and Nielsen, J. (2014). Metabolic engineering of *Saccharomyces cerevisiae* for production of fatty acid ethyl esters, an advanced biofuel, by eliminating non-essential fatty acid utilization pathways. *Appl. Energy* 115, 226–232. doi:10.1016/j.apenergy.2013.10.003
- Wang, W., and Lu, X. (2013). Microbial synthesis of alka(e)nes. *Front. Bioeng. Biotechnol.* 1:10. doi:10.3389/fbioe.2013.00010
- Willis, R. M., Wahlen, B. D., Seefeldt, L. C., and Barney, B. M. (2011). Characterization of a fatty acyl-CoA reductase from *Marinobacter aquaeolei* VT8: a bacterial enzyme catalyzing the reduction of fatty acyl-CoA to fatty alcohol. *Biochemistry* 50, 10550–10558. doi:10.1021/bi2008646
- Youngquist, J. T., Schumacher, M. H., Rose, J. P., Raines, T. C., Politz, M. C., Copeland, M. E., et al. (2013). Production of medium chain length fatty alcohols from glucose in *Escherichia coli*. *Metab. Eng.* 20, 177–186. doi:10.1016/j.ymben.2013.10.006
- Yu, K. O., Jung, J., Kim, S. W., Park, C. H., and Han, S. O. (2012). Synthesis of FAEEs from glycerol in engineered *Saccharomyces cerevisiae* using endogenously produced ethanol by heterologous expression of an unspecific bacterial acyltransferase. *Biotechnol. Bioeng.* 109, 110–115. doi:10.1002/bit.23311
- Zhang, F., Carothers, J. M., and Keasling, J. D. (2012). Design of a dynamic sensor-regulator system for production of chemicals and fuels derived from fatty acids. *Nat. Biotechnol.* 30, 354–359. doi:10.1038/nbt.2149
- Zhou, Y. J., Gao, W., Rong, Q., Jin, G., Chu, H., Liu, W., et al. (2012). Modular pathway engineering of diterpene synthases and the mevalonic acid pathway for mitratriene production. *J. Am. Chem. Soc.* 134, 3234–3241. doi:10.1021/ja2114486

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 31 May 2014; accepted: 18 August 2014; published online: 01 September 2014.

Citation: Zhou YJ, Buijs NA, Siewers V and Nielsen J (2014) Fatty acid-derived biofuels and chemicals production in *Saccharomyces cerevisiae*. *Front. Bioeng. Biotechnol.* 2:32. doi: 10.3389/fbioe.2014.00032

This article was submitted to *Synthetic Biology*, a section of the journal *Frontiers in Bioengineering and Biotechnology*.

Copyright © 2014 Zhou, Buijs, Siewers and Nielsen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.