



# Functional Pituitary Networks in Vertebrates

Yorgui Santiago-Andres<sup>1</sup>, Matan Golan<sup>2\*</sup> and Tatiana Fiordeliso<sup>1\*</sup>

<sup>1</sup> Laboratorio de Neuroendocrinología Comparada, Departamento de Ecología y Recursos Naturales, Biología, Facultad de Ciencias, Universidad Nacional Autónoma de México, Ciudad Universitaria, Ciudad de México, Mexico, <sup>2</sup> Department of Poultry and Aquaculture, Institute of Animal Sciences, Agricultural Research Organization, Rishon Lezion, Israel

## OPEN ACCESS

### Edited by:

Romain Fontaine,  
Norwegian University of Life Sciences,  
Norway

### Reviewed by:

Xavier Bonnefont,  
INSERM U1191 Institut de Génétique  
Fonctionnelle (IGF), France  
Takayoshi Ubuka,  
Waseda University, Japan

### \*Correspondence:

Tatiana Fiordeliso  
tfiorde@ciencias.unam.mx  
Matan Golan  
matan.golan@mail.huji.ac.il

### Specialty section:

This article was submitted to  
Neuroendocrine Science,  
a section of the journal  
Frontiers in Endocrinology

**Received:** 20 October 2020

**Accepted:** 07 December 2020

**Published:** 27 January 2021

### Citation:

Santiago-Andres Y, Golan M and  
Fiordeliso T (2021) Functional Pituitary  
Networks in Vertebrates.  
*Front. Endocrinol.* 11:619352.  
doi: 10.3389/fendo.2020.619352

The pituitary is a master endocrine gland that developed early in vertebrate evolution and therefore exists in all modern vertebrate classes. The last decade has transformed our view of this key organ. Traditionally, the pituitary has been viewed as a randomly organized collection of cells that respond to hypothalamic stimuli by secreting their content. However, recent studies have established that pituitary cells are organized in tightly wired large-scale networks that communicate with each other in both homo and heterotypic manners, allowing the gland to quickly adapt to changing physiological demands. These networks functionally decode and integrate the hypothalamic and systemic stimuli and serve to optimize the pituitary output into the generation of physiologically meaningful hormone pulses. The development of 3D imaging methods and transgenic models have allowed us to expand the research of functional pituitary networks into several vertebrate classes. Here we review the establishment of pituitary cell networks throughout vertebrate evolution and highlight the main perspectives and future directions needed to decipher the way by which pituitary networks serve to generate hormone pulses in vertebrates.

**Keywords:** pituitary, networks, plasticity, vertebrates, evolution

## INTRODUCTION

Biological variation is a defining characteristic in the process of species evolution and adaptation. Variation in morphology, physiology, or behavior within species is a prerequisite for evolutionary processes, like natural selection, to occur. In an ever-changing natural world, the organisms able to respond more efficiently to those changes are better fitted to their environment and could be selected (1). In this context, highly plastic systems are also adaptive since the information from both external and internal environments is better integrated. An example of such a highly plastic system is the pituitary gland, which secretes several key hormones that maintain homeostasis in all vertebrates (2).

The pituitary gland is embryonically derived from Rathke's pouch and is therefore considered to be of ectodermal origin (3, 4), although at least in fish, a small percentage of pituitary endocrine cells have endodermal origins (5). The pituitary of all vertebrates consists of three anatomically and developmentally distinct structures called the neurohypophysis or posterior pituitary, the adenohypophysis, or anterior lobe and the intermediate lobe, all of which function in close interdependence with the hypothalamus. The neurohypophysis comprises the neural lobe

(mainly oxytocin- and vasopressin-secreting neuroendocrine terminals in mammals and their homologous isotocin and vasotocin in non-mammalian vertebrates), the median eminence (the functional link between the hypothalamus and the anterior lobe), and the infundibulum. The intermediate lobe is mainly composed of melanotrophs that regulate skin pigmentation. The anterior lobe of the pituitary, which is the main focus of this review, is populated by five types of secretory cells: somatotrophs, gonadotrophs, lactotrophs, thyrotrophs, and corticotrophs, that may be identified by their hormone composition and their response to hypothalamic neurohormones. In mammals, these cell types follow a similar pattern of stimulus-secretion coupling that regulates the neuroendocrine axes and allows the preservation of homeostasis (6).

In all vertebrates, the main driver of pituitary secretion is the hypothalamic input. Factors released from hypothalamic neuron terminals into the pituitary portal system bind to their cognate receptors on target pituitary cells and elicit a signaling cascade that induces release (or inhibition) of hormone secretion from the pituitary (7) into the vascular system in a pulsatile manner, the pattern of which is critical in regulating the activity of the downstream target organs (2). Yet the individual cellular dynamics *in vitro* do not recapitulate the whole gland pulse generation needed for physiological actions. For example, *in vivo*, GH-secreting cells produce massive GH pulses (up to 1,000-fold rise) in response to physiological cues (8, 9), whereas these pulses are significantly weaker *in vitro* (10). The same dependence of hormone pulse generation on the tissue context has been observed in gonadotrophs (11–14), corticotrophs (15–17), and lactotrophs (18, 19). Thus, in order to generate this fine-tuned and coordinated secretion, a large-scale communication system must exist within the anterior pituitary that is critical for the ability to generate hormone surges.

In recent years, the use of high resolution imaging in combination with large-scale cell activity recordings has revealed the homotypic and heterotypic network arrangement of the mammalian pituitary cells (i.e., intercellular connectivity between the same secretory cell type or between different cell types, respectively). Research in this field has pointed out the relevance of the network organization in coordinating the synchronized and rhythmic secretion of hormones (16, 20). Intriguingly, the existence of cell networks has been identified in all major groups of vertebrates. As an example, in fish, which represent the most basal group of vertebrates, when the gonadotroph network connectivity is disrupted, the hormone secretion is compromised, as well as its biological significance (21, 22). It can be concluded from these facts that a) pulse generation is achieved thanks to cell networks, b) the importance of the pulse mode of secretion is revealed by studies in which its production is limited, and is followed by physiological perturbations in the target organs of regulation, and c) cell network formation, which is essential for the organization of these pulses, is conserved throughout vertebrate evolution (2).


Despite the clear importance of pituitary homotypic and heterotypic networks in the coordination of hormone

pulsatility, research into this subject has been limited. Here, we review the current knowledge and methodological approaches to assess the function of pituitary networks throughout vertebrate evolution. The mechanisms involved in pulse generation, as an adaptive and plastic process that promotes the rhythmic secretion of hormones in vertebrates, will be discussed from a cell network perspective. We then explore the evolution of these networks in different groups of vertebrates, from fish to mammals and discuss the importance of the cellular networks in the phenotypic plasticity. We conclude that a comparative approach to the study of pituitary networks is fundamental to the comprehension of the system both under normal and pathological conditions. It is important to clarify that factors such as the well-established hypothalamic drive, the peripheral tissues and the environment are fundamental in the generation of hormone pulses and should be integrated in order to understand the system as a whole. However, the pituitary gland has its own local and plastic modes of autonomic regulation and its interaction with the hypothalamus is a complex two-ways regulation.

## HYPOTHALAMIC CONTROL OF THE PITUITARY GLAND IN VERTEBRATES

Over the course of vertebrate evolution, the general morphology of the pituitary gland has remained constant, with three main lobes whose function depends on the integration of signals originating from hypothalamic nuclei and from the systemic circulation (23). While a general anatomic plan is observed in the pituitary gland of all vertebrates (**Figure 1**), the specific arrangement of the endocrine cells (**Figure 2**), as well as the way by which their communication with the brain is executed, show great variation across taxa.

Teleost pituitaries in particular exhibit an important deviation from the masterplan of the tetrapod pituitary in respect to the anatomy of the hypothalamus-pituitary interface. Studies of fish pituitaries found that in ancient forms such as cartilaginous fish, sturgeons, and gars the hypothalamic fibers terminate and release their content onto the basal membrane separating the neurohypophysis from the adenohypophysis. The secreted hypothalamic factors then diffuse through the connective tissue and are uptaken by the pituitary vascular system that delivers them to their pituitary target cells (24). This neuro-vascular mode of delivery was perpetuated in the tetrapod evolution and is the dominant mode by which the hypothalamus exerts its effect on pituitary cells in all known tetrapods (24) (see **Figure 1**). However, studies of teleost pituitaries revealed a unique characteristic that was not found in any other vertebrate: hypothalamic fibers crossed the basement membrane that separates the neuronal tract from the adenohypophysis and formed terminals adjacent to endocrine cells (see teleost fish in **Figure 3**) (25–31). This unique characteristic drew so much attention, that a direct, neuroglandular mode of regulation was considered the main mode operating in teleosts.




The development of transgenic fish with labeling of three components of the HP axis (GnRH3 cells, gonadotropes and blood vessels) enabled researchers to re-visit the existing dogma. Upon the examination of the GnRH tracts in pituitaries labeled for gonadotropes it became clear that the direct neuroglandular mode of regulation cannot be considered as the dominant route of regulation. This is because not all gonadotropes are contacted by GnRH terminals and because the vast majority of these terminals are not in the vicinity of the gonadotropes (32). By using transgenic fish with labeled blood vessels it became apparent that the GnRH terminals are in much closer association with the pituitary afferent vessels than with the gonadotropes (see the teleost fish in **Figures 3** and **5**). This organization strongly suggests that, at least in zebrafish, GnRH signals reach their pituitary targets through the blood or through diffusion rather than *via* direct neuro-glandular contacts. A dominant neuro-vascular interaction is also observed in other species and other hypothalamus-pituitary axes (31–36).

The distinction between a direct, neuroglandular mode of regulation *versus* an indirect, neuro-hemal mode of hypothalamic factor delivery has important consequences over the functional physiology of the hypothalamic pituitary axis. A direct, neuroglandular mode of communication would require considerably smaller concentrations of hypothalamic factors to be secreted since the secretagogues will be delivered directly to the target cell. Such a direct communication mode may also

affect the timescale of the regulation as a very temporally precise signal could be delivered to the target cells resembling the timescales of communication between neurons. Moreover, if the anatomical interactions between terminals and cells are very precise, it would make the variety of hypophysiotropic signals largely redundant as the specificity of the regulation could be achieved by controlling the type of neuron contacting a specific pituitary cell type rather than by different secretagogues. To date, there is no evidence to suggest that teleost pituitaries differ in any of these aspects from the more widely studied mammalian pituitary.

While it is admittedly difficult to evaluate the functional contribution of the different delivery modes, at least from a quantitative point of view, the direct, synaptic-like contacts between terminals and their pituitary targets seem to play a less important role than previously thought in the regulation of pituitary secretion. Instead, diffusion from terminals through the pituitary tissue to the target cells (either passive, or, like in tetrapods, assisted by local blood flow), appears to be the dominant way by which pituitary endocrine cells receive their hypothalamic signals. Therefore, the anterior neurohypophysis of teleosts may act as a functional median eminence that is embedded into the pituitary in teleosts. The evolution of the hypothalamo-pituitary axis has recently been thoroughly reviewed by Trudeau and Somoza (37) and interested readers are encouraged to refer to this excellent review for more detail.



In the next section we briefly cover the principal mechanisms involved in the maintenance of homotypic and heterotypic cellular networks in the pituitary, by means of cell-cell contacts or paracrine communication. Those mechanisms are present in all vertebrates but their contribution may vary depending on the communication with the hypothalamus, vascular arrangement of the pars distalis, distribution of cell types in the gland (regionalized or scattered), and formation of heterotypic networks (see **Figures 1** and **2**).


## MODES OF COMMUNICATION

### Gap Junctions

Gap-junctions (GJ) are transmembrane channels that allow the free cell-to-cell exchange of cytoplasmic molecules of <1,000 daltons molecular weight. The intercellular communication through GJ is an important source in the transport of cytoplasmic constituents (including water and ions), metabolic substrates (such as sugars, amino acids, nucleotides), and second messengers (mainly calcium,  $IP_3$ , and cAMP) between cells. In the pituitary gland, gap junctions are an important mode of communications in all studied vertebrates. For example, gap junctions have been shown to couple fish pituitary cells (38, 39). In the rat, cells show a high level of intercellular coupling through gap junctions, where lucifer yellow diffusion was

found up to 300-microns away from its site of injection (40). In addition, the transport between cells was primarily in lactotrophs, somatotrophs and folliculostellate cells (FS cells). Only a few LH, TSH, and ACTH cells were labeled (40). In all vertebrates gap junctions are considered a critical component in the coupling of pituitary cells and allow a coherent transmission of signals and the formation of cell networks (2, 19, 22, 41–48).

In terms of cell activity, Guéroux and collaborators have demonstrated the fundamental role of gap junctions in the synchronization of electrical activity (49). By multicellular measurement of spontaneous intracellular calcium mobilization in tissue preparations, the authors found the presence of clusters of pituitary cells that were in close proximity to each other and rhythmically coactive. As small molecules (Lucifer yellow, 457 Da), but not large molecules (Texas Red, 3 000 Da) spread between the co-active cells, in combination with the observation that a gap-junction blocker reduces the spread of synchronization, it was concluded that coordination between neighboring secretory cells is mediated by gap junctions. Of particular importance, the connected cells were principally of the same phenotype but not all the adjoining cells from the same phenotype were coupled. This fact indicated that the global response of secretory cells during hypothalamic stimulation involves, as we shall see, other mechanisms of cell communication that are not mutually exclusive.



**FIGURE 3** | The gonadotroph network and the relation with the hypophyseal portal system in vertebrates. Midsagittal view of the pituitary gland in relation to the hypothalamus (anterior at the left, dorsal to the top) using confocal microscopy. Shown are immunostainings for LH $\beta$  (green) in five different species of vertebrates (zebra fish, *Danio rerio*; axolotl, *Ambystoma mexicanum*; bullfrog, *Lithobates catesbeianus*; lizard, *Sceloporus aeneus*; mouse, *Mus musculus*). In contrast to tetrapods, where LH-containing gonadotrophs are well distributed in the pars distalis, in teleost these cells are confined to the rostral pars distalis. The merge detail column is a power magnification of the precedent, showing that gonadotrophs establish cell-cell contact to form an interdigitated network and reveal that this pattern of cell organization is extended to all vertebrates. Note that the vascular network pervades the pars distalis and gonadotrophs form process to the capillaries (red, lectin-rhodamine), with the exception of zebrafish. DAPI counterstain is shown in blue. Dashed lines mark the three main regions of the pituitary. HYP, hypothalamus; PD, pars distalis; PI, pars intermedia; PN, pars nervosa; ME, median eminence. Scale bars represent 400  $\mu$ m in whole-gland views and 20  $\mu$ m for merge detail.

Changes in gap junction expression patterns have been linked to plasticity in the pituitary gland. Modifications in connexins 43 and 26 expression (subunits of gap junctions), have been reported in rodents, following physiological challenges and

aging (50–54). For example, GJ expression is modified according to the stage of the estrous cycle, probably reflecting the plasticity of cell connectivity during the generation of hormone pulses. Indeed, other authors have reported that

estrogens play an important role in their regulation, since the absence of testosterone and estradiol downregulate the number of gap junctions by a quarter or more compared to intact rats (46, 55–57). Other experiments show that factors such as hydrocortisone, have a suppressive effect on the GJ formation between FS cells and between endocrine pituitary cells (58).

## Folliculostellate Cells

A population of glial-like cells, the FS cell network, is present in the pituitary gland of all major taxa of living vertebrates, and in mammals these non-endocrine cells comprise about 5–10% of the anterior pituitary cell mass (39, 59–62). These agranular cells have a characteristic stellate shape with long cytoplasmic processes, and arranged as a 3D anatomical network that extends over the whole gland in mammals (48, 53), aves (62), amphibians (63), and in fish (39). FS cells respond to a large variety of external and internal stimuli and produce growth factors and cytokines, such as fibroblast growth factor (FGF)-2, vascular endothelial growth factor (VEGF)-A, follistatin, and IL-6, all of which potentially have paracrine effects and regulate the activity of neighboring hormone-secreting pituitary cells (64).

The FS cell network has been suggested to wire the endocrine cells of the pituitary by relaying relevant signals throughout the gland. Secretory cells in the pituitary are intertwined through the vasculature and the FS network. The wave propagation of ions as  $\text{Ca}^{2+}$ , and small molecules as  $\text{IP}_3$  or cAMP, through GJ within the FS cell network, allow synchronized excitability within the anterior pituitary gland (48, 65). Although, the distance and velocity in which the information can propagate to parenchymal cells has not been investigated yet, there is evidence supporting that the anterior pituitary possesses an intrinsic system of communication, that allows long-distance transmission of information within its parenchyma (66, 67). FS cells may have a central role in this process due to their gap-junction mediated network properties, and the expression of a large variety of signaling molecules. Hence, distant endocrine cells might receive coordinated information from both systemic molecules through the vasculature and paracrine signals primarily coming from the FS-cell network (48).

In mammals, it has been reported that FS cells can generate different responses in groups of endocrine cells surrounded by their cytoplasmic processes, thus forming functional clusters (66–68). Interestingly, it has been observed that GnRH-neuron projections converge not only at the blood capillaries, they are also adjacent to FS cells in the pars tuberalis (68, 69). Still, gap junctions between the GnRH-neuron terminals and FS cells, as well as GnRH-induced calcium response in FS cells have been recognized (69–71). In this sense, in addition to the vascular system, cell signaling in the FS network generated by hypothalamic stimuli, may be a route to propagate signals throughout the pituitary (66–69, 72, 73). In this respect, the unique anatomy of the teleost pituitary may provide interesting insights, since stellate cells have been reported to form dense networks around the neurohypophysis that penetrates deep into the parenchyma of the adenohypophysis (39). As a hypothesis to be proved, some endocrine cells may receive the signals only

from the pituitary portal system and others could additionally integrate signals propagated by the FS-cell network, according to their distribution in the pituitary (70) (see **Figure 2**).

## Paracrine Communication

Paracrinicity is the process of short-distance communication between cells by substances released within the same tissue. The substance reaches its target by diffusion in the extracellular space or by direct contact formation (juxtacrine factors). More than 100 compounds in the pituitary have been identified as paracrine or autocrine factors (64). They include the neurotransmitters, growth factors, cytokines tissue factors such as annexin-1 and follistatin, hormones, ATP, NO, among others (64). The majority of paracrine or juxtacrine molecules, act through G-protein coupled receptors, with inhibitory ( $\text{Gi}$ ) or excitatory ( $\text{Gs}$ ) action, activating in most cases voltage gated channels activity and intracellular  $[\text{Ca}^{2+}]$ , as well as a whole host of other receptor types capable of affecting down-stream kinase and protein phosphorylation status (74). Paracrine signaling is a slow diffusion process, which occurs over seconds to minutes. This process of molecular diffusion is one of the main scenarios for possible organization and communication presumably sufficient to account for coordinated endocrine cell population activity and surge hormone output (74). However, this diffusion process seems to lack the directionality that would allow the formation of homo- or heterotypical networks, unless evidence is found to show that it could be a process directed through, for example, the interaction of molecules with the extracellular matrix. A more spatially precise intercellular signaling may be accomplished in the form of juxtacrine factors, as annexin 1 (formerly lipocortin-1), a phospholipid-binding protein which is predominantly expressed by the FS network (74).

In mammals, an extended review by Denef summarizes the paracrine interactions in the pituitary gland (64). The first evidence of the possible paracrine relationship between endocrine cells was established from gonadotrophs and lactotrophs, where the stimulation of gonadotrophs secretion with gonadotropin-releasing hormone (GnRH) triggers prolactin secretion (PRL) (75). More complex relationships were established between gonadotrophs and somatotrophs, in which it was observed that in aggregates of these two cell types, the GnRH stimulation produce an initial inhibition of GH release, whereas after GnRH stimulation, a rapid rebound secretion of GH was observed, suggesting that GnRH had both inhibitory and stimulatory actions on GH release (76). Gonadotrophs have a large number of secreted peptides who are potential candidates for a paracrine action. Some of these peptides modulate corticotroph activity when added exogenously to pituitary cell preparations (64). An inverse relation between lactotrophs and gonadotrophs was also established by the inhibition of LH secretion in hyperprolactinemia condition (77).

Other possible paracrine interactions imply coordination between different pituitary endocrine cells. For instance, the hypothalamic-pituitary-adrenal axis that is activated during stress. In this condition, there is a need to increase the metabolic rate in order to maintain body temperature, and this

process is regulated by the hypothalamic-pituitary-thyroid axis (78). Further example of crosstalk with the HPT axis is during pregnancy and lactation, in which energy consumption has to be adapted. The mechanisms involved in the regulation of these homeostatic changes are still largely unknown but paracrine interactions are likely to be involved in the heterotypic network communication (64).

In fish, paracrine interactions between pituitary cells have been shown to comprise an important regulatory layer of pituitary cell function and probably operate at a slower timescale than gap-junction mediated transmission. Perhaps one of the most interesting aspects of this communication mode in fish is the interplay between the growth and reproductive axis through the paracrine effects of LH on GH cells. Studies in grass carp have shown that both GH and LH have a stimulatory effect on GH expression and release whereas GH has an inhibitory effect on LH release (79). Cross communication between the growth and reproductive axes in fish is also achieved *via* the activin/inhibin/follistatin pathway as these genes are expressed in the pituitary and have known effects on gonadotropin expression (80–83) and the GH-LH interaction (84). The expression and secretion of somatolactin (SL), a member of the GH/prolactin family, has been shown to be autoregulated by the two SL isoforms (85, 86) as well as by kisspeptin (87), a hypothalamic hormone that in fish is also produced in the pituitary gland (88). Other proposed autocrine and paracrine effects between different axes in the fish pituitary have been proposed such as the effect of NKB produced in gonadotropes (89) on lactotrophs and somato-lactotrophs (90). Secretoneurin-a, a potent stimulator of gonadotropes in goldfish is produced in lactotrophs of the rostral pars distalis and may link the two axes *via* paracrine effects on the gonadotropes of the proximal pars distalis (91). Recent findings in long-term culture adult medaka female pituitary, showed that, Fsh cells which do not possess GnRH receptors start to express them, allowing Fsh cells to produce Lh. These observations suggest that *in vivo*, a paracrine signal inhibits GnRH receptor expression in Fsh cells and maintains their mono-hormonal identity (21, 35, 92).

It is also possible that paracrine factors not just mediate homotypic and heterotypic network formation, but also could have a priming effect on axis function because in PRL cells, the increased organization associated with lactation persists for months after weaning and leads to enhanced function (93). Despite a large body of evidence, we are still far from understanding the full effect of paracrinicity on network formation, because it is technically difficult to isolate this process from others such as communication between GJ or electrical conduction in FS cells.

## FUNCTIONAL PITUITARY NETWORKS IN VERTEBRATES

### Fish

One of the unique features of the fish pituitary is a high level of regionalization in the distribution of endocrine cells [(37, 94) and

Figures 2 and 3]. This compartmentalization of the cells results in homotypic clusters of cells that favor direct cell-cell interactions and the formation of homotypic cell networks. Yet despite this unique topology, the study of cell networks in fish pituitaries has lagged behind that of mammals. The first evidence for direct cell-cell communication in fish pituitaries was the discovery of electrotonic cell coupling in the tilapia pituitary (38). Using double-patch electrophysiology on live pituitary slices, this groundbreaking work has shown for the first time the existence of gap-junction mediated communication between teleost pituitary cells. However, at the time, the type of coupled cells was unknown. Since then, many of the new insights regarding pituitary networks in fish were drawn from three model species: the zebrafish (*Danio rerio*), the Japanese medaka (*Oryzias latipes*), and Nile tilapia (*Oreochromis niloticus*). In these species, the generation of transgenic fish with labeled pituitary cells (35, 95–98) was an important turning point as it provided the opportunity to study topological cells networks in 3D and opened the way to functional studies of cell-cell interactions. Interestingly, in all three species, the main study of pituitary networks has concentrated on the gonadotrope cells of the anterior pituitary, leaving future studies to focus on other cell types. One of the reasons for this specific focus on gonadotropes stems from the fact that in fish, as opposed to tetrapods, LH and FSH are secreted from discreet cell populations rather than from a single cell type. This feature makes teleosts a particularly attractive model to study the differential regulation of LH and FSH secretion.

Studies in these three teleost species confirmed the existence of functional gonadotrope networks in fish. In tilapia and zebrafish, we have shown that LH cells are connected by gap junctions (22). In medaka, Karigo and co-authors used calcium imaging to show that in the intact gland LH cells exhibit a high level of synchronization in their activity (98) although the mode by which the cells communicate was not investigated. More recently, working with pituitary slices from medaka, Hodne *et al.* have used precise calcium uncaging in individual LH cells to show that the calcium signal can propagate to neighboring cells in a both homo- and heterotypic manner, leading to the conclusion that LH cells act as a relay for GnRH signals to FSH cells (35). The importance of the functional coupling has been revealed by its effect on hormone secretion from tilapia pituitaries. Application of gap junction blockers decreased GnRH-induced LH release from perfused pituitary fragments seven-fold (22), underscoring the role of the network in mounting an efficient response to hypothalamic stimuli. Interestingly, while tilapia LH cell output is highly attenuated by application of gap-junction blockers, FSH release remains largely unaffected. In medaka, calcium activity of FSH cells also seems to be less synched compared to their LH counterparts suggesting that the lower level of coupling between FSH cells may be a common feature of fish FSH gonadotropes. The notion that FSH cells act individually while LH cells mount coordinated responses is also evident in their secretion patterns since the induction of gonadotropin secretion *in vivo* causes a considerably stronger fold increase over basal levels in LH compared to FSH (89, 99–101).

Long extensions have been described in fish gonadotropes both *in situ* and in culture (32, 95, 102, 103). It is difficult to postulate as to the function of these processes but two plausible options are that these appendages are used for cell-cell communication or as a bridge between the cells and the vasculature. One line of evidence from cultured medaka cells (102) suggests that these processes do not relay signals between cells but are instead used for cell motility which serve to bring the cells together thus promoting cell-cell contacts and the formation of a network. It is hard to evaluate whether the processes serve the same role *in vivo* as in the adult medaka pituitary the cells are highly clustered yet the projections are still observed (102). Findings from the zebrafish model suggest that these processes are more likely to operate as extensions of the cells toward the vasculature than for intercellular communication. In the developing zebrafish, FSH cell extensions are clearly oriented in one direction: toward the primary plexus of vessels entering the pituitary (32). The aforementioned cytoplasmic processes are all directed toward the afferent vessels and terminate on their endothelium. This finding is even less surprising when considering the fact that not only metabolites are acquired from the blood, but also GnRH signals. At this early developmental stage no contact between gonadotrope and GnRH terminal exists. However GnRH fibers do reach the primary plexus, thus their output is carried through the blood to reach the developing gonadotropes (32). In addition to receiving blood-borne signals through the processes, these projections can also be used to deposit hormones to the circulation. In fact, medaka LH cell projections have been shown to develop large varicosities around the process-vessel junctions (102). It therefore seems more likely that the gonadotrope extensions are used to access the circulation than to mediate cell communication, though further investigations into this aspect are required.

While the study of pituitary cell networks in fish is still in its early stages, it is clear today that these networks developed early during vertebrate evolution and their function was conserved in higher vertebrates. In order to fully appreciate the role and function of pituitary networks in fish, future studies will have to extend the focus onto other pituitary cell types and probe the plasticity of these networks in response to physiological challenges that require the modulation of pituitary output.

## Amphibians Metamorphosis

The process of metamorphosis in vertebrates is well represented by the post-embryonic morphological remodeling in amphibians, where changes in physiology and behavior also occur. During metamorphosis, the neuroendocrine system plays a major role orchestrating the modifications at cellular and molecular levels in tissues by thyroid hormones (104–106). Particularly, a peak of thyroid hormone secretion marks the beginning of the process and the thyrotrophs, from the hypothalamus-pituitary-thyroid axis, exert a direct action over the production of this peak (107). As thyroid hormone continues to stimulate tissue remodeling, a continuous activity of

thyrotrophs in the pituitary is also needed to maintain the secretion levels of the former (108, 109). Thus, thyrotrophs must shift plastically the secretion pattern by remodeling their structure and their relation with other cellular types within the pituitary and feedback loops in the axis (108, 110).

It is noteworthy that TSH secretion during amphibian metamorphosis is stimulated by CRF, the hypothalamic factor from the stress axis and not by TRH (111, 112). It is now understood that in amphibians, TRH activates TSH secretion in thyrotrophs of metamorphosed individuals but not in larval ones as reported in the bullfrog, reflecting that the thyrotroph plasticity is modified ontogenetically (113, 114). This phenomenon implies that thyrotrophs express receptors for TRH and/or that corticotrophs may regulate TSH secretion through paracrine communication (110, 115, 116). To date, the physiological and structural relationship between corticotrophs and thyrotrophs in the process of metamorphosis and whether a heterotypic network is formed remains to be elucidated, but it is possible to speculate about their intimate relationship, based on the shared main hypothalamic factor that regulates these cells types and also that the stress axis participates actively during the metamorphosis and its developmental plasticity (110, 117–120). Furthermore, cell plasticity in other pituitary cell types is also observed during metamorphosis. For example, when the salamander *Hynobius retardatus* was arrested prior to metamorphosis, an increase in the number of TSH-secreting cells was observed as well as an increase in bihormonal phenotypes such as TSH-PRL and TSH-GH, presumably involving transdifferentiation (121, 122).

The hypothalamus-pituitary-thyroid axis is also intertwined with the reproductive axis, where amphibian TSH in combination with gonadotropins has been demonstrated to trigger gonadal maturation (122, 123). Metamorphosis may modify gonadal maturation under environmental influence as demonstrated in urodels with neoteny, the heterochronic process where the larval phenotype is retained but organisms are sexually mature (124). Studies comparing the modification in the pituitary networks before and after metamorphosis in pedomorphic species (e.g., *Ambystoma mexicanum*; see **Figure 3**) with those with direct development (e.g., the bullfrog; see **Figure 3**) may reveal some of the molecular mechanisms involved in network plasticity, but also about its interaction with the hypothalamus (125–127).

## MSH Cells

Melanotrope cells are essential for amphibian camouflage *via* secretion of alpha-melanophore-stimulating hormones ( $\alpha$ -MSH), a hormone derived from the POMC protein (128, 129). The melanotrope cells are restricted to the pars intermedia; in contrast to other secretory cells in anterior pituitary that are heterogeneously distributed and where heterotypic contacts are formed. The cluster organization of melanotrophs, must permit a fine coordination in response and  $\alpha$ -MSH secretion, and probably involves gap junctions since expression of connexins (Cx43) has been reported (130, 131). The vascular system is a second candidate for melanotrope cells organization and

remodeling. In fact, in normal conditions, the pars intermedia is poorly vascularized compared to the adenohypophysis (see **Figure 3**), the capillary networks are present at the borders between the neural lobe and intermediate lobe (**Figure 1**). However, in transgenic frogs (*Xenopus laevis*) that overexpress VEGF-A in the pars intermedia, there is a complex formation of a vascular network (132). VEGF-A, the vascular endothelial growth factor, is involved in the formation of the vascular system during the development of the hypothalamus-pituitary system, and in the transgenic *Xenopus*, the increase of the vascular network has several consequences in the pars intermedia. First, these new capillaries are distributed around the lobules formed by melanotrophs. Second, the intermediate lobe is up to 3.5-fold greater than in control frogs as a consequence of melanotroph hypertrophy and hyperplasia, but the pars distalis is not reorganized. The implication of these results pointed to the active role of the vascular network in pituitary function and homeostasis. Further studies will be required to understand the effect of the vasculature in the pars distalis where cell networks are differentially disposed to the capillaries (see **Figure 3**). Specific modifications in the vascular system of animal models, as the example offered here, may also reveal how endocrine diseases emerge as vascular problems (2, 133).

## Reptiles and Birds

The establishment of cellular networks in the reptilian pituitary has not been studied directly, but the available literature indicates that homotypic and heterotypic networks may exist. For example, close contact between cells from the same lineage (including thyrotrophs), cluster formation, and cytoplasmic extensions have been reported by immunohistochemical studies (134–136). Due to their phylogenetic position between amphibians and mammals, reptiles gain relevance in the study of evolutionary transitions in the pituitary structure and function. Of particular importance may be the investigation of the chronological appearance of pituitary cell types, a process where ontogeny probably does not recapitulate phylogeny. Based on the work of Yamaki and collaborators, it has been demonstrated that the sequential development of endocrine cells in the mammalian pituitary varies largely in other vertebrates without a clear pattern, even close species show diversity in pituitary developmental programming (137). As an example, the establishment of the gonadotroph network upon the corticotroph scaffold, as described in mammals (138) may not be the case in turtles or fish, where corticotroph cell distribution is principally restricted to the rostral pars distalis, although gonadotrophs are widely distributed in the lobe (**Figures 2 and 3**). Comparative cell lineage in combination with other methods, such as lineage tracing, single RNA sequencing, and functional studies will reveal the mechanisms underlying the structure of pituitary networks, the heterochronic appearance in vertebrate development and their plasticity.

## Mammals

Of all vertebrates, pituitary networks in mammals, and particularly in mice, have been studied in the most

comprehensive manner and their existence has been demonstrated in lactotrophs, gonadotrophs, somatotrophs, corticotrophs, and folliculostellate cells (2, 16, 19, 20, 48). The modes by which mammalian pituitary cells communicate have been described earlier. Unlike in other vertebrate classes, works in mammals have focused not only on the existence of pituitary cell networks, but also on the way by which these networks modulate their function to meet the changing demands of the animal.

## Lactotrophs

The process of lactation requires a big amount of prolactin production and secretion from pituitary by lactotrophs. Milk production requires a 10–50-fold increase in prolactin secretion and involves a marked decrease of dopamine production by neurons in the hypothalamus (18, 139). Dopamine normally inhibits prolactin secretion through dopamine receptors coupled to inhibitory G-proteins that block the voltage gated calcium L-channel and change the membrane potential, but also through a short-loop feedback mediated by prolactin itself. However, during lactation DA-producing neurons switch from inhibiting prolactin secretion to promoting its release by secreting enkephalin (139). Although necessary, these hypothalamic changes do not account for the major alterations that support the increase in prolactin secretion (2).

In 2012, Hodson and collaborators reported the existence of physiological and morphological plasticity in the lactotroph network that reinforce the secretion activity of these cells and were not directly dependent on hypothalamic activity (19). First, from experiments of intracellular calcium measurements in lactotrophs, it was shown that during lactation, the network functional connectivity is more robust as both calcium coactivity between pairs of cells and correlation in calcium profiles increase. Remarkably, in virgin mice the network connectivity of lactotrophs is low and only few cells harbor the majority of these connections. In contrast, in lactotrophs from lactating mice there is an increase in the proportion of significantly correlated cells and new nodes emerge. This last characteristic of the network persists after weaning, even months after the lactation event is over. Second, increase in connectivity is not a direct result of the decrease in dopamine production, since in virgin mice treated with a dopamine-receptor antagonist, an increase in functional connectivity is not observed. Furthermore, the interruption or decrease in suckling stimulus during lactation significantly reduces this functional network at values to those observed in virgin mice. Third, the coordinated functional activity in the network is directly dependent on structural connectivity due to lactotroph hypertrophy and gap junctions. When a gap junction blocker was applied to the lactotroph network from a lactating female, the connectivity resembled the network of virgin mice. Finally, the plastic transition of this network remains assembled and is crucial in subsequent lactation events that are characterized by a prolactin secretion improvement (19). The effects of the lactotroph network reconfiguration on other networks, for example gonadotrophs and reproductive axis, and the heterotypic interactions remain open questions, but their physiological relevance is clear.

## Gonadotrophs

Another process where the hypothalamic regulation of pituitary does not account for the shift in hormone release is offered by gonadotrophs and LH secretion throughout the estrous cycle. LH and FSH secretion are necessary for follicle maturation and ovulation in females but they are differentially secreted through the reproductive cycle (140). For example, in proestrus the pattern of LH pulse secretion turns into a surge mechanism that has been largely seen as a consequence of the GnRH surge, the main stimulating factor of gonadotrophs in the hypothalamus. However, GnRH stimulus *per se* is not the causal phenomena for the significant increase in responsiveness and secretion by gonadotrophs (140, 141).

*In vitro*, it has been reported that during diestrus, the gonadotroph response to GnRH is limited and just a small proportion of the population present intracellular calcium activity (142). High concentrations of GnRH (100 to 1,000 nM) are necessary in this state of the cycle to produce the whole-population response, but the amount of LH secretion does not increase significantly. On the other hand, in proestrus, low GnRH concentrations are sufficient to trigger a whole population response and, more importantly, the majority of gonadotrophs secrete a larger amount of LH. Despite the increase in gonadotroph responsiveness and LH secretion during proestrus that depend upon GnRH receptor expression, these processes also involve the gonadotroph network reconfiguration and its relation with vasculature (143).

From experiments *ex vivo*, where tissue interactions are preserved, the gonadotroph network reveals dynamic and plastic adaptation in proestrus through an increase in cell number (probably involving cell proliferation and transdifferentiation). Gonadotrophs also increase cell-cell contacts (involving cell motility), and the increase of protrusions toward other gonadotrophs and to the vascular systems (141, 143). Taken together, these cell modifications and changes in responsiveness to GnRH during the estrous cycle, account for the regulation in the rhythm and mode of LH secretion at different phases of the reproductive cycle. The plasticity in gonadotrope cells at the molecular, cellular and population levels and a comparison of shared mechanisms between fish and mammals has been recently reviewed and for a more in-depth description the reader can refer to (21, 144).


Plasticity in gonadotrophs is canalized in a context-dependent manner: inputs from the reproductive axis as well as from the environment are interpreted within the network to activate or attenuate the system output. For example, testosterone, estradiol, and GnRH have been reported to stimulate the LH secretion, cell proliferation, and GnRH receptor expression when present at certain concentrations and rhythmicity, but may desensitize the network and inhibit the LH production and secretion when present for prolonged periods of time or higher concentrations (21, 145).

Most factors involved in the network establishment are necessary but not sufficient for a complete explanation of the mechanisms involved in the gonadotroph function as a network. GnRH is an element that triggers the network response since the

intrinsic activity of gonadotrophs is not particularly synchronized and is not associated with gonadotropin secretion (see **Figure 4**) (147). Furthermore, the amount of gonadotropin secretion is influenced by the balance of hormones in the reproductive axis and includes a reconfiguration in the network connectivity and this change is associated but is not a consequence of the GnRH stimulus (141, 143). Such connectivity is principally constructed through soma-soma contact and cytoplasmic extensions in which gap junction-mediated intercellular communication or cytosolic connection are involved. However, if the gonadotroph network is wired only by cell-cell contact, one could expect that cells in close proximity to each other, should present a similar pattern of response and form clusters where degrees of synchronization may be highest (19, 47). Interestingly, the analysis of clusters by correlation in the patterns of calcium mobilization shows that distant cells respond to GnRH similarly, but cells in contact or surrounded by capillaries may exhibit different responses, indicating that other mechanisms such as paracrine communication or connectivity with FS cells play an important role in the network function (64, 148, 149) (**Figure 4**). In addition, the functional and structural connection in the gonadotroph network is plastic and remodels throughout the estrous cycle, probably following the physiological demand during the proestrus of an LH surge (143). The variety of molecules and processes involved in the network formation, their relative contribution, as well as their degree of plasticity requires further investigation.

## Heterotypic Interactions

The homotypic network organization of gonadotrophs and lactotrophs discussed before provide evidence for its physiological relevance and plasticity. GH-secreting cells are the third population in pituitary that are organized in a homotypic network. This network is reconfigured during puberty with a sexually dimorphic pattern, probably as a result of different and dynamic feedback exerted by gonadal steroids (8, 150, 151). To date, the homotypic GH-cell network is the most extensively studied population in the anterior pituitary. There is evidence demonstrating the intimate interaction of the somatotroph network with other cellular types forming exquisitely arranged heterotypic networks, and when these interactions are disrupted, there are important repercussions at the gland-level hormone secretion (150, 152–155). For example, the ablation of somatotrophs at different degrees, compromises the pituitary production of TSH, PRL, LH, and ACTH, resulting in a modified ultrastructural composition of the cells and pituitary hypoplasia (152, 156, 157). Importantly, GH-cell ablation also impairs cell-cell communication between this homotypic network and other cell types, instead of causing GH reduction. The mechanisms of buffering the complete loss of somatotrophs in this model suggests an overproduction of the lineage from progenitor cells and a reduction in the generation of other lineages (152, 156, 157). Other possibilities could reflect transdifferentiation from different cell types since lactotrophs have been reported to decrease in number in these conditions, and it may also reflect the plastic function of multiresponsive



**FIGURE 4** | Reconfiguration of the gonadotroph network throughout the reproductive cycle. **(A)** Intracellular calcium recordings in *ex vivo* preparations of mouse pituitary loaded with the calcium sensor Fluo 4-AM (green) and rhodamine for vasculature (red). Monitoring of calcium activity was performed on the ventral side of the gland under an epifluorescence stereomicroscope and continuously perfused with Ringer's solution [methods as in (146)]. Color traces show the intracellular calcium activity of individual gonadotrophs in response to 10mM GnRH, indicating clusters of gonadotrophs surrounded by vasculature with a similar (pink and yellow boxes), or heterogenous (blue box) calcium response. **(B)** Heatmap and network organization of basal calcium activity of correlated pairs of gonadotrophs from a male mouse. Gonadotrophs stimulated with GnRH in male mouse **(C)**, female mouse in diestrus **(D)**, and proestrus **(E)** showed an increase in synchronized calcium activity compared to basal activity. Notice that the overall correlation values are similar between diestrus and proestrus, as well as their connectivity. Bars in the heatmaps indicate sorting of gonadotrophs by correlation but not by spatial proximity, while groups of gonadotrophs surrounded by vasculature are represented with the same color in bars. Network maps were plotted using qgraph and significant pairs (edges) of correlated cells (nodes) are shown ( $P < 0.05$ ).

cells that have been observed to increase under several physiological challenges (158, 159). However, the mechanisms involved in this process remain to be explored.

Less understood is the ontogeny of the homotypic and heterotypic networks in the pituitary gland. The observation of the complex arrangement of networks leads to the question regarding the developmental origin of cellular interactions and two options are possible. First, it is possible that at the point of cell proliferation of one lineage, cells never lose the contacts (such as cytonemas) and form a continuum allowing high synchronicity of activity and identity prevalence of the homotypic networks, as observed in the first stages of spermatogenesis (160). Alternatively, the cells might appear isolated, establish a position in the gland, and after that, the lineages could acquire the connectivity observed in adulthood. The available literature revealed that the second option is observed during development in the corticotroph and gonadotroph networks (138). Corticotrophs and gonadotrophs first appear in the ventral side of the pituitary as small cells without cytoplasmic projections, and isolated from other cells of the same lineage. Then, corticotrophs reshape their morphology, penetrate into the adenohypophysis, and interconnect with other corticotrophs and with the vasculature through cytoplasmic projections (138). Gonadotrophs follow the same pattern and it is reasonable to speculate that the organization of other networks in the gland also appears just after terminal differentiation (16, 138).

The sequential differentiation of cell types in the pituitary also suggests that homotypic interactions establish first and the heterotypic networks appear later, although the lineages that are developmentally older have a repercussion over the newborn cells and lineages (156, 161). For example, LH-positive cells first appear in the ventral surface of the pituitary at embryonic day 17.5 when the homotypic network of corticotrophs is already established. LH cells then form a homotypic network using the corticotroph network as a scaffold. In fact, blocking cell differentiation of the POMC lineage leads to an increase in the number of LH-expressing gonadotrophs that are mainly located where corticotrophs should be distributed (138, 162).


One may ask whether the increase of the gonadotroph population after the interruption of corticotroph establishment during development is due to a physical or morphogenetic constraints exerted from corticotrophs to the other populations. The idea that corticotrophs and gonadotrophs probably arise from a common precursor, suggests that the extended gonadotrophs lineage in deficient POMC cell differentiation is more than expansion of the former in an “empty niche,” reflecting a complex feedback between networks. For example, the transcription factor *Tpit*, expressed exclusively by POMC cells, exerts a negative effect over *SF1*, the transcription factor associated with the establishment of the gonadotroph phenotype (162, 163). This and other molecular interactions (e.g., *pit-1* blocking the expression of the gonadotroph-like phenotype in thyrotrophs throughout *GATA-2*), suggests that a molecular interplay exists between cell lineages and results in an equilibrium of cell populations (16,

138, 164). The dynamic equilibrium achieved during ontogeny could persist to adulthood and shift to different points of equilibrium in a context-dependent manner, with the possibility of return to previous states, revealing the substrate of a highly plastic gland with cell populations that are able to navigate toward alternative phenotypes [see for example *Sox2*-expressing cells in adult organisms (156, 159, 165, 166)].

To date, other heterotypic interactions between the five mammalian adenohypophyseal populations have not been investigated but there are reasons to speculate about their relation and interdependence. First, the interlocked cell arrangement of gonadotrophs and corticotrophs was uncovered recently only by 3D image analysis that is hidden under classical 2D microscopy analysis (138). In this sense, thyrotrophs, the less abundant secretory population, may reveal a 3D continuous distribution in the pituitary [but see (130)]. Second, under diverse physiological challenges, transdifferentiation between cell types occurs independently of their ontogenetic proximity. For these reasons, the proportion of multi-responsive and multi-hormonal cells are constantly fluctuating, and different phenotypes such as lactosomatotrophs, somato-gonadotrophs, gonado-lactotrophs have been observed (144). Finally, as we have mentioned before in this work, paracrine communication may play an important role in heterotypic communication, although its full implications are not yet clear (64).

## CONCLUSIONS

Despite variations in the hypothalamus-pituitary communication and the organization of the cell populations within the gland, pituitary cell networks have persisted throughout evolution, revealing their pivotal role in the ability of the pituitary to mount effective responses to hypothalamic and systemic stimuli. Two major elements allow the formation of networks in the pituitary gland: cell-cell contacts and diffusible factors, both present in vertebrate species, although with different preponderance. For example, in fish, where cell types are present in clusters and hypothalamic innervation invades the pituitary gland (see **Figure 5**), the secreted factors by these neurons near the vasculature reach their pituitary targets by diffusion. However, as not all cells in a cluster (see gonadotrophs from **Figure 5**) are in close proximity to neuron terminals and capillaries, a synchronized response is also achieved by gap junctions. Furthermore, the relevance of the vascular systems may reside in the delivery of nutrient and oxygen supply as well transport of pituitary outputs to the blood circulation, meanwhile paracrine and FS cells could produce a balance between distinct secretory cells and neuroendocrine axes. In contrast, in mammals, where the endocrine cells are distributed along the gland and the hypothalamic neurons secrete to the vasculature in the median eminence, a smaller proportion of the network could receive hypothalamic factors by diffusion (**Figure 5**). It seems that homo and heterotypic communication allow coordinated propagation of signals



**FIGURE 5** | A multimodal communication between the hypothalamus and pituitary cell networks. Shown are sagittal sections (20  $\mu$ m) of 4% paraformaldehyde fixed tissue, anterior at the left, dorsal to the top. Immunocytochemistry for gonadotrophs containing the LH $\beta$  subunit (green) and GnRH neurons (gray), species as in **Figure 3**. The distribution of gonadotrophs in the pars distalis, proximal to the hypothalamus, have remarkable cell-cell contacts in the tetrapod species, in teleost these cells are exclusively found in the rostral pars distalis forming a clear-cut cluster. The gonadotroph network arrangement is highly influenced by the vascular system (red; lectin-rhodamine), since these cells have cytoplasmic extensions to the capillaries. In tetrapods, the GnRH-producing neurons extend their projections to the median eminence where GnRH is secreted to the capillaries. In contrast, the GnRH neurons in teleost unfold throughout the pituitary and establish at the boundaries of the gonadotroph patch, following the vascular system. Note the close proximity between gonadotrophs and GnRH at the median eminence of tetrapods. DAPI is shown in blue for counterstain. Scale bars represent 200  $\mu$ m.

between distant cell subpopulations, that may be enhanced through FS cells, paracrine factors and gap junction in small spaces (100  $\mu\text{m}$ ). Moreover, in other cell types that are less proximal to the median eminence (e.g., thyrotrophs), probably the hypothalamic signals are initially delivered through the vasculature and propagated by FS afterward. In summary, plasticity and evolution of the pituitary gland is the tinkering of existing parts combined in new ways rather than the creation of new parts.

In all vertebrate species, different physiological changes such as development, metamorphosis, reproductive cycles, stress, and metabolism require different pituitary outputs, not limited to large changes in the amount of secreted hormones, but also in timing and pulse frequency. Therefore, the gland and its interactions with the hypothalamus, organs, and tissues of the body are constantly being remodeled. The integration of all signals and the generation of hormone pulses, is the result of the plasticity of pituitary cell homotypic and heterotypic networks. Therefore, the interaction between these endocrine networks, are an important driver of the organism homeostasis.

The plasticity of hormone secretion in the pituitary gland has largely been seen as the result of changes in the hypothalamic input and pituitary-intrinsic modulations such as cell proliferation, cell size modifications, transdifferentiation, and multi-responsiveness. However, this review and other authors have contributed to the discussion of the elements that support the complexity of hypophysial plasticity. Several elements are involved to achieve this plasticity and have been discussed in detail (16, 20), these are: a) *gain of function*, by integration of stimuli from neuroendocrine axis and systemic information in the interaction of axes, as GH and PRL-networks, whose coordinated response to GHRH and TRH, respectively, is not seen when cells are dispersed; b) *plasticity for hormone secretion*, during changes in physiological demand, as seen in puberty when the GH axis is highly active and there is an increase in male somatotroph clustering associated with large amplitude of GH pulses; c) *experience-dependent response*, that is improved as a consequence of repeat demand, as mentioned in lactotrophs during the process of lactation; d) *network support of transcriptional synchronization*, when gene transcription follows a rhythmic pattern at the population level that is lost in dispersed cells, as it happens in lactotrophs; e) *redundancy through modularity*, although all the cells in a homotypic network are connected, there are subgroups of cells connected more strongly than other, this fact promotes certain independence of activity, and perturbations in a given region not necessarily affecting the whole network given the possibility of local homeostasis.

It is noteworthy that, despite their enormous functional importance, the study of pituitary networks is still in its infancy. In the last years there has been more literature focusing on the study of the pituitary as a system with functional networks. The term “network(s)” associated with the pituitary gland, as a main topic, first appeared in scientific literature in 1990, with the study of avian adenohypophysis and the effect of a serotonin precursor on thyrotrophs and the

follicle-stellate network (167). In the next years, the concept of pituitary networks became popular and just in the last 10 years, 890 publications have been reported using the term network as a topic, according to The Web of Science Core Collection. Research of pituitary networks refers to different levels of interaction including protein-protein, protein-DNA, protein-metabolite, and cell homotypic and heterotypic networks (20). Accordingly, this young research field is highly productive and growing rapidly, showing the relevance of the cellular interactions when studying plasticity of the gland under diverse physiological demand.

So far research has focused mainly on the mammalian model of mice and, more recently, on fish. Research in both of these organisms has relied heavily on transgenic animal models that allow the identification, monitoring and manipulation of specific cell populations in live tissue as well as 3D imaging of the cell networks. Extending the study of functional pituitary networks to other vertebrate classes will be greatly assisted by the generation of transgenic animals in these taxa.

The development of genetically encoded calcium indicators (168) and *in vivo* imaging techniques (169) are expected to play an important role in advancing our understanding of the function and plasticity of pituitary cell networks in the coming years. These methods will allow researchers to extend the study of these networks from slices to freely moving and behaving animals and will therefore significantly increase the physiological relevance of our insights. Furthermore, single-cell transcriptomics (5, 170) and precise genome editing through viral delivery (171) will allow researchers to reveal the molecular machinery that drives the function and plasticity of pituitary cell networks, the relative role played by the elements that establish these networks and the relevance of the multimodal communication in the hypothalamus-pituitary system.

## AUTHOR CONTRIBUTIONS

TF and MG: Conception of the line of work, structure of ideas, and writing. YS-A: Writing, immunocytochemistry, calcium recordings, and figures. All authors contributed to the article and approved the submitted version.

## FUNDING

Authors were supported by grants from CONACyT and the Agence Nationale de la Recherche (273513, Repro-Net) and PAPIIT (IN222613 and IN227416). YS-A was supported by CONACyT master funding.

## ACKNOWLEDGMENTS

We would like to thank Edgar Jiménez-Díaz and the LANSBIODYT-UNICUA-UNAM (CONACYT) for equipment

and the technical support in the acquisition of confocal images. The amphibian and lizard for immunohistochemistry shown in this manuscript were kindly provided by Sebastian Zúñiga-Lagunes and Oscar Flores-Villela, the authors are deeply grateful. We would like

to thank Berta-Levavi Sivan for her permission to use the zebrafish pituitary images. YS-A would like to thank the Posgrado en Ciencias Biológicas (UNAM) for the academic formation through its Master's degree program in Biological Sciences.

## REFERENCES

- West-Eberhard MJ. Developmental plasticity and the origin of species differences. *Proc Natl Acad Sci USA* (2005) 102 Suppl 1:6543–9. doi: 10.1073/pnas.0501844102
- Le Tissier P, Campos P, Lafont C, Romano N, Hodson DJ, Mollard P. An updated view of hypothalamic-vascular-pituitary unit function and plasticity. *Nat Rev Endocrinol* (2017) 13:257–67. doi: 10.1038/nrendo.2016.193
- Schlosser G. From so simple a beginning – what amphioxus can teach us about placode evolution. *Int J Dev Biol* (2017) 61:633–48. doi: 10.1387/ijdb.170127gs
- Rathke H. Ueber die Entstehung der Glandula pituitaria. *Arch Anat Physiol Sci Med* (1838) 40:482–5.
- Fabian P, Tseng K-C, Smeeton J, Lancman JJ, Duc Si Dong P, Cerny R, et al. Lineage analysis reveals an endodermal contribution to the vertebrate pituitary. *Science* (2020) 370:463–7. doi: 10.1126/science.aba4767
- Melmed S. *The Pituitary*. China: Elsevier Academic Press (2010). Available at: <https://play.google.com/store/books/details?id=OFFW68kMpFIC>.
- Harris GW. Neural Control Of The Pituitary Gland. *Physiol Rev* (1948) 28:139–79. doi: 10.1152/physrev.1948.28.2.139
- Bonnefont X, Lacampagne A, Sanchez-Hormigo A, Fino E, Creff A, Mathieu M-N, et al. Revealing the large-scale network organization of growth hormone-secreting cells. *Proc Natl Acad Sci USA* (2005) 102:16880–5. doi: 10.1073/pnas.0508202102
- Painson JC, Tannenbaum GS. Sexual dimorphism of somatostatin and growth hormone-releasing factor signaling in the control of pulsatile growth hormone secretion in the rat. *Endocrinology* (1991) 128:2858–66. doi: 10.1210/endo-128-6-2858
- Sartor O, Bowers CY, Chang D. Parallel studies of His-DTrp-Ala-Trp-DPhe-Lys-NH<sub>2</sub> and human pancreatic growth hormone-releasing factor-44-NH<sub>2</sub> in rat primary pituitary cell monolayer culture. *Endocrinology* (1985) 116:952–7. doi: 10.1210/endo-116-3-952
- Wen S, Schwarz JR, Niculescu D, Dinu C, Bauer CK, Hirdes W, et al. Functional characterization of genetically labeled gonadotropes. *Endocrinology* (2008) 149:2701–11. doi: 10.1210/en.2007-1502
- Navratil AM, Knoll JG, Whitesell JD, Tobet SA, Clay CM. Neuroendocrine plasticity in the anterior pituitary: gonadotropin-releasing hormone-mediated movement in vitro and in vivo. *Endocrinology* (2007) 148:1736–44. doi: 10.1210/en.2006-1153
- Campos P, Herbison AE. Optogenetic activation of GnRH neurons reveals minimal requirements for pulsatile luteinizing hormone secretion. *Proc Natl Acad Sci USA* (2014) 111:18387–92. doi: 10.1073/pnas.1415226112
- Levine JE, Pau KY, Ramirez VD, Jackson GL. Simultaneous measurement of luteinizing hormone-releasing hormone and luteinizing hormone release in unanesthetized, ovariectomized sheep. *Endocrinology* (1982) 111:1449–55. doi: 10.1210/endo-111-5-1449
- Romano N, McClafferty H, Walker JJ, Le Tissier P, Shipston MJ. Heterogeneity of Calcium Responses to Secretagogues in Corticotrophs From Male Rats. *Endocrinology* (2017) 158:1849–58. doi: 10.1210/en.2017-00107
- Mollard P, Hodson DJ, Lafont C, Rizzotti K, Drouin J. A tridimensional view of pituitary development and function. *Trends Endocrinol Metab* (2012) 23:261–9. doi: 10.1016/j.tem.2012.02.004
- Walker JJ, Terry JR, Lightman SL. Origin of ultradian pulsatility in the hypothalamic–pituitary–adrenal axis. *Proc R Soc B: Biol Sci* (2010) 277:1627–33. doi: 10.1098/rspb.2009.2148
- Le Tissier PR, Hodson DJ, Martin AO, Romano N, Mollard P. Plasticity of the prolactin (PRL) axis: mechanisms underlying regulation of output in female mice. *Adv Exp Med Biol* (2015) 846:139–62. doi: 10.1007/978-3-319-12114-7\_6
- Hodson DJ, Schaeffer M, Romano N, Fontanaud P, Lafont C, Birkenstock J, et al. Existence of long-lasting experience-dependent plasticity in endocrine cell networks. *Nat Commun* (2012) 3:605. doi: 10.1038/ncomms1612
- Le Tissier PR, Hodson DJ, Lafont C, Fontanaud P, Schaeffer M, Mollard P. Anterior pituitary cell networks. *Front Neuroendocrinol* (2012) 33:252–66. doi: 10.1016/j.yfrne.2012.08.002
- Fontaine R, Ciani E, Haug TM, Hodne K, Ager-Wick E, Baker DM, et al. Gonadotrope plasticity at cellular, population and structural levels: A comparison between fishes and mammals. *Gen Comp Endocrinol* (2020) 287:113344. doi: 10.1016/j.ygcen.2019.113344
- Golan M, Martin AO, Mollard P, Levavi-Sivan B. Anatomical and functional gonadotrope networks in the teleost pituitary. *Sci Rep* (2016) 6:23777. doi: 10.1038/srep23777
- Green JD. The comparative anatomy of the hypophysis, with special reference to its blood supply and innervation. *Am J Anat* (1951) 88:225–311. doi: 10.1002/aja.1000880204
- Holmes RL, Ball JN. *The Pituitary Gland: A Comparative Account*. Cambridge, Great Britain: Cambridge University Press: CUP Archive (1974). Available at: [https://books.google.com/books/about/The\\_Pituitary\\_Gland.html?hl=&id=UZs8AAAIAAJ](https://books.google.com/books/about/The_Pituitary_Gland.html?hl=&id=UZs8AAAIAAJ).
- Abraham M. The ultrastructure of the cell types and of the neurosecretory innervation in the pituitary of Mugil cephalus L. from fresh water, the sea, and a hypersaline lagoon. *Gen Comp Endocrinol* (1971) 17:334–50. doi: 10.1016/0016-6480(71)90144-4
- Abraham M. Ultrastructure of the cell types and of the neurosecretory innervation in the pituitary of Mugil cephalus L. from freshwater, the sea, and a hypersaline lagoon II. The proximal pars distalis. *Gen Comp Endocrinol* (1974) 24:121–32. doi: 10.1016/0016-6480(74)90164-6
- Abraham M, Dinari-Lavie V, Lotan R. The pituitary of Aphanis dispar (Rüppell) from hypersaline marshes and freshwater II. Ultrastructure of the rostral pars distalis. *Cell Tissue Res* (1977) 179(3):317–30. doi: 10.1007/bf00221103
- Kah O, Dubourg P, Onteniente B, Geffard M, Calas A. The dopaminergic innervation of the goldfish pituitary. An immunocytochemical study at the electron-microscope level using antibodies against dopamine. *Cell Tissue Res* (1986) 244:577–82. doi: 10.1007/BF00212536
- Batten TF, Ball JN. Ultrastructure of the neurohypophysis of the teleost *Poecilia latipinna* in relation to neural control of the adenohypophysial cells. *Cell Tissue Res* (1977) 185:409–33. doi: 10.1007/BF00220300
- Batten T, Ball JN, Benjamin M. Ultrastructure of the adenohypophysis in the teleost *Poecilia latipinna*. *Cell Tissue Res* (1975) 161:239–61. doi: 10.1007/bf00220372
- Batten TF. Ultrastructural characterization of neurosecretory fibres immunoreactive for vasotocin, isotocin, somatostatin, LHRH and CRF in the pituitary of a teleost fish, *Poecilia latipinna*. *Cell Tissue Res* (1986) 244:661–72. doi: 10.1007/BF00212547
- Golan M, Zelinger E, Zohar Y, Levavi-Sivan B. Architecture of GnRH-Gonadotrope-Vasculature Reveals a Dual Mode of Gonadotropin Regulation in Fish. *Endocrinology* (2015) 156:4163–73. doi: 10.1210/en.2015-1150
- Abraham M, Kieselstein M, Lisson-Begon S. The extravascular channel system in the rostral pituitary of Mugil cephalus (Teleostei) as revealed by use of horseradish peroxidase. *Cell Tissue Res* (1976) 167:289–96. doi: 10.1007/BF00219143
- Abraham M. Ultrastructure of the cell types and of the neurosecretory innervation in the pituitary of Mugil cephalus L. from freshwater, the sea, and a hypersaline lagoon. *Gen Comp Endocrinol* (1976) 29:511–21. doi: 10.1016/0016-6480(76)90035-6
- Hodne K, Fontaine R, Ager-Wick E, Weltzien F-A. GnRH1-Induced Responses Are Indirect in Female Medaka Fish Cells, Generated Through Cellular Networks. *Endocrinology* (2019) 160:3018–32. doi: 10.1210/en.2019-00595
- Shainer I, Buchshtab A, Hawkins TA, Wilson SW, Cone RD, Gothilf Y. Novel hypophysiotropic AgRP2 neurons and pineal cells revealed by BAC transgenesis in zebrafish. *Sci Rep* (2017) 7:44777. doi: 10.1038/srep44777

37. Trudeau VL, Somoza GM. Multimodal hypothalamo-hypophysial communication in the vertebrates. *Gen Comp Endocrinol* (2020) 293:113475. doi: 10.1016/j.ygcen.2020.113475
38. Levavi-Sivan B, Bloch CL, Gutnick MJ, Fleiderovich IA. Electrotonic coupling in the anterior pituitary of a teleost fish. *Endocrinology* (2005) 146:1048–52. doi: 10.1210/en.2004-1415
39. Golan M, Hollander-Cohen L, Levavi-Sivan B. Stellate Cell Networks in the Teleost Pituitary. *Sci Rep* (2016) 6:24426. doi: 10.1038/srep24426
40. Morand I, Fonlupt P, Guerrier A, Trouillas J, Calle A, Remy C, et al. Cell-to-cell communication in the anterior pituitary: evidence for gap junction-mediated exchanges between endocrine cells and folliculostellate cells. *Endocrinology* (1996) 137:3356–67. doi: 10.1210/endo.137.8.8754762
41. Fletcher WH, Anderson NC, Everett JW. Intercellular communication in the rat anterior pituitary gland: an in vivo and in vitro study. *J Cell Biol* (1975) 67:469–76. doi: 10.1083/jcb.67.2.469
42. Abraham M, Sandri C, Akert K. Freeze-etch study of the teleostean pituitary. *Cell Tissue Res* (1979) 199:397–407. doi: 10.1007/BF00236078
43. Herbert DC. Intercellular junctions in the rhesus monkey pars distalis. *Anat Rec* (1979) 195:1–6. doi: 10.1002/ar.1091950101
44. Wilfinger WW, Larsen WJ, Downs TR, Wilbur DL. An in vitro model for studies of intercellular communication in cultured rat anterior pituitary cells. *Tissue Cell* (1984) 16:483–97. doi: 10.1016/0040-8166(84)90026-0
45. Soji T, Herbert DC. Intercellular communication between rat anterior pituitary cells. *Anatomical Record* (1989) 224:523–33. doi: 10.1002/ar.1092240410
46. Soji T, Herbert DC. Intercellular communication within the rat anterior pituitary Gland. II. castration effects and changes after injection of luteinizing hormone-releasing hormone (LH-RH) or testosterone. *Anatomical Record* (1990) 226:342–6. doi: 10.1002/ar.1092260311
47. Göngrich C, García-González D, Le Magueresse C, Roth LC, Watanabe Y, Burks DJ, et al. Electrotonic Coupling in the Pituitary Supports the Hypothalamic-Pituitary-Gonadal Axis in a Sex Specific Manner. *Front Mol Neurosci* (2016) 9(65):1–12. doi: 10.3389/fnmol.2016.00065
48. Fauquier T, Guérineau NC, McKinney RA, Bauer K, Mollard P. Folliculostellate cell network: a route for long-distance communication in the anterior pituitary. *Proc Natl Acad Sci USA* (2001) 98:8891–6. doi: 10.1073/pnas.151339598
49. Guérineau NC, Bonnefont X, Stoeckel L, Mollard P. Synchronized spontaneous Ca<sup>2+</sup> transients in acute anterior pituitary slices. *J Biol Chem* (1998) 273:10389–95. doi: 10.1074/jbc.273.17.10389
50. Meda P, Pepper MS, Traub O, Willecke K, Gros D, Beyer E, et al. Differential expression of gap junction connexins in endocrine and exocrine glands. *Endocrinology* (1993) 133:2371–8. doi: 10.1210/endo.133.5.8404689
51. Soji T, Nishizono H, Yashiro T, Herbert DC. Intercellular communication within the rat anterior pituitary gland. III. Postnatal development and periodic changes of cell-to-cell communications in female rats. *Anat Rec* (1991) 231:351–7. doi: 10.1002/ar.1092310309
52. Soji T, Shirasawa N, Kurono C, Yashiro T, Herbert DC. Immunohistochemical study of the post-natal development of the folliculo-stellate cells in the rat anterior pituitary gland. *Tissue Cell* (1994) 26:1–8. doi: 10.1016/0040-8166(94)90078-7
53. Soji T, Mabuchi Y, Kurono C, Herbert DC. Folliculo-stellate cells and intercellular communication within the rat anterior pituitary gland. *Microsc Res Tech* (1997) 39:138–49. doi: 10.1002/(SICI)1097-0029(19971015)39:2<138::AID-JEMT5>3.0.CO;2-H
54. Soji T, Yashiro T, Herbert DC. Intercellular communication within the rat anterior pituitary gland: IV. Changes in cell-to-cell communications during pregnancy. *Anat Rec* (1992) 233:97–102. doi: 10.1002/ar.1092330113
55. Nishizono H, Soji T, Herbert DC. Intercellular communication within the rat anterior pituitary gland: V. Changes in cell-to-cell communications as a function of the timing of castration in male rats. *Anat Rec* (1993) 235:577–82. doi: 10.1002/ar.1092350410
56. Kurono C. Intercellular communication within the rat anterior pituitary gland: VI. Development of gap junctions between folliculo-stellate cells under the influence of ovariectomy and sex steroids in the female rat. *Anat Rec* (1996) 244:366–73. doi: 10.1002/(SICI)1097-0185(199603)244:3<366::AID-AR8>3.0.CO;2-W
57. Soji T, Yashiro T, Herbert DC. Intercellular communication within the rat anterior pituitary Gland. I. Postnatal development and changes after injection of luteinizing hormone-releasing hormone (LH-RH) or testosterone. *Anatomical Record* (1990) 226:337–41. doi: 10.1002/ar.1092260310
58. Sakuma E, Soji T, Herbert DC. Effects of hydrocortisone on the formation of gap junctions and the abnormal growth of cilia within the rat anterior pituitary gland: possible role of gap junctions on the regulation of cell development. *Anat Rec* (2001) 262:169–75. doi: 10.1002/1097-0185(20010201)262:2<169::AID-AR1026>3.0.CO;2-Z
59. Wada I, Sakuma E, Shirasawa N, Wakabayashi K, Otsuka T, Hattori K, et al. Intercellular communications within the rat anterior pituitary. XVI: postnatal changes of distribution of S-100 protein positive cells, connexin 43 and LH-RH positive sites in the pars tuberalis of the rat pituitary gland. An immunohistochemical and electron microscopic study. *Tissue Cell* (2014) 46:33–9. doi: 10.1016/j.tice.2013.10.001
60. Borst JGG, Lodder JC, Roubos EW, Kits KS. In situ recordings of presumed folliculo-stellate cells in the intermediate lobe of the pituitary gland of *Xenopus laevis*. *Neurosci Lett* (1996) 209:61–4. doi: 10.1016/0304-3940(96)12602-1
61. Allaerts W, Vankelecom H. History and perspectives of pituitary folliculo-stellate cell research. *Eur J Endocrinol* (2005) 153:1–12. doi: 10.1530/eje.1.01949
62. Nishimura S, Yamashita M, Kaneko T, Kawabata F, Tabata S. Cytokeratin-positive folliculo-stellate cells in chicken adenohypophysis. *Anim Sci J* (2017) 88:1835–41. doi: 10.1111/asj.12866
63. Gracia-Navarro F, Gonzalez-Reyes JA, Guerrero-Callejas F, Garcia-Herdugo G. An electron microscopic study of stellate cells and cavities in the pars distalis of frog pituitary. *Tissue Cell* (1983) 15:729–36. doi: 10.1016/0040-8166(83)90046-0
64. Deneff C. Paracrinicity: the story of 30 years of cellular pituitary crosstalk. *J Neuroendocrinol* (2008) 20:1–70. doi: 10.1111/j.1365-2826.2007.01616.x
65. Sato Y, Hashitani H, Shirasawa N, Sakuma E, Naito A, Suzuki H, et al. Intercellular communications within the rat anterior pituitary XII: immunohistochemical and physiological evidences for the gap junctional coupling of the folliculo-stellate cells in the rat anterior pituitary. *Tissue Cell* (2005) 37:281–91. doi: 10.1016/j.tice.2005.03.005
66. Shirasawa N, Sakuma E, Wada I, Naito A, Horiuchi O, Mabuchi Y, et al. Intercellular Communication Within the Rat Anterior Pituitary: XIV Electron Microscopic and Immunohistochemical Study on the Relationship between the Agranular Cells and GnRH Neurons in the Dorsal Pars Tuberalis of the Pituitary Gland. *Anatomical Record: Adv Integr Anat Evol Biol* (2007) 290:1388–98. doi: 10.1002/ar.20596
67. Shirasawa N, Kihara H, Yamaguchi S, Yoshimura F. Pituitary folliculo-stellate cells immunostained with S-100 protein antiserum in postnatal, castrated and thyroidectomized rats. *Cell Tissue Res* (1983) 231:235–49. doi: 10.1007/BF00222177
68. Foster WG, Younglai EV. An immunohistochemical study of the GnRH neuron morphology and topography in the adult female rabbit hypothalamus. *Am J Anat* (1991) 191:293–300. doi: 10.1002/aja.1001910308
69. Hattori K, Shirasawa N, Suzuki H, Otsuka T, Wada I, Yashiro T, et al. Intercellular Communication within the Rat Anterior Pituitary Gland. XV. Properties of Spontaneous and LHRH-Induced Ca<sup>2+</sup> Transients in the Transitional Zone of the Rat Anterior Pituitary in Situ. *Endocrinology* (2013) 154:400–9. doi: 10.1210/en.2012-1501
70. Shirasawa N, Mabuchi Y, Sakuma E, Horiuchi O, Yashiro T, Kikuchi M, et al. Intercellular communication within the rat anterior pituitary gland: X. Immunohistochemistry of S-100 and connexin 43 of folliculo-stellate cells in the rat anterior pituitary gland. *Anat Rec A Discovery Mol Cell Evol Biol* (2004) 278:462–73. doi: 10.1002/ar.a.20040
71. Mabuchi Y, Shirasawa N, Sakuma E, Hashimoto Y, Kuno M, Coombs RJ, et al. Intercellular communication within the rat anterior pituitary: relationship between LH-RH neurons and folliculo-stellate cells in the pars tuberalis. *Cell Tissue Res* (2004) 317:79–90. doi: 10.1007/s00441-003-0851-9
72. Soji T, Taya K, Igarashi M, Yoshimura F. Acute and subacute effect of LH-RH upon LH- and FSH-gonadotrophs in castrated female rats with short-term estrogen-progesterone pretreatment. *Endocrinol Jpn* (1974) 21:407–28. doi: 10.1507/endocrj1954.21.407
73. Naik DV. Immunoreactive LH-RH neurons in the hypothalamus identified by light and fluorescent microscopy. *Cell Tissue Res* (1975) 157:423–36. doi: 10.1007/BF00222597

74. Hodson DJ, Romanò N, Schaeffer M, Fontanaud P, Lafont C, Fiordelisio T, et al. Coordination of calcium signals by pituitary endocrine cells in situ. *Cell Calcium* (2012) 51:222–30. doi: 10.1016/j.ceca.2011.11.007
75. De Paul AL, Bonaterra M, Aoki A, Torres AI. Cellular and functional interactions between gonadotrophs and lactotrophs in pituitary cell cultures. *Med Electron Microsc* (2000) 33:231–40. doi: 10.1007/s007950000023
76. Robberecht W, Andries M, Denef C. Angiotensin II Is Retained in Gonadotrophs of Pituitary Cell Aggregates Cultured in Serum-Free Medium but Does Not Mimic the Effects of Exogenous Angiotensins and Luteinizing-Hormone-Releasing Hormone on Growth Hormone Release. *Neuroendocrinology* (1992) 56:550–60. doi: 10.1159/000126273
77. McNeilly AS. Lactational control of reproduction. *Reprod Fertil Dev* (2001) 13:583–90. doi: 10.1071/rd01056
78. Hefco E, Krulich L, Illner P, Larsen PR. Effect of acute exposure to cold on the activity of the hypothalamic-pituitary-thyroid system. *Endocrinology* (1975) 97:1185–95. doi: 10.1210/endo-97-5-1185
79. Zhou H, Wang X, Ko KW, Wong AOL. Evidence for a novel intrapituitary autocrine/paracrine feedback loop regulating growth hormone synthesis and secretion in grass carp pituitary cells by functional interactions between gonadotrophs and somatotrophs. *Endocrinology* (2004) 145:5548–59. doi: 10.1210/en.2004-0362
80. Lin S-W, Ge W. Differential regulation of gonadotropins (FSH and LH) and growth hormone (GH) by neuroendocrine, endocrine, and paracrine factors in the zebrafish—an in vitro approach. *Gen Comp Endocrinol* (2009) 160:183–93. doi: 10.1016/j.ygcen.2008.11.020
81. Yuen C-W, Ge W. Follistatin suppresses FSH $\beta$  but increases LH $\beta$  expression in the goldfish - evidence for an activin-mediated autocrine/paracrine system in fish pituitary. *Gen Comp Endocrinol* (2004) 135:108–15. doi: 10.1016/j.ygcen.2003.08.012
82. Aroua S, Maugars G, Jeng S-R, Chang C-F, Weltzien F-A, Rousseau K, et al. Pituitary gonadotropins FSH and LH are oppositely regulated by the activin/follistatin system in a basal teleost, the eel. *Gen Comp Endocrinol* (2012) 175:82–91. doi: 10.1016/j.ygcen.2011.10.002
83. Yaron Z, Gur G, Melamed P, Rosenfeld H, Levavi-Sivan B, Elizur A. Regulation of gonadotropin subunit genes in tilapia. *Comp Biochem Physiol B Biochem Mol Biol* (2001) 129:489–502. doi: 10.1016/s1096-4959(01)00345-1
84. Fung RSK, Bai J, Yuen KWY, Wong AOL. Activin/follistatin system in grass carp pituitary cells: - Regulation by local release of growth hormone and luteinizing hormone and its functional role in growth hormone synthesis and secretion. *PLoS One* (2017) 12:e0179789. doi: 10.1371/journal.pone.0179789
85. Jiang Q, Wong AOL. Signal transduction mechanisms for autocrine/paracrine regulation of somatolactin- $\alpha$  secretion and synthesis in carp pituitary cells by somatolactin- $\alpha$  and - $\beta$ . *Am J Physiol Endocrinol Metab* (2013) 304:E176–86. doi: 10.1152/ajpendo.00455.2012
86. Jiang Q, Ko KW, Wong AOL. Insulin-like growth factor as a novel stimulator for somatolactin secretion and synthesis in carp pituitary cells via activation of MAPK cascades. *Am J Physiol Endocrinol Metab* (2011) 301: E1208–19. doi: 10.1152/ajpendo.00347.2011
87. Jiang Q, He M, Ko KW, Wong AOL. Kisspeptin induction of somatolactin- $\alpha$  release in goldfish pituitary cells: functional role of cAMP/PKA-, PLC/PKC-, and Ca(2+)/calmodulin-dependent cascades. *Am J Physiol Endocrinol Metab* (2014) 307:E872–84. doi: 10.1152/ajpendo.00321.2014
88. Espigares F, Zanuy S, Gómez A. Kiss2 as a Regulator of Lh and Fsh Secretion via Paracrine/Autocrine Signaling in the Teleost Fish European Sea Bass (*Dicentrarchus labrax*). *Biol Reprod* (2015) 93:114. doi: 10.1095/biolreprod.115.131029
89. Biran J, Golan M, Mizrahi N, Ogawa S, Parhar IS, Levavi-Sivan B. LPXRFa, the piscine ortholog of GnIH, and LPXRF receptor positively regulate gonadotropin secretion in Tilapia (*Oreochromis niloticus*). *Endocrinology* (2014) 155:4391–401. doi: 10.1210/en.2013-2047
90. Hu G, He M, Ko KW, Lin C, Wong AOL. Novel pituitary actions of TAC3 gene products in fish model: receptor specificity and signal transduction for prolactin and somatolactin  $\alpha$  regulation by neurokinin B (NKB) and NKB-related peptide in carp pituitary cells. *Endocrinology* (2014) 155:3582–96. doi: 10.1210/en.2014-1105
91. Zhao E, Grey CL, Zhang D, Mennigen JA, Basak A, Chang JP, et al. Secretoneurin is a potential paracrine factor from lactotrophs stimulating gonadotropin release in the goldfish pituitary. *Am J Physiol Regul Integr Comp Physiol* (2010) 299:R1290–7. doi: 10.1152/ajpregu.00407.2010
92. Fontaine R, Ager-Wick E, Hodne K, Weltzien F-A. Plasticity in medaka gonadotropes via cell proliferation and phenotypic conversion. *J Endocrinol* (2020) 245:21–37. doi: 10.1530/JOE-19-0405
93. Le Tissier P, Fiordelisio Coll T, Mollard P. The Processes of Anterior Pituitary Hormone Pulse Generation. *Endocrinology* (2018) 159:3524–35. doi: 10.1210/en.2018-00508
94. Kasper RS, Shved N, Takahashi A, Reinecke M, Eppler E. A systematic immunohistochemical survey of the distribution patterns of GH, prolactin, somatolactin, beta-TSH, beta-FSH, beta-LH, ACTH, and alpha-MSH in the adenohypophysis of *Oreochromis niloticus*, the Nile tilapia. *Cell Tissue Res* (2006) 325:303–13. doi: 10.1007/s00441-005-0119-7
95. Golan M, Biran J, Levavi-Sivan B. A novel model for development, organization, and function of gonadotropes in fish pituitary. *Front Endocrinol* (2014) 5:182. doi: 10.3389/fendo.2014.00182
96. Golan M, Levavi-Sivan B. Social dominance in tilapia is associated with gonadotropin hyperplasia. *Gen Comp Endocrinol* (2013) 192:126–35. doi: 10.1016/j.ygcen.2013.04.032
97. Hildahl J, Sandvik GK, Lifjeld R, Hodne K, Nagahama Y, Haug TM, et al. Developmental tracing of luteinizing hormone  $\beta$ -subunit gene expression using green fluorescent protein transgenic medaka (*Oryzias latipes*) reveals a putative novel developmental function. *Dev Dyn* (2012) 241:1665–77. doi: 10.1002/dvdy.23860
98. Karigo T, Aikawa M, Kondo C, Abe H, Kanda S, Oka Y. Whole brain-pituitary in vitro preparation of the transgenic medaka (*Oryzias latipes*) as a tool for analyzing the differential regulatory mechanisms of LH and FSH release. *Endocrinology* (2014) 155:536–47. doi: 10.1210/en.2013-1642
99. Biran J, Golan M, Mizrahi N, Ogawa S, Parhar IS, Levavi-Sivan B. Direct regulation of gonadotropin release by neurokinin B in tilapia (*Oreochromis niloticus*). *Endocrinology* (2014) 155:4831–42. doi: 10.1210/en.2013-2114
100. Maruska KP, Levavi-Sivan B, Biran J, Fernald RD. Plasticity of the Reproductive Axis Caused by Social Status Change in an African Cichlid Fish: I. Pituitary Gonadotropins. *Endocrinology* (2011) 152:281–90. doi: 10.1210/en.2010-0875
101. Aizen J, Kasuto H, Golan M, Zakay H, Levavi-Sivan B. Tilapia Follicle-Stimulating Hormone (FSH): Immunochemistry, Stimulation by Gonadotropin-Releasing Hormone, and Effect of Biologically Active Recombinant FSH on Steroid Secretion. *Biol Reprod* (2007) 76:692–700. doi: 10.1095/biolreprod.106.055822
102. Grønlien HK, Fontaine R, Hodne K, Tysseng I, Ager-Wick E, Weltzien F-A, et al. Gonadotropes in medaka grow long extensions with varicosity-like swellings, projecting towards each other and blood vessels. *bioRxiv* (2019). doi: 10.1101/777151
103. Strandabø RAU, Hodne K, Ager-Wick E, Sand O, Weltzien F-A, Haug TM. Signal transduction involved in GnRH2-stimulation of identified LH-producing gonadotropes from lhb-GFP transgenic medaka (*Oryzias latipes*). *Mol Cell Endocrinol* (2013) 372:128–39. doi: 10.1016/j.mce.2013.03.022
104. Buchholz DR, Paul BD, Fu L, Shi Y-B. Molecular and developmental analyses of thyroid hormone receptor function in *Xenopus laevis*, the African clawed frog. *Gen Comp Endocrinol* (2006) 145:1–19. doi: 10.1016/j.ygcen.2005.07.009
105. Shi Y-B. *Amphibian Metamorphosis: From Morphology to Molecular Biology*. New York, NY: Wiley-Liss, Inc (1999). Available at: [https://books.google.com/books/about/Amphibian\\_Metamorphosis.html?hl=&id=WjVmQgAACAAJ](https://books.google.com/books/about/Amphibian_Metamorphosis.html?hl=&id=WjVmQgAACAAJ).
106. Denver RJ. Neuroendocrinology of amphibian metamorphosis. *Curr Top Dev Biol* (2013) 103:195–227. doi: 10.1016/B978-0-12-385979-2.00007-1
107. Sachs LM, Buchholz DR. Insufficiency of Thyroid Hormone in Frog Metamorphosis and the Role of Glucocorticoids. *Front Endocrinol* (2019) 10:287. doi: 10.3389/fendo.2019.00287
108. Huang H, Cai L, Remo BF, Brown DD. Timing of metamorphosis and the onset of the negative feedback loop between the thyroid gland and the pituitary is controlled by type II iodothyronine deiodinase in *Xenopus laevis*. *Proc Natl Acad Sci USA* (2001) 98:7348–53. doi: 10.1073/pnas.131198998
109. Buckbinder L, Brown DD. Expression of the *Xenopus laevis* prolactin and thyrotropin genes during metamorphosis. *Proc Natl Acad Sci USA* (1993) 90:3820–4. doi: 10.1073/pnas.90.9.3820
110. Denver RJ. Stress hormones mediate environment-genotype interactions during amphibian development. *Gen Comp Endocrinol* (2009) 164:20–31. doi: 10.1016/j.ygcen.2009.04.016

111. Norris DO. Evolution of Endocrine Regulation of Metamorphosis in Lower Vertebrates. *Am Zoologist* (1983) 23:709–18. doi: 10.1093/icb/23.3.709
112. Jacobs GF, Michielsen RP, Kühn ER. Thyroxine and triiodothyronine in plasma and thyroids of the neotenic and metamorphosed axolotl *Ambystoma mexicanum*: influence of TRH injections. *Gen Comp Endocrinol* (1988) 70:145–51. doi: 10.1016/0016-6480(88)90103-7
113. Denver RJ, Licht P. Neuropeptide stimulation of thyrotropin secretion in the larval bullfrog: Evidence for a common neuroregulator of thyroid and interrenal activity in metamorphosis. *J Exp Zool* (1989) 252:101–4. doi: 10.1002/jez.1402520114
114. Okada R, Yamamoto K, Koda A, Ito Y, Hayashi H, Tanaka S, et al. Development of radioimmunoassay for bullfrog thyroid-stimulating hormone (TSH): effects of hypothalamic releasing hormones on the release of TSH from the pituitary in vitro. *Gen Comp Endocrinol* (2004) 135:42–50. doi: 10.1016/j.ygcen.2003.09.001
115. De Groef B, Van der Geyten S, Darras VM, Kühn ER. Role of corticotropin-releasing hormone as a thyrotropin-releasing factor in non-mammalian vertebrates. *Gen Comp Endocrinol* (2006) 146:62–8. doi: 10.1016/j.ygcen.2005.10.014
116. Denver RJ. Endocrinology of Complex Life Cycles: Amphibians. *Hormones Brain Behav* (2017) 2:145–68. doi: 10.1016/b978-0-12-803592-4.00024-9
117. Krain LP, Denver RJ. Developmental expression and hormonal regulation of glucocorticoid and thyroid hormone receptors during metamorphosis in *Xenopus laevis*. *J Endocrinol* (2004) 181:91–104. doi: 10.1677/joe.0.1810091
118. Kühn ER, De Groef B, Grommen SVH, Van der Geyten S, Darras VM. Low submetamorphic doses of dexamethasone and thyroxine induce complete metamorphosis in the axolotl (*Ambystoma mexicanum*) when injected together. *Gen Comp Endocrinol* (2004) 137:141–7. doi: 10.1016/j.ygcen.2004.03.005
119. Boorse GC, Denver RJ. Acceleration of *Ambystoma tigrinum* metamorphosis by corticotropin-releasing hormone. *J Exp Zool* (2002) 293:94–8. doi: 10.1002/jez.10115
120. Denver RJ. Evolution of the corticotropin-releasing hormone signaling system and its role in stress-induced phenotypic plasticity. *Ann N Y Acad Sci* (1999) 897:46–53. doi: 10.1111/j.1749-6632.1999.tb07877.x
121. Furuta K, Kanki K, Wakahara M. Differentiation behavior of pituitary cells in normal and metamorphosis-arrested larvae of the salamander *Hynobius retardatus*. *Int J Dev Biol* (2001) 45:903–6.
122. Yamaguchi M, Tanaka S, Wakahara M. Immunohisto- and Immunocytochemical Studies on the Dynamics of TSH and GTH Cells in Normally Metamorphosing, Metamorphosed, and Metamorphosis-Arrested *Hynobius retardatus*. *Gen Comp Endocrinol* (1996) 104:273–83. doi: 10.1006/gcen.1996.0172
123. Kanki K, Wakahara M. Spatio-temporal expression of TSH beta and FSH beta genes in normally metamorphosing, metamorphosed, and metamorphosis-arrested *Hynobius retardatus*. *Gen Comp Endocrinol* (2000) 119:276–86. doi: 10.1006/gcen.2000.7502
124. Laudet V. The Origins and Evolution of Vertebrate Metamorphosis. *Curr Biol* (2011) 21:R726–37. doi: 10.1016/j.cub.2011.07.030
125. Wakahara M. Heterochrony and neotenic salamanders: possible clues for understanding the animal development and evolution. *Zool Sci* (1996) 13:765–76. doi: 10.2108/zsj.13.765
126. Childs G. Gonadotropes and Lactotropes. *Knobil Neill's Physiol Reprod* (2006) 1:1483–579. doi: 10.1016/b978-0-12515400-0/50034-8
127. García-Navarro S, Malagón MM, Gracia-Navarro F. Immunohistochemical localization of thyrotropic cells during amphibian morphogenesis: a stereological study. *Gen Comp Endocrinol* (1988) 71:116–23. doi: 10.1016/0016-6480(88)90302-4
128. Stojilkovic SS, Tabak J, Bertram R. Ion Channels and Signaling in the Pituitary Gland. *Endocrine Rev* (2010) 31:845–915. doi: 10.1210/er.2010-0005
129. Baker BI. Biological Role of the Pars Intermedia in Lower Vertebrates. *Ciba Foundation Symposium 81 - Peptides Pars Intermedia* (2008) 81:166–79. doi: 10.1002/9780470720646.ch10
130. Hodson DJ, Legros C, Desarménien MG, Guérineau NC. Roles of connexins and pannexins in (neuro)endocrine physiology. *Cell Mol Life Sci* (2015) 72:2911–28. doi: 10.1007/s00018-015-1967-2
131. Kehl SJ, McBurney RN. The firing patterns of rat melanotrophs recorded using the patch clamp technique. *Neuroscience* (1989) 33:579–86. doi: 10.1016/0306-4522(89)90410-7
132. Tanaka S, Nakakura T, Jansen EJR, Unno K, Okada R, Suzuki M, et al. Angiogenesis in the intermediate lobe of the pituitary gland alters its structure and function. *Gen Comp Endocrinol* (2013) 185:10–8. doi: 10.1016/j.ygcen.2013.01.009
133. Turner HE, Nagy Z, Gatter KC, Esiri MM, Harris AL, Wass JAH. Angiogenesis in Pituitary Adenomas and the Normal Pituitary Gland. *J Clin Endocrinol Metab* (2000) 85:1159–62. doi: 10.1210/jcem.85.3.6485
134. Yip DY, Lofts B. Adenohypophysial cell-types in the pituitary gland of the soft-shelled turtle, *Trionyx sinensis*. I. Seasonal cycles. *Cell Tissue Res* (1976) 170:523–37. doi: 10.1007/BF00361710
135. Doerr-Schott J. Immunohistochemical detection, by light and electron microscopy, of pituitary hormones in cold-blooded vertebrates. *Gen Comp Endocrinol* (1976) 28:513–29. doi: 10.1016/0016-6480(76)90159-3
136. Ávila-Mendoza J, Carranza M, Pérez-Rueda E, Luna M, Arámburo C. Characterization of pituitary growth hormone and its receptor in the green iguana (*Iguana iguana*). *Gen Comp Endocrinol* (2014) 203:281–95. doi: 10.1016/j.ygcen.2014.04.009
137. Saga T, Yamaki K-I. A Comparative Study of the Chronological Appearance of Adenohypophysial Cells in Vertebrates with Emphasis on the Japanese Soft-Shelled Turtle (*Pelodiscus sinensis japonicus*). *BioMed Res* (2006) 17:81–94.
138. Budry L, Lafont C, El Yandouzi T, Chauvet N, Conéjero G, Drouin J, et al. Related pituitary cell lineages develop into interdigitated 3D cell networks. *Proc Natl Acad Sci USA* (2011) 108:12515–20. doi: 10.1073/pnas.1105929108
139. Yip SH, Romanò N, Gustafson P, Hodson DJ, Williams EJ, Kokay IC, et al. Elevated Prolactin during Pregnancy Drives a Phenotypic Switch in Mouse Hypothalamic Dopaminergic Neurons. *Cell Rep* (2019) 26:1787–99.e5. doi: 10.1016/j.celrep.2019.01.067
140. Herbison AE, Porteous R, Pape J-R, Mora JM, Hurst PR. Gonadotropin-releasing hormone neuron requirements for puberty, ovulation, and fertility. *Endocrinology* (2008) 149:597–604. doi: 10.1210/en.2007-1139
141. Edwards BS, Clay CM, Ellsworth BS, Navratil AM. Functional Role of Gonadotrope Plasticity and Network Organization. *Front Endocrinol* (2017) 8:223. doi: 10.3389/fendo.2017.00223
142. Leong DA, Thorner MO. A potential code of luteinizing hormone-releasing hormone-induced calcium ion responses in the regulation of luteinizing hormone secretion among individual gonadotropes. *J Biol Chem* (1991) 266:9016–22.
143. Alim Z, Hartshorn C, Mai O, Stitt I, Clay C, Tobet S, et al. Gonadotrope plasticity at cellular and population levels. *Endocrinology* (2012) 153:4729–39. doi: 10.1210/en.2012-1360
144. Childs GV, MacNicol AM, MacNicol MC. Molecular Mechanisms of Pituitary Cell Plasticity. *Front Endocrinol* (2020) 11:656. doi: 10.3389/fendo.2020.00656
145. Durán-Pastén ML, Fiordelisio T. GnRH-Induced Ca<sup>2+</sup> Signaling Patterns and Gonadotropin Secretion in Pituitary Gonadotrophs. Functional Adaptations to Both Ordinary and Extraordinary Physiological Demands. *Front Endocrinol* (2013) 4(127):1–13. doi: 10.3389/fendo.2013.00127
146. Durán-Pastén ML, Fiordelisio-Coll T, Hernández-Cruz A. Castration-induced modifications of GnRH-elicited [Ca<sup>2+</sup>]<sub>i</sub> signaling patterns in male mouse pituitary gonadotrophs in situ: studies in the acute pituitary slice preparation. *Biol Reprod* (2013) 88:38. doi: 10.1095/biolreprod.112.103812
147. Stojilkovic SS, Bjelobaba I, Zemkova H. Ion Channels of Pituitary Gonadotrophs and Their Roles in Signaling and Secretion. *Front Endocrinol* (2017) 8:126. doi: 10.3389/fendo.2017.00126
148. Bilezikjian LM, Leal AMO, Blount AL, Corrigan AZ, Turnbull AV, Vale WW. Rat Anterior Pituitary Folliculostellate Cells Are Targets of Interleukin-1 $\beta$  and a Major Source of Intrapituitary Follistatin. *Endocrinology* (2003) 144:732–40. doi: 10.1210/en.2002-220703
149. Childs GV, Unabia G, Lee BL, Rougeau D. Heightened secretion by small and medium-sized luteinizing hormone (LH) gonadotropes late in the cycle suggests contributions to the LH surge or possible paracrine interactions. *Endocrinology* (1992) 130:345–52. doi: 10.1210/endo.130.1.1727708
150. Sanchez-Cardenas C, Fontanaud P, He Z, Lafont C, Meunier A-C, Schaeffer M, et al. Pituitary growth hormone network responses are sexually dimorphic and regulated by gonadal steroids in adulthood. *Proc Natl Acad Sci* (2010) 107:21878–83. doi: 10.1073/pnas.1010849107

151. Schaeffer M, Hodson DJ, Meunier A-C, Lafont C, Birkenstock J, Carmignac D, et al. Influence of Estrogens on GH-Cell Network Dynamics in Females: A Live in Situ Imaging Approach. *Endocrinology* (2011) 152:4789–99. doi: 10.1210/en.2011-1430
152. Waite E, Lafont C, Carmignac D, Chauvet N, Coutry N, Christian H, et al. Different degrees of somatotroph ablation compromise pituitary growth hormone cell network structure and other pituitary endocrine cell types. *Endocrinology* (2010) 151:234–43. doi: 10.1210/en.2009-0539
153. Lafont C, Desarménien MG, Cassou M, Molino F, Lecoq J, Hodson D, et al. Cellular in vivo imaging reveals coordinated regulation of pituitary microcirculation and GH cell network function. *Proc Natl Acad Sci USA* (2010) 107:4465–70. doi: 10.1073/pnas.0902599107
154. Chauvet N, El-Yandouzi T, Mathieu M-N, Schlernitzauer A, Galibert E, Lafont C, et al. Characterization of adherens junction protein expression and localization in pituitary cell networks. *J Endocrinol* (2009) 202:375–87. doi: 10.1677/JOE-09-0153
155. Mullis PE. Genetics of Growth Hormone Deficiency. *Endocrinol Metab Clinics North America* (2007) 36:17–36. doi: 10.1016/j.ecl.2006.11.010
156. Fauquier T, Rizzoti K, Dattani M, Lovell-Badge R, Robinson ICAF. SOX2-expressing progenitor cells generate all of the major cell types in the adult mouse pituitary gland. *Proc Natl Acad Sci USA* (2008) 105:2907–12. doi: 10.1073/pnas.0707886105
157. Vankelecom H. Pituitary stem /progenitor cells: embryonic players in the adult gland? *Eur J Neurosci* (2010) 32:2063–81. doi: 10.1111/j.1460-9568.2010.07523.x
158. Borrelli E, Heyman RA, Arias C, Sawchenko PE, Evans RM. Transgenic mice with inducible dwarfism. *Nature* (1989) 339:538–41. doi: 10.1038/339538a0
159. Núñez L, Villalobos C, Senovilla L, García-Sancho J. Multifunctional Cells of Mouse Anterior Pituitary Reveal a Striking Sexual Dimorphism. *J Physiol* (2003) 549:835–43. doi: 10.1113/jphysiol.2003.040758
160. de Rooij DG, Mizrak SC. Deriving multipotent stem cells from mouse spermatogonial stem cells: a new tool for developmental and clinical research. *Development* (2008) 135:2207–13. doi: 10.1242/dev.015453
161. Suga H, Kadoshima T, Minaguchi M, Ohgushi M, Soen M, Nakano T, et al. Self-formation of functional adenohypophysis in three-dimensional culture. *Nature* (2011) 480:57–62. doi: 10.1038/nature10637
162. Pulichino A-M, Vallette-Kasic S, Tsai JP-Y, Couture C, Gauthier Y, Drouin J. Tpit determines alternate fates during pituitary cell differentiation. *Genes Dev* (2003) 17:738–47. doi: 10.1101/gad.1065703
163. Zhao L, Bakke M, Krimkevich Y, Cushman LJ, Parlow AF, Camper SA, et al. Steroidogenic factor 1 (SF1) is essential for pituitary gonadotrope function. *Development* (2001) 128:147–54.
164. Dasen JS, O'Connell SM, Flynn SE, Treier M, Gleiberman AS, Szeto DP, et al. Reciprocal interactions of Pit1 and GATA2 mediate signaling gradient-induced determination of pituitary cell types. *Cell* (1999) 97:587–98. doi: 10.1016/s0092-8674(00)80770-9
165. Andoniadou CL, Matsushima D, Gharavy SNM, Signore M, Mackintosh AI, Schaeffer M, et al. Sox2 Stem/Progenitor Cells in the Adult Mouse Pituitary Support Organ Homeostasis and Have Tumor-Inducing Potential. *Cell Stem Cell* (2013) 13:433–45. doi: 10.1016/j.stem.2013.07.004
166. Rizzoti K, Akiyama H, Lovell-Badge R. Mobilized adult pituitary stem cells contribute to endocrine regeneration in response to physiological demand. *Cell Stem Cell* (2013) 13:419–32. doi: 10.1016/j.stem.2013.07.006
167. Van Nassauw L, Harrisson F. 5-hydroxytryptophan-induced accumulation of secondary lysosomes in the avian adenohypophysis. *Eur Arch Biol* (1990) 101:219–35.
168. Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, et al. High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. *Nat Methods* (2019) 16:649–57. doi: 10.1038/s41592-019-0435-6
169. Hoa O, Lafont C, Fontanaud P, Guillou A, Kemkem Y, Kineman RD, et al. Imaging and Manipulating Pituitary Function in the Awake Mouse. *Endocrinology* (2019) 160:2271–81. doi: 10.1210/en.2019-00297
170. Zhang S, Cui Y, Ma X, Yong J, Yan L, Yang M, et al. Single-cell transcriptomics identifies divergent developmental lineage trajectories during human pituitary development. *Nat Commun* (2020) 11(1):5275. doi: 10.1038/s41467-020-19012-4
171. Ran FA, Ann Ran F, Cong L, Yan WX, Scott DA, Gootenberg JS, et al. In vivo genome editing using Staphylococcus aureus Cas9. *Nature* (2015) 520:186–91. doi: 10.1038/nature14299

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Santiago-Andres, Golan and Fiordelisio. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.