



Negative Frequency-Dependent Selection Is Frequently Confounding

*Dustin Brisson**

Biology Department, University of Pennsylvania, Philadelphia, PA, United States

Persistent genetic variation within populations presents an evolutionary problem, as natural selection and genetic drift tend to erode genetic diversity. Models of balancing selection were developed to account for the maintenance of genetic variation observed in natural populations. Negative frequency-dependent selection is a powerful type of balancing selection that maintains many natural polymorphisms, but it is also commonly misinterpreted. This review aims to clarify the processes underlying negative frequency-dependent selection, describe classes of polymorphisms that can and cannot result from these processes, and discuss the empirical data needed to accurately identify processes that generate or maintain diversity in nature. Finally, the importance of accurately describing the processes affecting genetic diversity within populations as it relates to research progress is considered.

OPEN ACCESS

Edited by:

Norman A. Johnson,
University of Massachusetts Amherst,
United States

Reviewed by:

Rama Shankar Singh,
McMaster University, Canada
Cock Van Oosterhout,
University of East Anglia,
United Kingdom

*Correspondence:

Dustin Brisson
dbrisson@sas.upenn.edu

Specialty section:

This article was submitted to
Evolutionary and Population Genetics,
a section of the journal
Frontiers in Ecology and Evolution

Received: 01 December 2017

Accepted: 19 January 2018

Published: 21 February 2018

Citation:

Brisson D (2018) Negative
Frequency-Dependent Selection Is
Frequently Confounding.
Front. Ecol. Evol. 6:10.
doi: 10.3389/fevo.2018.00010

Keywords: negative frequency dependent selection, balancing selection, killing the winner hypothesis, multiple niche polymorphism, density dependent selection

INTRODUCTION

Natural diversity—the “endless forms most beautiful and most wonderful” (Darwin, 2012)—is an enduring focus of both evolutionary biologists and nature lovers. The evolutionary processes that have generated or are maintaining many examples of diversity in nature, however, remain obscure and often controversial (Chesson, 2000). The processes that result in persistent polymorphisms within populations demand a special explanation as both directional natural selection and genetic drift should eliminate alleles and thus erode genetic diversity (Lewontin, 1974; Charlesworth and Hughes, 2000; Nielsen, 2005). Nevertheless, many examples of persistent polymorphisms occur in nature (Hedrick, 1986; Mallet and Joron, 1999; Richman, 2000; Carius et al., 2001; Hedrick et al., 2002; Delph and Kelly, 2014). Models of balancing selection—including negative frequency-dependent selection, spatial or temporal habitat heterogeneity, and heterozygote advantage—provide theoretical frameworks describing the processes that can account for persistent polymorphisms within populations. A core tenet of each balancing selection model is that the selective value of an allele—whether it is beneficial or detrimental—is dependent on the environmental context (Dobzhansky, 1982; Clarke et al., 1988). That is, alleles are advantageous and deleterious in different circumstances.

Negative frequency-dependent selection has been called the most powerful selective force maintaining balanced polymorphisms (Ayala and Campbell, 1974; Turelli and Barton, 2004; Fitzpatrick et al., 2007; Kazancıoğlu and Arnqvist, 2014), with some proposing that a large proportion of natural genetic polymorphisms are maintained by selection favoring rare alleles (Kojima and Yarbrough, 1967). Negative frequency-dependent selection occurs when the selective value of a variant (relative to other variants) is a function of its abundance in the population (relative to other variants) such that its relative fitness increases as the relative abundance, or frequency,

of the variant decreases (Wright, 1939) (please see Clarke, 1979; Levin, 1988 for foundational mathematical descriptions and assumptions of this process). That is, rare variants have a selective advantage specifically because of their rarity while common variants are disadvantaged because of their commonness. Negative frequency-dependent selection has the potential to maintain polymorphisms within populations because relatively rare variants have a selective advantage over more common variants and thus tend to increase in frequency and avoid local extinction. Negative frequency-dependent selection models are a narrow subset of a broad field of models describing the impact of variant frequency on natural selection; the overwhelming majority of this broad field is beyond the scope of the concepts addressed here. Here, I focus on natural polymorphisms that can be explained by negative frequency-dependent selection, where genetic diversity is maintained when a variant becomes disadvantageous as it becomes more frequent, and polymorphisms that are more accurately explained by other process.

Numerous ecological interactions can result in a selective advantage for relatively rare alleles including sexual selection, parasite or predator preferences, and resource competition. In fact, each of these mechanisms has been shown to create a selective advantage for rare alleles that has resulted in persistent polymorphisms in multiple natural populations (Fisher, 1930; Wright, 1939; Harvey et al., 1975; Gigord et al., 2001; Delph and Kelly, 2014). While ecological context and natural history determine the proximate ecological mechanism affecting the differential survival or reproduction of variants in a population, changes in relative survival or reproduction must be negatively correlated with variant frequency for negative frequency-dependent selection to maintain natural polymorphisms. In a classic example, color polymorphisms are maintained in natural populations of *Cepaea nemoralis* snails by negative frequency-dependent selection because their predators, the song thrush (*Turdus philomelos*), form a search image for the most common morph resulting in much greater predation pressure on the common than the rare morph (Harvey et al., 1975; Allen, 1988). The rare morph can increase in frequency due to the relaxed predation pressure until it becomes common, resulting in a search image switch that now targets the new common morph, a process that maintains this polymorphism in *C. nemoralis* populations. Two luminaries in population genetics—R. Fisher and S. Wright—have also demonstrated the power of negative frequency-dependent selection to maintain diversity in natural systems. Wright famously demonstrated that self-incompatibility alleles, a genetic mechanism in plants to prevent inbreeding, are incredibly diverse because pollen containing a rare allele is more likely to find a receptive mate than pollen containing a common allele (Wright, 1964, 1969; Castric and Vekemans, 2004). Thus, plants with rare alleles have a selective advantage (Figure S1 and Appendix 2). Similarly, Fisher's principle demonstrates that human males and females are equally frequent because, if one sex were more frequent, parents producing the alternate sex would enjoy an advantage resulting in more grandchildren (Fisher, 1930; Edwards, 1998).

The many incontrovertible demonstrations of the power of negative frequency-dependent selection to maintain polymorphisms in nature have led some to suggest that it is a “pervasive” force maintaining natural diversity (Clarke, 1979). The pervasiveness of negative frequency-dependent selection has been further supported by the perception that “nearly every [selective agent] works in a way liable to produce frequency-dependent selection of the kind that favors rare phenotypes and hinders common ones” (Clarke, 1979). Although negative frequency-dependent selection may be a “powerful, perhaps a dominant, factor maintaining genetic diversity” within populations (Clarke, 1979), many natural polymorphisms are maintained by other evolutionary processes (Allison, 1954; Barton, 1986; Smith, 1990; Weatherall, 1997; Schmidt et al., 2000; Brisson and Dykhuizen, 2004). Nevertheless, many natural polymorphisms have been assumed to result from negative frequency-dependent selection even when empirical data from the system are inconsistent with the theoretical framework in which selection favors relatively rare variants. In this essay, I describe several patterns of allele dynamics that are commonly described in the literature as resulting from negative frequency-dependent selection despite data demonstrating that other causative processes. These processes include allelic diversity resulting from directional selection within a changing ecological context, density-dependent population regulation, other models of balancing selection, and aspects of community ecology. I will discuss concepts and experiments that can aid in identifying the processes underlying patterns of allele dynamics and suggest that accurately identifying the evolutionary process underlying natural patterns facilitates the development of hypotheses and future experiments to determine the ecological interactions or molecular mechanisms at the root of the process.

DIRECTIONAL SELECTION DESCRIBED AS NEGATIVE FREQUENCY-DEPENDENT SELECTION

Conceptually, negative frequency-dependent selection may be the “most intuitively obvious explanation” of polymorphisms in nature (Trotter and Spencer, 2007). However, the original concept becomes ambiguous, complex, and even controversial as a result of differing definitions and applications in both theoretical and empirical work (Heino et al., 1998). Even some of the greatest thinkers in evolutionary biology have used negative frequency-dependent selection to explain scenarios in which the selective values of alleles are independent of their relative abundance. A prominent example comes from an influential essay by JBS Haldane outlining mechanisms by which infectious diseases drive natural selection in metazoans (Haldane, 1949). Although most of these ideas have been “followed profitably” (very profitably indeed), the negative frequency-dependent selection framework described in this essay appears to be one of the few unsound lines of thought. In this framework, Haldane suggested that a host with a rare defensive phenotype has a selective advantage in the face of highly-adapted pathogens, “For just because of its rarity it will be resistant

to diseases which attack the majority of its fellows.” That is, the adapted pathogen has evolved mechanisms to overcome the common defensive phenotypes in host populations but cannot overcome the rare defensive phenotypes. Thus, hosts expressing rare but effective defensive phenotypes, or *escape variants*, enjoy a selective advantage over hosts expressing common but exploitable defenses.

The scenario described by Haldane, however, confounds natural selection favoring a specific (*effective*) phenotype in the current environment with a selective advantage resulting from rarity. Haldane’s escape variants have a selective advantage because they cannot be subverted by the pathogen, not because they are rare. Although both rarity and novelty can result in a selective advantage, the novel defensive phenotype maintains its efficacy against the pathogen not because it is rare, but because it is novel. This point can be illustrated by extending this line of thought to allow migration of many individuals expressing a novel and effective defensive phenotype. These migrants would enjoy the same selective advantage over the previously common resident phenotype, regardless of frequency of the novel phenotype in the population immediately following the mass-migration event. The evolutionary dynamics occurring in this framework do not occur because of rare advantage and, in most cases, will not result in a balanced polymorphism. These evolutionary dynamics are more likely the result of directional selection in a continuously changing environment (Levins, 1968; Lande and Shannon, 1996; Orr, 2005; Collins et al., 2007; Bell, 2010). These two processes—negative frequency-dependent selection and selection in a changing environment—can potentially be distinguished by artificially manipulating variant frequencies or by introducing a previously common but now extinct variant into a controlled population.

The genetic diversity of haemagglutinin (HA) glycoproteins in the influenza virus is another conspicuous example of selection in a changing environment that is often confounded with negative frequency-dependent selection. The dynamics of HA alleles change over time such that rare alleles enter the population, rise to high population sizes, and subsequently decline toward extinction (Earn et al., 2002; Andreasen, 2003; Lin et al., 2003). The strains expressing a numerically common allele have relatively low fitness and decline in frequency because there are few hosts still susceptible to this strain, as hosts acquire immunity to strains with which they have been previously infected (Pease, 1987; Stegeman et al., 2004; Virseda et al., 2010). By contrast, strains expressing numerically rare alleles have many susceptible hosts available and enjoy high rates of secondary infections per infected host causing a numerical increase (Stegeman et al., 2004; Virseda et al., 2010). While there is undoubtedly strong selection at the HA locus, the selective advantage is derived not from relative rarity but from antigenic novelty (Plotkin and Dushoff, 2003; Nelson and Holmes, 2007; Cherry et al., 2009; Virseda et al., 2010), similar to Haldane’s example. The presence or frequency of alternative HA alleles does not affect the fitness (growth rate) or temporal dynamics of the alleles. That is, the population dynamics of a numerically rare allele is the same if the host population is already plagued by other numerically common strains (0.0001% when one novel allele enters a population of


106 infected hosts) and if it enters a host population in which no other influenza strain is circulating (100% when one novel allele enters a previous uninfected host population) (**Figure 1**). As the selective value of the allele is conditioned on the absolute abundance—but not the relative abundance—of the allele, it is unlikely that negative frequency-dependent selection is the evolutionary process underlying the polymorphism commonly observed at the HA locus. More likely, the common variant is changing its own environment such that there are few susceptible hosts in which new infections can establish, but it is not affecting the environment of alternative variants.

DENSITY-DEPENDENT FITNESS DYNAMICS DESCRIBED AS NEGATIVE FREQUENCY-DEPENDENT SELECTION

A preeminent evolutionary biologist, Lewontin suggested that negative frequency-dependent selection should be pervasive because, whenever “a genotype is its own worst enemy, its fitness will decrease as it becomes more common” (Lewontin, 1974). As similar variants occupy similar ecological niches and are commonly their own worst enemy, this logic suggests that negative frequency-dependent selection should indeed be pervasive. However, “common” in this case refers not to relative abundance but absolute abundance. For example, the fitness (growth rate) of individuals within a monomorphic population, one in which the frequency of a genotype is always at 100%, decreases as it “becomes more common” in absolute abundance as it approaches a carrying capacity. Further, relatively rare variants suffer negative fitness effects in proportion to the absolute abundance of their numerically common competitors such that relative rarity may not provide a selective advantage.

There is an extensive literature describing fitness (growth rate) as a function of the absolute abundance of each variant in a population (Birch, 1955; MacArthur, 1962; MacArthur et al., 1967; Roughgarden, 1971; Emlen, 1985). The above scenario can be characterized using classical Logistic growth models that include competition among variants such that “a genotype is its own worst enemy” (Lotka-Volterra models) (Equation 1). The growth rates of the variants in these models are a function of the absolute abundance of each variant—discounted by their competitive abilities (α_{ij})—with respect to the carrying capacity (K), but are not explicitly conditioned on the abundance of the variants relative to each other. An interesting body of literature uses this modeling framework to describe the generation and maintenance of polymorphisms not through negative frequency-dependent selection mechanisms but through disruptive selection conditioned on the strength of competitive interactions and the abundance of each variant (ex Kisdi, 1999).

$$\begin{aligned} \frac{dN_1}{dt} &= r_1 N_1 \left(1 - \frac{N_1 + \alpha_{12} N_2}{K_1} \right) \\ \frac{dN_2}{dt} &= r_2 N_2 \left(1 - \frac{N_2 + \alpha_{21} N_1}{K_2} \right) \end{aligned} \quad (1)$$



It is often challenging to distinguish the effect of numerical rarity from relative rarity on the selective value of an allele through observations of patterns of allelic diversity. Experimental manipulations of the carrying capacity (K), potentially through resource supplementation, can assuage the reductions in relative fitness experienced by common variants that result from high densities without altering relative frequencies. In these experiments, the relative fitness of common variants should increase if the effects are associated with density while the relative fitness of the common and rare variants should not be altered if the allelic diversity is maintained by negative frequency-dependent selection.

MULTIPLE NICHE POLYMORPHISMS DESCRIBED AS NEGATIVE FREQUENCY-DEPENDENT SELECTION

In the multiple niche selection model of balancing selection, the selective value of a trait is conditioned on its ability to exploit different environmental features in a heterogeneous habitat (Levene, 1953; Ravigne et al., 2004). Multi-niche selection maintains multiple variants in a population if each variant has a selective advantage in some available habitats while other variants are superior in other habitats. This idea—that environmentally variable selection can result in balanced polymorphisms—has a long history in the literature in which the foundational idea is stated by Dobzhansky (1982). Although incontrovertible examples of multi-niche selection maintaining polymorphism in natural populations are relatively rare, correct inference of the process resulting in balancing selection is necessary to generate hypotheses and design experiments to determine the ecological interactions or molecular mechanisms underlying the process.

The study of pattern, in isolation from the evolutionary processes that generated it, is not likely to advance general theories nor an understanding of specific systems (Cale et al., 1989). However, determining the processes responsible for balanced polymorphism patterns observed in nature is a difficult task (Barrett, 1988; Chaboudez and Burdon, 1995; Laine et al., 2011; Kazancıoğlu and Arnqvist, 2014). The balanced polymorphism at the outer surface protein C (*ospC*) locus in populations of *Borrelia burgdorferi*, the cause of human Lyme disease, provides a fitting example. Although the function of *OspC* remains unclear (Pal et al., 2004; Tilly et al., 2006, 2013; Xu et al., 2007; Onder et al., 2012; Carrasco et al., 2015), the within-population diversity at this locus bears all the hallmarks of balancing selection—large numbers of alleles in all local populations; allele frequencies that are more even than expected at neutrally evolving loci; and genetic evidence of an ancient polymorphism (Charlesworth et al., 1997; Qiu et al., 1997, 2002; May et al., 1999; Wang et al., 1999; Brisson and Dykhuizen, 2004).

Negative frequency-dependent selection and multi-niche selection have both been proposed as processes maintaining *ospC* polymorphisms, and both frameworks have empirical support (Qiu et al., 1997; Wang et al., 1999; Haven et al., 2011;

Brisson et al., 2012; Seifert et al., 2015). The negative frequency-dependent selection model suggests that the polymorphism can be maintained if previously infected hosts are immune to subsequent infections by the same *OspC* variant but susceptible to novel variants, a molecular mechanism that has been demonstrated in laboratory animals (Gilmore et al., 1996; Probert et al., 1997; but see, Devevey et al., 2015). However, in this scenario the frequency or even presence of alternative *OspC* variants does not affect the number of susceptible hosts for the common strain, similar to the influenza example, arguing against negative frequency-dependent selection as an evolutionary process maintaining *ospC* polymorphisms. Further, negative frequency-dependent selection is most effective when few hosts remain susceptible to the common *ospC* variants, a pattern that is not observed in natural data sets (Brisson and Dykhuizen, 2004; Hanincova et al., 2006; Ogden et al., 2008; States et al., 2014; Vuong et al., 2014). Studies investigating allelic diversity at *ospC* from natural hosts consistently demonstrate that most natural reservoir hosts, those that are regularly infected with *B. burgdorferi*, are rarely infected with all of the common *ospC* variants (Brisson and Dykhuizen, 2004; Hanincova et al., 2006; Vuong et al., 2014; Mechai et al., 2016). Most hosts are, however, infected with a subset of the *ospC* variants, as expected if each host species represented a different ecological niche (Brisson and Dykhuizen, 2004; Hanincova et al., 2006; Vuong et al., 2014; Mechai et al., 2016). Further, host individuals of the same species, including humans, are often infected by the same subset of *ospC* variants across both time and geography (Seinost et al., 1999; Brisson and Dykhuizen, 2004; Hanincova et al., 2006; Dykhuizen et al., 2008; Wormser et al., 2008; Vuong et al., 2014; Mechai et al., 2016). The collective evidence suggests that the balanced *ospC* polymorphisms are more likely maintained by multi-niche selection—with each host species representing multiple niches (Brisson et al., 2011), one for each *ospC* variant by which it can be infected—than by negative frequency-dependent selection. These results suggest that the mechanisms causing the balanced polymorphism are more likely to involve *ospC* variant-by-host species interactions than to involve a memory immune response mechanism that is conserved across vertebrate species.


It has been argued that “Selection in multiple niches is not an alternative to [negative] frequency-dependent selection... but a way of generating it” (Clarke, 1979). However, scenarios in which balanced polymorphisms can be maintained without a selective advantage favoring relatively rare variants are not uncommon, suggesting that these are two distinct evolutionary processes in at least some cases. To illustrate this point, imagine two variants occupying a heterogeneous habitat where each variant has a selective advantage in one niche but is disadvantaged in another, a classical multi-niche selection scenario (Levene, 1953; Ravigne et al., 2004). Here we assume that the carrying capacity in niche A is much lower than the carrying capacity in niche B ($K_A = 10$; $K_B = 10^5$). In this scenario, variant B—which has a competitive advantage in niche B—can retain a fitness advantage (a greater *per capita* growth rate) even when it is more common than variant A—which has a competitive advantage in niche A. For example, in a population with 90 variant B individuals and 10 variant A individuals, variant B has a rapid

per capita rate of increase while variant A does not increase (Figure 2). Here, the relatively common variant B has a “selective advantage” over the relatively rare variant A due to multi-niche selection, which is independent of negative frequency-dependent selection. Depending on the parameter values in this model, a balanced polymorphism can be maintained in the absence of rare advantage.

COMMUNITY DIVERSITY DESCRIBED AS NEGATIVE FREQUENCY-DEPENDENT SELECTION

Prominent population geneticists including Williams and Maynard Smith, among many others, have demonstrated that the efficacy of natural selection decreases at increasing levels of biological organization such that selection among individuals within populations is much more efficient than selection among species within communities (Maynard Smith, 1964, 1976; Williams, 1966). Additionally, selection at higher levels of organization (i.e., among species within communities) “tends to be undermined by natural selection at lower levels” (i.e., among individuals with populations) (Wilson and Wilson, 2007). Nevertheless, several studies have suggested that negative frequency-dependent selection maintains species diversity within ecological communities. There is a rich empirical and theoretical history describing the causes and consequences of species diversity within ecological communities (Connell and Orias, 1964; Schoener, 1974; Lubchenco, 1978; Ricklefs and Schluter, 1993; Chesson, 2000; Wright, 2002). Mechanisms of coexistence function in two major ways: *equalizing* mechanisms minimize the average fitness differences between species while *stabilizing* mechanisms increase negative *intraspecific* interactions relative to negative *interspecific* interactions (Chesson, 2000). Stabilizing mechanisms promote species coexistence and include mechanisms such as resource partitioning and frequency-dependent predation, as well as mechanisms that depend on spatial or temporal fluctuations in population densities or environmental factors. Equalizing mechanisms contribute to stable coexistence when they reduce large average fitness inequalities which might negate the effects of stabilizing mechanisms (Chesson, 2000). While some natural forces that affect the maintenance of community diversity have frequency-dependent mechanisms, this should not be mistaken for negative frequency-dependent selection which maintains polymorphisms within populations. Applying models of natural selection to levels of biological organization above the population level should be exercised only with the greatest of caution (Williams, 1966).

The Killing the Winner (KtW) hypothesis is a recent endeavor to understand patterns of diversity within communities using a negative frequency-dependent selection framework (Thingstad and Lignell, 1997; Thingstad, 2000). The KtW hypothesis suggests that predators target species that maximize reproductive effort over those that invest heavily in predator defense. Recent extensions of the KtW hypothesis suggest that this predator functional response promotes community diversity through



negative frequency dependent selection. However, the functional response in this hypothesis is often not conditioned on the frequency of the prey species but on the presence or absence of character traits in the prey species (Thingstad and Lignell,

1997; Suttle, 2007; Winter et al., 2010; Koskella and Meaden, 2013). The “winner” in this hypothesis refers to species that invest resource into reproduction at the expense of investing in predator defenses, which may or may not correspond to the most frequent species (Winter et al., 2010). In these cases, neither the relative nor the absolute abundance of the prey species affects the functional responses of the predator.

CONCLUDING REMARKS AND FUTURE PERSPECTIVES

Understanding the processes that produce or maintain diversity in natural populations is a central challenge in evolutionary biology. Negative frequency-dependent selection maintains many noted and striking polymorphisms in nature (Kojima and Tobar, 1969; Gigord et al., 2001; Charlesworth, 2006; Loisel et al., 2006; Fitzpatrick et al., 2007; Mitchell-Olds et al., 2007; Mokkonen et al., 2011), and many polymorphisms exist in the absence of a selective advantage favoring rare variants (Allison, 1954; Barton, 1986; Smith, 1990; Weatherall, 1997; Schmidt et al., 2000; Brisson and Dykhuizen, 2004). Ideally, one could unequivocally determine the causative process through observations of the patterns of variation in nature. Unfortunately, many processes result in identical patterns, especially when those patterns are observed over short time scales. In some cases, long-term observations of allelic dynamics can distinguish polymorphisms caused by mutation-selection balance or selection in a changing environment from a stable polymorphism resulting from balancing selection (Roy, 1998; Schmidt et al., 2000; Schmidt, 2001; Siemens and Roy, 2005; Olendorf et al., 2006; Koskella and Lively, 2009). Evidence suggesting negative frequency-dependent selection—such as allelic cycles where each allele gains a selective advantage as it becomes more rare—may also be observed from long-term observational studies (Gigord et al., 2001; Thrall et al., 2012). More directly, the patterns resulting from specific evolutionary processes can be tested through controlled and natural experiments such as manipulating allele frequencies in sub-populations (Roy, 1998; Schmidt et al., 2000; Olendorf et al., 2006; Koskella and Lively, 2009).

Ecological and molecular mechanisms are rarely deducible from patterns (Kershaw, 1963), but accurate identification of the evolutionary processes causing the pattern can generate hypotheses about these mechanisms. For example, the northern acorn barnacle, *Semibalanus balanoides*, shows clear evidence of a balanced polymorphism at the mannose-6-phosphate isomerase (*mpi*) locus (Hoffmann, 1981; McDonald, 1991). The pattern of *mpi* genotype frequencies among intertidal microhabitats, where one allele is common in high intertidal zones but rare in low intertidal zones, suggests that multi-niche selection maintains this polymorphism (Schmidt and Rand, 1999). Experimental manipulations of genotypes among microhabitats confirmed that multi-niche selection is the process responsible for the allelic variation (Schmidt et al., 2000; Schmidt and Rand, 2001). The molecular mechanism linking mannose utilization with survivorship in high intertidal zones, where

temperature and desiccation stress is high, was subsequently elucidated through controlled laboratory experiments (Schmidt, 2001). As this and many other examples demonstrate, the ecological interaction or molecular mechanism underlying an evolutionary process can best be understood when the evolutionary process is accurately determined.

AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and approved it for publication.

REFERENCES

- Allen, J. A. (1988). Frequency-dependent selection by predators. *Philos. Trans. R. Soc. B* 319, 485–503. doi: 10.1098/rstb.1988.0061
- Allison, A. C. (1954). Protection afforded by sickle-cell trait against subtertian malarial infection. *Br. Med. J.* 1, 290–294. doi: 10.1136/bmj.1.4857.290
- Andreasen, V. (2003). Dynamics of annual influenza epidemics with immunoselection. *J. Math. Biol.* 46, 504–536. doi: 10.1007/s00285-002-0186-2
- Ayala, F. J., and Campbell, C. A. (1974). Frequency-dependent selection. *Annu. Rev. Ecol. Syst.* 5, 115–138. doi: 10.1146/annurev.es.05.110174.000555
- Barrett, J. A. (1988). Frequency-dependent selection in plant fungal interactions. *Philos. Trans. R. Soc. B* 319, 473–483. doi: 10.1098/rstb.1988.0060
- Barton, N. (1986). The maintenance of polygenic variation through a balance between mutation and stabilizing selection. *Genet. Res.* 47, 209–216. doi: 10.1017/S0016672300023156
- Bell, G. (2010). Fluctuating selection: the perpetual renewal of adaptation in variable environments. *Philos. Trans. R. Soc. B* 365, 87–97. doi: 10.1098/rstb.2009.0150
- Birch, L. C. (1955). Selection in *Drosophila pseudoobscura* in relation to crowding. *Evolution* 9, 389–399. doi: 10.1111/j.1558-5646.1955.tb01549.x
- Brisson, D., Baxamusa, N., Schwartz, L., and Wormser, G. P. (2011). Biodiversity of *Borrelia burgdorferi* strains in tissues of Lyme disease patients. *PLoS ONE* 6:e22926. doi: 10.1371/journal.pone.0022926
- Brisson, D., Drecktrah, D., Eggers, C. H., and Samuels, D. S. (2012). Genetics of *Borrelia burgdorferi*. *Annu. Rev. Genet.* 46, 515–536. doi: 10.1146/annurev-genet-011112-112140
- Brisson, D., and Dykhuizen, D. E. (2004). *ospC* diversity in *Borrelia burgdorferi*: different hosts are different niches. *Genetics* 168, 713–722. doi: 10.1534/genetics.104.028738
- Cale, W., Henebry, G., and Yeakley, J. (1989). Inferring process from pattern in natural communities. *Bioscience* 39, 600–605. doi: 10.2307/1311089
- Carius, H. J., Little, T. J., and Ebert, D. (2001). Genetic variation in a host-parasite association: potential for coevolution and frequency-dependent selection. *Evolution* 55, 1136–1145. doi: 10.1111/j.0014-3820.2001.tb00633.x
- Carrasco, S. E., Troxell, B., Yang, Y., Brandt, S. L., Li, H., Sandusky, G. E., et al. (2015). Outer surface protein OspC is an antiphagocytic factor that protects *Borrelia burgdorferi* from phagocytosis by macrophages. *Infect. Immun.* 83, 4848–4860. doi: 10.1128/IAI.01215-15
- Castric, V., and Vekemans, X. (2004). Plant self-incompatibility in natural populations: a critical assessment of recent theoretical and empirical advances. *Mol. Ecol.* 13, 2873–2889. doi: 10.1111/j.1365-294X.2004.02267.x
- Chaboudez, P., and Burdon, J. J. (1995). Frequency-dependent selection in a wild plant-pathogen system. *Oecologia* 102, 490–493. doi: 10.1007/BF00341361
- Charlesworth, B., and Hughes, K. A. (2000). “The maintenance of genetic variation in life-history traits,” in *Evolutionary Genetics: from Molecules to Morphology*, eds R. S. Singh and C. B. Krimbas (Cambridge, UK: Cambridge University Press), 369–392.
- Charlesworth, B., Nordborg, M., and Charlesworth, D. (1997). The effects of local selection, balanced polymorphism and background selection on equilibrium patterns of genetic diversity in subdivided populations. *Genet. Res.* 70, 155–174. doi: 10.1017/S0016672397002954
- Charlesworth, D. (2006). Balancing selection and its effects on sequences in nearby genome regions. *PLoS Genet.* 2:e64. doi: 10.1371/journal.pgen.0020064
- Cherry, J. L., Lipman, D. J., Nikolskaya, A., and Wolf, Y. (2009). Evolutionary dynamics of N-glycosylation sites of influenza virus hemagglutinin. *PLoS Curr.* 1:RRN1001. doi: 10.1371/currents.RRN1001
- Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* 31, 343–366. doi: 10.1146/annurev.ecolsys.31.1.343
- Clarke, B. C. (1979). The evolution of genetic diversity. *Proc. R. Soc. B* 205, 453–474. doi: 10.1098/rspb.1979.0079
- Clarke, B. C., Shelton, P. R., and Mani, G. S. (1988). Frequency-dependent selection, metrical characters and molecular evolution. *Philos. Trans. R. Soc. B* 319, 631–640. doi: 10.1098/rstb.1988.0070
- Collins, S., De Meaux, J., and Acquisti, C. (2007). Adaptive walks toward a moving optimum. *Genetics* 176, 1089–1099. doi: 10.1534/genetics.107.072926
- Connell, J. H., and Orias, E. (1964). The ecological regulation of species diversity. *Am. Nat.* 98, 399–414. doi: 10.1086/282335
- Darwin, C. (2012). *The Origin of Species*, Dover Thrift Edition. New York, NY: Courier Corporation.
- Delph, L. F., and Kelly, J. K. (2014). On the importance of balancing selection in plants. *New Phytol.* 201, 45–56. doi: 10.1111/nph.12441
- Devevey, G., Dang, T., Graves, C. J., Murray, S., and Brisson, D. (2015). First arrived takes all: inhibitory priority effects dominate competition between co-infecting *Borrelia burgdorferi* strains. *BMC Microbiol.* 15:61. doi: 10.1186/s12866-015-0381-0
- Dobzhansky, T. (1982). *Genetics and the origin of Species*. New York, NY: Columbia University Press.
- Dykhuizen, D. E. D. E., Brisson, D., Sandigursky, S., Wormser, G. P. G. P., Nowakowski, J., Nadelman, R. B., et al. (2008). The propensity of different *Borrelia burgdorferi* sensu stricto genotypes to cause disseminated infections in humans. *Am. J. Trop. Med. Hyg.* 78, 806–810. doi: 10.4269/ajtmh.2008.78.806
- Earn, D. J. D., Dushoff, J., and Levin, S. A. (2002). Ecology and evolution of the flu. *Trends Ecol. Evol.* 17, 334–340. doi: 10.1016/S0169-5347(02)02502-8
- Edwards, A. W. F. (1998). Natural selection and the sex ratio: Fisher's sources. *Am. Nat.* 151, 564–569.
- Emlen, J. M. (1985). The assessment of frequency- and density-dependent influences on fitness in natural populations. *Am. Nat.* 125, 507–520. doi: 10.1086/284359
- Fisher, R. A. (1930). *The Genetical Theory of Natural Selection*. New York, NY: Oxford University Press.
- Fitzpatrick, M. J., Feder, E., Rowe, L., and Sokolowski, M. B. (2007). Maintaining a behaviour polymorphism by frequency-dependent selection on a single gene. *Nature* 447, 210–212. doi: 10.1038/nature05764
- Gigord, L. D., MacNair, M. R., and Smithson, A. (2001). Negative frequency-dependent selection maintains a dramatic flower color polymorphism in the rewardless orchid *Dactylorhiza sambucina* (L.). *Soo. Proc. Natl. Acad. Sci. U.S.A.* 98, 6253–6255. doi: 10.1073/pnas.111162598
- Gilmore, R. D., Kappel, K. J., Dolan, M. C., Burkot, T. R., and Johnson, B. J. B. (1996). Outer surface protein C (OspC), but not P39, is a protective immunogen against a tick-transmitted *Borrelia burgdorferi* challenge: evidence for a conformational protective epitope in OspC. *Infect. Immun.* 64, 2234–2239.
- Haldane, J. B. S. (1949). Disease and evolution. *La Ric. Sci.* 19(Suppl.), 1–11.

FUNDING

Burroughs Wellcome Fund (1012376), the National Institutes of Health (NIH AI076342 and AI097137), and National Science Foundation (DEB 1354184).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fevo.2018.00010/full#supplementary-material>

- Hanincova, K., Kurtenbach, K., Diuk-Wasser, M., Brei, B., and Fish, D. (2006). Epidemic spread of Lyme borreliosis, northeastern United States. *Emerg. Infect. Dis.* 12, 604–611. doi: 10.3201/eid1204.051016
- Harvey, P. H., Birley, N., and Blackstock, T. H. (1975). The effect of experience on the selective behaviour of song thrushes feeding on artificial populations of *Cepaea*. *Genetica* 45, 211–216. doi: 10.1007/BF01517197
- Haven, J., Vargas, L. C., Mongodin, E. F., Xue, V., Hernandez, Y., Pagan, P., et al. (2011). Pervasive recombination and sympatric genome diversification driven by frequency-dependent selection in *Borrelia burgdorferi*, the Lyme disease bacterium. *Genetics* 189, 951–966. doi: 10.1534/genetics.111.130773
- Hedrick, P. W. (1986). Genetic-polymorphism in heterogeneous environments - a decade later. *Annu. Rev. Ecol. Syst.* 17, 535–566. doi: 10.1146/annurev.es.17.110186.002535
- Hedrick, P. W., Lee, R. N., and Garrigan, D. (2002). Major histocompatibility complex variation in red wolves: evidence for common ancestry with coyotes and balancing selection. *Mol. Ecol.* 11, 1905–1913. doi: 10.1046/j.1365-294X.2002.01579.x
- Heino, M., Metz, J. A. J., and Kaitala, V. (1998). The enigma of frequency-dependent selection. *Trends Ecol. Evol.* 13, 367–370. doi: 10.1016/S0169-5347(98)01380-9
- Hoffmann, R. J. (1981). Evolutionary genetics of *Metridium senile*. I. Kinetic differences in phosphoglucose isomerase allozymes. *Biochem. Genet.* 19, 129–144. doi: 10.1007/BF00486143
- Kazancıoğlu, E., and Arnqvist, G. (2014). The maintenance of mitochondrial genetic variation by negative frequency-dependent selection. *Ecol. Lett.* 17, 22–27. doi: 10.1111/ele.12195
- Kershaw, K. A. (1963). Pattern in vegetation and its causality. *Ecology* 44, 377–388. doi: 10.2307/1932185
- Kisdi, E. (1999). Evolutionary branching under asymmetric competition. *J. Theor. Biol.* 197, 149–162. doi: 10.1006/jtbi.1998.0864
- Kojima, K., and Tobari, Y. (1969). Selective modes associated with karyotypes in *Drosophila ananassae*. II. Heterosis and frequency-dependent selection. *Genetics* 63, 639–651.
- Kojima, K., and Yarbrough, K. M. (1967). Frequency-dependent selection at the *esterase-6-locus* in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U.S.A.* 57, 645–649. doi: 10.1073/pnas.57.3.645
- Koskella, B., and Lively, C. M. (2009). Evidence for negative frequency-dependent selection during experimental coevolution of a freshwater snail and a sterilizing trematode. *Evolution* 63, 2213–2221. doi: 10.1111/j.1558-5646.2009.00711.x
- Koskella, B., and Meaden, S. (2013). Understanding bacteriophage specificity in natural microbial communities. *Viruses* 5, 806–823. doi: 10.3390/v5030806
- Laine, A. L., Burdon, J. J., Dodds, P. N., and Thrall, P. H. (2011). Spatial variation in disease resistance: from molecules to metapopulations. *J. Ecol.* 99, 96–112. doi: 10.1111/j.1365-2745.2010.01738.x
- Lande, R., and Shannon, S. (1996). The role of genetic variation in adaptation and population persistence in a changing environment. *Evolution* 56, 429.
- Levene, H. (1953). Genetic equilibrium when more than one ecological niche is available. *Am. Nat.* 87, 331–333. doi: 10.1086/281792
- Levin, B. R. (1988). Frequency dependent selection in bacterial populations. *Philos. Trans. R. Soc. B* 319, 459–472. doi: 10.1098/rstb.1988.0059
- Levins, R. (1968). *Evolution in Changing Environments*. Princeton, NJ: Princeton University Press.
- Lewontin, R. C. (1974). *The Genetic Basis of Evolutionary Change*. New York, NY: Columbia University Press.
- Lin, J., Andreasen, V., Casagrandi, R., and Levin, S. A. (2003). Traveling waves in a model of influenza A drift. *J. Theor. Biol.* 222, 437–445. doi: 10.1016/S0022-5193(03)00056-0
- Loisel, D. A., Rockman, M. V., Wray, G. A., Altmann, J., and Alberts, S. C. (2006). Ancient polymorphism and functional variation in the primate MHC-DQA1 5' cis-regulatory region. *Proc. Natl. Acad. Sci. U.S.A.* 103, 16331–16336. doi: 10.1073/pnas.0607662103
- Lubchenco, J. (1978). Plant species diversity in a marine intertidal community: importance of herbivore food preference and algal competitive. *Am. Nat.* 112, 23–39. doi: 10.1086/283250
- MacArthur, R. H. (1962). Some generalized theorems of natural selection. *Genetics* 48, 1893–1897
- MacArthur, R. H., Wilson, E. O., and MacArthur, W. (1967). *The Theory of Island Biogeography*. Princeton, NJ: Princeton University Press.
- Mallet, J., and Joron, M. (1999). Evolution of diversity in warning color and mimicry: polymorphisms, shifting balance, and speciation. *Annu. Rev. Ecol. Syst.* 30, 201–233. doi: 10.1146/annurev.ecolsys.30.1.201
- May, G., Shaw, F., Badrane, H., and Vekemans, X. (1999). The signature of balancing selection: fungal mating compatibility gene evolution. *Proc. Natl. Acad. Sci. U.S.A.* 96, 9172–9177. doi: 10.1073/pnas.96.16.9172
- Maynard Smith, J. (1964). Group selection and kin selection. *Nature* 201, 1145–1147. doi: 10.1038/2011145a0
- Maynard Smith, J. (1976). Commentary - group selection. *Q. Rev. Biol.* 51, 277–283.
- McDonald, J. H. (1991). Contrasting amounts of geographical variation as evidence for direct selection: the *mpi* and *pgm* loci in eight crustacean species. *Heredity* 67, 215–219. doi: 10.1038/hdy.1991.82
- Mechai, S., Margos, G., Feil, E. J., Barairo, N., Lindsay, L. R., Michel, P., et al. (2016). Evidence for host-genotype associations of *Borrelia burgdorferi* sensu stricto. *PLoS ONE* 11:e0149345. doi: 10.1371/journal.pone.0149345
- Mitchell-Olds, T., Willis, J. H., and Goldstein, D. B. D. B. (2007). Which evolutionary processes influence natural genetic variation for phenotypic traits? *Nat. Rev. Genet.* 8, 845–856. doi: 10.1038/nrg2207
- Mokkonen, M., Kokko, H., Koskela, E., Lehtonen, J., Mappes, T., Martiskainen, H., et al. (2011). Negative frequency-dependent selection of sexually antagonistic alleles in *Myodes glareolus*. *Science* 334, 972–974. doi: 10.1126/science.1208708
- Nelson, M. I., and Holmes, E. C. (2007). The evolution of epidemic influenza. *Nat. Rev. Genet.* 8, 196–205. doi: 10.1038/nrg2053
- Nielsen, R. (2005). Molecular signatures of natural selection. *Annu. Rev. Genet.* 39, 197–218. doi: 10.1146/annurev.genet.39.073003.112420
- Ogden, N. H., Lindsay, L. R., Hanincova, K., Barker, I. K., Bigras-Poulin, M., Charron, D. F., et al. (2008). Role of migratory birds in introduction and range expansion of *Ixodes scapularis* ticks and of *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in Canada. *Appl. Environ. Microbiol.* 74, 1780–1790. doi: 10.1128/AEM.01982-07
- Olendorf, R., Rodd, F. H., Punzalan, D., Houde, A. E., Hurt, C., Reznick, D. N., et al. (2006). Frequency-dependent survival in natural guppy populations. *Nature* 441, 633–636. doi: 10.1038/nature04646
- Onder, O., Humphrey, P. T., McOmber, B., Korobova, F., Francella, N., Greenbaum, D. C., et al. (2012). OspC is potent plasminogen receptor on surface of *Borrelia burgdorferi*. *J. Biol. Chem.* 287, 16860–16868. doi: 10.1074/jbc.M111.290775
- Orr, H. A. (2005). Theories of adaptation: what they do and don't say. *Genetica* 123, 3–13. doi: 10.1007/s10709-004-2702-3
- Pal, U., Yang, X., Chen, M., Bockenstedt, L. K., Anderson, J. F., Flavell, R. A., et al. (2004). OspC facilitates *Borrelia burgdorferi* invasion of *Ixodes scapularis* salivary glands. *J. Clin. Invest.* 113, 220–230. doi: 10.1172/JCI200419894
- Pease, C. M. (1987). An evolutionary epidemiological mechanism, with applications to type A influenza. *Theor. Popul. Biol.* 31, 422–452. doi: 10.1016/0040-5809(87)90014-1
- Plotkin, J. B., and Dushoff, J. (2003). Codon bias and frequency-dependent selection on the hemagglutinin epitopes of influenza A virus. *Proc. Natl. Acad. Sci. U.S.A.* 100, 7152–7157. doi: 10.1073/pnas.1132114100
- Probert, W. S., Crawford, M., Cadiz, R. B., and LeFebvre, R. B. (1997). Immunization with outer surface protein (Osp) A, but not OspC, provides cross-protection of mice challenged with North American isolates of *Borrelia burgdorferi*. *J. Infect. Dis.* 175, 400–405. doi: 10.1093/infdis/175.2.400
- Qiu, W. G., Bosler, E. M., Campbell, J. R., Ugine, G. D., Wang, I. N., Luft, B. J., et al. (1997). A population genetic study of *Borrelia burgdorferi* sensu stricto from eastern Long Island, New York, suggested frequency-dependent selection, gene flow and host adaptation. *Heredity* 127, 203–216. doi: 10.1111/j.1601-5223.1997.00203.x
- Qiu, W. G., Dykhuizen, D. E., Acosta, M. S., and Luft, B. J. (2002). Geographic uniformity of the Lyme disease spirochete (*Borrelia burgdorferi*) and its shared history with tick vector (*Ixodes scapularis*) in the northeastern United States. *Genetics* 160, 833–849.
- Ravigne, V., Olivieri, I., and Dieckmann, U. (2004). Implications of habitat choice for protected polymorphisms. *Evol. Ecol. Res.* 6, 125–145.
- Richman, A. (2000). Evolution of balanced genetic polymorphism. *Mol. Ecol.* 9, 1953–1963. doi: 10.1046/j.1365-294X.2000.01125.x

- Ricklefs, R. E., and Schluter, D. (1993). *Species Diversity in Ecological Communities: Historical and Geographical Perspectives*. Chicago, IL: University of Chicago Press.
- Roughgarden, J. (1971). Density-dependent natural selection. *Ecology* 52, 453–468. doi: 10.2307/1937628
- Roy, B. A. (1998). Differentiating the effects of origin and frequency in reciprocal transplant experiments used to test negative frequency-dependent selection hypotheses. *Oecologia* 115, 73–83. doi: 10.1007/s004420050493
- Schmidt, P. S. (2001). The effects of diet and physiological stress on the evolutionary dynamics of an enzyme polymorphism. *Proc. R. Soc. B* 268, 9–14. doi: 10.1098/rspb.2000.1323
- Schmidt, P. S., Bertness, M. D., and Rand, D. M. (2000). Environmental heterogeneity and balancing selection in the acorn barnacle, *Semibalanus balanoides*. *Philos. Trans. R. Soc. B* 267, 379–384. doi: 10.1098/rspb.2000.1012
- Schmidt, P. S., and Rand, D. M. (1999). Intertidal microhabitat and selection at *Mpi*: interlocus contrasts in the northern acorn barnacle, *Semibalanus glandula*. *Evolution* 53, 135–146.
- Schmidt, P. S., and Rand, D. M. (2001). Adaptive maintenance of genetic polymorphism in an intertidal barnacle: habitat- and life-stage-specific survivorship of *Mpi* genotypes. *Evolution* 55, 1336–1344. doi: 10.1111/j.0014-3820.2001.tb00656.x
- Schoener, T. W. (1974). Resource partitioning in ecological communities research on how similar species divide resources helps reveal the natural regulation of species diversity. *Science* 185, 27–39. doi: 10.1126/science.185.4145.27
- Seifert, S. N., Khatchikian, C. E., Zhou, W., and Brisson, D. (2015). Evolution and population genomics of the Lyme borreliosis pathogen, *Borrelia burgdorferi*. *Trends Genet.* 31, 201–207. doi: 10.1016/j.tig.2015.02.006
- Seinost, G., Dykhuizen, D. E., Dattwyler, R. J., Golde, W. T., Dunn, J. J., Wang, I. N., et al. (1999). Four clones of *Borrelia burgdorferi* sensu stricto cause invasive infection in humans. *Infect. Immun.* 67, 3518–3524.
- Siemens, D. H., and Roy, B. A. (2005). Tests for parasite-mediated frequency-dependent selection in natural populations of an asexual plant species. *Evol. Ecol.* 19, 321–338. doi: 10.1007/s10682-005-6639-5
- Smith, T. B. (1990). Natural selection on bill characters in the two bill morphs of the African finch, *Pyrenestes ostrinus*. *Evolution* 44, 832–842. doi: 10.1111/j.1558-5646.1990.tb03808.x
- States, S. L., Brinkerhoff, R. J., Carpi, G., Steeves, T. K., Folsom-O'Keefe, C., DeVeaux, M., et al. (2014). Lyme disease risk not amplified in a species-poor vertebrate community: Similar *Borrelia burgdorferi* tick infection prevalence and OspC genotype frequencies. *Infect. Genet. Evol.* 27, 566–575. doi: 10.1016/j.meegid.2014.04.014
- Stegeman, A., Bouma, A., Elbers, A. R. W., de Jong, M. C. M., Nodelijk, G., de Klerk, F., et al. (2004). Avian influenza A virus (H7N7) epidemic in the Netherlands in 2003: course of the epidemic and effectiveness of control measures. *J. Infect. Dis.* 190, 2088–2095. doi: 10.1086/425583
- Suttle, C. A. (2007). Marine viruses—major players in the global ecosystem. *Nat. Rev. Microbiol.* 5, 801–812. doi: 10.1038/nrmicro1750
- Thingstad, T. F. (2000). Elements of a theory for the mechanisms controlling abundance, diversity, and biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnol. Oceanogr.* 45, 1320–1328. doi: 10.4319/lo.2000.45.6.1320
- Thingstad, T. F., and Lignell, R. (1997). Theoretical models for the control of bacterial growth rate, abundance, diversity and carbon demand. *Aquat. Microb. Ecol.* 13, 19–27. doi: 10.3354/ame013019
- Thrall, P. H., Laine, A. L., Ravensdale, M., Nemri, A., Dodds, P. N., Barrett, L. G., et al. (2012). Rapid genetic change underpins antagonistic coevolution in a natural host-pathogen metapopulation. *Ecol. Lett.* 15, 425–435. doi: 10.1111/j.1461-0248.2012.01749.x
- Tilly, K., Bestor, A., and Rosa, P. A. (2013). Lipoprotein succession in *Borrelia burgdorferi*: similar but distinct roles for OspC and VlsE at different stages of mammalian infection. *Mol. Microbiol.* 89, 216–227. doi: 10.1111/mmi.12271
- Tilly, K., Krum, J. G., Bestor, A., Jewett, M. W., Grimm, D., Bueschel, D., et al. (2006). *Borrelia burgdorferi* OspC protein required exclusively in a crucial early stage of mammalian infection. *Infect. Immun.* 74, 3554–3564. doi: 10.1128/IAI.01950-05
- Trotter, M. V., and Spencer, H. G. (2007). Frequency-dependent selection and the maintenance of genetic variation: exploring the parameter space of the multi-allelic pairwise interaction model. *Genetics* 176, 1729–1740. doi: 10.1534/genetics.107.073072
- Turelli, M., and Barton, N. H. (2004). Polygenic variation maintained by balancing selection: pleiotropy, sex-dependent allelic effects and G × E interactions. *Genetics* 166, 1053–1079. doi: 10.1534/genetics.166.2.1053
- Virseda, S., Restrepo, M. A., Arranz, E., Magan-Tapia, P., Fernandez-Ruiz, M., de la Camara, A. G., et al. (2010). Seasonal and pandemic A (H1N1) 2009 influenza vaccination coverage and attitudes among health-care workers in a Spanish university hospital. *Vaccine* 28, 4751–4757. doi: 10.1016/j.vaccine.2010.04.101
- Vuong, H. B., Canham, C. D., Fonseca, D. M., Brisson, D., Morin, P. J., Ostfeld, R. S., et al. (2014). Occurrence and transmission efficiencies of *Borrelia burgdorferi* ospC types in avian and mammalian wildlife. *Infect. Genet. Evol.* 27, 594–600. doi: 10.1016/j.meegid.2013.12.011
- Wang, I. N., Dykhuizen, D. E., Qiu, W., Dunn, J. J., Bosler, E. M., and Luft, B. J. (1999). Genetic diversity of ospC in a local population of *Borrelia burgdorferi* sensu stricto. *Genetics* 151, 15–30.
- Weatherall, D. J. (1997). Thalassaemia and malaria, revisited. *Ann. Trop. Med. Parasitol.* 91, 885–890. doi: 10.1080/00034983.1997.11813215
- Williams, G. C. (1966). *Adaptation and Natural Selection*. Princeton, NJ: Princeton University Press.
- Wilson, D. S., and Wilson, E. O. (2007). Rethinking the theoretical foundation of sociobiology. *Q. Rev. Biol.* 82, 327–348. doi: 10.1086/522809
- Winter, C., Bouvier, T., Weinbauer, M. G., and Thingstad, T. F. (2010). Trade-offs between competition and defense specialists among unicellular planktonic organisms: the “killing the winner” hypothesis revisited. *Microbiol. Mol. Biol. Rev.* 74, 42–57. doi: 10.1128/MMBR.00034-09
- Wormser, G. P., Brisson, D., Liveris, D., Hanincova, K., Sandigursky, S., Nowakowski, J., et al. (2008). *Borrelia burgdorferi* genotype predicts the capacity for hematogenous dissemination during early Lyme disease. *J. Infect. Dis.* 198, 1358–1364. doi: 10.1086/592279
- Wright, S. (1939). The distribution of self-sterility alleles in populations. *Genetics* 24, 538–552.
- Wright, S. (1964). The distribution of self-incompatibility alleles in populations. *Evolution* 18, 609–619. doi: 10.1111/j.1558-5646.1964.tb01675.x
- Wright, S. (1969). *Evolution and the Genetics of Populations, Vol. 2, The Theory of Gene Frequencies*. Chicago, IL: University of Chicago Press.
- Wright, S. J. (2002). Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia* 130, 1–14. doi: 10.1007/s004420100809
- Xu, Q., McShan, K., and Liang, F. T. (2007). Identification of an ospC operator critical for immune evasion of *Borrelia burgdorferi*. *Mol. Microbiol.* 64, 220–231. doi: 10.1111/j.1365-2958.2007.05636.x

Conflict of Interest Statement: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Brisson. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.