



# **Corrigendum:** *In situ* **Pumping Rate** of 20 Marine Demosponges Is a Function of Osculum Area

Teresa Maria Morganti<sup>1,2,3\*</sup>, Marta Ribes<sup>2</sup>, Raz Moskovich<sup>4,5</sup>, Jeremy Brian Weisz<sup>6</sup>, Gitai Yahel<sup>5</sup> and Rafel Coma<sup>3\*</sup>

<sup>1</sup> Max Planck Institute for Marine Microbiology, Bremen, Germany, <sup>2</sup> Institut de Ciències del Mar, Consejo Superior de Investigaciones Cientificas (ICM-CSIC), Barcelona, Spain, <sup>3</sup> Centre d'Estudis Avançats de Blanes, Consejo Superior de Investigaciones Cientificas (CEAB-CSIC), Girona, Spain, <sup>4</sup> School of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel, <sup>5</sup> Faculty of Marine Sciences, Ruppin Academic Center, Michmoret, Israel, <sup>6</sup> Department of Biology, Linfield University, McMinnville, OR, United States

Keywords: size, allometric scaling, osculum, HMA-LMA sponges, sponge (Porifera), pumping rate

#### A Corrigendum on

#### In situ Pumping Rate of 20 Marine Demosponges Is a Function of Osculum Area

by Morganti, T. M., Ribes, M., Moskovich, R., Weisz, J. B., Yahel, G., and Coma, R. (2021). Front. Mar. Sci. 8:583188. doi: 10.3389/fmars.2021.583188

In the published article, there was an error regarding the affiliation for Raz Moskovich. As well as having affiliation 5, she should also have 4 "School of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel".

Murk, A. J., et al. (2020). The effect of depth on the morphology, bacterial clearance, and

respiration of the Mediterranean sponge Chondrosia reniformis (Nardo, 1847). Mar. Drugs 18:358.

doi: 10.3390/md18070358" was not cited in the article. The citation has now been inserted in

total OSA (i.e., the total osculum cross-sectional area) and sponge volume conformed to a power

function allometric scaling with a similar exponent (b = 0.6 - 0.7), except for tropical HMAs (b =

0.99). A theoretical allometric scaling of the increase in OSA with sponge volume has been recently

suggested to be: OSA  $\sim V^{2/3}$  and this suggestion was confirmed in small specimens in laboratory experiments (*Halicondria panicea*, b = 0.66, Goldstein et al., 2019). Our results agree with those and

previous estimates (*Aphrocallistes vastus*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, b = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Leys et al., 2011; *Cinachyrella* cf. *cavernosa*, *b* = 0.84, Cinachyrella cf. *cavernosa*, *b* = 0.84, Cinachyrella cf. *cavernosa*, *b* = 0.84, Cinachyrella cf. *cavernosa*, *cavernosa*,

0.50, Dahihande and Thakur, 2019) and provide an in situ confirmation of the suggested theoretical

exponent based on a very limited size range, on several species that encompass the natural range of

both parameters (Table 4). An increase of total OSA with sponge volume can be achieved either by

the increase of the number of oscula and/or by the increase of OSA with sponge volume. Due to the

high morphological plasticity of sponges, changes in the number of oscula and aquiferous modules

may be attributed to different environmental conditions (Plotkin et al., 1999; Ereskovskii, 2003). A

recent study observed a depth induced change in osculum morphology in Chondrosia reniformis.

This change was attributed to wave action and sediment loading and did not affect total OSA,

bacterial clearance, respiration and growth (Gökalp et al., 2020a; but see Lesser et al., 2020). As the

sampled specimens inhabited the same area under similar environmental conditions, we postulate

that the observed variability within each species should be attributed to inherent properties such as

sponge size, metabolism, and internal structure, rather than to abiotic factors such as temperature

"In both the temperate and tropical species analyzed in this study, the relationships between the

Discussion, Oscula Number and Their Cross-Sectional Area, paragraph 1 and should read:

In the original article "Gökalp, M., Kooistra, T., Rocha, M. S., Silva, T. H., Osinga, R.,

## **OPEN ACCESS**

#### Approved by:

Frontiers Editorial Office, Frontiers Media SA, Switzerland

### \*Correspondence:

Teresa Maria Morganti tmorgant@mpi-bremen.de Rafel Coma coma@ceab.csic.es

#### Specialty section:

This article was submitted to Aquatic Physiology, a section of the journal Frontiers in Marine Science

> **Received:** 21 May 2021 **Accepted:** 24 May 2021 **Published:** 05 July 2021

#### Citation:

Morganti TM, Ribes M, Moskovich R, Weisz JB, Yahel G and Coma R (2021) Corrigendum: In situ Pumping Rate of 20 Marine Demosponges Is a Function of Osculum Area. Front. Mar. Sci. 8:712856. doi: 10.3389/fmars.2021.712856

Frontiers in Marine Science | www.frontiersin.org

1

and current (Riisgård et al., 1993)."

Additionally, in the original article the reference for "Gökalp et al., 2020" was incorrectly given as "Gokalp, M., Kuehnhold, H., de Goeij, J. M., and Osinga, R. (2020). Depth and turbidity affect *in situ* pumping activity of the Mediterranean sponge Chondrosia reniformis (Nardo, 1847). *bioRxiv* 2020.03.30.009290. doi: 10.1101/2020.03.30.009290". It should be "Gökalp, M., Kuehnhold, H., de Goeij, J. M., and Osinga, R. (2020b). Depth and turbidity affect *in situ* pumping activity of the Mediterranean sponge Chondrosia reniformis (Nardo, 1847). *bioRxiv* 2020.03.30.009290. doi: 10.1101/2020.03.30.009290". A correction has been made to **Introduction, paragraph 3**:

"Direct relationship between the osculum cross-sectional area and the amount of water pumped by a sponge has been recently demonstrated in laboratory experiments with small sponge explants (Strehlow et al., 2016; Kumala et al., 2017; Goldstein et al., 2019; Kealy et al., 2019) and in one *in situ* study (Gökalp et al., 2020b), suggesting that osculum cross-sectional area and the number of oscula may be more practical predictors of sponge pumping."

The authors apologize for these errors and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2021 Morganti, Ribes, Moskovich, Weisz, Yahel and Coma. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.