

Cytokines induce effector T-helper cells during invasive aspergillosis; what we have learned about T-helper cells?

Raman Thakur¹, Rajesh Anand², Shraddha Tiwari¹, Agam P. Singh²,
Bhupendra N. Tiwary³ and Jata Shankar^{1*}

¹ Department of Biotechnology and Bioinformatics, Jaypee University of Information Technology, Solan, India, ² Infectious Diseases Laboratory, National Institute of Immunology, New Delhi, India, ³ Department of Biotechnology, Guru Ghasidas Vishwavidyalaya, Bilaspur, India

OPEN ACCESS

Edited by:

Frederic Lamoth,
Lausanne University Hospital,
Switzerland

Reviewed by:

Nina Khanna,
University of Basel, Switzerland
Lars Tramsen,
University Hospital Frankfurt,
Germany

*Correspondence:

Jata Shankar,
Department of Biotechnology
and Bioinformatics, Jaypee University
of Information Technology,
Waknaghat, Solan 173234,
Himachal Pradesh, India
jata.shankar@juit.ac.in,
jata_s@yahoo.com

Specialty section:

This article was submitted to
Fungi and Their Interactions,
a section of the journal
Frontiers in Microbiology

Received: 02 January 2015

Accepted: 22 April 2015

Published: 11 May 2015

Citation:

Thakur R, Anand R, Tiwari S,
Singh AP, Tiwary BN and Shankar J
(2015) Cytokines induce effector
T-helper cells during invasive
aspergillosis; what we have learned
about T-helper cells?
Front. Microbiol. 6:429.
doi: 10.3389/fmicb.2015.00429

Invasive aspergillosis caused by *Aspergillus* species (*Aspergillus fumigatus*, *A. flavus*, and *A. terreus*) is life-threatening infections in immunocompromised patients. Understanding the innate and adaptive immune response particularly T-helper cells (T_H-cells) against these *Aspergillus* species and how the different sub-set of T_H-cells are regulated by differentiating cytokines at primary target organ site like lung, kidney and brain is of great significance to human health. This review focuses on presentation of *Aspergillus* through Antigen presenting cells (APCs) to the naive CD4⁺ T-cells in the host. The production of differentiating/effector cytokines that activate following T_H-cells, e.g., T_H1, T_H2, T_H9, and T_H17 has been reported in association or alone in allergic or invasive aspergillosis. Chemokines (CXCL1, CXCL2, CCL1, and CCL20) and their receptors associated to these T_H-cells have also been observed in invasive aspergillosis. Thus, further study of these T_H-cells in invasive aspergillosis and other elements of adaptive immune response with *Aspergillus* species are required in order to have a better understanding of host response for safer and effective therapeutic outcome.

Keywords: cytokines, T-helper cells, dendritic cells, *Aspergillus*, antigen presenting cells, invasive aspergillosis

Introduction

Fungi are the most common microorganisms and have clinical importance. Few of them are pathogenic or opportunistic pathogen and results in morbidity and mortality to human beings. There is a rise in opportunistic fungal infections in recent years due to increased incidence of immunocompromised host (Chamilos et al., 2006; Romani, 2008). After *Candida albicans*, the leading causes of fungal infections in immunocompromised individuals are from *Aspergillus* species. *Aspergillus* is one of the most ubiquitous medically important opportunistic fungi (Weaver et al., 2007). The genus *Aspergillus*, contains about 40 species that can cause infection (Verweij and Brandt, 2007), among them *A. fumigatus*, *A. flavus*, and *A. terreus* are the leading cause of invasive Aspergillosis in immunocompromised individuals. These species produce conidia at a concentration of around 1–100 conidia per m³ (Barnes and Marr, 2006). Human routinely inhale hundreds of these conidia per day, despite these exposure to *Aspergillus* conidia, human do not develop any disease due to the clearance of conidia from lung by innate immunity especially phagocytic cells (Chamilos et al., 2006; Romani, 2008). However, due to rise in immunocompromised host, e.g., patients receiving organ

transplant, immunosuppressive therapy for autoimmune or neoplastic disease and HIV patients, inhaled conidia if not cleared in these host, colonization of *Aspergillus* occurs (Stevens et al., 2000). The adaptive immune response in human responsible for conidia clearance is not well understood in immunocompetent host as well as where conidia colonize in immunocompromised host. It is worth to note that secondary metabolites (e.g., Gliotoxins, Aflatoxins) excreted by *Aspergillus* especially have been recognized to modulate immunological responses (Shankar, 2013). Thus, we reviewed recent advances made in immune responses against *Aspergillus* species in mice model studies and clinical aspergillosis patients. *A. fumigatus* is the prominent species, which cause 90% of Aspergillosis followed by *A. flavus* and *A. terreus*. Studies have showed that *Aspergillus* species associated with infection after hematopoietic stem cell transplantation include *A. fumigatus* with 56% followed by *A. flavus* (18.7%) and *A. terreus* (16%) (Steinbach et al., 2004; Morgan et al., 2005). The involvement of *Aspergillus* infection in pulmonary tuberculosis and in asthmatic patients has been reported by Denning et al. It has been estimated annually at least 372,385 patients developed chronic pulmonary aspergillosis worldwide following treated pulmonary tuberculosis (Denning et al., 2011). Similarly, around 4,837,000 patients develop Allergic bronchopulmonary aspergillosis out of 193 million adults with active asthma (Denning et al., 2013). However, in a recent study of Indian population by Agarwal et al. (2014) the estimated ABPA burden was 1.38 million out of 27.6 million adults with asthma. This review touches different aspect of antifungal immunity against aspergillosis that include antigen presenting cells (APCs), dendritic cells (DCs), fungal pattern recognition receptors (PRRs), T_H-cells with their subsets profile during infection associated to *Aspergillus* species at different site of infection, e.g., lung, kidney, and brain.

Recognition of *Aspergillus* by the Host

Presentation of Pathogen via Soluble Receptors and Cell Bound Receptors

After the inhalation of *A. fumigatus* conidia, they are entrapped by the lung alveoli and if they are not efficiently cleared from lung, they germinate and establish lung infection termed invasive pulmonary aspergillosis and it may also disseminates to other organs if not treated (Park and Mehrad, 2009). The recognition of *A. fumigatus* conidia and hyphae occurs by PRRs those include soluble receptors and cell-bound receptors. Conidial germination starts with hydrophobic layer degradation and exposure of inner cell wall components mainly polysaccharides, which includes chitin, β -glucan, mannan, and galactomannan. These are termed as pathogen associated molecular patterns (PAMP), are recognized by PRRs (Netea et al., 2006; Inoue and Shinohara, 2014). PRRs soluble receptor such as pulmonary collectins, family of C-type lectins, pentraxin-3, pulmonary surfactant proteins-A and D have been reported in aspergillosis. Further, the cell-bound receptors in association with aspergillosis include Toll like receptor-2 (TLR), TLR-4 and TLR-9, which potentially induce the production of pro-inflammatory cytokines and reactive oxygen species through MyD88 signaling pathway (Willment and Brown, 2008).

Antigen Presenting Cells and T-cell Differentiation

Antigen Presenting Cells Triggers Cytokines Production

The activation of the innate immunity through PRRs present on the APCs that regulate the development of T-cell. APCs express wide-array of PRRs that provides the link between adaptive and innate immunity (Park and Mehrad, 2009). APCs, dominantly DCs, are responsible for antigen monitoring and then shaping T-cell response by secreting cytokines and chemokines. DCs express PRRs on their cell surface and endosomal compartments, which serve to recognize PAMPs. After interaction with DCs, naive T-cells are activated. The activation of T-cell response is regulated by the cytokines milieu predominantly framed by DCs (Akdis et al., 2011). Chemokines secreted by DCs recruit the phagocytic cells to infected areas to clear the *Aspergillus* components. APC cells, e.g., monocytes differentiate into distinct sub-populations of CD14⁺ and CD16⁺ cells after *A. fumigatus* conidia infection (Serbina et al., 2009). Monocytes interact with *Aspergillus* antigens resulting in maturation of monocytes into macrophages or DCs. Macrophages and DCs interact with antigens and secrete effector cytokines (Osugi et al., 2002; Ramirez-Ortiz and Means, 2012). Major sub-populations of DCs are myeloid DCs, plasmacytoid DC (pDCs) and monocyte-derived DCs (Bozza et al., 2002; Osugi et al., 2002). pDC recognize the nucleic acids from *A. fumigatus* via TLR-9 and lead to resistance to *A. fumigatus* infection in mice (Ramirez-Ortiz et al., 2008). Further, monocytes migrate toward the lung to differentiate into either DCs or alveolar macrophages during invasive aspergillosis (Cramer et al., 2011; Morton et al., 2012). Monocytes express different chemokines receptor predominantly CCR2, which help in migration of monocytes from bone marrow toward lung in response to *A. fumigatus* infection (Serbina et al., 2009). It has been shown that monocytes expressing CCR2 in the lung involved in conidial uptake and killing (Espinosa et al., 2014). Furthermore, alveolar macrophages induce APCs to release IL-1 β in pulmonary invasive infection (Park and Mehrad, 2009). IL-18 has also been observed in lung during invasive aspergillosis mice model (Akdis et al., 2011). Recently, it has been observed that *A. fumigatus* pulmonary challenge induces expression of the inflammasome-dependent cytokines IL-1 β and IL-18 within the first 12 h, while IL-1 α expression continually increases over at least the first 48 h (Caffrey et al., 2015). Moretti et al. (2014) showed in a pulmonary invasive aspergillosis model that mice injected with IL-37 prior to *A. fumigatus* infection has significant reduction in IL- β production and recruitment of neutrophils and resulted in diminution in lung inflammation and damage. The anti-inflammatory activity of IL-37 has been observed as an inhibitor of the innate response. Thus, cytokines play a vital role in modulation of immune response and coordinate the innate as well as adaptive responses. APCs secrete cytokines that act on naive T-cells leading to the differentiation of naive T-cells. These differentiated T-cells further secrete effector cytokines and regulate the function of T_H-cells. The profile of cytokine depends on the type of *Aspergillus* antigens, route of infection, immunological status of the host and cytokines milieu present during the interaction (Romani, 2008; Chai et al., 2010b). CD4⁺

T-cells can be divided into distinct subtypes according to cytokine profile, and they can differentiate to T_H1 , T_H2 , T_H17 , T_H9 , and T-follicular effector cells (Kerzerho et al., 2013; Kara et al., 2014). On the basis of the cytokine profile, these T_H -cells perform distinct functions. However, it is not clear how T-follicular effector cells respond during *Aspergillus* infection (Wüthrich et al., 2012).

Cytokines Associated with T_H1 Type of Response

Aspergillus fumigatus challenged intranasally in mice interacts with DCs and alveolar macrophages in the lung. Secretion of T_H1 associated pro-inflammatory cytokines IL-12, IFN- γ , TNF- α , IL-18 has been observed after the challenge (Chotirmall et al., 2013). Among these, IL-12 is the prominent cytokine released from activated monocytes and macrophages in lung that help in shaping T-cell immune response. IL-12 is a heterodimeric cytokine composed of IL-12p35 and IL-12p40 polypeptides that form the bioactive IL-12p70. The heterodimer binds to the IL-12 receptor composed of IL-12R β 1 and IL-12R β 2 chains and signals through STAT-4 (Shao et al., 2005). IL-12 acts on NK cells to promote IFN- γ secretion and differentiate $CD4^+$ T-cells into T_H1 -cells, once $CD4^+$ cells differentiate to T_H1 -cells, they increase the secretion of IFN- γ , which suppress T_H17 and T_H2 response in the lung (Espinosa and Rivera, 2012; Camargo and Husain, 2014). IL-12, hence, is the most important regulator of T_H1 response during lung infection. IL-12 deficient mice failed to generate a T_H1 response, leading to increased secretion of IL-4 and IL-10 cytokines, which shifts the immune response toward T_H2 pathway (Cenci et al., 1998). In *A. fumigatus* induced neutropenic aspergillosis in mice, NK cells can be the primary source of IFN- γ responsible for activating phagocytic cells and direct antifungal effectors cells against *A. fumigatus* (Park et al., 2009). Further, patients with invasive *Candida* and/or *Aspergillus* infections, recombinant treatment of IFN- γ in combination with antifungal drug partially restored immune function (Delsing et al., 2014). In intravenous infection of *A. flavus* mice model studies, lung and brain homogenate showed pro-inflammatory cytokines IL-12 and IFN- γ and relative absence of IL-4, IL-23, and IL-17 suggesting a T_H1 response (Anand et al., 2013, 2015). *A. terreus* induced invasive aspergillosis showed the presence of IL-1 β , IL-6, and reduced level of IL-10 in mice model studies. Although there is activation of T_H17 type of adaptive immune response through IL-1 β but the later is suppressed by T_H1 cytokines particularly IFN- γ (Vyas, 2011; Lass-Florl, 2012). The T_H1 response is thus also mounted by *A. terreus* and there is a lack of T_H2 response in contrast to *A. fumigatus* infection where T_H2 promoting cytokines are observed.


Cytokines Associated with T_H17 Type Response

Aspergillus fumigatus mediated infections in lung induce T_H17 and T_H1 -cells. These cells play an important role in protection and induction of inflammation (Chai et al., 2010b). Activation of T_H17 -cell depends on Dectin-1 signaling pathway. Various studies have suggested that dectin-1 deficient mice entirely activate T_H1 -cells. So Dectin-1 signaling not only serves as a positive factor to promote T_H17 differentiation but rather act to balance T_H1 versus T_H17 differentiation. Activation of the APCs by Dectin-1, release the proinflammatory cytokines IL-1 β , IL-6, IL-23, and

IL-22 which differentiates $CD4^+$ T-cells to T_H17 -cells, which further secretes IL-17A and IL-17F cytokines and maintain T_H17 response (Werner et al., 2011). IL-23 is a member of IL-12 family, produced by phagocytic cells, macrophages and activated DCs in lung. IL-23 contains two subunits IL-12p40 and IL-23p19 and it binds to heterodimeric receptors IL-12R β 1, expressed by activated T-cells (Zelante et al., 2007). IL-6 is another important cytokine involved in regulation of T_H17 response. IL-6 is a multifunctional cytokine, promote T_H17 -cells differentiation, inflammation and acute response (Akdis et al., 2011). During T_H17 differentiation, human naïve T-cells are exposed to IL-1 β , IL-6, and IL-23 (Zelante et al., 2009; Gresnigt and van de Veer-donk, 2014). T_H17 promoting cytokine IL-17 binds to IL-17RA and IL-17RC receptors expressed in lung cells, like fibroblast, epithelial cells and T-cells. After release of IL-17 from T_H17 -cells, it activates the neutrophils migration toward infected area and increases inflammation (Wilson et al., 2007). In *A. flavus* and *A. terreus*, the role of T_H17 -cells during lung infection is yet to be established.

Cytokines Associated with T_H2 Type of Response

Aspergillus fumigatus is associated with both invasive and allergic form of aspergillosis. In case, if conidia are not cleared, they germinate to produce hyphae, which are responsible for invasion in host tissues that leads to inflammation. In a healthy human T-cells response, *A. fumigatus* not only evoke pro-inflammatory type of immune response via T_H1 and T_H17 -cells but also anti-inflammatory type of immune response mediated by T_H2 -cells (Chaudhary et al., 2010). Immune response initiated by IL-4 and IL-10 inhibits T_H1 and T_H17 response and increased secretion of IL-4 and IL-10 inhibits IFN- γ and IL-12 production. T_H2 -cells differentiation depends on IL-4 and IL-10 and after differentiation in to T_H2 cells, these cells further secretes IL-5 and IL-13 which maintain T_H2 response. T_H2 immune response is triggered in acute bronchopulmonary aspergillosis and also in invasive pulmonary infection during some time point of infection. IL-4 and IL-10 deficient mice show lower *A. fumigatus* burden and increased survival rates compared to wild type mouse in invasive pulmonary aspergillosis (Cenci et al., 2000). It has been shown that ESTs (L3 ribosomal protein, L7A ribosomal protein, Histone -H2A) have high sequence similarity with human counter parts suggesting molecular mimicry between human and pathogen protein (Shankar et al., 2004). However, role of these genes in eliciting allergic immune response needs investigations. In *A. flavus* mediated infection, lung homogenate showed the absence of T_H2 response in a limited cytokine profile study (Anand and Tiwary, 2010). However, T_H2 type response may get activated in later stages of infection in lung due to rise in IL-4 and IL-10, which suppress the T_H1 response but consistent expression IFN- γ overcomes T_H2 response. In addition to T_H2 , the role of T_H9 -cells has been shown during infection with a Virus, bacteria, parasites and fungi. T_H9 response contributes to allergic inflammation during allergic aspergillosis due to *A. fumigatus* in mice model (Kerzerho et al., 2013). T_H9 -cells develop in the presence of IL-1 α and TGF- β along with T_H2 -cells. However, role of T_H9 and T_H2 response during invasive aspergillosis remains unclear.



Is Their Co-evolution of T-helper Cells During Invasive Aspergillosis?

In immunocompromised mice model studies, repeated exposure of *A. fumigatus* conidia in hosts lead to co-evolution of T_H1, T_H2, and T_H17 response in infected lung (Murdock et al., 2011). They have observed the presence of IFN-γ and IL-17 in infected lung of mice along with T_H2 response. These mixed responses might be occurring at different time points during progression of infection and leads to either protection or infection. T_H1 and T_H17 response probably leads to protection whereas T_H2 response further complicates the disease. T_H2 response helps in evasion of *A. fumigatus* from immune cells and further increases the IgE level, which leads to high inflammation at the site of infection. Mice model of ABPA demonstrated the T_H2 cytokine profile consisting of IL-4, IL-10, and IL-5 (Latge, 1999).

Interplay of Cytokines; T_H1 or T_H2 or T_H17 Type of Response

T_H-cells response during invasive *Aspergillus* depends on differentiating cytokines. T_H1 response is activated by differentiating

cytokine IL-12 followed by secretion of IFN-γ. Secretion of IFN-γ further stimulates T_H1 cells, if IFN-γ dominates initially it suppresses the other cytokines of T_H2 and T_H17, i.e., IL-4 and IL-17 (Harrington et al., 2005). If IL-4 dominates during the initial period of *Aspergillus* infection, it suppresses the protective T_H1 type immune response by inhibiting differential cytokine IL-12 and IFN-γ (Harrington et al., 2005). Recognition of *Aspergillus* antigens by Dectin-1 signaling pathway inhibits the production of IFN-γ and IL-12 receptors suppressing T_H1, which leads to differentiation of T_H17-cells and production of IL-17. In this way Dectin-1 signal balances the T_H1 and T_H17 response through the regulation of their respective cytokines (Rivera et al., 2011; Figure 1). The development of effective CD4⁺ T_H-cells response not only depends upon cytokines, but also on chemokines and their receptors. Chemokines help in recruitment of leukocytes, i.e., neutrophils, monocytes and NK cells toward lung during *Aspergillus* infection. These cells express chemokine receptors; neutrophils contain CXCR2 chemokine receptor for ligand CXCL1 and CXCL2, monocytes contain CCR2 and CCR6 receptor for CCL2 and CCL20 ligands, whereas NK cells contain CCR2 receptor for CCL2 ligand. So these chemokine ligands attract monocytes, neutrophils and NK cells to clear the *Aspergillus* hyphae during

lung infection (Park and Mehrad, 2009). These chemokines receptors are also present on DCs, regulatory T-cells (Tregs) and T_H-cells and help in their trafficking (Bendall, 2005). In this way, there is an interdependent relationship between chemokines and cytokines that help in evolution of effector T_H-cells response. CCL17, a chemokine, help in trafficking of DCs, Tregs and T_H1-cells toward infected area during invasive aspergillosis in response to CCR4 chemokine receptor present on these cells. Further, CCR6 receptor present on DCs and T_H17-cells help in migration of these cells in response to chemokine CCL20 (Bendall, 2005; Wüthrich et al., 2012).

Conclusion

Cytokines are important in the development of CD4⁺ T_H-cells. Understanding of trafficking of CD4⁺ T_H-cells and their regulation through differentiating/effector cytokines during invasive

aspergillosis will be crucial for the targeted immunotherapy. Overall, cytokines and chemokines may serve as prognostic biomarkers that could be followed to assess the effectiveness of treatment response during invasive aspergillosis. Measurement of selected cytokines in the blood samples of aspergillosis patients may be a promising tool for the monitoring of treatment responses. Also, manipulation of cytokine response e. g, IFN- γ or IFN- γ in combination with antifungal drug, IL-37, may be a future avenue for the development of better therapeutic against invasive aspergillosis.

Acknowledgments

Authors are thankful to Department of Biotechnology and Bioinformatics, Jaypee University of Information Technology, Solan, Himachal Pradesh, India, for financial support to Ph.D. students RT and ST.

References

- Agarwal, R., Denning, D. W., and Chakrabarti, A. (2014). Estimation of the burden of chronic and allergic pulmonary aspergillosis in India. *PLoS ONE* 9:e114745. doi: 10.1371/journal.pone.0114745
- Akdis, M., Burgler, S., Cramer, R., Eiwegger, T., Fujita, H., Gomez, E., et al. (2011). Interleukins, from 1 to 37, and interferon-gamma: receptors, functions, and roles in diseases. *J. Allergy Clin. Immunol.* 127, 701–721 e701–e770. doi: 10.1016/j.jaci.2010.11.050
- Anand, R., Shankar, J., Singh, A. P., and Tiwary, B. N. (2013). Cytokine milieu in renal cavities of immunocompetent mice in response to intravenous challenge of *Aspergillus flavus* leading to aspergillosis. *Cytokine* 61, 63–70. doi: 10.1016/j.cyto.2012.08.024
- Anand, R., Shankar, J., Tiwary, B. N., and Singh, A. P. (2015). *Aspergillus flavus* induces granulomatous cerebral aspergillosis in mice with display of distinct cytokine profile. *Cytokine* 72, 166–172. doi: 10.1016/j.cyto.2015.01.006
- Anand, R., and Tiwary, B. N. (2010). Th1 and Th2 cytokines in a self-healing primary pulmonary *Aspergillus flavus* infection in BALB/c mice. *Cytokine* 52, 258–264. doi: 10.1016/j.cyto.2010.07.428
- Barnes, P. D., and Marr, K. A. (2006). Aspergillosis: spectrum of disease, diagnosis, and treatment. *Infect. Dis. Clin. North Am.* 20, 545–561, vi. doi: 10.1016/j.idc.2006.06.001
- Bendall, L. (2005). Chemokines and their receptors in disease. *Histol. Histopathol.* 20, 907–926.
- Bozza, S., Gaziano, R., Spreca, A., Bacci, A., Montagnoli, C., Di Francesco, P., et al. (2002). Dendritic cells transport conidia and hyphae of *Aspergillus fumigatus* from the airways to the draining lymph nodes and initiate disparate Th responses to the fungus. *J. Immunol.* 168, 1362–1371. doi: 10.4049/jimmunol.168.3.1362
- Caffrey, A. K., Lehmann, M. M., Zickovich, J. M., Espinosa, V., Shepardson, K. M., Watschke, C. P., et al. (2015). IL-1 α signaling is critical for leukocyte recruitment after pulmonary *Aspergillus fumigatus* challenge. *PLoS Pathog.* 11:e1004625. doi: 10.1371/journal.ppat.1004625
- Camargo, J. F., and Husain, S. (2014). Immune correlates of protection in human invasive aspergillosis. *Clin. Infect. Dis.* 59, 569–577. doi: 10.1093/cid/ciu337
- Cenci, E., Mencacci, A., Bacci, A., Bistoni, F., Kurup, V. P., and Romani, L. (2000). T cell vaccination in mice with invasive pulmonary aspergillosis. *J. Immunol.* 165, 381–388. doi: 10.4049/jimmunol.165.1.381
- Cenci, E., Mencacci, A., Fe D'ostiani, C., Del Sero, G., Mosci, P., Montagnoli, C., et al. (1998). Cytokine- and T helper-dependent lung mucosal immunity in mice with invasive pulmonary aspergillosis. *J. Infect. Dis.* 178, 1750–1760. doi: 10.1086/314493
- Chai, L., Netea, M. G., Teerenstra, S., Earnest, A., Vonk, A. G., Schlamm, H. T., et al. (2010a). Early proinflammatory cytokines and C-reactive protein trends as predictors of outcome in invasive Aspergillosis. *J. Infect. Dis.* 202, 1454–1462. doi: 10.1086/656527
- Chai, L. Y., Van De Veerdonk, F., Marijnissen, R. J., Cheng, S. C., Khoo, A. L., Hectors, M., et al. (2010b). Anti-*Aspergillus* human host defence relies on type 1 T helper (Th1), rather than type 17 T helper (Th17), cellular immunity. *Immunology* 130, 46–54. doi: 10.1111/j.1365-2567.2009.03211.x
- Chamilos, G., Luna, M., Lewis, R. E., Bodey, G. P., Chemaly, R., Tarrand, J. J., et al. (2006). Invasive fungal infections in patients with hematologic malignancies in a tertiary care cancer center: an autopsy study over a 15-year period (1989–2003). *Haematologica* 91, 986–989.
- Chaudhary, N., Staab, J. F., and Marr, K. A. (2010). Healthy human T-Cell Responses to *Aspergillus fumigatus* antigens. *PLoS ONE* 5:e9036. doi: 10.1371/journal.pone.0009036
- Chotirmall, S. H., Al-Alawi, M., Mirkovic, B., Lavelle, G., Logan, P. M., Greene, C. M., et al. (2013). *Aspergillus*-associated airway disease, inflammation, and the innate immune response. *Biomed. Res. Int.* 2013, 723129. doi: 10.1155/2013/723129
- Cramer, R. A., Rivera, A., and Hohl, T. M. (2011). Immune responses against *Aspergillus fumigatus*: what have we learned? *Curr. Opin. Infect. Dis.* 24, 315–322. doi: 10.1097/QCO.0b013e328348b159
- Delsing, C. E., Gresnigt, M. S., Leentjens, J., Preijers, F., Frager, F. A., Kox, M., et al. (2014). Interferon-gamma as adjunctive immunotherapy for invasive fungal infections: a case series. *BMC Infect. Dis.* 14:166. doi: 10.1186/1471-2334-14-166
- Denning, D. W., Pleuvry, A., and Cole, D. C. (2011). Global burden of chronic pulmonary aspergillosis as a sequel to pulmonary tuberculosis. *Bull. World Health Organ.* 89, 864–872. doi: 10.2471/blt.11.089441
- Denning, D. W., Pleuvry, A., and Cole, D. C. (2013). Global burden of allergic bronchopulmonary aspergillosis with asthma and its complication chronic pulmonary aspergillosis in adults. *Med. Mycol.* 51, 361–370. doi: 10.3109/13693786.2012.738312
- Espinosa, V., Jhingran, A., Dutta, O., Kasahara, S., Donnelly, R., Du, P., et al. (2014). Inflammatory monocytes orchestrate innate antifungal immunity in the lung. *PLoS Pathog.* 10:e1003940. doi: 10.1371/journal.ppat.1003940
- Espinosa, V., and Rivera, A. (2012). Cytokines and the regulation of fungus-specific CD4 T cell differentiation. *Cytokine* 58, 100–106. doi: 10.1016/j.cyto.2011.11.005
- Gresnigt, M. S., and van de Veerdonk, F. L. (2014). The role of interleukin-1 family members in the host defence against *Aspergillus fumigatus*. *Mycopathologia* 178, 395–401. doi: 10.1007/s11046-014-9776-y
- Harrington, L. E., Hatton, R. D., Mangan, P. R., Turner, H., Murphy, T. L., Murphy, K. M., et al. (2005). Interleukin 17-producing CD4⁺ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nat. Immunol.* 6, 1123–1132. doi: 10.1038/ni1254
- Inoue, M., and Shinohara, M. L. (2014). Clustering of pattern recognition receptors for fungal detection. *PLoS Pathog.* 10:e1003873. doi: 10.1371/journal.ppat.1003873
- Kara, E. E., Comerford, I., Fenix, K. A., Bastow, C. R., Gregor, C. E., Mckenzie, D. R., et al. (2014). Tailored immune responses: novel effector helper T cell

- subsets in protective immunity. *PLoS Pathog.* 10:e1003905. doi: 10.1371/journal.ppat.1003905
- Kerzerho, J., Maazi, H., Speak, A. O., Szely, N., Lombardi, V., Khoo, B., et al. (2013). Programmed cell death ligand 2 regulates TH9 differentiation and induction of chronic airway hyperreactivity. *J. Allergy Clin. Immunol.* 131, 1048–1057, 1057 e1041–e1042. doi: 10.1016/j.jaci.2012.09.027
- Korn, T., Bettelli, E., Oukka, M., and Kuchroo, V. K. (2009). IL-17 and Th17 Cells. *Annu. Rev. Immunol.* 27, 485–517. doi: 10.1146/annurev.immunol.021908.132710
- Lass-Flörl, C. (2012). *Aspergillus terreus*: how inoculum size and host characteristics affect its virulence. *J. Infect. Dis.* 205, 1192–1194. doi: 10.1093/infdis/jis185
- Latge, J. P. (1999). *Aspergillus fumigatus* and aspergillosis. *Clin. Microbiol. Rev.* 12, 310–350.
- Moretti, S., Bozza, S., Oikonomou, V., Renga, G., Casagrande, A., Iannitti, R. G., et al. (2014). IL-37 inhibits inflammasome activation and disease severity in murine aspergillosis. *PLoS Pathog.* 10:e1004462. doi: 10.1371/journal.ppat.1004462
- Morgan, J., Wannemuehler, K. A., Marr, K. A., Hadley, S., Kontoyiannis, D. P., Walsh, T. J., et al. (2005). Incidence of invasive aspergillosis following hematopoietic stem cell and solid organ transplantation: interim results of a prospective multicenter surveillance program. *Med. Mycol.* 43(Suppl. 1), S49–S58. doi: 10.1080/13693780400020113
- Morton, C. O., Bouzani, M., Loeffler, J., and Rogers, T. R. (2012). Direct interaction studies between *Aspergillus fumigatus* and human immune cells; what have we learned about pathogenicity and host immunity? *Front. Microbiol.* 3:413. doi: 10.3389/fmicb.2012.00413
- Murdock, B. J., Shreiner, A. B., McDonald, R. A., Osterholzer, J. J., White, E. S., Toews, G. B., et al. (2011). Coevolution of TH1, TH2, and TH17 responses during repeated pulmonary exposure to *Aspergillus fumigatus* conidia. *Infect. Immun.* 79, 125–135. doi: 10.1128/iai.00508-10
- Netea, M. G., Ferwerda, G., Van Der Graaf, C. A., Van Der Meer, J. W., and Kullberg, B. J. (2006). Recognition of fungal pathogens by toll-like receptors. *Curr. Pharm. Des.* 12, 4195–4201. doi: 10.2174/138161206778743538
- Osugi, Y., Vuckovic, S., and Hart, D. N. (2002). Myeloid blood CD11c⁺ dendritic cells and monocyte-derived dendritic cells differ in their ability to stimulate T lymphocytes. *Blood* 100, 2858–2866. doi: 10.1182/blood.V100.8.2858
- Park, S. J., Hughes, M. A., Burdick, M., Strieter, R. M., and Mehrad, B. (2009). Early NK cell-derived IFN- γ is essential to host defense in neutropenic invasive aspergillosis. *J. Immunol.* 182, 4306–4312. doi: 10.4049/jimmunol.0803462
- Park, S. J., and Mehrad, B. (2009). Innate immunity to *Aspergillus* species. *Clin. Microbiol. Rev.* 22, 535–551. doi: 10.1128/cmr.00014-09
- Ramirez-Ortiz, Z. G., and Means, T. K. (2012). The role of dendritic cells in the innate recognition of pathogenic fungi (*A. fumigatus*, *C. neoformans*, and *C. albicans*). *Virulence* 3, 635–646. doi: 10.4161/viru.22295
- Ramirez-Ortiz, Z. G., Specht, C. A., Wang, J. P., Lee, C. K., Bartholomeu, D. C., Gazzinelli, R. T., et al. (2008). Toll-like receptor 9-dependent immune activation by unmethylated CpG motifs in *Aspergillus fumigatus* DNA. *Infect. Immun.* 76, 2123–2129. doi: 10.1128/iai.00047-08
- Rivera, A., Hohl, T. M., Collins, N., Leiner, I., Gallegos, A., Saijo, S., et al. (2011). Dectin-1 diversifies *Aspergillus fumigatus*-specific T cell responses by inhibiting T helper type 1 CD4 T cell differentiation. *J. Exp. Med.* 208, 369–381. doi: 10.1084/jem.20100906
- Romani, L. (2008). Cell mediated immunity to fungi: a reassessment. *Med. Mycol.* 46, 515–529. doi: 10.1080/13693780801971450
- Serbina, N. V., Cherny, M., Shi, C., Bleau, S. A., Collins, N. H., Young, J. W., et al. (2009). Distinct responses of human monocyte subsets to *Aspergillus fumigatus* conidia. *J. Immunol.* 183, 2678–2687. doi: 10.4049/jimmunol.0803398
- Shankar, J. (2013). An overview of toxins in *Aspergillus* associated with pathogenesis. *Int. J. Life Sci. Biotechnol. Pharm. Res.* 2, 16–31.
- Shankar, J., Nigam, S., Saxena, S., Madan, T., and Sarma, P. U. (2004). Identification and assignment of function to the genes of *Aspergillus fumigatus* expressed at 37 degrees C. *J. Eukaryot. Microbiol.* 51, 428–432. doi: 10.1111/j.1550-7408.2004.tb00390.x
- Shao, C., Qu, J., He, L., Zhang, Y., Wang, J., Zhou, H., et al. (2005). Dendritic cells transduced with an adenovirus vector encoding interleukin-12 are a potent vaccine for invasive pulmonary aspergillosis. *Genes Immun.* 6, 103–114. doi: 10.1038/sj.gene.6364167
- Steinbach, W. J., Benjamin, D. K. Jr., Kontoyiannis, D. P., Perfect, J. R., Lutsar, I., et al. (2004). Infections due to *Aspergillus terreus*: a multicenter retrospective analysis of 83 cases. *Clin. Infect. Dis.* 39, 192–198. doi: 10.1086/421950
- Stevens, D. A., Kan, V. L., Judson, M. A., Morrison, V. A., Dummer, S., Denning, D. W., et al. (2000). Practice guidelines for diseases caused by *Aspergillus*. Infectious Diseases Society of America. *Clin. Infect. Dis.* 30, 696–709. doi: 10.1086/313756
- Verweij, P. E., and Brandt, M. E. (2007). *Aspergillus, Fusarium and Other Opportunistic Moniliaceous Fungi*. Washington, DC: ASM Press.
- Vyas, J. M. (2011). The duality of *Aspergillus terreus*: differential immune responses to distinct conidia. *Virulence* 2, 181–184. doi: 10.4161/viru.2.3.16613
- Weaver, C. T., Hatton, R. D., Mangan, P. R., and Harrington, L. E. (2007). IL-17 family cytokines and the expanding diversity of effector T cell lineages. *Annu. Rev. Immunol.* 25, 821–852. doi: 10.1146/annurev.immunol.25.022106.141557
- Werner, J. L., Gessner, M. A., Lilly, L. M., Nelson, M. P., Metz, A. E., Horn, D., et al. (2011). Neutrophils produce interleukin 17A (IL-17A) in a dectin-1- and IL-23-dependent manner during invasive fungal infection. *Infect. Immun.* 79, 3966–3977. doi: 10.1128/IAI.05493-11
- Willment, J. A., and Brown, G. D. (2008). C-type lectin receptors in antifungal immunity. *Trends Microbiol.* 16, 27–32. doi: 10.1016/j.tim.2007.10.012
- Wilson, N. J., Boniface, K., Chan, J. R., McKenzie, B. S., Blumenschein, W. M., Mattson, J. D., et al. (2007). Development, cytokine profile and function of human interleukin 17-producing helper T cells. *Nat. Immunol.* 8, 950–957. doi: 10.1038/ni1497
- Wüthrich, M., Deepe, G. S., Jr., and Klein, B. (2012). Adaptive immunity to fungi. *Annu. Rev. Immunol.* 30, 115–148. doi: 10.1146/annurev-immunol-020711-074958
- Zelante, T., Bozza, S., De Luca, A., D'angelo, C., Bonifazi, P., Moretti, S., et al. (2009). Th17 cells in the setting of *Aspergillus* infection and pathology. *Med. Mycol.* 47(Suppl. 1), S162–S169. doi: 10.1080/13693780802140766
- Zelante, T., De Luca, A., Bonifazi, P., Montagnoli, C., Bozza, S., Moretti, S., et al. (2007). IL-23 and the Th17 pathway promote inflammation and impair anti-fungal immune resistance. *Eur. J. Immunol.* 37, 2695–2706. doi: 10.1002/eji.200737409

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Thakur, Anand, Tiwari, Singh, Tiwary and Shankar. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.