



Extrolites of *Aspergillus fumigatus* and Other Pathogenic Species in *Aspergillus* Section *Fumigati*

Jens C. Frisvad * and Thomas O. Larsen

Section of Eukaryotic Biotechnology, Department of Systems Biology, Technical University of Denmark, Kongens Lyngby, Denmark

OPEN ACCESS

Edited by:

Frederic Lamothe,
Lausanne University Hospital,
Switzerland

Reviewed by:

Vito Valiante,
Leibniz-Institute for Natural Product Research and Infection Biology - Hans Knöll Institute, Germany
William Charles Nierman,
J. Craig Venter Institute, USA
Miroslav Kolarik Kolarik,
Institute of Microbiology,
Czech Republic

***Correspondence:**

Jens C. Frisvad
jcf@bio.dtu.dk

Specialty section:

This article was submitted to
Fungi and Their Interactions,
a section of the journal
Frontiers in Microbiology

Received: 05 August 2015

Accepted: 09 December 2015

Published: 07 January 2016

Citation:

Frisvad JC and Larsen TO (2016)
Extrolites of *Aspergillus fumigatus* and
Other Pathogenic Species in
Aspergillus Section *Fumigati*.
Front. Microbiol. 6:1485.
doi: 10.3389/fmicb.2015.01485

Aspergillus fumigatus is an important opportunistic human pathogen known for its production of a large array of extrolites. Up to 63 species have been described in *Aspergillus* section *Fumigati*, some of which have also been reliably reported to be pathogenic, including *A. felis*, *A. fischeri*, *A. fumigatiaffinis*, *A. fumisynnematus*, *A. hiratsukae*, *A. laciniosus*, *A. lentulus*, *A. novofumigatus*, *A. parafelis*, *A. pseudofelis*, *A. pseudoviridinutans*, *A. spinosus*, *A. thermomutatus*, and *A. udagawae*. These species share the production of hydrophobins, melanins, and siderophores and ability to grow well at 37°C, but they only share some small molecule extrolites, that could be important factors in pathogenicity. According to the literature gliotoxin and other exometabolites can be contributing factors to pathogenicity, but these exometabolites are apparently not produced by all pathogenic species. It is our hypothesis that species unable to produce some of these metabolites can produce proxy-exometabolites that may serve the same function. We tabulate all exometabolites reported from species in *Aspergillus* section *Fumigati* and by comparing the profile of those extrolites, suggest that those producing many different kinds of exometabolites are potential opportunistic pathogens. The exometabolite data also suggest that the profile of exometabolites are highly specific and can be used for identification of these closely related species.

Keywords: *Aspergillus*, gliotoxin, fumagillin, extrolites, proxy-exometabolites

INTRODUCTION

The genus *Aspergillus* comprises 344 species (Samson et al., 2014), and some of these can cause human diseases. *A. fumigatus* is the most important species (Latgé, 1999), but several other species in *Aspergillus* section *Fumigati* have been shown to be pathogenic in humans and animals with an inefficient immune system, including *A. lentulus* (Balajee et al., 2005a; Alhambra et al., 2008; Alcazar-Fuoli et al., 2014; Howard, 2014), *A. fumisynnematus* (Alcazar-Fuoli et al., 2014), *A. fumigatiaffinis* (Alcazar-Fuoli et al., 2014), *A. novofumigatus* (Peláez et al., 2013), *A. felis* (Barrs et al., 2013), *A. fischeri* (Kano et al., 2015), *A. viridinutans* (Vinh et al., 2009a; Coelho et al., 2011; Alcazar-Fuoli et al., 2014), *A. pseudofelis*, *A. pseudoviridinutans*, and *A. parafelis* (Sugui et al., 2014), *A. thermomutatus* (Toskova et al., 2013; Alcazar-Fuoli et al., 2014; Howard, 2014; Khare et al., 2014), *A. laciniosus* (Malejczyk et al., 2013), *A. hiratsukae* (Guarro et al., 2002; Alcazar-Fuoli et al., 2014), *A. spinosus* (Sutton et al., 2002); and *A. udagawae* (Kano et al., 2008; Vinh et al., 2009b; Sugui et al., 2010; Posteraro et al., 2011; Gyotoku et al., 2012; Kano et al., 2013). The taxonomy and identification of the causing Aspergilli is not always clear-cut and some isolates have been

misidentified (Balajee et al., 2005a,b, 2006; Álvarez-Pérez et al., 2014; Howard, 2014). For example pathogenic isolates identified as *A. viridinutans* (Varga et al., 2000; Vinh et al., 2009a; Kano et al., 2013) proved to be *A. felis*, *A. pseudoviridinutans*, *A. parafelis*, or *A. pseudofelis* (Barrs et al., 2013; Novaková et al., 2014; Sugui et al., 2014). *Aspergillus* species in subgenus *Circumdati* have also been reported as pathogenic including *Aspergillus terreus* in section *Terrei*, *A. flavus* in section *Flavi* and *A. tubingensis* in section *Nigri*, *A. persii*, and *A. tanneri* in section *Circumdati*, *A. nidulans* in section *Nidulantes*, (Sugui et al., 2012, 2015; Howard, 2014; Visagie et al., 2014) and *Aspergillus* section *Phialosimplex* [*Ph. caninus* = *Aspergillus caninus* and *Ph. salinarum* = *Aspergillus salinarus* (Sigler et al., 2010; Greiner et al., 2014)]. Small molecule extrolites (secondary metabolites) have been shown to be involved in the infection process (Kamei and Watanabe, 2005; Abad et al., 2010), so it might be expected that the pathogenic Aspergilli produce the same extrolites. In this review we examine whether the closely related pathogenic species in *Aspergillus* section *Fumigati* produce the same extrolites.

ASPERGILLUS TAXONOMY

Since 2011, all ascomycetous species can only have one name (Hawksworth et al., 2011; Hawksworth, 2012; McNeill et al., 2012). All species formerly included in *Dichotomomyces*, *Cristaspora*, *Phialosimplex*, *Polypaecilum*, in addition to *Penicillium inflatum*, have been formally combined into *Aspergillus* (Houbraken et al., 2014; Samson et al., 2014). Furthermore, all species of *Eurotium*, *Emericella*, *Chaetosartorya*, *Fennellia*, *Neocarpenteles*, *Neopetromyces*, *Neosartorya*, *Petromyces*, *Saitoa*, and *Stilbothamnium* have also been transferred to *Aspergillus* (Samson et al., 2014). Ascomata producing species in section *Fumigati* were originally described under the name *Neosartorya* (Samson et al., 2006, 2007), but have now all been transferred to *Aspergillus* (Samson et al., 2014). Several of the species originally thought to produce only the asexual state have later been shown to be able to produce mature ascocarps when crossed with the opposite mating type, for example *A. fumigatus* (O’Gorman et al., 2009) and *A. lentulus* (Swilaiman et al., 2013). Other opportunistically pathogenic species such as *A. flavus* (Horn et al., 2009), *A. tubingensis* (Horn et al., 2013), and *A. terreus* (Samson et al., 2011; Arabatsis and Velegraki, 2013) can also produce mature ascocarps when crossed with the opposite mating type. All species in *Aspergillus* and *Penicillium* have now been placed in the family *Aspergillaceae* (Houbraken and Samson, 2011). Species in *Aspergillus* section *Fumigati* are both phenotypically and genotypically distinct (Raper and Fennell, 1965; Geiser et al., 1998; Hong et al., 2005, 2006, 2008; Katz et al., 2005; Geiser et al., 2007; Samson et al., 2007; Yaguchi et al., 2007). *Aspergillus lentulus* was originally claimed to be a sibling species of *A. fumigatus*, but was later shown to be phenotypically very different from *A. fumigatus*, especially concerning extrolite profiles (Larsen et al., 2007; Tamiya et al., 2015). The species *A. pseudofelis*, *A. parafelis*, and *A. pseudoviridinutans* have not been examined chemically, but they are very close phylogenetically and morphologically to *A. felis* and may be real sibling species with no phenotypic

differences (Sugui et al., 2014). The 63 species listed in **Table 1** are all those that have been described in *Aspergillus* section *Fumigati* and *Neosartorya*, but some of them are not yet available for the scientific community, so their identity and probably synonymy with other species is unknown. Samson et al. (2007) indicated that several species were synonyms of already known species in *Aspergillus* section *Fumigati* and *Neosartorya*. Thus the total number of species in *Fumigati* may be less than 63.

CHEMOTAXONOMY OF ASPERGILLUS SUBGENUS FUMIGATI

Species in subgenus *Fumigati* can produce many different extrolites (Frisvad and Samson, 1990; Samson et al., 2007; Stack et al., 2007; Varga et al., 2007; Frisvad et al., 2009; Sanchez et al., 2012; Kang et al., 2013; Frisvad and Larsen, 2015) of which some are specific to section *Fumigati*, while others are shared with the closely related section *Clavati* and the *Dichotomomyces* clade. *Aspergillus cepii* in the *Dichotomomyces* clade produces gliotoxin, acetylgliotoxin, acetylgliotoxin G, bis(dethio)bis(methylthio)gliotoxin, fiscalin B, xanthocillin X monomethylether, tryptoquivalones, emindole SB, emindole SB β-mannoside, and 27-O-methylasporyzin (Varga et al., 2007; Harms et al., 2014; Rodrigues et al., 2015) possibly in addition to asporyzin A-C, emeniveol, JBIR-03, and asporyergosterol and other sterols (Qiao et al., 2010a,b). The producing strain of the latter exometabolites was probably misidentified as *A. oryzae*, since none of these exometabolites have ever been found in *A. oryzae* (Rank et al., 2012). Apart from some few other shared extrolites with *Aspergillus* species in other sections, most extrolites are unique to section *Fumigati*.

Aspergillus section *Clavati* contains species mostly associated to dung, and have not been reported to cause infections of vertebrate lungs (Varga et al., 2007). Species in *Aspergillus* section *Clavati* produce several bioactive extrolites, but few of these are found in *Aspergillus* section *Fumigati*. Examples of such *Aspergillus* section *Clavati* specific extrolites include patulin, cytochalasin E and K, antafumicins, expansolides, and clavatols, and these extrolites may be important for competition in a dung habitat, rather than in the compost habitats in which species of *Aspergillus* section *Fumigati* thrives. Some similar extrolites are in common between species in *Aspergillus* sections *Fumigati* and *Clavati*, however. Ribotoxins like the sarcins in *Aspergillus* section *Clavati* (Varga and Samson, 2008) are closely related to mitogillin and restrictocin in *Aspergillus* section *Fumigati* (Kao et al., 2001; Schwienbacher et al., 2005; Virág et al., 2014). Furthermore, some tryptoquivalins are produced by species in both *Aspergillus* sections.

Like other filamentous fungi, *A. fumigatus* isolates produce extrolites in a species specific manner (Larsen et al., 2005; Frisvad et al., 2008), but some strains do not produce all the extrolites expected. This weaker exometabolic vigor is most pronounced in isolates directly isolated from patients (Frisvad and Samson, 1990; Tamiya et al., 2015). These isolates are often floccose and less strongly sporulating. However, isolates from natural habitats, such as compost, always sporulate heavily and

TABLE 1 | Species in Aspergillus section Fumigati and their extrolite production (species written in bold are known to be pathogenic to humans and/or other mammals).

Aspergillus arcoverdensis: N.E. (Matsusawa et al., 2015)
Aspergillus assulatus: aszonapyrone A, indole alkaloids and apolar metabolites (Samson et al., 2007)
Aspergillus auratus: helvolic acid (Samson et al., 2007)
Aspergillus aureolus: fiscalins, fumagillin, fumiquinazolines, helvolic acid, pseurotin A, tryptoquivalines, tryptoquivalones, viriditoxin (Samson et al., 2007; Kaur et al., 2013)
Aspergillus australensis: aszonalenins, wortmannins (Samson et al., 2007)
Aspergillus beijingensis: N.E. (Li et al., 1998)
Aspergillus botucatensis (= A. spinosus) (Horie et al., 1995; Samson et al., 2007)
Aspergillus brevipes: roquefortine C, cf. meleagrin, viriditoxin (trace) (Lillehoj and Milburn, 1973; Samson et al., 2007)
Aspergillus brevistipitatus: N.E. (Novaková et al., 2014)
Aspergillus caatingaensis: N.E. (Matsusawa et al., 2014b)
Aspergillus conversis: N.E. (Novaková et al., 2014)
Aspergillus "coreanus" (<i>Neosartorya coreana</i>): aszonalenins (Samson et al., 2007)
Aspergillus delicatus (= <i>A. tatenoi</i>) (Samson et al., 2007)
Aspergillus denticulatus: gliotoxin, viriditoxin (Samson et al., 2007)
Aspergillus duricaulis: asperdurin, asperpentyn, cyclopaldic acid, duricaulic acid, fumagillin, 3-O-methylcyclopolic acid, furochromanols and phthalides, pseurotin A (Brillinger et al., 1978; Achenbach et al., 1982a,b, 1985a,b; Mühlendorf and Achenbach, 1988a,b; Samson et al., 2007)
Aspergillus felis: fumagillin, fumioclavine C, fumitremorgin A and C, helvolic acid, monomethylsulochrin, pyripyropene A, E, O, S, trypacidin (reported as "A. viridinutans", Tamiya et al., 2015, but <i>A. viridinutans</i> has a very different profile of extrolites, and many isolates reported as <i>A. viridinutans</i> have been shown to be <i>A. felis</i> ; Barrs et al., 2013)
Aspergillus fennelliae: asperfuran, aszonalenins, fumioclavines, viridicatumtoxin (Samson et al., 2007)
Aspergillus ferenczii: asperfuran, aszonalenins, fumioclavine, fumigatins, cf. gliotoxin, viridicatumtoxin (Samson et al., 2007)
Aspergillus fischeri: 5-N-acetylardeemin, 5-N-acetyl-15b-didehydroardeemin, 5-N-acetyl-16-hydroxyardeemin, acetylaszonalenin, ardeemin, aszonalenin, aszonapyrone A, B, cottoquinazolin E & F, cyclotryprostatin B, 12a,13a-dihydroxyfumitremorgin C, rel-(8S)-19,20-dihydroxy-8-methoxy-9,18-epifumitremorgin C, fiscalin A, B, C, fischerin, 1-formyl-5-hydroxyazonalenin, fumitremorgin A, B, C, helvolic acid, 6-hydroxyazonalenin, 15b-β-hydroxy-5-N-ardeemin, isoterrein, neofipiperazine A, B, C, neosartorin, nortryptoquivalone, 13-oxofumitremorgin B, pyripyropene A, pyripyrone S, sarcins, sartopyrone B & D, sesterfischeric acid, sesterfischroler, terrein, TR-2, trypacidin, tryptoquivalines, verruculogen (Samson et al., 1990; Wong et al., 1993; Wakana et al., 2006; Samson et al., 2007; Yin et al., 2009; Eamvijarn et al., 2013a; Lee et al., 2013; Gomes et al., 2014; Shan et al., 2014; Sodngam et al., 2014) (as "Xylaria humosa") (Zhang et al., 2014; Zheng et al., 2014; Kaifu et al., 2015; Shan et al., 2015; Ye et al., 2015). There are indications that <i>A. fischeri</i> can also produce fumagillin (Lin et al., 2013; Wiemann et al., 2013)
[fiscalin B, helvolic acid, helvolinic acid, 27-epi-nortryptoquivalone, setosusin, 2-(1-oxo-2-hydroxyethyl)furan, 27-epi-tryptoquivalone, was found in "Corynascus setosus," which is probably an <i>Aspergillus fischeri</i> or alternatively the <i>Corynascus</i> culture was overgrown by <i>A. fischeri</i> ; Fujimoto et al., 1996]
[cladoquinazoline, epi-cladoquinazoline, CS-C, deoxynortryptoquivalone, deoxytryptoquivalone, glyantrypine, 3-hydroxyglyantrypine, norquindoline A, oxoglyantrypine, prelaptin B, quinadoline A, B, tryptoquivalone was reported from a <i>Cladosporium</i> sp., but the culture may have been overgrown with an <i>Aspergillus fischeri</i> ; Peng et al., 2013]
Aspergillus fumigatiaffinis: auranthine, cycloechinuline, fumioclavines, helvolic acid, neosartorin, palitantin, pyripyropene A, E, O, S, tryptoquivalins (Samson et al., 2007; Ola et al., 2014)
Aspergillus fumigatus: Wang compound 1,2, 3, Zhao compound 1, 2, 3, Zuck compound 1,2,3, N-acetyltyramine, asperfumigatin, asperfumin, asperfumoid, azasiprene, bisdechloroegedin, chaetominine, bisdethio(bismethylthio)gliotoxin, brevianamide F, 4-carboxy-5,5'-dihydroxy-3,3'-dimethylidiphenylether, cephalimycin A, B, C, D, 2-chloro-1,3,8-trihydroxy-6-methyl-9-anthrone, cyclo-(Ala-Val), cyclotryptostatin A, B, C, D, cyclo(L-4-hydroxyproline-L-leucine), cyclo(L-4-hydroxyproline-L-phenylalanine), cyclo(L-Pro-L-Pro), cyclo(L-Pro-L-Gly), cyclo(L-Pro-L-Leu), cyclo(L-Pro-L-Pro), cyclo(L-Pro-L-Val), cyclo(L-Val-L-Leu), cyclotryptostatin C, 9-deacetoxyfumigoclavine C, 9-deacetylumigoclavine C, 13-dehydroxycyclotryptostatin C, demethoxyfumitremorgin C, (4S,5S,6S,8S,9S,10R,13R,14S,16S,17Z)-6,16-diacetoxy-25-hydroxy-3,7-dioxy-29-nordammara-1,17(20)-dien-21-oic acid, didehydrobisdethiobis(methylthio)gliotoxin, difructosedianhydride, 1,2-dihydrohelvolic acid, 12,13-dihydroxyfumitremorgin C = TR-3, 2,3-dihydroxy-5-methyl-1,4-benzoquinone, 5,8-dihydroxy-9,12-octadecadienoic acid, 2,6-dihydroxyphenylacetic acid, dimethoxyfumitremorgin C, emodin, emodin 1,6-dimethylether, epoxysuccinic acid, ferrichrome C, festuclavine, FD-889, FK-463, fumagillin, fumagiringillin, fumigefungin, fumioclavine A, B, C,D, E, F, G, H (fumioclavine A reported also from <i>A. tamarii</i> , but this was an <i>A. fumigatus</i> , Janardhanan et al., 1984), fumigatin, fumigatin chlorhydrin, fumigatin oxide, fumigatin quinol, fumigatonin (identity of producer not verified), fumigatoside B, C, and D, fumipyrrole, fumiquinazolin A, B, C, D, E, F, G, J, and K, fumiquinone A and B, fumitremorgin A, B, C, and D, (GERI-BP002-A), fusarinine C, glionitrin A and B, gliotoxin E and G, helvolic acid, helvolinic acid, hexahydroproprenol-18, 19, 20, 21, 22, 23, 24, 3-β-hydroxy-cyclo-L-tryptophyl-L-proline, 2-hydroxy-3-methoxy-5-methyl-1,4-benzoquinone, N-(2-(4-hydroxyphenyl)ethenyl)formamide, 14-hydroxyterezine D, 3-hydroxytoluquinone, 20-hydroxytryptostatin B, isochaetominin, isorhodoptilometrin, LL-S490β, 6-methoxyspirotryprostatin B, 8'-O-methylasterric acid id, 11-O-methylpseurotin, monomethylsulochrin, orsellinic acid, 13-oxofumitremorgin B, 18-oxotryptostatin A, 13-oxo-verruculogen, 14-norpseurotin A, N-prenyl-cyclo-L-tryptophyl-L-proline, pseurotin A, A1, A2, D, F1, F2, pyripyropene A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, and S, questin, RK-95113, cis and trans-ruakuric acid (Cutler et al., 1996, identity of fungus could not be checked), Sch-528647, sphingofungin A, B, C, D, spinulosin, spinulosin hydrate, spinulosin quinol hydrate, spirotryprostatin A, B, C, D, and E, synerazol, terezine D, TR-2, 1,2,3,4-tetrahydroxy-5-methylbenzene, 4,8,10,14-tetramethyl-6-acetoxy-14-[16-acetoxy-19-(20,21-dimethyl)-18-ene]phenanthrene-1-ene-3,7-dione, 3-thiomethyl-cyclo(Ser, Phe), triacetyl fusarinine, tryptostatin A, B, trypacidin, 1,2-seco-tryprostatin, tryptoquivaline F, G, H, I, J, L, M, and N, tryptoquivalin R?, S?, verruculogen (Samson et al., 1990; Land et al., 1993; Cui et al., 1996; Tepsic et al., 1997; Furtado et al., 2005; Han et al., 2007a,b; Samson et al., 2007; Wang et al., 2008; Zhang et al., 2008 (as " <i>A. sydowii</i> "); Frisvad et al., 2009; Zhao et al., 2010; Afifyatullov et al., 2012; Zhang et al., 2012; Cano et al., 2013; Ding et al., 2013; Zhou et al., 2013; Alcazar-Fuoli et al., 2014; Haas, 2014; Kim et al., 2014; Owens et al., 2014; Wiekmann et al., 2014; Xu et al., 2014; Liang et al., 2015; Liu et al., 2015; MacHelein et al., 2015; Tamiya et al., 2015; Xie et al., 2015)

(Continued)

TABLE 1 | Continued

- [*Aspergillus fumigatus* (fungus misidentified): antafumicin A and B, cytochalasin E, expansolide A and B Macías et al., 2003, the strain used was misidentified and was an *Aspergillus clavatus*; isosclerone Li et al., 2014]
- Aspergillus fumisynnematus:** cyclopiazonic acid, fumimycin, neosartorin, pyripyropes (Kwon et al., 2007; Samson et al., 2007)
- Aspergillus galapagensis:* gregatins (Samson et al., 2007)
- Aspergillus hiratsukae:** avenaciolide (Samson et al., 2007)
- Aspergillus huiyanae:* N.E. (Matsusawa et al., 2014a)
- Aspergillus indohii:* N.E. (Horie et al., 2003)
- Aspergillus lacinosus:** aszonalenins, aszonapyrone A and B, 3'-(4-oxoquinazolin-3-yl)spiro[1H-indol-3,5'-oxolane]2,2'-dione, 4(3H)-quinazoline, tryptoquivaline L & T (Samson et al., 2007; Eamvijarn et al., 2013a; Gomes et al., 2014)
- Aspergillus lentulus:** auranthine, cyclopiazonic acid, fumifungin, fumigaclavine A, B, C, fumiquinazoline F or G, monomethylsulochrin, neosartorin, pyripyropes A, E, O, S, sphingofungin A, B, C, D, terrein, trypacidin (Larsen et al., 2007; Samson et al., 2007; Frisvad et al., 2009; Tamiya et al., 2015)
- Aspergillus marvanovae:* apolar indoloterpenes (Hubka et al., 2013)
- Aspergillus multiplicatus:* aszonapyrone A, helvolic acid (Samson et al., 2007)
- Aspergillus neoglaber:* asperpentyn, avenaciolide, glabramycin A, B, C, Mer-NF8054A, Mer-NF8054X, NK-372135A, B, C, sartoryglabrin A, B, C, wortmannins (Ellis et al., 1964; Morino et al., 1994; Samson et al., 2007; Jayasuriya et al., 2009; Kijjoa et al., 2011)
- Aspergillus nishimurae:* Anishidiol, 4-hydroxybenzaldehyde, 4-methylbenzylalcohol, monochaetin (Hosoe et al., 2011)
- Aspergillus novofumigatus:** Dihydroterrein, epi-aszonalenin A, B, C, ent-cycloechinulin, dihydroterrein, fiscalins, helvolic acid, neosartorin, novoamauromin, novobenzomalonin A, B, C, novofumigatamide, novofumigatonin, palitantin, terrein, territrem B (Rank et al., 2006; Samson et al., 2007; Rank et al., 2008; Hosoe et al., 2009; Ishikawa et al., 2010a,b, 2011)
- Aspergillus otanii* = *A. fennelliae* (Takeda et al., 2001; Samson et al., 2007)
- Aspergillus papuensis:* wortmannins (Samson et al., 2007)
- Aspergillus parafelis:** N.E. (Sugui et al., 2014)
- Aspergillus paulistensis* [= *A. spinosus* according to Samson et al. (2007)]: 3'-(4-oxoquinazolin-3-yl)spiro[1H-indol-3,5'-oxolane]2,2'-dione, 4(3H)-quinazoline, sartorypyrone C, tryptoquivaline L (Horie et al., 1995; Gomes et al., 2014)
- Aspergillus pernambucoensis:* N.E. (Matsusawa et al., 2014b)
- Aspergillus primulinus* = *A. quadricinctus* (Samson et al., 2007)
- Aspergillus pseudofelis:** N.E. (Sugui et al., 2014)
- Aspergillus pseudoviridinutans:* N.E. (Sugui et al., 2014)
- Aspergillus quadricinctus:* aszonalenins, PF1223, quinolactacin (Ozoe et al., 2004; Samson et al., 2007)
- Aspergillus qizutongii:** N.E. (Li et al., 1998)
- Aspergillus shendawai:* N.E. (Yaguchi et al., 2010)
- Aspergillus slamensis:* chevalone B, C, 4-dihydroxy-3-methylacetophenone, fiscalin A, C, epi-fiscalin A, C, neofiscalin A, epi-neofiscalin A, 3'-(4-oxoquinazolin-3-yl)spiro[1H-indole-3,5'-oxolane]-2,2'-dione, sartorymensin, tryptoquivaline, tryptoquivaline F, H, L, O (Buttachon et al., 2012; Eamvijarn et al., 2013b)
- Aspergillus similanensis:* chevalone, B, C, E, 6,8-dihydroxy-3,7-dimethylisocoumarin, 6,8-dihydroxy3-methylisocoumarin, p-hydroxybenzaldehyde, 5-hydroxy-8-methyl-2H,6H-pyrano[3,4-g]chromene-2,6-dione, pyripyropes E, S, and T, reticulol, S14-95, similanamide, similanpyrone C (Prompanya et al., 2014, 2015)
- Aspergillus solicola:* aszonalenins, chromanols, tryptoquivalines, tryptoquivalones, wortmannins (Samson et al., 2007, 2014)
- Aspergillus spathulatus:* aszonalenins, xanthocillins (Samson et al., 2007)
- Aspergillus spinosus:** aszonalenins, pseurotins, 2-pyrovoylaminobenzamide, fumigachlorin (Atsumi et al., 1970; Samson et al., 2007)
- Aspergillus stramenius:* avenaciolide, quinolactacin (Samson et al., 2007)
- Aspergillus sublevisporus:* N.E. (Someya et al., 1999)
- Aspergillus takakii* = *A. spinosus* (?) (Samson et al., 2007): acetylaszonalenin, aszonalenin, aszonapyrone A, chevalone B, 6-hydroxymellein, 3'-(4-oxoquinazolin-3-yl)spiro[1H-indole-3,5'-oxolane]-2,2'-dione, tryptoquivaline F, H, L, U (Zin et al., 2015)
- Aspergillus tatenoi:* aszonalenin, aszonapyrone A, B, tatenoi acid (Samson et al., 2007; Yim et al., 2014)
- Aspergillus thermomutatus:** 6-acetylbis(methylthio)gliotoxin, acetylgliotoxin, asperfuran, bisdethiobis(methylthio)gliotoxin, bis-N-norgliovictin, brasiliannamide B, cadinene, CJ-12662?, 3,8-diacyl-4-(3-methoxy-4,5-methyleneoxy)benzyl-7-phenyl-6-oxa-3,8-diazabicyclo[3.2.1]octane, didehydrobisdethiobis(methylthio)gliotoxin, euchevalierine, fiscalins (?), fischerindoline, gliotoxin, helvolic acid, 3-hydroxy-5-methylphenyl-2,4-dihydroxy-6-methylbenzoate, N-methyl-1H-indole-2-carboxamide, neosartorin A, B, C, pseudofischerine, pyripyropes A, E, O, S, 1,2,3,4-tetrahydro-2,3-dimethyl-1,4-dioxopyrazino[1,2-a]indole, 1,2,3,4-tetrahydro-2-methyl-1,3,4-trioxopyrazino[1,2-a]indole, (tryptoquivalin R, S ?) [maybe: misidentified as "Eurotium chevalieri": cadinene, chevalone A, B, C, D, aszonapyrone A, B, euchevalierine, CJ-12662 Kanokmedhakul et al., 2011] (Samson et al., 2007; Eamvijarn et al., 2012; Masi et al., 2013; Xu et al., 2013; Liang et al., 2014)
- Aspergillus tsunodae:* helvolic acid, sartorypyrone A and B (Yaguchi et al., 2010; Eamvijarn et al., 2013a; Gomes et al., 2014)
- Aspergillus tsurutae:* N.E. (Horie et al., 2003)
- Aspergillus turcosus:* aszonalenins, gliotoxin, kotanins (Samson et al., 2007; Hubka et al., 2013)

(Continued)

TABLE 1 | Continued

Aspergillus udagawae: fumagillin, fumigaclavine A and C, fumigatins, fumiquinazolin F or G, helvolic acid, monomethylsulochrin, pyripyropene A, E, trypacidin, tryptoquivalines, tryptoquivalones (Samson et al., 2007; Tamiya et al., 2015)

Aspergillus unilateralis: aszonapyrones, mycophenolic acid (Samson et al., 2007; Hubka et al., 2013)

Aspergillus viridinutans: 4-acetyl-6,8-dihydroxy-5-methyl-2-benzopyran-1-1 A, 13-O-methylviriditin, phomaligin A, SC-28763, SC-30532, semiviriditoxin, viriditoxin, viritin, viriditin, wasabidienone B0 and B1 (Omolo et al., 2000; Samson et al., 2007)

Aspergillus waksmanii: apolar indoloterpenes (Hubka et al., 2013)

Aspergillus wangduanglpii: N.E. (Li et al., 1998)

Aspergillus wyomingensis: N.E. (Novaková et al., 2014)

N.E.: Not Examined.

produce most of the expected species specific extrolites (Frissvad and Samson, 1990; Tepsic et al., 1997; Hong et al., 2010a,b). Production of small molecule extrolites is depending on the growth conditions and the growth media (Nielsen et al., 2011; Frissvad, 2012; Brakhage, 2013), and some of these extrolites may need biological / chemical stimulants of the producing fungus to be expressed (Brakhage and Schroeckh, 2011; Zuck et al., 2011; Netzker et al., 2015).

Being species specific, the difference between the extrolites profiles of different species of *Aspergillus* section *Fumigati* can be used in identification of the species in *Aspergillus* section *Fumigati* as an alternative to sequence-based or MALDI-TOF based identification (Panda et al., 2015), or used together with morphology and physiology in a polyphasic identification approach (Samson et al., 2007). For example *A. fumigatus* can be distinguished from *A. lentulus* by exometabolite profiling (Larsen et al., 2007), MALDI-TOF (Verwer et al., 2014), and sequencing (Balajee et al., 2005a; Samson et al., 2007), but only partially by morphology and Raman spectroscopy (Verwer et al., 2014).

EXTROLITES PRODUCED BY ASPERGILLUS FUMIGATUS AND OTHER PATHOGENIC SPECIES IN FUMIGATI

A. fumigatus has been reported to produce many different extrolites that are bioactive and may contribute to infection in humans and other animals (Amitani et al., 1995; Tomee and Kauffman, 2000; Reeves et al., 2006; Cramer et al., 2009; Abad et al., 2010; Coleman et al., 2011). Melanins are polyketide derived conidium pigments that may have an influence on the infection process (Tsai et al., 1998; Jahn et al., 2000; Tsai et al., 2001; Langfelder et al., 2003). Since all species in *Aspergillus* section *Fumigati* produce green conidia, it is expected that they all contain melanin (Perrin et al., 2007). Another more general small molecule pathogenicity factor is siderophores, of which *A. fumigatus* produces fusarinine C and triacetyl fusarinine C extracellularly (Haas, 2014; Petrik et al., 2014). Furthermore hydrophobins are also present in all species of *Aspergillus* section *Fumigati* (Geiser et al., 1998; Pedersen et al., 2011). These proteins will protect conidia from being recognized by the immune system in mammals (Aimanianda et al., 2009). Other proteins, especially proteases also play a role in the infection process and may be expected to be produced by many pathogenic

species (Tomee and Kauffman, 2000; Abad et al., 2010; Dhingra et al., 2012). Small molecule siderophores are also considered to be important pathogenicity factors, and given the general importance for fungi they can be expected to be produced by all pathogenic species of *Aspergillus* (Fedorova et al., 2008; Abad et al., 2010; Haas, 2014), but probably also by non-pathogenic species.

However, other extrolites are not produced by all species in *Aspergillus* section *Fumigati*. Gliotoxin has long been known to be important for the infection process by inhibiting the immune response, phagocytosis and angiogenesis (Watanabe et al., 2003, 2004; Tsunawaki et al., 2004; Bok et al., 2005; Lewis et al., 2005; Stanzani et al., 2005; Coméra et al., 2007; Sugui et al., 2007; Ben-Ami et al., 2009; Abad et al., 2010). Gliotoxin has been reported from the pathogenic species *A. fumigatus* and *A. thermomutatus*, but also from *A. denticulatus*, *A. ferenczii* and *A. turcosus* (Table 1) the latter three not yet known to be pathogenic. Annotation of the genomes of *A. fumigatus* and *A. fischeri* indicates that the latter species can also produce gliotoxin given the right conditions (Inglis et al., 2013). However, many other *Aspergillus* section *Fumigati* extrolites appear to be involved in pathogenesis. Verruculogen, produced by *A. fumigatus* and *A. fischeri*, modifies electrophysical properties of the human nasal epithelial cells (Khoufache et al., 2007) but is also a potent tremorgen (Land et al., 1993; Kelman et al., 2004). Verruculogen and fumitremorgin C (Rabindran et al., 2000) are produced by *A. fumigatus* and *A. fischeri* (Table 1) in section *Fumigati*. Fumagillin suppresses the immune response, neutrophil function and angiogenesis (Fallon et al., 2010, 2011) and is produced by the pathogenic species *A. felis*, *A. fumigatus*, and *A. udagawae*, but also by species in *Aspergillus* section *Fumigati*, such as *A. aureolus* and *A. viridinutans* that have not been reported as yet to be pathogenic (Table 1). Pseurotin A is an inhibitor of immunoglobulin E and is responding to hypoxia (Schmedemann-Hirschmann et al., 2008; Ishikawa et al., 2009; Vödisch et al., 2011). Pseurotins are produced by the pathogenic *A. fumigatus* and *A. spinosus*, but are also produced by *A. duricalis* and *A. aureolus* (Table 1). Sulochrin inhibits eosinophil activation (Ohashi et al., 1997, 1998) and is produced by four pathogenic species in section *Fumigati*: *A. felis*, *A. fumigatus*, *A. lentulus*, and *A. udagawae* (Table 1). The related asterric acid is produced by the same species and this extrolite inhibits vascular endothelial growth factor induced tube formation (Lee et al., 2013). Another related extrolite is trypacidin, which is cytotoxic (Gauthier

et al., 2012), but was originally isolated as an antiprotozoan metabolite (Balan et al., 1963). The fumiquinazolins are also cytotoxic (Lim et al., 2014), and are produced consistently by *A. fumigatus* (Frisvad et al., 2009). The fumiquinazolines (Takahashi et al., 1995) are produced by the pathogenic *A. fumigatus* and *A. lentulus*, but are also produced by *A. aureolus* (**Table 1**). The chemically similar fiscalins and cottoquinazolins (norfumiquinazolins; Ames and Walsh, 2010; Shan et al., 2015) are produced by *A. fischeri*, indicating that these metabolites are of importance for the competitiveness of these fungi. The pyripyropenes have antiangiogenic activity (Hayashi et al., 2009) and are produced by nearly all the known pathogenic species in section *Fumigati*: *A. fumigatus*, *A. fumigatiaffinis*, *A. fumisynnematus*, *A. lentulus*, *A. thermomutatus*, and *A. udagawae* (**Table 1**). In addition pyripyropens are produced by *A. similansensis*, a species that has not yet been tested for pathogenicity or isolated from any animal tissues.

Helvolic acid has been reported as an antibiotic and antifungal extrolite (Rementeria et al., 2005), but it also has been reported to affect human respiratory epithelium (Amitani et al., 1995) and the metabolism of macrophages (Shinohara et al., 1992). Helvolic acid has been reported from *Aspergillus auratus*, *A. aureolus*, *A. felis*, *A. fischeri*, *A. fumigatiaffinis*, *A. fumigatus*, *A. multiplicatus*, *A. novofumigatus*, *A. thermomutatus*, *A. tsunodae*, and *A. udagawae*. It is upregulated with gliotoxin in *A. fumigatus* (O'Keeffe et al., 2014). Thus helvolic acid may also be a pathogenicity factor, but of the species listed above *A. auratus*, *A. aureolus*, *A. multiplicatus*, and *A. tsunodae* have not been reported as pathogenic. Among bioactive proteins it seems that mitogillin is playing a role in the infection process (Schwienbacher et al., 2005; Abad et al., 2010), but these ribotoxins have not been screened in the other 62 species in *Aspergillus* section *Fumigati*.

Several small molecule extrolites have not yet been claimed to be involved in pathogenesis. The fumigaclavines are produced by the pathogenic species *A. felis*, *A. fumigatus*, *A. fumigatiaffinis*, and *A. lentulus*, but are also produced by *A. fennelliae* and *A. ferenczii* (**Table 1**). Even though these ergot alkaloids are associated with conidiation in *A. fumigatus* (Coyle et al., 1981), their role in animal pathogenesis is unknown. The fumigatins have mostly been found in soil-borne strains of *A. fumigatus* (Frisvad and Samson, 1990), and may rather have a role in competitiveness in compost and soil, than in animal pathogenesis.

PREDICTION OF OTHER POTENTIAL OPPORTUNISTIC PATHOGENIC SPECIES IN ASPERGILLUS SECTION FUMIGATI BASED ON EXTROLITES

Among the 63 species described in *Aspergillus* section *Fumigati*, 17 have until now been reported to be opportunistic pathogens of vertebrate animals (in bold, **Table 1**). Several extrolites have been shown to have a certain role in the infection process, but these extrolites may have a different role in the natural habitats of these fungi, of which plant compost may be the

primary habitat (Latgé, 1999; Abad et al., 2010). It appears that when growing on plant compost these fungi need a certain profile of extrolites (small molecule extrolites and exoproteins), while as vertebrate opportunistic pathogens they may need quite a different profile of extrolites (Abad et al., 2010). For example cellulases would be important in the compost situation (Srivastava et al., 2014; Miao et al., 2015), while hemolysins are probably only important for the vertebrate infection process (Abad et al., 2010). The same would be the case for antifungals and antibiotics, especially anti-streptomycte metabolites, as *A. fumigatus* and other members of *Aspergillus* section *Fumigati* are thermotolerant / thermophilic species competing with other thermotolerant and thermophilic species of fungi and bacteria (Langarica-Fuentes et al., 2014). Several species, such as *A. assulatus*, *A. australensis*, *A. brevipes*, *A. "coreanus"*, *A. duricaulis*, *A. fennelliae*, *A. galapagensis*, *A. neoglaber*, *A. marvanovae*, *A. nishimurae*, *A. papuensis*, *A. quadricinctus*, *A. solicola*, *A. spathulatus*, *A. tatenoi*, *A. unilateralis*, *A. viridinutans*, and *A. waksmanii* produce few if any of the extrolites suspected to play a role in the infection process, and so would not be predicted to be potential opportunistic pathogens of vertebrates. Some species, such as *A. auratus*, *A. denticulatus*, *A. similansensis*, *A. tsunodae*, and *A. turcosus* only produce one of the extrolites believed to play a role in pathogenesis, and may or may not be prospective vertebrate pathogens. Finally *A. aureolus*, *A. ferenczii*, and *A. siamensis* produce several of the extrolites potentially involved in pathogenesis, and thus may be predicted to be potential opportunistic vertebrate pathogens.

The many extrolites that have been suspected to be pathogenicity factors and are produced by species in *Aspergillus* section *Fumigati* are biosynthetically derived from polyketides, amino acids, terpenes, shikimic acid or are of mixed biosynthetic origin. The formula of some of the most important extrolites common in *Aspergillus* section *Fumigati* are shown in **Figure 1**. Some of the extrolites are not produced in the same patterns in different species in *Aspergillus* section *Fumigati*. While *A. fumigatus* produces fumiquinazolins A–G, J, and K, *A. fischeri* produces the related norfumiquinazolins (Shan et al., 2015). These extrolites may have the same function, even though they are chemically somewhat different. Whether such proxy-extrolites have the same function for pathogenicity in vertebrates is unknown. It is known, however, that other opportunistic pathogenic aspergilli in other sections of *Aspergillus* produce secondary metabolites that are biosynthetically and functionally closely related. While *A. fumigatus*, *A. thermomutatus*, and other species in section *Fumigati* produce gliotoxin, *A. flavus* in *Aspergillus* section *Flavi* can produce aspirochlorine and *A. terreus* in *Aspergillus* section *Terrei* can produce acetylalaranotin (Frisvad and Larsen, 2015). While not identical to gliotoxin, these epidithiodioxopiperazines could be predicted to play a role in pathogenicity of *A. flavus* and *A. terreus*. The reports that *A. niger*, *A. flavus*, and *A. terreus* could produce gliotoxin (Lewis et al., 2005; Kupfahl et al., 2008) have not been confirmed (Samson et al., 2011; Varga et al., 2011a,b).

Close phylogenetic relationships seem to be less suited pathogenicity predictors. For example *A. viridinutans* seems to be non-pathogenic, while the closely related *A. felis* is pathogenic




FIGURE 1 | Structures of the most important extrolites from *Aspergillus* section *Fumigati* potentially involved in pathogenesis.

(Barrs et al., 2013; Novaková et al., 2014; Sugui et al., 2014). Good growth at 37°C also seems to be a contributing factor to pathogenicity, and for example *A. brevis*, *A. duricaulis* and *A. viridinutans* grow relatively poorly at 37°C, and in addition are not considered potentially pathogenic *Aspergillus* species in section *Fumigati*, based on extrolite evidence and absence of reports of pathogenicity. However, while there are many data on the involvement of exometabolites for *A. fumigatus* (Abad et al., 2010), there are few data on production of exoproteins for other opportunistic pathogenic species such as *A. thermomutatus*.

Genome sequencing and systematic comparison of the genomes and transcriptomes of other members of *Aspergillus* section *Fumigati* may help in predicting which pathogenicity factors are especially important (Galaghan et al., 2005; Nierman et al., 2005; Wortman et al., 2006; Fedorova et al., 2008; McDonagh et al., 2008; Chooi et al., 2013; Inglis et al., 2013; Cerqueira et al., 2014; Kusuya et al., 2015; Lind et al., 2015). These data should be compared to phenotypic data such as profiles of

large and small molecule extrolites, growth temperatures, carbon dioxide tolerance etc.

Altogether, approximately one third of the species in *Aspergillus* section *Fumigati* are common pathogenic species, one third are rare species of unknown pathogenicity and one third are predicted to be non-pathogenic, based on their production of relatively few exometabolites. Exometabolite pathogenicity factors found in the successful opportunistic pathogenic fungus *A. fumigatus* may have proxy-exometabolites with the same function in other species in that section, but also in less closely related pathogenic Aspergilli, especially species from sections *Nigri*, *Terrei*, *Circumdati*, and *Flavi*.

ACKNOWLEDGMENTS

We thank Agilent technologies for Agilent Thought Leader Award # 2871 and the Novo Nordisk Foundation for grant NNF 13OC0005201.

REFERENCES

- Abad, A., Fernández-Molina, J. V., Bikandi, J., Ramírez, A., Margareto, J., Sendino, J., et al. (2010). What makes *Aspergillus fumigatus* a successful pathogen? Genes and molecules involved in invasive aspergillosis. *Rev. Iberoam. Micol.* 27, 155–182. doi: 10.1016/j.riam.2010.10.003
- Achenbach, H., Mühlenfeld, A., and Brillinger, G. U. (1985a). Stoffwechselprodukte von mikroorganismen. XXX. phthalide und chromanole aus *Aspergillus duricaulis*. *Liebigs Annalen der Chemie* 1985, 1596–1628.
- Achenbach, H., Mühlenfeld, A., Kohl, W., and Brillinger, G. U. (1985b). Stoffwechselprodukte von mikroorganismen, XXXI [1]. Duricaulinsäure, ein neuer Naturstoff vom phthalimidin-Typ aus *Aspergillus duricaulis*. *Z. Naturfors.* 40b, 1219–1225.
- Achenbach, H., Mühlenfeld, A., Weber, B., and Brillinger, G. U. (1982b). Highly substituted chromanols of *Aspergillus duricaulis*. *Tetrahedron Lett.* 23, 4659–4660.
- Achenbach, H., Mühlenfeld, A., Weber, B., Kohl, W., and Brillinger, G. U. (1982a). Stoffwechselprodukte von mikroorganismen. XXVII [1]. Cyclopaldsäure und 3-O-methylcyclopolsäure, zwei antibiotisch worksame Substanzen aus *Aspergillus duricaulis*. *Z. Naturfors.* 37b, 1091–1097.
- Afiyatullov, S. S., Zhuravleva, O. I., Antonov, A. S., Kalinovsky, A. I., Pivkin, M. V., Mentinskaya, E. S., et al. (2012). New metabolites from the marine-derived fungus *Aspergillus fumigatus*. *Nat. Prod. Commun.* 7, 497–500.
- Aimanianda, V., Bayry, J., Bozza, S., Kniemeyer, O., Perrucio, K., Elluru, S. R., et al. (2009). Surface hydrophobin prevents immune recognition of airborne fungal spores. *Nature* 460, 1117–1121. doi: 10.1038/nature08264
- Alcazar-Fuoli, L., Cairns, T., Lopen, J. F., Zonja, B., Pérez, J., Barceló, D., et al. (2014). A modified recombining protocol for the genetic manipulation of gene clusters in *Aspergillus fumigatus*. *PLoS ONE* 9:e111875. doi: 10.1371/journal.pone.0111875
- Alhambra, A., Catalán, M., Moragues, M. D., Brena, S., Pontón, J., Montejo, J. C., et al. (2008). Isolation of *Aspergillus lentulus* in Spain from a critically ill patient with chronic obstructive pulmonary disease. *Rev. Iberoam. Micol.* 25, 246–249. doi: 10.1016/S1130-1406(08)70058-5
- Álvarez-Pérez, S., Mellado, E., Serrano, P., Blanco, J. L., García, M. E., Kwon, M., et al. (2014). Polyphasic characterization of fungal isolates from a published case of invasive aspergillosis reveals misidentification of *Aspergillus felis* as *Aspergillus viridinutans*. *J. Med. Microbiol.* 63, 617–619. doi: 10.1099/jmm.0.068502-0
- Ames, B. D., and Walsh, C. T. (2010). Anthranilate-activating modules from fungal non-ribosomal peptide assembly lines. *Biochemistry* 49, 3351–3365. doi: 10.1021/bi100198y
- Amitani, R., Taylor, G., Elezis, E. N., Llewellyn-Jones, C., Mitchell, J., Kuza, F., et al. (1995). Purification and characterization of factors produced by *Aspergillus fumigatus* which affect human ciliated respiratory epithelium. *Infect. Immun.* 63, 3266–3271.
- Arabatis, M., and Velegakis, A. (2013). Sexual reproduction cycle in the opportunistic human pathogen *Aspergillus terreus*. *Mycologia* 105, 71–79. doi: 10.3852/11-426
- Atsumi, K., Takada, M., Mizuno, K., and Ando, T. (1970). Fumigachlorin, a new antifungal antibiotic. *J. Antibiot.* 23, 223–224. doi: 10.7164/antibiotics.23.223
- Balajee, S. A., Gribskov, J., Brandt, M., Ito, J., Fothergill, A., and Marr, K. A. (2005b). Mistaken identity: *Neosartorya pseudofischeri* and its anamorph masquerading as *Aspergillus fumigatus*. *J. Clin. Microbiol.* 43, 5996–5999. doi: 10.1128/JCM.43.12.5996-5999.2005
- Balajee, S. A., Gibbskov, J. L., Hanley, E., Nickle, C., and Marr, K. A. (2005a). *Aspergillus lentulus* sp. nov., a new sibling species of *A. fumigatus*. *Eukaryotic Cell* 4, 625–632. doi: 10.1128/EC.4.3.625-632.2005
- Balajee, S. A., Nickle, D., Varga, J., and Marr, K. A. (2006). Molecular studies reveal frequent misidentification of *Aspergillus fumigatus* by morphotyping. *Eukaryotic Cell* 5, 1705–1712. doi: 10.1128/EC.00162-06
- Balan, J., Ebringer, L., Nemec, P., Kovac, S., and Dobias, J. (1963). Antiprotozoal antibiotics. II. Isolation and characterization of trypacidin, a new antibiotic, active against *Trypanosoma cruzi* and *Toxoplasma gondii*. *J. Antibiot.* 16, 157–160.
- Barris, V. R., van Doorn, T. M., Houbraken, J., Kidd, S. E., Martin, P., Pinheiro, M. D., et al. (2013). *Aspergillus felis* sp. nov., an emerging agent of aspergillosis in humans, cats, and dogs. *PLOS ONE* 8:e64871. doi: 10.1371/journal.pone.0064871
- Ben-Ami, R., Lewis, R. E., Leventakos, K., and Kontoyiannis, D. P. (2009). *Aspergillus fumigatus* inhibits angiogenesis through the production of gliotoxin and other secondary metabolites. *Blood* 114, 5393–5399. doi: 10.1182/blood-2009-07-231209
- Bok, J. W., Chung, D., Balajee, S. A., Marr, K. A., Andes, D., Nielsen, K. F., et al. (2005). Gliz, a transcriptional regulator of gliotoxin biosynthesis, contributes to *Aspergillus fumigatus* virulence. *Infect. Immun.* 74, 6761–6768. doi: 10.1128/IAI.00780-06
- Brakhage, A. A. (2013). Regulation of fungal secondary metabolism. *Nat. Rev. Microbiol.* 11, 21–32. doi: 10.1038/nrmicro2916
- Brakhage, A. A., and Schroeck, V. (2011). Fungal secondary metabolites – strategies to activate silent gene clusters. *Fungal Genet. Biol.* 48, 15–22. doi: 10.1016/j.fgb.2010.04.004
- Brillinger, G. U., Heberle, W., Weber, B., and Achenbach, H. (1978). Metabolic products of microorganisms. 167. Cyclopaldic acid from *Aspergillus duricaulis*. 1. Production, isolation and biological properties. *Arch. Microbiol.* 116, 245–252. doi: 10.1007/BF00417847
- Buttachon, S., Chandrapatya, A., Manoch, L., Silva, A., Gales, L., Bruylants, C., et al. (2012). Sartorymensin, a new indole alkaloid, and new analogues of tryptoquinoline and fiscalins produced by *Neosartorya siamensis* (KUFC 6349). *Tetrahedron* 68, 3253–3262. doi: 10.1016/j.tet.2012.02-024
- Cano, P. M., Jamin, E. L., Tadrist, S., Bourdau'hui, P., Pean, M., Debrauwer, L., et al. (2013). New untargeted metabolite profiling combining mass spectrometry and isotopic labelling: application to *Aspergillus fumigatus* grown on wheat. *Anal. Chem.* 85, 8412–8420. doi: 10.1021/ac401872f
- Cerdeira, G. C., Arnaud, M. B., Inglis, D. O., Skrzypek, M. S., Binkley, G., Simison, M., et al. (2014). The *Aspergillus* genome database: multispecies curation and incorporation of RNA-Seq data to improve structural gene annotations. *Nucleic Acids Res.* 42, D705–D710. doi: 10.1093/nar/gkt1029
- Chooi, Y.-H., Fang, J., Liu, H., Filler, S. G., Wang, P., and Tang, Y. (2013). Genome mining of a prenylated and immunosuppressive polyketide from pathogenic fungi. *Org. Lett.* 15, 780–783. doi: 10.1021/ol303435y
- Coelho, D., Silva, S., Vale-Silva, L., Gomes, H., Pinto, E., Sarmento, A., et al. (2011). *Aspergillus viridinutans*: an agent of adult chronic invasive aspergillosis. *Med. Mycol.* 49, 755–759. doi: 10.3109/13693786.2011.556672
- Coleman, J. J., Ghosh, S., Okoli, I., and Mylonakis, E. (2011). Antifungal activity of microbial secondary metabolites. *PLoS ONE* 6:e25321. doi: 10.1371/journal.pone.0025321
- Coméra, C., André, K., Lafitte, J., Collet, X., Galtier, P., and Maridonneau-Parini, I. (2007). Gliotoxin from *Aspergillus fumigatus* affects phagocytosis and the organization of the actin cytoskeleton by distinct signaling pathways in human neutrophils. *Microb. Infect.* 9, 47–54. doi: 10.1016/j.micinf.2006.10.009
- Coyle, C. M., Kenaley, S. C., Rittenour, W. R., and Panaccione, D. G. (1981). Association of ergot alkaloids with conidiation in *Aspergillus fumigatus*. *Mycologia* 99, 804–811. doi: 10.3852/mycologia.99.6.804
- Cramer, R. A., Keats Schwab, E. Jr., and Keller, N. P. (2009). “Genetic regulation of *Aspergillus* secondary metabolites and their role in pathogenesis,” in *Aspergillus fumigatus and Aspergillosis* eds J. P. Latgé and W. J. Steinbach (Washington, DC: ASM Press), 185–199. ISBN 978-1-55581-438-0
- Cui, C. B., Kakeya, H., Okada, G., Onose, R., and Osada, H. (1996). Novel mammalian cell cycle inhibitor, tryprostatins A, B and other diketopiperazines produced by *Aspergillus fumigatus*. II. Physico-chemical properties and structures. *J. Antibiot.* 49, 527–533. doi: 10.7164/antibiotics.49.527
- Cutler, H. G., Lauren, D. R., Wilkins, A. L., Holland, P. T., Hill, R. A., and Dugan, F. M. (1996). Ruakuric acid: a natural product from *Aspergillus fumigatus*. *Phytochemistry* 43, 209–214. doi: 10.1016/0031-9422(96)00224-5
- Dhingra, S., Andes, D., and Calvo, A. M. (2012). VeA regulated conidiation, Gliotoxin production, and protease activity in the opportunistic human pathogen *Aspergillus fumigatus*. *Eukaryotic Cell* 11, 1531–1543. doi: 10.1128/EC.00222-12
- Ding, G.-Z., Liu, J., Wang, J.-M., Faing, L., and Yi, S.-S. (2013). Secondary metabolites from the endophytic fungi *Penicillium polonicum* and *Aspergillus fumigatus*. *J. Asian Nat. Prod. Res.* 15, 446–452. doi: 10.1080/10286020.2013.780349
- Eamvijarn, A., Gomes, N. M., Dethou, T., Buaruang, J., Manoch, L., Manoch, L., et al. (2013a). Bioactive meroterpenes and indole alkaloids from the

- soil fungus *Neosartorya fischeri* (KUFC 6344), and the marine-derived fungi *Neosartorya laciniosa* (KUFC 7896) and *Neosartorya tsunodae* (KUFC 9213). *Tetrahedron* 69, 8583–8591. doi: 10.1016/j.tet.2013.07.078
- Eamvijarn, A., Kijjoa, A., Bruyère, C., Mathieu, V., Manoch, L., Lefranc, F., et al. (2012). Secondary metabolites from a culture of the fungus *Neosartorya pseudofischeri* and their *in vitro* cytostatic activity in human cancer cells. *Planta Med.* 78, 1767–1776. doi: 10.1055/s-0032-1315301
- Eamvijarn, A., Manoch, L., Chamswarn, C., Piasai, O., Visarathanonth, N., Langsa-ard, J. J., et al. (2013b). *Aspergillus siamensis* sp. nov. from soil in Thailand. *Mycoscience* 54, 401–405. doi: 10.1016/j.myc.2013.01.005
- Ellis, J. J., Stodola, F. H., Vesonder, R. F., and Glass, C. A. (1964). $C_{15}H_{22}O_4$ compound produced by the fungus *Aspergillus fischeri* var. *glaber*. *Nature* 203:1382. doi: 10.1038/2031382a0
- Fallon, J. P., Reeves, E. P., and Kavanagh, K. (2010). Inhibition of neutrophil function following exposure to the *Aspergillus fumigatus* toxin fumagillin. *J. Med. Microbiol.* 59, 625–633. doi: 10.1099/jmm.0.018192-0
- Fallon, J. P., Reeves, E. P., and Kavanagh, K. (2011). The *Aspergillus fumigatus* toxin fumagillin suppresses the immune response of *Galleria mellonella* larvae by inhibiting the action of haemocytes. *Microbiology* 157, 1481–1488. doi: 10.1099/mic.0.043786-0
- Fedorova, N. D., Khaldi, N., Joarder, V. S., Maiti, R., Amedeo, P., Anderson, M. J., et al. (2008). Genomic islands in the pathogenic filamentous fungus *Aspergillus fumigatus*. *PLoS Genet.* 4:e1000046. doi: 10.1371/journal.pgen.1000046
- Frisvad, J. C. (2012). "Media and growth conditions for induction of secondary metabolites," in *Fungal Secondary Metabolism: Methods and Protocols* eds N. P. Keller and G. Turner (New York, NY: Humana Press), 47–58.
- Frisvad, J. C., Andersen, B., and Thrane, U. (2008). The use of secondary metabolite profiling in fungal taxonomy. *Mycol. Res.* 112, 231–240. doi: 10.1016/j.mycres.2007.08.018
- Frisvad, J. C., and Larsen, T. O. (2015). Chemodiversity in the genus *Aspergillus*. *Appl. Microbiol. Biotechnol.* 99, 7859–7877. doi: 10.1007/s00253-015-6839-z
- Frisvad, J. C., Rank, C., Nielsen, K. F., and Larsen, T. O. (2009). Metabolomics of *Aspergillus fumigatus*. *Med. Mycol.* 47, S53–S71. doi: 10.1080/13693780802307720
- Frisvad, J. C., and Samson, R. A. (1990). "Chemotaxonomy and morphology of *Aspergillus fumigatus* and related species," in *Modern Concepts in Penicillium and Aspergillus Classification* eds R. A. Samson and J. I. Pitt (New York, NY: Plenum Press), 201–208.
- Fujimoto, H., Negishi, E., Yamaguchi, K., Nishi, N., and Yamazaki, M. (1996). Isolation of new tremorgenic metabolites from an Ascomycete, *Corynascus setosus*. *Chem. Pharm. Bull.* 44, 1843–1848. doi: 10.1248/cpb.44.1843
- Furtado, N. A. J. C., Pupo, M. T., Carvalho, I., Campo, V. L., Duarte, M. C. T., and Bastos, J. K. (2005). Diketopiperazines produced by an *Aspergillus fumigatus* Brazilian strain. *J. Braz. Chem. Soc.* 16, 1448–1453. doi: 10.1590/S0103-50532005000800026
- Galaghan, J. E., Calvo, S. E., Cuomo, C., Ma, L. J., Wortman, J. R., Batzoglou, S., et al. (2005). Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* 438, 1105–1115. doi: 10.1038/nature04341
- Gauthier, T., Wang, X., Dos Santos, J. S., Fysikopoulos, A., Tadrist, S., Canlet, C., et al. (2012). Trypacidin, a spore-borne toxin from *Aspergillus fumigatus*, is cytotoxic to lung cells. *PLoS ONE* 7:e29906. doi: 10.1371/journal.pone.0029906
- Geiser, D. M., Frisvad, J. C., and Taylor, J. W. (1998). Evolutionary relationships in *Aspergillus* section *Fumigati* inferred from partial beta-tubulin and hydrophobin DNA sequences. *Mycologia* 90, 832–846. doi: 10.2307/3761325
- Geiser, D. M., Klich, M. A., Frisvad, J. C., Peterson, S. W., Varga, J., and Samson, R. A. (2007). The current status of species recognition and identification in *Aspergillus*. *Stud. Mycol.* 59, 1–10. doi: 10.3114/sim.2007.59.01
- Gomes, N. M., Bessa, L. J., Buttacon, S., Costa, P. M., Buaruang, J., Dethoup, T., et al. (2014). Antibacterial and antibiofilm activities of tryptoquinolines and meroterpenes isolated from the marine-derived fungi *Neosartorya paulistensis*, *N. laciniosa*, *N. tsunodae*, and the soil fungi *N. fischeri* and *N. siamensis*. *Mar. Drugs* 12, 822–839. doi: 10.3390/MD12020822
- Greiner, K., Peršoh, D., Weig, A., and Rambold, G. (2014). *Phialosimplex salinarum*, a new species of *Eurotiomycetes* from a hypersaline habitat. *IMA Fungus* 5, 161–172. doi: 10.5598/imapfungus.2014.05.02.01
- Guarro, J., Kallas, E. G., Godoy, P., Karenina, A., Gene, J., Stchigel, A., et al. (2002). Cerebral aspergillosis caused by *Neosartorya hiratsukae* infection. *Emerging Infect. Dis.* 8, 989–991. doi: 10.3201/eid0809.020073
- Gyotoku, H., Izumikawa, K., Ikeda, H., Takazono, T., Morinaga, Y., Nakamura, S., et al. (2012). A case of bronchial aspergillosis caused by *Aspergillus udagawae* and its mycological features. *Med. Mycol.* 50, 631–636. doi: 10.3109/13693786.2011.639036
- Haas, H. (2014). Fungal siderophore metabolism with a focus on *Aspergillus fumigatus*. *Nat. Prod. Rep.* 31, 1266–1276. doi: 10.1039/C4NP00071D
- Han, X., Xu, X., Cui, C., and Gu, Q. (2007a). Alkaloidal compounds produced by a marine-derived fungus, *Aspergillus fumigatus* H1-04, and their antitumor activities. *Chin. J. Med. Chem.* 17, 232–237.
- Han, X., Xu, X., Cui, C., and Gu, Q. (2007b). Diketopiperazines produced by a marine-derived fungus, *Aspergillus fumigatus* H1-04, and their antitumor activities. *Chin. J. Med. Chem.* 17, 155–159.
- Harms, H., Rempel, V., Kehraus, S., Kaiser, M., Hufendick, P., Müller, C. E., et al. (2014). Indoloterpenes from a marine-derived fungal strain of *Dichotomomyces cepii* with an antagonistic activity at GPR18 and cannabinoid receptors. *J. Nat. Prod.* 77, 673–677. doi: 10.1021/np400850g
- Hawksworth, D. L. (2012). Managing and coping with names of pleomorphic fungi in a period of transition. *IMA Fungus* 3, 15–24. doi: 10.5598/imapfungus.2012.03.01.03
- Hawksworth, D. L., Crous, P. W., Redhead, S. A., Reynolds, D. R., Samson, R. A., Seifert, K. A., et al. (2011). The Amsterdam declaration on fungal nomenclature. *IMA Fungus* 2, 105–112. doi: 10.5598/imapfungus.2011.02.01.14
- Hayashi, A., Artai, M., Fujita, M., and Kobayashi, M. (2009). Pyripyropens, fungal sesquiterpenes conjugated with alpha-pyrone and pyridine moieties, exhibits anti-angiogenicity against human umbilical vein endothelial cells. *Biol. Pharm. Bull.* 32, 1261–1265. doi: 10.1248/bpb.32.1261
- Hong, S. B., Cho, H. S., Shin, H. D., Frisvad, J. C., and Samson, R. A. (2006). New *Neosartorya* species isolated from soil in Korea. *Int. J. Syst. Evol. Microbiol.* 56, 439–442. doi: 10.1099/ijss.0.63980-0
- Hong, S. B., Go, S. J., Shin, H. D., Frisvad, J. C., and Samson, R. A. (2005). Polyphasic taxonomy of *Aspergillus fumigatus* and related species. *Mycologia* 97, 1316–1329. doi: 10.3852/mycologia.97.6.1316
- Hong, S. B., Kim, D. H., Park, I. C., Choi, Y. J., Shin, H. D., and Samson, R. A. (2010a). Re-identification of *Aspergillus fumigatus sensu lato* based on a new concept of species delimitation. *J. Microbiol.* 48, 607–615. doi: 10.1007/s12275-010-0084-z
- Hong, S. B., Kim, D.-H., Park, I. C., Samson, R. A., and Shin, H. D. (2010b). Isolation and identification of *Aspergillus* section *Fumigati* strains from arable soil in Korea. *Mycobiology* 38, 1–6. doi: 10.4489/MYCO.2010.38.1.001
- Hong, S. B., Shin, H. D., Hong, J., Frisvad, J. C., Nielsen, P. V., et al. (2008). New taxa of *Neosartorya* and *Aspergillus* in *Aspergillus* section *Fumigati*. *Antonie van Leeuwenhoek* 93, 87–98. doi: 10.1007/s10482-007-9183-1
- Horie, Y., Abliz, P., Fukushima, K., Okada, K., and Takaki, G. M. C. (2003). Two new species of *Neosartorya* from Amazonian soil, Brazil. *Mycoscience* 44, 397–402. doi: 10.1007/S10267-003-0132-1
- Horie, Y., Miyaji, M., Nishimura, K., Franco, M. F., and Colho, K. I. R. (1995). Two new species of *Neosartorya* from Brazilian soil. *Mycoscience* 36, 159–165. doi: 10.1007/BF02268552
- Horn, B. W., Moore, G. G., and Carbone, I. (2009). Sexual reproduction in *Aspergillus flavus*. *Mycologia* 101, 174–183. doi: 10.3852/10-115
- Horn, B. W., Olarte, R. A., Peterson, S. W., and Carbone, I. (2013). Sexual reproduction in *Aspergillus tubingensis* from section *Nigri*. *Mycologia* 105, 1153–1163. doi: 10.3852/13-101
- Hosoe, T., Mori, N., Kamano, K., Itabashi, T., Yaguchi, T., and Kawai, K. (2011). A new antifungal yellow pigment from *Aspergillus nishimurae*. *J. Antibiot.* 64, 211–212. doi: 10.1038/ja.2010.132
- Hosoe, T., Moriyama, H., Wakana, D., Itabashi, T., Kawai, K., Yaguchi, T., et al. (2009). Inhibitory effects of dihydroterrein and terrain isolated from *Aspergillus novofumigatus* on platelet aggregation. *Mycotoxins* 59, 75–82. doi: 10.2520/myco.59.75
- Houbraek, J., de Vries, R. P., and Samson, R. A. (2014). Modern taxonomy of biotechnologically important *Aspergillus* and *Penicillium* species. *Adv. Appl. Microbiol.* 86, 199–249. doi: 10.1016/B978-0-12-800262-9.00004-4

- Houbraken, J., and Samson, R. A. (2011). Phylogeny of *Penicillium* and the segregation of *Trichocomaceae* into three families. *Stud. Mycol.* 70, 1–51. doi: 10.3114/sim.2011.70.01
- Howard, S. J. (2014). Multiresistant aspergillosis due to cryptic species. *Mycopathologia* 178, 427–433. doi: 10.1007/s11046-014-9774-0
- Hubka, V., Peterson, S. W., Frissvad, J. C., Yaguchi, T., Kubátová, A., Kolaøik, M., et al. (2013). *Aspergillus waksmanii* sp. nov. and *Aspergillus marvanova* sp. nov., two new closely related species in section *Fumigati*. *Int. J. Syst. Evol. Microbiol.* 63, 763–789. doi: 10.1099/ij.s.0.047076-0
- Inglis, D. O., Binkley, J., Skrzypek, M. S., Arnaud, M. B., Cerqueira, G. C., Shah, P., et al. (2013). Comprehensive annotation of secondary biosynthetic genes and gene clusters of *Aspergillus nidulans*, *A. fumigatus*, *A. niger* and *A. oryzae*. *BMC Microbiol.* 13:91. doi: 10.1186/1471-2180-13-91
- Ishikawa, K., Hosoe, T., Itabashi, T., Sato, F., Wachi, H., Nagase, H., et al. (2011). Quinazolinobenzodiazepine derivatives, novobenzomalvins A-C: Fibronectin expression regulators from *Aspergillus novofumigatus*. *Sci. Pharm.* 79, 937–950. doi: 10.3797/scipharm.1106-21
- Ishikawa, K., Hosoe, T., Itabashi, T., Takizawa, K., Yaguchi, T., and Kawai, K. (2010b). Novofumigatamide, new cyclic tripeptide from *Aspergillus novofumigatus*. *Heterocycles* 81, 2143–2148. doi: 10.3987/COM-10-12005
- Ishikawa, K., Hosoe, T., Itabashi, T., Wakana, D., Takizawa, K., Yuguchi, T., et al. (2010a). Novoamauromine and ent-cycloechinulin: two new diketopiperazine derivatives from *Aspergillus novofumigatus*. *Chem. Pharm. Bull.* 58, 717–719. doi: 10.1248/cpb.58.717
- Ishikawa, M., Nionomiya, T., Akabane, H., Kushida, N., Tsujiuchi, G., Ohyama, M., et al. (2009). Pseurotin A and its analogues as inhibitors of immunoglobulin E production. *Bioorg. Med. Chem. Lett.* 19, 1457–1460. doi: 10.1016/j.bmcl.2009.01.029
- Jahn, B., Boukhallouk, F., Lotz, J., Langfelder, K., Wanner, G., and Brakhage, A. A. (2000). Interaction of human phagocytes with pigmentless *Aspergillus* conidia. *Infect. Immun.* 68, 3736–3739. doi: 10.1128/IAI.68.6.3736-3739.2000
- Janardhanan, K. K., Satta, A., and Husain, A. (1984). Production of fumigaclavine by *Aspergillus tamarii* Kita. *Can. J. Microbiol.* 30, 247–250. doi: 10.1139/m84-036
- Jayasuriya, H., Zink, D., Basilio, A., Vicente, F., Collado, J., Bills, G., et al. (2009). Discovery and antibacterial activity of glabramycin A-C from *Neosartorya glabra* by an antisense strategy. *J. Antibiot.* 62, 265–269. doi: 10.1038/ja.2009.26
- Kaifuchi, S., Mori, M., Nonaka, K., Musauma, R., Omura, S., Shiomi, K., et al. (2015). Sartopyrone D: a new NADH-fumarate reductase inhibitor produced by *Neosartorya fischeri* FO-5897. *J. Antibiot.* 68, 403–405. doi: 10.1038/ja.2014.167
- Kamei, K., and Watanabe, A. (2005). *Aspergillus* mycotoxins and their effects on the host. *Med. Mycol.* 43, S95–S99. doi: 10.1080/13693780500051547
- Kang, D., Son, G. H., Park, H. M., Kim, J., Choi, J. N., Kim, H. Y., et al. (2013). Culture condition-dependant metabolite profiling of *Aspergillus fumigatus* with antifungal activity. *Fungal Biol.* 117, 211–219. doi: 10.1016/j.funbio.2013.01.009
- Kano, R., Itamoto, K., Okuda, M., Inokuma, H., Hasegawa, A., and Balajee, S. A. (2008). Isolation of *Aspergillus udagawae* from a fatal case of feline orbital aspergillosis. *Mycoses* 51, 360–361. doi: 10.1111/j.1439-0507.2008.01493.x
- Kano, R., Shinahashi, A., Fujino, Y., Sakai, H., Mori, T., Tsujimoto, H., et al. (2013). Two cases of feline orbital aspergillosis due to *Aspergillus udagawae* and *A. viridinutans*. *J. Vet. Med. Sci.* 75, 7–10. doi: 10.1292/jvms.12-0119
- Kano, R., Takahashi, T., Hayawaka, T., Yamayaka, Y., Hasegawa, A., and Kamata, H. (2015). The first case of feline sinonasal aspergillosis due to *Aspergillus fischeri* in Japan. *J. Vet. Med. Sci.* 77, 1183–1185. doi: 10.1292/jvms.14-0454
- Kanokmedhakul, K., Kanokmedhakul, S., Suwannatrat, R., Soytong, K., Prabpais, S., and Kongsaeree, P. (2011). Bioactive meroterpenoids and alkaloids from the fungus *Eurotium chevalieri*. *Tetrahedron* 67, 5461–5468. doi: 10.1016/j.tet.2011.05.066
- Kao, R., Martinez-Ruiz, A., del Poozo, A. M., Cramer, R., and Davies, J. (2001). Mitogillin and related fungal ribotoxins. *Meth. Enzymol.* 34, 324–335. doi: 10.1016/S0076-6879(01)41161-X
- Katz, M. E., Dougall, A. M., Weeks, K., and Cheetham, B. F. (2005). Multiple genetically distinct groups revealed among clinical isolates identified as atypical *Aspergillus fumigatus*. *J. Clin. Microbiol.* 43, 551–555. doi: 10.1128/JCM.43.2.551-555.2005
- Kaur, A., Swenson, D. C., Wicklow, D. T., and Gloer, J. B. (2013). New fiscalin, tryptoquinavine, and fumiquinazoline analogues from an endophytic isolate of *Neosartorya aureola*. *Planta Med.* 79, 852. doi: 10.1055/s-0033-1348666
- Kelman, B. J., Robbins, C. A., Swenson, L. J., and Hardin, B. D. (2004). Risk from inhaled mycotoxins in indoor office and residential environments. *Int. J. Toxicol.* 23, 3–10. doi: 10.1080/10915810490265423
- Khare, R., Gupta, S., Arif, S., Jentoft, M. E., Denzil, P. H., Roden, A. C., et al. (2014). Misidentification of *Neosartorya pseudofischeri* as *Aspergillus fumigatus* in a lung transplant patient. *J. Clin. Microbiol.* 52, 2722–2725. doi: 10.1128/JCM.00216-14
- Khoufache, K., Puel, O., Loiseau, N., Delaforge, M., Rivollet, D., Coste, A., et al. (2007). Verruculogen associated with *Aspergillus fumigatus* hyphae and conidia modifies the electrophysiological properties of human nasal epithelial cells. *BMC Microbiol.* 25:5. doi: 10.1186/1471-2180-7-5
- Kijjoa, A., Santos, S., Dethoup, T., Manoch, L., Almeida, A. P., Vasconcelos, M. H., et al. (2011). Sartoryglabrins, analogs of ardeemins, from *Neosartorya glabra*. *Nat. Prod. Commun.* 6, 807–812.
- Kim, Y.-J., Park, H. B., Yoo, J.-H., Kwon, H. C., Kim, J., and Yang, H. O. (2014). Glionitrin A, a new diketopiperazine disulphide, activates ATM-ATR-Chk1/2 via 53BP1 phosphorylation in DU145 cells and shows antitumor effect in xenograft model. *Biol. Pharm. Bull.* 37, 378–386. doi: 10.1248/bpb.b13-00719
- Kupfahl, C., Michalka, A., Lass-Flörl, C., Fischer, G., Haase, G., Ruppert, T., et al. (2008). Gliotoxin production by clinical and environmental *Aspergillus fumigatus* strains. *Int. J. Med. Microbiol.* 298, 319–327. doi: 10.1016/j.ijmm.2007.04.006
- Kusuya, Y., Takahashi-Nakaguchi, A., Takahashi, H., and Yaguchi, T. (2015). Draft genome sequence of the pathogenic filamentous fungus *Aspergillus udagawae* strain IFM 46973^T. *Genome Announc.* 3, e00834–e00815. doi: 10.1128/genomeA.00834-15
- Kwon, Y.-J., Sohn, M.-J., Zheng, C.-J., and Kim, W.-G. (2007). Fumimycin: a peptide deformylase inhibitor with an unusual skeleton produced by *Aspergillus fumisynnematus*. *Org. Lett.* 9, 2449–2451. doi: 10.1021/o10703231
- Land, C. J., Lundström, H., and Werner, S. (1993). Production of tremorgenic mycotoxins by isolates of *Aspergillus fumigatus* from sawmills in Sweden. *Mycopathologia* 124, 87–93. doi: 10.1007/BF01103107
- Langarica-Fuentes, A., Handley, P. S., Houlden, A., Fox, G., and Robson, G. D. (2014). An investigation of the biodiversity of thermophilic and thermotolerant fungal species in composts using culture-based and molecular techniques. *Fungal Ecol.* 11, 132–144. doi: 10.1016/j.funeco.2014.05.007
- Langfelder, K., Streibel, M., Jahn, B., Haase, G., and Brakhaage, A. A. (2003). Biosynthesis of fungal melanins and their importance for human pathogenic fungi. *Fungal Genet. Biol.* 38, 143–158. doi: 10.1016/S1087-1845(02)00526-1
- Larsen, T. O., Smedsgaard, J., Nielsen, K. F., Hansen, M. E., and Frissvad, J. C. (2005). Phenotypic taxonomy and metabolite profiling in microbial drug discovery. *Nat. Prod. Rep.* 22, 672–695. doi: 10.1039/b404943h
- Larsen, T. O., Smedsgaard, J., Nielsen, K. F., Hansen, M. E., Samson, R. A., and Frissvad, J. C. (2007). Production of mycotoxins by *Aspergillus lentulus* and other medically important and closely related species in section *Fumigati*. *Med. Mycol.* 45, 225–232. doi: 10.1080/13693780601185939
- Latgé, J. P. (1999). *Aspergillus fumigatus* and aspergillosis. *Clin. Microbiol. Rev.* 12, 310–350.
- Lee, M. Y., Park, H. M., Son, G. H., and Lee, C. H. (2013). Liquid chromatography-mass spectrometry based chemotaxonomic classification of *Aspergillus* species and evaluation of the biological activity of its unique metabolite neosartorin. *J. Microbiol. Biotechnol.* 23, 932–941. doi: 10.4014/jmb.1212.12068
- Lewis, R. E., Wiederhold, N. P., Lionakis, M. S., Prince, R. A., and Kontoyiannis, D. P. (2005). Frequency and species distribution of gliotoxin-producing *Aspergillus* isolates recovered from patients at a tertiary-care cancer center. *J. Clin. Microbiol.* 43, 6120–6122. doi: 10.1128/JCM.43.12.6120-6122.2005
- Li, D. M., Horie, Y., Wang, Y., and Li, R. (1998). Three new *Aspergillus* species isolated from clinical sources as a causal agent of human aspergillosis. *Mycoscience* 39, 299–305. doi: 10.1007/BF02464012
- Li, Y.-X., Himaya, S. W. A., Dewapriya, P., Kim, H. J., and Kim, S. K. (2014). Antiproliferative effects of isosclerone from marine fungus *Aspergillus fumigatus* in MCF-7 human breast cancer cells. *Process Biochem.* 49, 2292–2298. doi: 10.1016/j.procbio.2014.08.016
- Liang, W.-L., Le, X., Li, H.-J., Yang, X.-L., Chen, J.-X., Xu, J., et al. (2014). Exploring the chemodiversity and biological activities of the secondary metabolites from

- the marine fungus *Neosartorya pseudofischeri*. *Mar. Drugs* 12, 5657–5676. doi: 10.3390/mdl2115657
- Liang, Z., Zhang, T., Zhang, X., Zhang, J., and Zhao, C. (2015). An alkaloid and a steroid from the endophytic fungus *Aspergillus fumigatus*. *Molecules* 20, 1424–1433. doi: 10.3390/molecules20011424
- Lillehoj, E. B., and Milburn, M. S. (1973). Viriditoxin production by *Aspergillus viridi-nutans* and related species. *Appl. Microbiol.* 26, 202–205.
- Lim, F. Y., Ames, B., Walsch, C. T., and Keller, N. P. (2014). Co-ordination between BrlA regulation and secretion of the oxidoreductase FmqD directs selective accumulation of fumiquinazoline C to conidial tissues in *Aspergillus fumigatus*. *Cell. Microbiol.* 16, 1267–1283. doi: 10.1111/cmi.12284
- Lin, H.-C., Chooi, Y.-H., Dhingra, S., Xu, W., Calvo, A. M., and Tang, Y. (2013). The fumagillin biosynthetic gene cluster in *Aspergillus fumigatus* encodes a cryptic terpene cyclase involved in the formation of β-trans-bergamotene. *J. Am. Chem. Soc.* 135, 4616–4619. doi: 10.1021/ja312503y
- Lind, A. L., Wisecaver, J. H., Smith, T. D., Feng, X., Calvo, A. M., and Rokas, A. (2015). Examining the evolution of the regulatory circuit controlling secondary metabolism and development in the fungal genus *Aspergillus*. *PLoS Genet.* 11:e1005096. doi: 10.1371/journal.pgen.1005096
- Liú, J., Wei, X., Kim, E. L., Lin, X., Yang, X.-W., Zhou, X., et al. (2015). New glucosidated pyrazinoquinazoline indole alkaloids from the fungus *Aspergillus fumigatus* derived from a jellyfish. *Tetrahedron* 71, 271–275. doi: 10.1016/j.tet.2014.11.063
- MacHeleidt, J., Scherlach, K., Neuwirth, T., Schmidt-Heck, W., Strassburger, M., Spraker, J., et al. (2015). Transcriptome analysis of cyclic AMP-dependent protein kinase A-regulated genes reveals the production of the novel natural compound fumipyrrole by *Aspergillus fumigatus*. *Mol. Microbiol.* 96, 148–162. doi: 10.1111/mmi.12926
- Macías, F. A., Varela, R. M., Simonet, A. M., Cutler, H. G., Cutler, S. J., and Hill, R. A. (2003). Absolute configuration of bioactive expansolides A and B from *Aspergillus fumigatus* Fresenius. *Tetrahedron Lett.* 44, 941–943. doi: 10.1016/S0040-4039(02)02778-8
- Malejczyk, K., Sigler, L., Gibas, G. F. C., and Smith, S. W. (2013). Invasive sino-orbital mycosis in an aplastic anaemia patient caused by *Neosartorya laciniosa*. *J. Clin. Microbiol.* 51, 1316–1319. doi: 10.1128/JCM.02919-12
- Masi, M., Andolfi, A., Mathieu, V., Boari, A., Cimmino, A., Y., Banuls, L. M. et al. (2013). Fischerindoline, a pyrrolindole sesquiterpenoid isolated from *Neosartorya pseudofischeri*, with *in vitro* growth inhibitory activity in human cancer cell lines. *Tetrahedron* 69, 7466–7470. doi: 10.1016/j.tet.2013.06.031
- Matsusawa, T., Horie, Y., Abilz, P., Gono, T., and Yaguchi, T. (2014a). *Aspergillus huiyaniae* sp. nov., a teleomorphic species in sect. *Fumigati* isolated from desert soil in China. *Mycoscience* 55, 213–220. doi: 10.1016/j.myc.2013.08.007
- Matsusawa, T., Takaki, G. M. C., Yaguchi, T., Okada, K., Abilz, P., Gono, T., et al. (2015). *Aspergillus arcoverdensis*, a new species of *Aspergillus* section *Fumigati* isolated from caatinga soil in the state of Pernambuco, Brazil. *Mycoscience* 56, 123–131. doi: 10.1016/j.myc.2014.04.006
- Matsusawa, T., Takaki, G. M. C., Yaguchi, T., Okada, K., Gono, T., and Horie, Y. (2014b). Two new species of *Aspergillus* section *Fumigati* isolated from caatinga soil in the state of Pernambuco, Brazil. *Mycoscience* 55, 79–88. doi: 10.1016/j.myc.2013.04.001
- McNeill, J., Barrie, F. R., Buck, W. R., Demoulin, V., Greuter, W., Gams, W., et al. (2012). *International Code of Nomenclature for Algae, Fungi and Plants (Melbourne Code)* Adopted by the 18th International Botanical Congress, Melbourne, Australia, Jul 2011. *Regnum Vegetabile* 154. Koenigstein: Koeltz Scientific Books.
- McDonagh, A., Fedorova, N. D., Crabtree, J., Yu, Y., Kim, S., Chen, D., et al. (2008). Sub-telomere directed gene expression during initiation of invasive aspergillosis. *PLoS Pathog.* 4:e1000154. doi: 10.1371/journal.ppat.1000154
- Miao, Y.-Z., Liu, D.-Y., Li, G.-Q., Li, P., Xu, Y.-C., Shen, Q.-R., et al. (2015). Genome-wide transcriptomic analysis of a superior biomass-degrading strain of *Aspergillus fumigatus* revealed active lignocellulose-degrading genes. *BMC Genomics* 16:459. doi: 10.1186/s12864-015-1658-2
- Morino, T., Nishimoto, M., Itou, N., and Nishikiori, T. (1994). NK372135s, novel antifungal agents produced by *Neosartorya fischeri*. *J. Antibiot.* 47, 1546–1548. doi: 10.7164/antibiotics.47.1546
- Mühlenfeld, A., and Achenbach, H. (1988a). Asperpentyn, a novel acetylenic cyclohexene epoxide from *Aspergillus duricaulis*. *Phytochemistry* 27, 3853–3855.
- Mühlenfeld, A., and Achenbach, H. (1988b). Stoffwechselprodukte von mikroorganismen. 35 mitt. Asperdurin, ein neues antimykotisches phthalide aus *Aspergillus duricaulis*. *Archiv der Pharmazie (Weinheim)* 321, 803–805.
- Netzker, T., Fischer, J., Weber, J., Mattem, D. J., Konig, C. C., Valiante, V., et al. (2015). Microbial communication leading to the activation of silent fungal secondary metabolite gene clusters. *Front. Microbiol.* 6:299. doi: 10.3389/fmicb.2015.00299
- Nielsen, M. L., Nielsen, J. B., Rank, C., Klejnstrup, M. L., Holm, D. M. K., Brogaard, K. H., et al. (2011). A genome-wide polyketide synthase deletion library uncovers novel genetic links to polyketides and meroterpenoids in *Aspergillus nidulans*. *FEMS Microbiol. Lett.* 321, 157–166. doi: 10.1111/j.1574-6968.2011.02327.x
- Nierman, W. C., Pain, A., Anderson, M. J., Wortman, J. R., Kim, H. S., Arroyo, J., et al. (2005). Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus fumigatus*. *Nature* 438, 1151–1156. doi: 10.1038/nature04332
- Novaková, A., Hubka, V., Dudová, Z., Matsuzawa, T., Kubatová, A., Yaguchi, T., et al. (2014). New species in *Aspergillus* section *Fumigati* from reclamation sites in Wyoming (USA) and revision of *A. viridinutans* complex. *Fungal Divers.* 64, 253–274. doi: 10.1007/s13225-013-0262-5
- O’Gorman, C. M., Fuller, H. T., and Dyer, P. S. (2009). Discovery of a sexual cycle in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Nature* 457, 471–474. doi: 10.1038/nature07528
- Ohashi, H., Ishikawa, M., Ito, J., Ueno, A., Gleich, G. J., Kita, H., et al. (1997). Sulochrin inhibits eosinophil degranulation. *J. Antibiot.* 50, 972–974. doi: 10.7164/antibiotics.50.972
- Ohashi, H., Motegi, Y., Kitas, H., Gleich, G. J., Miura, T., Ishikawa, M., et al. (1998). Sulochrin inhibits eosinophil activation and chemotaxis. *Inflamm. Res.* 47, 409–415. doi: 10.1007/s000110050352
- O’Keeffe, G., Hammel, S., Owens, R. A., Keane, T. M., FitzPatrick, O. A., Jones, G. W., et al. (2014). RNA-sequencing reveals the pan-transcriptomic impact of attenuating the gliotoxin self-protection mechanism in *Aspergillus fumigatus*. *BMC Genomics* 15:894. doi: 10.1186/1471-2164-15-894
- Ola, A. R. B., Debba, A., Aly, A. H., Mandi, A., Zerfass, I., Hamacher, A., et al. (2014). Absolute configuration and antibiotic activity of neosartorin from the endophytic fungus *Aspergillus fumigataffinis*. *Tetrahedron Lett.* 55, 1020–1023. doi: 10.1016/j.tetlet.2013.12.070
- Omolo, J. O., Anke, H., Chhabara, S., and Sternner, O. (2000). New variotin analogues from *Aspergillus viridi-nutans*. *J. Nat. Prod.* 63, 975–977. doi: 10.1021/np990509b
- Owens, K. A., Hammel, S., Sheridan, K. J., Jones, G. W., and Doyle, S. (2014). A proteomic approach to investigating gene cluster expression and secondary metabolite functionality in *Aspergillus fumigatus*. *PLoS ONE* 9:e106942. doi: 10.1371/journal.pone.0106942
- Ozoe, Y., Kuriyama, T., Tachibana, Y., Hariyama, K., Takahashi, N., Yuguchi, T., et al. (2004). Isocoumarin derivative as a novel GABA receptor ligand from *Neosartorya quadricincta*. *J. Pestic. Sci.* 29, 328–333. doi: 10.1584/jpestics.29.328
- Panda, A., Ghosh, A. K., Mirdha, B. R., Xess, I., Paul, S., Samanaray, J. C., et al. (2015). MALDI-TOF mass spectrometry for rapid identification of clinical fungal isolates based on ribosomal protein markers. *J. Microbiol. Methods* 109, 93–105. doi: 10.1016/j.mimet.2014.12.014
- Pedersen, M. H., Borodina, I., Moresco, J. L., Svendsen, W. E., Frisvad, J. C., and Sondergaard, I. (2011). High-yield production of hydrophobins RodA and RodB from *Aspergillus fumigatus* in *Pichia pastoris*. *Appl. Microbiol. Biotechnol.* 90, 1923–1932. doi: 10.1007/s00253-011-3235-1
- Peláez, T., Álvarez-Perez, S., Mellado, E., Serrano, D., Valerio, M., Blanco, J. L., et al. (2013). Invasive aspergillosis caused by cryptic *Aspergillus* species: a report of two consecutive episodes in a patient with leukemia. *J. Med. Mycol.* 62, 474–478. doi: 10.1099/jmm.0.044867-0
- Peng, J., Lin, T., Wang, W., Xin, Z., Zhu, T., Gu, Q., et al. (2013). Antiviral alkaloids produced by the mangrove-derived fungus *Cladosporium* sp. PJX-41. *J. Nat. Prod.* 76, 1133–1140. doi: 10.1021/np400200k
- Perrin, R. M., Fedorova, N. D., Bok, J. W., Cramer, R. A. Jr., Wortman, J. R., Kim, H. S. et al. (2007). Transcriptional regulation of chemical diversity in *Aspergillus fumigatus* by LaeA. *PLoS Pathog.* 3:e50. doi: 10.1371/journal.ppat.0030050
- Petrik, M., Haas, J. H., Laverman, P., Schrettl, M., Franssen, G. M., Blatzer, M., et al. (2014). ⁶⁸Ga-triacytlyfusarinine C and ⁶⁸Ga-ferrioxamine E for *Aspergillus* infection imaging: Uptake specificity in various microorganisms. *Mol. Imag. Biol.* 16, 102–108. doi: 10.1007/s11307-013-0654-7

- Posteraro, B., Mattei, R., Trivella, F., Maffei, A., Torre, A., De Carolis, E., et al. (2011). Uncommon *Neosartorya udagawae* fungus as a causative agent of severe corneal infection. *J. Clin. Microbiol.* 49, 2357–2360. doi: 10.1128/JCM.00134-11
- Prompanya, C., Dethoup, T., Bessa, L. J., Pinto, M. M. M., Gales, L., Costa, P. M., et al. (2014). New isocoumarin derivatives and meroterpenoids from the marine sponge-associated fungus *Aspergillus similanensis*. *Mar. Drugs* 12, 5160–5173. doi: 10.3390/md12105160
- Prompanya, C., Fernandes, C., Cravo, S., Pinto, M. M. M., Dethoup, T., Silva, A. M. S., et al. (2015). A new cyclic hexapeptide and a new isocoumarin derivative from the marine sponge-associated fungus *Aspergillus similanensis* KUFA 0013. *Mar. Drugs* 13, 1432–1450. doi: 10.3390/md13031432
- Qiao, M.-F., Ji, N.-Y., Liu, X.-H., Li, F., and Xue, Q.-Z. (2010b). Asporgerosterol, a new steroid from an algicolous isolate of *Aspergillus oryzae*. *Nat. Prod. Commun.* 5, 1575–1578.
- Qiao, M.-F., Ji, N.-Y., Liu, X.-H., Li, K., Zhu, Q.-M., Xue, Q.-Z., et al. (2010a). Indoloterpenes from an algicolous isolate of *Aspergillus oryzae*. *Biorg. Med. Chem. Lett.* 20, 5677–5680. doi: 10.1016/j.bmcl.2010.08.024
- Rabindran, S. K., Ross, D. D., Doyle, L. A., Yang, W., and Greenberger, L. M. (2000). Fumitremorgin C reverses multidrug resistance in cells transfected with the cancer resistance protein. *Cancer Res.* 60, 47–50.
- Rank, C., Phipps, R. K., Harris, P., Frisvad, J. C., Gotfredsen, C. H., and Larsen, T. O. (2006). Epi-aszonalenin A, B, and C from *Aspergillus novofumigatus*. *Tetrahedron Lett.* 47, 6099–6102. doi: 10.1016/j.tetlet.2006.06.086
- Rank, C., Phipps, R. K., Harris, P., Fristrup, P., Larsen, T. O., and Gotfredsen, C. H. (2008). Novofumigatonin, a new orthoester meroterpenoid from *Aspergillus novofumigatus*. *Org. Lett.* 10, 401–404. doi: 10.1021/o107206834
- Rank, C., Klejnstrup, M. L., Petersen, L. M., Kildegaard, S., Frisvad, J. C., Gotfredsen, C. H., et al. (2012). Comparative chemistry of *Aspergillus oryzae* (RIB40) and *A. flavus* (NRRL 3357). *Metabolites* 2, 39–56. doi: 10.3390/metabo2010039
- Raper, K. B., and Fennell, D. I. (1965). *The Genus Aspergillus*. Baltimore, MD: Williams and Wilkins.
- Reeves, E. P., Reiber, K., Neville, C., Schreibner, O., Kavanagh, K., and Doyle, S. (2006). A nonribosomal peptide synthetase (Pes1) confers protection against oxidative stress in *Aspergillus fumigatus*. *FEBS J.* 273, 3038–3053. doi: 10.1111/j.1742-4658.2006.05315.x
- Rementeria, A., Lopez-Molina, N., Ludwig, A., Vivanco, A. B., Belen, A., Bikandi, J., et al. (2005). Genes and molecules involved in *Aspergillus fumigatus* virulence. *Rev. Iberoam. Micol.* 22, 1–23. doi: 10.1016/S1130-1406(05)70001-2
- Rodrigues, B. S. F., Sahm, B. D. B., Jimenez, P. C., Pinto, F. C. L., Mafezoli, J., Mattos, M. C., et al. (2015). Bioprospection of cytotoxic compounds in fungal strains recovered from sediments of the Brazilian coast. *Chem. Biodivers.* 12, 432–442. doi: 10.1002/cbdv.201400193
- Samson, R. A., Hong, S.-B., and Frisvad, J. C. (2006). Old and new concepts of species differentiation in *Aspergillus*. *Med. Mycol.* 44, S133–S144. doi: 10.1080/13693780600913224
- Samson, R. A., Hong, S.-B., Peterson, S. W., Frisvad, J. C., and Varga, J. (2007). Polyphasic taxonomy of *Aspergillus* section *Fumigati* and its teleomorph *Neosartorya*. *Stud. Mycol.* 59, 147–203. doi: 10.3114/sim.2007.59.14
- Samson, R. A., Nielsen, P. V., and Frisvad, J. C. (1990). “The genus *Neosartorya*: differentiation by scanning electron microscopy and mycotoxin profiles,” in *Modern Concepts in Penicillium and Aspergillus Classification*, R. A. Samson and J. I. Pitt (New York, NY: Plenum Press), 455–467. doi: 10.1007/978-1-4899-3579-3_40
- Samson, R. A., Peterson, S. W., Frisvad, J. C., and Varga, J. (2011). New species in *Aspergillus* section *Terrei*. *Stud. Mycol.* 69, 39–55. doi: 10.3114/sim.2011.69.04
- Samson, R. A., Visagie, C. M., Houbraken, J., Hong, S.-B., Hubka, V., Klaasen, C. H. W., et al. (2014). Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Stud. Mycol.* 78, 141–173. doi: 10.1016/j.simco.2014.07.004
- Sanchez, J. F., Somoza, A. D., Keller, N. P., and Wang, C. C. (2012). Advances in *Aspergillus* secondary metabolite research in the post-genomic era. *Nat. Prod. Rep.* 29, 351–371. doi: 10.1039/c2np00084a
- Schmeda-Hirschmann, G., Hormazabal, E., Rodriguez, J. A., and Theoduloz, C. (2008). Cycloaspeptide A and pseurotin A from the endophytic fungus *Penicillium janczewskii*. *Z. Naturforsch. C* 63, 383–388. doi: 10.1515/znc-2008-5-612
- Schwienbacher, M., Weig, M., Thies, S., Regula, J. T., Heeseman, J., and Ebel, F. (2005). Analysis of the major proteins secreted by the human opportunistic pathogen *Aspergillus fumigatus* under *in vitro* conditions. *Med. Mycol.* 43, 623–630. doi: 10.1080/13693780500089216
- Shan, W., Wang, S., Ying, Y., Ma, L., and Zhan, Z. (2014). Indole-benzodiazepine-2,5-dione derivatives from *Neosartorya fischeri*. *J. Chem. Res.* 38, 692–694. doi: 10.3184/174751914X1414003469581
- Shan, W.-G., Wang, S.-L., Lang, H. Y., Chen, S. M., Ying, Y.-M., and Zhan, Z.-J. (2015). Cottoquinazolins E and F from *Neosartorya fischeri* NRRL 181. *Helv. Chim. Acta* 98, 552–556. doi: 10.1002/hlca.201400270
- Shinohara, C., Hasumi, K., and Endo, A. (1992). Inhibition of oxidized low-density lipoprotein metabolism in macrophage J774 by helvolic acid. *Biochim. Biophys. Acta* 1167, 303–306. doi: 10.1016/0005-2760(93)90233-Y
- Sigler, L., Sutton, D. A., Gibas, C. F. C., Summerbell, R. C., Noel, R. K., and Iwen, P. C. (2010). *Phialosimplex*, a new anamorphic genus associated with infections in dogs and having phylogenetic affinity to the *Trichocomaceae*. *Med. Mycol.* 48, 335–345. doi: 10.3109/13693780903225805
- Sodngam, S., Sawadsitang, S., Suwananasai, N., and Mongkoltharanuk, W. (2014). Chemical constituents, and their cytotoxicity, of the rare wood decaying fungus *Xylaria humosa*. *Nat. Prod. Commun.* 9, 157–158.
- Someya, A., Yaguchi, T., and Udagawa, S. (1999). *Neosartorya sublevispora*, a new species of soil-borne Eurotiales. *Mycoscience* 40, 405–409. doi: 10.1007/BF02464395
- Srivastava, N., Rawal, R., Sharma, R., Oberoi, H. S., Srivastava, M., and Singh, J. (2014). Effect of nickel-cobaltite nanoparticles on production and thermostability of cellulases from newly isolated thermotolerant *Aspergillus fumigatus* NS (Class: Eurotiomycetes). *Appl. Biochem. Biotechnol.* 174, 1092–1103. doi: 10.1007/s12010-014-0940-0
- Stack, D., Neville, C., and Doyle, S. (2007). Nonribosomal peptide synthesis in *Aspergillus fumigatus* and other fungi. *Microbiology* 153, 1297–1306. doi: 10.1099/mic.0.2006/006908-0
- Stanzani, M., Orciuolo, E., Lewis, R., Kontoyiannis, D. P., Martins, S. L. R., St. John, L. S., et al. (2005). *Aspergillus fumigatus* suppresses the human cellular immune response via gliotoxin-mediated apoptosis of monocytes. *Blood* 105, 2258–2265. doi: 10.1182/blood-2004-09-3421
- Sugui, J. A., Pardo, J., Chang, Y. C., Zaremba, K. A., Nardone, G., Galvez, E. M., et al. (2007). Gliotoxin is a virulence factor of *Aspergillus fumigatus*; *gliP* deletion attenuates virulence in mice immunocompromised with hydrocortisone. *Eukaryotic Cell* 6, 1562–1569. doi: 10.1128/EC.00141-07
- Sugui, J. A., Kwon-Chung, K. J., Juvvadi, P. R., Latgé, J. P., and Steinbach, W. J. (2015). *Aspergillus fumigatus* and related species. *Cold Spring Harb. Perspect. Med.* 5: a019786. doi: 10.1101/cshperspect.a019786
- Sugui, J. A., Peterson, S. W., Clark, L. P., Nardone, G., Folio, L., Riedlinger, G., et al. (2012). *Aspergillus tanneri* sp. nov., a new pathogen that causes invasive disease refractory to antifungal therapy. *J. Clin. Microbiol.* 50, 3309–3317. doi: 10.1128/JCM.01509-12
- Sugui, J. A., Peterson, S. W., Figat, A. B., Hansen, B., Samson, R. A., Mellado, E., et al. (2014). Genetic relatedness versus biological compatibility between *Aspergillus fumigatus* and related species. *J. Clin. Microbiol.* 52, 3707–3721. doi: 10.1128/JCM.01704-14
- Sugui, J. A., Vinh, D. C., Nardone, G., Sheat, Y. R., Chang, Y. C., Zelazny, A. M., et al. (2010). *Neosartorya udagawae* (*Aspergillus udagawae*), an emerging agent of aspergillosis: how different is it from *Aspergillus fumigatus*? *J. Clin. Microbiol.* 48, 220–228. doi: 10.1128/JCM.01556-09
- Sutton, D. A., Summerbell, R. C., Samson, R., and Rinalde, M. F. (2002). “First report of *Neosartorya spinosa* (Raper & Fennell) Kozakiewicz inciting human disease,” in *Abstracts of the General Meeting of ASM*, Vol. 102 (Salt Lake City, UT), 212.
- Swilaiman, S. S., O’Gorman, C. M., Balajee, S. A., and Dyer, P. S. (2013). Discovery of a sexual cycle in *Aspergillus lentulus*, a close relative of *A. fumigatus*. *Eukaryotic Cell* 12, 962–969. doi: 10.1128/EC.00040-13
- Takahashi, C., Matsushida, T., Doi, M., Minoura, K., Shingu, T., Komeda, Y., et al. (1995). Fumiquinazolions A-G, novel metabolites of a fungus separated from a *Pseudolabrus* marine fish. *J. Chem. Soc. Perkin Trans. I* 1995, 2345–2353. doi: 10.1039/p19950002345
- Takeda, M., Horie, Y., and Abliz, P. (2001). Two new heterothallic *Neosartorya* from African soil. *Mycoscience* 42, 361–367. doi: 10.1007/BF02461219
- Tamiya, H., Ochiai, E., Kikuchi, K., Yahiro, M., Toyotome, T., Watanabe, A., et al. (2015). Secondary metabolite profiles and antifungal drug susceptibility of *Aspergillus fumigatus* and closely related species, *A. lentulus*, *A. udagawae*,

- and *A. viridinutans*. *Journal of Infection and Chemotherapy*. 21, 385–391. doi: 10.1016/j.jiac.2015.01.005
- Tepsic, K., Gunde-Cimerman, N., and Frissvad, J. C. (1997). Growth and mycotoxin production by *Aspergillus fumigatus* strains isolated from a saltern. *FEMS Microbiol. Lett.* 157, 9–12. doi: 10.1111/j.1574-6968.1997.tb12745.x
- Tomee, J. F., and Kauffman, H. F. (2000). Putative virulence factors of *Aspergillus fumigatus*. *Clin. Exp. Allergy* 30, 476–484. doi: 10.1046/j.1365-2222.2000.00796.x
- Toskova, M., Palousova, D., Kocmanova, I., Pavlovsky, Z., Timilsinas, S., Langerova, M., et al. (2013). Invasive mould disease involving the gastrointestinal tract caused by *Neosartorya pseudofischeri* in a haematological patient. *Mycoses* 56, 385–388. doi: 10.1111/myc.12038
- Tsai, H.-F., Chang, Y. C., Washburn, R. G., Wheeler, M. H., and Kwon-Chung, K. J. (1998). The developmentally regulated *alb1* gene of *Aspergillus fumigatus*: its role in modulation of conidial morphology and virulence. *J. Bacteriol.* 180, 3031–3038.
- Tsai, H.-F., Fujii, I., Watanabe, A., Wheeler, M. H., Chang, Y. C., Yasuaka, Y., et al. (2001). Pentaketide melanin biosynthesis in *Aspergillus fumigatus* requires chain-length shortening of a heptaketide precursor. *J. Biol. Chem.* 276, 29292–29298. doi: 10.1074/jbc.M101998200
- Tsunawaki, S., Yoshida, L. S., Nishida, S., Kobayashi, T., and Shimoyama, T. (2004). Fungal metabolite gliotoxin inhibits assembly of the human respiratory burst NADPH oxidase. *Infect. Immun.* 72, 3373–3382. doi: 10.1128/IAI.72.6.3373-3382.2004
- Varga, J., Due, M., Frissvad, J. C., and Samson, R. A. (2007). Taxonomic revision of *Aspergillus* section *Clavati* based on molecular, morphological and physiological data. *Stud. Mycol.* 59, 89–106. doi: 10.3114/sim.2007.59.11
- Varga, J., Frissvad, J. C., Kocsimbé, S., Brancovics, B., Szegedi, G., and Samson, R. A. (2011a). New and revisited species in *Aspergillus* section *Nigri*. *Stud. Mycol.* 69, 1–17. doi: 10.3114/sim.2011.69.01
- Varga, J., Frissvad, J. C., and Samson, R. A. (2011b). Two new aflatoxin producing species, and an overview of *Aspergillus* section *Flavi*. *Stud. Mycol.* 69, 57–80. doi: 10.3114/sim.2011.69.05
- Varga, J., and Samson, R. A. (2008). Ribotoxin genes in *Aspergillus* section *Clavati*. *Antonie van Leeuwenhoek* 94, 481–485. doi: 10.1007/s10482-008-9266-7
- Varga, J., Tóth, B., Rigó, K., Debets, F., and Kozakiewicz, Z. (2000). Genetic variability within the *Aspergillus viridinutans* species. *Folia Microbiol.* 45, 423–428. doi: 10.1007/BF02817615
- Verwer, P. E. B., van Leuwen, W. B., Girard, V., van Belkum, A., Staab, J. F., Verbrugh, H. A., et al. (2014). Diuscrimination of *Aspergillus lentulus* from *Aspergillus fumigatus* by Raman spectroscopy and MALDI-TOF MS. *Eur. J. Clin. Microbiol.* 33, 245–251. doi: 10.1007/s10096-013-1951-4
- Vinh, D. C., Shea, Y. R., Jones, P. A., Freeman, A. F., Zelazny, A., and Holland, S. M. (2009a). Chronic invasive aspergillosis caused by *Aspergillus viridinutans*. *Emerging Infect. Dis.* 15, 1292–1294. doi: 10.3201/eid1508.090251
- Vinh, D. C., Shea, Y. R., Sogui, J. A., Parrilla-Castellar, E. R., Freeman, A. F., Campbell, J. W., et al. (2009b). Invasive aspergillosis due to *Neosartorya udagawae*. *Clin. Infect. Dis.* 49, 102–111. doi: 10.1086/599345
- Virág, M., Vörös, D., Kele, Z., Kovács, L., Fizilo, Á., Lakatos, G., et al. (2014). Production of a defensin-like antifungal protein NFAP from *Neosartorya fischeri* in *Pichia pastoris* and its antifungal activity against filamentous fungal isolates from human infections. *Protein Expr. Purif.* 94, 78–84. doi: 10.1016/j.pep.2013.11.003
- Visagie, C., Varga, J., Houbraken, J., Meijer, M., Yilmaz, N., Fotedar, R., et al. (2014). Ochratoxin production and taxonomy of the yellow aspergilli (*Aspergillus* section *Circumdati*). *Stud. Mycol.* 78, 1–61. doi: 10.1016/j.simyco.2014.07.001
- Vödisch, M., Scherlach, K., Winkler, R., Hertweck, C., Braun, H. P., Roth, M., et al. (2011). Analysis of the *Aspergillus fumigatus* proteome reveals metabolic changes and the activation of the pseurotin A biosynthesis gene cluster in response to hypoxia. *J. Proteome Res.* 10, 2508–2524. doi: 10.1021/pr1012812
- Wakana, D., Hosoe, T., Itabashi, T., Nozawa, K., Okada, K., Campos Takaki, G. M., et al. (2006). Isolation of isoterrein from *Neosartorya fischeri*. *Mycotoxins* 56, 3–6. doi: 10.2520/myco.56.3
- Wang, F., Fang, Y., Zhu, T., Zhang, M., Lin, A., Gu, Q., et al. (2008). Seven new prenylated indole diketopiperazine alkaloids from holothurian-derived fungus *Aspergillus fumigatus*. *Tetrahedron* 64, 7986–7991. doi: 10.1016/j.tet.2008.06.013
- Watanabe, A., Kamei, K., Sekine, T., Hiraguchi, H., Ochiai, E., Hashimoto, Y., et al. (2004). Cytotoxic substances from *Aspergillus fumigatus* in oxygenated or poorly oxygenated environment. *Mycopathologia* 158, 1–7. doi: 10.1023/B:MYCO.0000038439.56108.3c
- Watanabe, A., Kamei, K., Sekine, T., Waku, M., Nishimura, K., Miyaji, M., et al. (2003). Immunosuppressive substances in *Aspergillus fumigatus* culture filtrate. *J. Infect. Chemother.* 9, 114–121. doi: 10.1007/s10156-002-0227-1
- Wiekmann, I., Lechner, B. E., Baccile, J. A., Velk, T. A., Yin, W.-B., Bok, J. B., et al. (2014). Perturbations in small molecule synthesis uncovers an iron-responsive secondary metabolite network in *Aspergillus fumigatus*. *Front. Microbiol.* 5:530. doi: 10.3389/fmicb.2014.00530
- Wiemann, P., Guo, C. J., Palmer, J. M., Sekonyela, R. R., Wang, C. C. C., and Keller, N. P. (2013). Prototype of an intertwined secondary-metabolite supercluster. *Proc. Natl. Acad. Sci. U.S.A.* 110, 17065–17070. doi: 10.1073/pnas.1313258110
- Wong, S. M., Musza, L. L., Kydd, G. C., Kullnig, R., Gillum, A. M., and Cooper, R. (1993). Fiscalins: new substance P inhibitors produced by the fungus *Neosartorya fischeri*. Taxonomy, fermentation, structures, and biological properties. *J. Antibiot.* 46, 545–553. doi: 10.7164/antibiotics.46.545
- Wortman, J. R., Fedorova, N., Crabtree, J., Joardar, V., Maiti, R., Haas, B. J., et al. (2006). Whole genome comparison of the *Aspergillus fumigatus* family. *Med. Mycol.* 44, S3–S7. doi: 10.1080/13693780600835799
- Xie, F., Li, X.-B., Zhou, J.-C., Xu, Q.-Q., Wang, X.-N., Yuan, H.-Q., et al. (2015). Secondary metabolites from *Aspergillus fumigatus*, an endophytic fungus from the liverwort *Heteroscyphus tener* (STEPH.) SCHIFFN. *Chem. Biodivers.* 12, 1313–1321. doi: 10.1002/cbdv.201400317
- Xu, N., Cao, Y., Wang, L., Chen, G., and Pei, Y.-H. (2013). New alkaloids from a marine-derived fungus *Neosartorya* sp. HN-M-3. *J. Asian Nat. Prod. Res.* 15, 731–736. doi: 10.1080/10286020.2013.797895
- Xu, J., Song, Y. C., Guo, Y., Mei, Y. N., and Tan, R. X. (2014). Fumigaclavine D-H, new ergot alkaloids from endophytic *Aspergillus fumigatus*. *Planta Med.* 80, 1131–1137. doi: 10.1055/s-0034-1382958
- Yaguchi, T., Horie, Y., Tanaka, R., Matsuzawa, T., Ito, J., and Nishimura, K. (2007). Molecular phylogenetics of multiple genes on *Aspergillus* section *Fumigati* isolated from clinical specimens in Japan. *Jpn. J. Med. Mycol.* 48, 37–46. doi: 10.3314/jjmm.48.37
- Yaguchi, T., Matsuzawa, T., Tanaka, R., Abliz, P., Hui, Y., and Horie, Y. (2010). Two new species of *Neosartorya* isolated from soil in Xinjiang, China. *Mycoscience* 51, 253–262. doi: 10.1007/S10267-010-0037-8
- Ye, Y., Minami, A., Mandi, A., Liu, C., Taniguchi, T., Kuzuyama, T., et al. (2015). Genome mining for sesquiterpenes using bifunctional terpene synthases reveals a unified intermediate of di/sesquiterpenes. *J. Am. Chem. Soc.* 137, 11846–11853. doi: 10.1021/jacs.5b08319
- Yim, T., Kanokmedhakul, K., Kanokmedhakul, S., Sanmanoch, W., and Boonlue, S. (2014). A new meroterpenoid tatenoi acid from the fungus *Neosartorya tatenoi*. *Nat. Prod. Res.* 28, 1847–1852. doi: 10.1080/14786419.2014.951353
- Yin, W.-B., Grundmann, A., Cheng, Y., and Li, S.-M. (2009). Acetylaszonalein biosynthesis in *Neosartorya fischeri*. Identification of the biosynthetic gene cluster by genomic mining and functional proof of the genes by biochemical characterization. *J. Biol. Chem.* 284, 100–109. doi: 10.1074/jbc.M80760200
- Zhang, H.-W., Ying, C., and Tang, Y.-F. (2014). Four ardeemin analogs from endophytic *Aspergillus fumigatus* SPS-02 and their reversal effects on multi-drug resistant tumor cells. *Chem. Biodivers.* 11, 85–91. doi: 10.1002/cbdv.201300220
- Zhang, M., Wang, W.-L., Fang, Y.-C., Zhu, T.-J., Gu, Q.-Q., and Zhu, W.-M. (2008). Cytotoxic alkaloids and antibiotic nordammarane triterpenoids from the marine-derived fungus *Aspergillus sydowi*. *J. Nat. Prod.* 71, 985–989. doi: 10.1021/np700737g
- Zhang, H.-C., Ma, Y.-M., Liu, R., and Zhou, F. (2012). Endophytic fungus *Aspergillus tamarii* from *Ficus carica* L., a new source of indolyl dikeropiperazines. *Biochem. Syst. Ecol.* 45, 31–33. doi: 10.1016/j.bse.2012.07.020
- Zhao, W. Y., Zhu, T. J., Fan, G. T., Liu, H. B., Fang, Y. C., Gu, Q. Q., et al. (2010). Three new dioxopiperazine metabolites from a marine-derived fungus *Aspergillus fumigatus* Fres. *Nat. Prod. Res.* 24, 953–957. doi: 10.1080/14786410902726134
- Zheng, Z. Z., Shan, W. C., Wang, S.-L., Ying, Y.-M., Ma, L.-F., and Zhan, Z.-J. (2014). Three new prenylated dikeropiperazines from *Neosartorya fischeri*. *Helv. Chim. Acta* 97, 1020–1026. doi: 10.1002/hclca.201300416
- Zhou, F., Zhang, H., Liu, R., and Zhang, D. (2013). Isolation and biological evaluation of secondary metabolites from the endophytic fungus *Aspergillus*

- fumigatus* from *Astragalus membranaceus*. *Chem. Nat. Comp.* 49, 568–570. doi: 10.1007/s10600-013-0675-0
- Zin, W. W. M., Buttachon, S., Buaruang, J., Gales, L., Pereira, J. A., Pinto, M. M. M., et al. (2015). A new meroditerpene and a new tryptoquivaline analog from the algicolous fungus *Neosartorya takakii* KUFC 7898. *Mar. Drugs* 13, 3776–3790. doi: 10.3390/md13063776
- Zuck, K. M., Shipley, S., and Newman, D. J. (2011). Induced production of N-formyl alkaloids from *Aspergillus fumigatus* by co-culture with *Streptomyces peucetius*. *J. Nat. Prod.* 74, 1653–1657. doi: 10.1021/np200255f

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Frisvad and Larsen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.