



# The Complete Genome Sequence of the Fish Pathogen *Tenacibaculum maritimum* Provides Insights into Virulence Mechanisms

David Pérez-Pascual<sup>1</sup>, Aurelie Lunazzi<sup>1</sup>, Ghislaine Magdelenat<sup>2</sup>, Zoe Rouy<sup>3</sup>, Alain Roulet<sup>4,5</sup>, Celine Lopez-Roques<sup>4,5</sup>, Robert Larocque<sup>6</sup>, Tristan Barbeyron<sup>6</sup>, Angélique Gobet<sup>6</sup>, Gurvan Michel<sup>6</sup>, Jean-François Bernardet<sup>1</sup> and Eric Duchaud<sup>1\*</sup>

<sup>1</sup> Virologie et Immunologie Moléculaires, Institut National de la Recherche Agronomique, Université Paris-Saclay, Jouy-en-Josas, France, <sup>2</sup> Commissariat à l'Énergie Atomique et aux Énergies Alternatives, Institut de Génétique, Genoscope, Evry, France, <sup>3</sup> Laboratoire d'Analyses Bioinformatiques en Génétique et Métabolisme, Centre National de la Recherche Scientifique (UMR-8030), Commissariat à l'Énergie Atomique et aux Énergies Alternatives, Institut de Génétique, Genoscope, Evry, France, <sup>4</sup> Genotoul Genome & Transcriptome (GeT-PlaGe), Institut National de la Recherche Agronomique, Castanet-Tolosan, France, <sup>5</sup> Institut National de la Recherche Agronomique (UAR1209), Castanet-Tolosan, France, <sup>6</sup> Laboratoire de Biologie Intégrative des Modèles Marins (UMR 8227), Centre National de la Recherche Scientifique, Université Pierre et Marie Curie, Station Biologique de Roscoff, Sorbonne Universités, Roscoff, France

## OPEN ACCESS

### Edited by:

Senjie Lin,  
University of Connecticut,  
United States

### Reviewed by:

Jose M. Gonzalez,  
Universidad de La Laguna, Spain  
Edward Fell,  
University of Bath, United Kingdom

### \*Correspondence:

Eric Duchaud  
eric.duchaud@inra.fr

### Specialty section:

This article was submitted to  
Aquatic Microbiology,  
a section of the journal  
Frontiers in Microbiology

**Received:** 28 February 2017

**Accepted:** 31 July 2017

**Published:** 16 August 2017

### Citation:

Pérez-Pascual D, Lunazzi A, Magdelenat G, Rouy Z, Roulet A, Lopez-Roques C, Larocque R, Barbeyron T, Gobet A, Michel G, Bernardet J-F and Duchaud E (2017) The Complete Genome Sequence of the Fish Pathogen *Tenacibaculum maritimum* Provides Insights into Virulence Mechanisms. *Front. Microbiol.* 8:1542. doi: 10.3389/fmicb.2017.01542

*Tenacibaculum maritimum* is a devastating bacterial pathogen of wild and farmed marine fish with a broad host range and a worldwide distribution. We report here the complete genome sequence of the *T. maritimum* type strain NCIMB 2154<sup>T</sup>. The genome consists of a 3,435,971-base pair circular chromosome with 2,866 predicted protein-coding genes. Genes encoding the biosynthesis of exopolysaccharides, the type IX secretion system, iron uptake systems, adhesins, hemolysins, proteases, and glycoside hydrolases were identified. They are likely involved in the virulence process including immune escape, invasion, colonization, destruction of host tissues, and nutrient scavenging. Among the predicted virulence factors, type IX secretion-mediated and cell-surface exposed proteins were identified including an atypical sialidase, a sphingomyelinase and a chondroitin AC lyase which activities were demonstrated *in vitro*.

**Keywords:** *Tenacibaculum maritimum*, fish pathogen, virulence factors, genome, toxins

## INTRODUCTION

*Tenacibaculum maritimum* (formerly *Flexibacter maritimus*), a member of the family Flavobacteriaceae, phylum Bacteroidetes (Suzuki et al., 2001), is the etiological agent of tenacibaculosis, a very serious bacterial disease of many commercial marine fish species (for a review, see Avendaño-Herrera et al., 2006b), responsible for considerable economic losses in all major areas of marine finfish aquaculture worldwide (i.e., Japan, Europe including the Atlantic, Channel and Mediterranean coasts, North America, Australia, and the Red Sea). Moreover, *T. maritimum* can affect a large number of feral, captive, and cultured fish species such as: Dover sole (*Solea solea*), Senegalese sole (*Solea senegalensis*), wedge sole (*Dicologlossa cuneata*), turbot (*Scophthalmus maximus*), Atlantic salmon (*Salmo salar*), Japanese flounder (*Paralichthys olivaceus*), yellowtail (*Seriola quinqueradiata*), red sea bream (*Pagrus major*), black sea bream (*Acanthopagrus schlegelii*), gilthead sea bream (*Sparus aurata*), European sea bass (*Dicentrarchus labrax*), puffer fish (*Takifugu rubripes*), Pacific sardine (*Sardinops sagax*), lump sucker

(*Cyclopterus lumpus*), and sand tiger shark (*Carcharias taurus*) (Bernardet et al., 1990; Avendaño-Herrera et al., 2006a; López et al., 2009; AbdEl-Galil and Hashiem, 2011; Rahman et al., 2014; Florio et al., 2016; Småge et al., 2016; and references therein). Affected fish usually display a variety of external signs including eroded mouth, skin ulcers, fin necrosis, and tail-rot. Skin lesions are often colonized by opportunistic pathogens such as *Vibrio* spp. So far, only one specific vaccine is commercially available to prevent tenacibaculosis in turbot. Hence, in all other fish species, the control of tenacibaculosis outbreaks remains restricted to the use of antibiotics, sometimes combined with external disinfectants (Avendaño-Herrera et al., 2008).

So far, three serotypes have been documented that show varying degrees of association with host fish species (Avendaño-Herrera et al., 2005a). This serological diversity could have important consequences for the development of an efficient vaccine. Recently, multilocus sequence analysis (MLSA) of *T. maritimum* isolates representative of the worldwide diversity revealed that this species constitutes a cohesive group, exhibiting moderate levels of nucleotide diversity and recombination [average pairwise nucleotide diversity ( $\pi$ ) estimated to be 0.44% and  $r/m$  ratio estimated to be 2.7]. Moreover, the population structure of *T. maritimum* did not reveal dominant genotypes or clonal complexes but rather suggested an endemic colonization of fish farms by local strains with no contribution of long-distance contamination related to fish movements. In addition, the same MLSA genotype was identified in different host species in the same geographical area, suggesting host versatility (Habib et al., 2014).

Despite the significance of tenacibaculosis outbreaks in the aquaculture industry, little is known about the virulence mechanisms of *T. maritimum* (Avendaño-Herrera et al., 2006b). Adhesion to hydrophobic surfaces (Burchard et al., 1990) or fish skin mucus (Magariños et al., 1995), hemagglutination (Pazos, 1997), extracellular products including proteolytic activity (Baxa et al., 1988; Handlinger et al., 1997; Pazos, 1997; van Gelderen et al., 2009), and iron uptake mechanisms (Avendaño-Herrera et al., 2005b) have been suggested to play roles in virulence. However, the molecular factors involved remain to be identified. Loss-of-function studies for experimental validation of genes as virulence factors are still inaccessible due to the absence of genetic tools.

In the present work, we sequenced and analyzed the complete genome of *T. maritimum* NCIMB 2154<sup>T</sup> to forecast the genes relevant to the bacterial lifestyle, in particular those linked to virulence. These *in silico* predictions paved the way for assessing for the first time the functional role of some relevant components. This genome will serve as a reference for future whole genome-based molecular epidemiology surveys aimed at analyzing disease emergence and propagation (Bayliss et al., 2017).

## MATERIALS AND METHODS

### Bacterial Growth Conditions

Several batches of the *T. maritimum* type strain (i.e., NCIMB 2154<sup>T</sup>, ATCC 43398<sup>T</sup>, CIP 103528<sup>T</sup>, and DSM 17995<sup>T</sup>),

*Tenacibaculum discolor* LL04 11.1.1<sup>T</sup>, *Tenacibaculum jejuense* CNURIC013<sup>T</sup>, and *Tenacibaculum soleae* LL04 12.1.7<sup>T</sup> were routinely grown in marine broth and agar 2216 (Difco) at 28°C and 170 rpm.

### Genome Sequencing

*Tenacibaculum maritimum* NCIMB 2154<sup>T</sup> was sequenced with a combination of PacBio RSII (N50 reads 7.4 kb, estimated coverage 234 x) and Illumina (HiSeq 2x100 pair-end reads with 300 bp insert size, 54,259,876 filtered sequences, estimated coverage 1500 x) reads and assembled with MHAP to completion to obtain a circular molecule. The final, quiver polished assembly was validated by optical mapping using *NcoI*.

### Annotation and Genome Comparisons

Genome annotation, including manual curation, and comparisons were performed using the web interface MicroScope (Vallenet et al., 2013) which allows graphic visualization enhanced by a synchronized representation of synteny groups<sup>1</sup>. Predictions of repeated sequences were performed using Repseek (Achaz et al., 2007) and those of genomic islands (GIs) using SIGI-HMM (Waack et al., 2006) and Alien hunter (Vernikos and Parkhill, 2006). The dbCAN database was used to identify carbohydrate active enzymes (CAZymes)<sup>2</sup> (Yin et al., 2012). The genomic sequence reported in this article has been deposited in the EMBL database under the accession number LT634361.

### MLST on Selected Strains

The four above-mentioned batches of the *T. maritimum* type strain were genotyped using the MLST scheme described in Habib et al. (2014).

### Chondroitin AC Lyase and Sphingomyelinase Cloning, Expression, and Enzymatic Activity

The genes encoding the chondroitin AC lyase (*csIA*, locus identifier: *MARIT\_2107*) and sphingomyelinase (*sph*, locus identifier: *MARIT\_1748*) were cloned according to Groisillier et al. (2010). Briefly, primers were designed to amplify the coding region corresponding to the catalytic module of CslA (forward primer 5'-TTTTTTAGATCTACTTCTCTAACTT TGGATGTAAATTCG-3'; reverse primer 5'-TTTTTTGAA TTCTTATATTTTAAGAACTTCTCTGTTATTAG-3') and *sph* (forward primer 5'-AAAAAAGGATCCAATGATGACGTTTCC CTTGGAGAAA-3'; reverse primer 5'-TTTTTTCAATTGTT AGTAGCTAAAGTAAAAAGTTTGCTTG-3') by PCR from *T. maritimum* genomic DNA. After digestion with the restriction enzymes *Bgl*III and *Eco*RI, and *Bam*HI and *Mfe*I respectively, the purified PCR products were ligated using the T4 DNA ligase into the expression vector pFO4 predigested by *Bam*HI and *Eco*RI (referred to as the plasmid pCslA and psph), resulting in a recombinant protein with a N-terminal hexa-histidine tag for each construct. The obtained plasmids were transformed

<sup>1</sup><http://www.genoscope.cns.fr/agc/mage/>

<sup>2</sup><http://csbl.bmb.uga.edu/dbCAN/index.php>

into *Escherichia coli* DH5 $\alpha$  for storage and in *E. coli* BL21(DE3) for protein expression. *E. coli* BL21(DE3) cells harboring the plasmid pCslA or psph were cultivated at 20°C in a 3 mL auto-induction ZYP 5052 medium (Studier, 2005) supplemented with 100  $\mu$ g/mL ampicillin. Cultures were stopped after 72 h and centrifuged for 35 min at 4°C, 3,000 g. The cells were resuspended in 500  $\mu$ L of buffer A (20 mM sodium phosphate pH 7.4, 500 mM NaCl, 10 mM imidazole). An anti-proteases mixture (cOmplete<sup>TM</sup> EDTA-free, Roche) and 0.1 mg/mL of DNase were added. The cells were disrupted by sonication. After centrifugation at 12,500 g for 2 h at 4°C the supernatant was loaded onto a His spin trap column (GE Healthcare Life Science) equilibrated with buffer A. After extensive washing with buffer A, the recombinant proteins were eluted with 400  $\mu$ L of buffer B (20 mM sodium phosphate pH 7.4, 500 mM NaCl, 500 mM imidazole). The results were analyzed by 12 % sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

### Chondroitin AC Lyase *In Vitro* Activity

The native chondroitin AC lyase activity of *T. maritimum* was assayed *in vitro* according to Li et al. (2015). Briefly, 6  $\mu$ L of mid-log-phase bacterial cultures (OD<sub>600</sub> = 0.6) were spotted on marine agar 2216 supplemented with 0.2% chondroitin sulfate A or C (Sigma) and 2% bovine serum albumin (BSA, Sigma), and incubated for 48 h at 28°C. Chondroitin lyase activity was visualized as a clear halo surrounding the bacterial growth after the plates were flooded with 0.35 N HCl.

The activity of the recombinant CslA was determined by adding 1  $\mu$ L of the recombinant protein solution to a 1-mL cuvette containing 600  $\mu$ L of 50 mM Tris-HCl, pH 8.0 supplemented by 1 mg/mL of chondroitin A or C at 30°C. Product formation was monitored as an increase in absorbance at 232 nm as a function of time (Michel et al., 2004). The assay was performed in triplicate.

### Sphingomyelinase Activity

The activity of the recombinant Sph was determined using a coupled assay Amplex Red Sphingomyelinase assay kit (Life Technologies, Invitrogen) following the manufacturer's instructions. The reactions were performed in 96-well special optics flat clear bottom black polystyrene Microplates (Corning). The reaction mixture (200  $\mu$ L) contained 100  $\mu$ L of 1.3 and 13 ng of recombinant protein and 100  $\mu$ L of 100  $\mu$ M Amplex red reagent (containing 2 U/mL horseradish peroxidase, 0.2 U/mL choline oxidase, 8 U/mL alkaline phosphatase, and 0.5 mM sphingomyelin). The fluorescence was measured every minute at excitation and emission wavelengths of 530 nm and 590 nm, respectively, using the TECAN Infinite<sup>®</sup> Pro200 microplate reader at 28°C for 15 min. The background fluorescence was corrected by subtracting the negative control (i.e., without recombinant protein). The positive control of each experiment was performed with *Bacillus cereus* sphingomyelinase provided by the manufacturer. The assay was performed in triplicate.

### Sialidase Activity

The fluorogenic substrate 2'-(4-methylumbelliferyl)- $\alpha$ -D-N-acetylneuraminic acid sodium salt hydrate (MUAN) was used

to determine sialidase activity according to Mally et al. (2008). Briefly, 10<sup>7</sup> mid-log phase bacteria were incubated with 0.1 mM of MUAN (Sigma) in 100 mM sodium acetate buffer pH 7.4 at 28°C. The reaction was stopped by the addition of a 0.5 M Na<sub>2</sub>CO<sub>3</sub> solution at pH 10. Released 4-methylumbelliferone was measured by fluorescence in a TECAN Infinite<sup>®</sup> Pro200 microplate reader at an excitation wavelength of 360 nm and an emission wavelength of 440 nm. The background fluorescence was corrected by subtracting the negative control (i.e., without bacteria). The assay was performed in triplicate.


## RESULTS AND DISCUSSION

### General Genome Features

The genome of *T. maritimum* NCIMB 2154<sup>T</sup> consists of a circular chromosome of 3,435,971 bp with a 32.01% GC content. No plasmid was identified. The chromosome is predicted to contain 2,866 protein-coding genes, 6 rDNA operons, and 57 tRNA (Figure 1). The minimal gene set includes well-conserved housekeeping genes for basic metabolism and macromolecular synthesis, many of which are essential (the list of these genes was taken from Gil et al., 2004). Accordingly, we formally identified 203 out of the 206 protein-coding genes proposed by these authors to represent this core minimal gene set. Strikingly, the genome is rich in repeat sequences, encompassing 10.35% of the genome. Twenty-five GIs were predicted (Supplementary Table S1), two of which (GI2 and GI9) are likely from phage origin (i.e., containing elements such as a tRNA border and motility genes, and displaying a compositional bias). Genes encoding rhs family proteins and rhs (rearrangement hot-spot) associated vgr proteins (or their remnants) were frequently identified in the predicted islands. Genes encoding Cas1, Cas2, and Cas9 were identified in GI19. However, the unconventional gene organization of the *cas* genes and the lack of short direct repeats interspersed with spacer sequences strongly argues for a functionally defective CRISPR system.

### The Genome of *T. maritimum* NCIMB 2154<sup>T</sup> Reveals Errors in Culture Collections

While this work was in progress, a WGS assembly of the *T. maritimum* type strain retrieved from the NBRC collection (NBRC 15946<sup>T</sup>; RefSeq assembly accession: GCF\_000509405.1) was released but not published. Its comparison with the complete genome of strain NCIMB 2154<sup>T</sup> presented in this study revealed unexpected sequence discrepancies as both cultures were supposed to represent the type strain. Using MLST (Habib et al., 2014), we confirmed that strains NCIMB 2154<sup>T</sup>, ATCC 43398<sup>T</sup>, and CIP 103528<sup>T</sup> all share identical sequences for the seven loci [corresponding to Sequence Type 1 (ST1)] and are, as predicted, most likely of the same origin. In contrast, strain NBRC 15946<sup>T</sup> and its derivative strain DSM 17995<sup>T</sup>, both possess the same sequences for the seven loci (corresponding to ST32) that differ from the sequences of ST1 at four loci (i.e., *atpA*,



**FIGURE 1 |** Circular representation of the *Tenacibaculum maritimum* NCIMB 2154<sup>T</sup> genome. Circles display (from the outside): (1) GC percent deviation (GC window – mean GC) in a 1,000-bp window. (2) Predicted CDSs transcribed in the clockwise direction. (3) Predicted CDSs transcribed in the counterclockwise direction. Genes displayed in (2) and (3) are color-coded according to the following categories: red and blue, MaGe validated annotations; purple, primary/automatic annotations. (4) GC skew (G + C/G – C) in a 1,000-bp window. (5) rRNA (blue), tRNA (green), miscellaneous other RNA (orange), transposable elements (pink), and pseudogenes (gray).

*dnaK*, *glyA*, and *gyrB*). According to the dates and order of deposition in the different culture collections, NCIMB 2154<sup>T</sup>, ATCC 43398<sup>T</sup>, and CIP 103528<sup>T</sup> represent the *bona fide* type strain (Supplementary Figure S1).

## Metabolism

Genome analysis indicated the presence of a complete Embden–Meyerhof–Parnas pathway, of the tricarboxylic acid cycle and of genes encoding NADH dehydrogenase subunits, cytochrome *c*, cytochrome *c* oxidase, components of ATP synthase genes as well as enzymes needed to synthesize amino acids, nucleotides, fatty acids, heme, vitamins, and coenzymes (e.g., biotin, farnesyl diphosphate, coenzyme A, NAD, FAD, dihydrofolate, mevalonate, and thiamin). However, and in contrast with *T. soleae*, the cobalamin biosynthesis encoding genes are absent in the *T. maritimum* genome. Nitrate reduction (Wakabayashi et al., 1986) is likely performed by the periplasmic, cytochrome *c*-linked, nitrate reductase complex NapAB (*MARIT\_1701-1700*).

A relevant characteristic of the lifestyle of pathogens is nutrient acquisition from their host. *T. maritimum* is able to degrade proteinaceous compounds (e.g., gelatin and casein) and to grow on casamino acids or tryptone as a sole carbon and nitrogen source (Wakabayashi et al., 1986). Accordingly, the *T. maritimum* genome encodes predicted secreted proteases (see below), peptide/amino acid transporters and peptide/amino acid catabolic pathways likely involved in protein degradation and uptake from host tissues. *T. maritimum* has been reported to be unable to degrade most simple and more complex carbohydrates (Wakabayashi et al., 1986; Avendaño-Herrera et al., 2006b). However, sugar transporters and CAZymes were predicted. Overall, the genome of *T. maritimum* encodes 59 CAZymes encompassing 18 glycosyl hydrolase, 30 glycosyl transferases, one polysaccharide lyase, six carbohydrate binding modules containing proteins, and four carbohydrate esterases (Supplementary Table S2).

Polysaccharide utilization loci (PUL), restricted to and very common within the phylum Bacteroidetes, are gene clusters involved in the capture, degradation, and import of complex carbohydrates. PUL-encoded proteins encompass a SusD-family cell-surface lipoprotein that binds the oligosaccharide, and a SusC-family TonB-dependent receptor for its transport across the bacterial outer membrane (Anderson and Salyers, 1989). Moreover, genes encoding SusC- and SusD-family proteins are usually organized in tandem in the genomes of Bacteroidetes (Terrapon et al., 2015). In full accordance with its inability to use carbohydrates, the *T. maritimum* genome presents a very low amount of *susC/susD* pairs compared to other members of the phylum Bacteroidetes (Barbeyron et al., 2016). Moreover, among the six identified *susC/susD* containing loci, only one harbors a typical PUL structure (*MARIT\_2678 - 2679*) and is predicted to be involved in glycan harvesting from host glycoproteins (see below).

## Iron Acquisition and Utilization

Iron acquisition from host plays an important role in virulence of many pathogenic bacteria. In biological systems, high-affinity iron-binding proteins can chelate iron, and pathogens have developed efficient mechanisms to obtain iron from their hosts (Ratledge and Dover, 2000). In the *T. maritimum* genome, we identified a siderophore biosynthesis gene cluster (*MARIT\_0169-0174*) highly similar to the *mbs* locus from a deep-sea metagenome (Fujita et al., 2012). As this gene cluster is predicted to be involved in the production of the macrocyclic hydroxamate class bisucaberin siderophore, we named the genes *tbs* for *Tenacibaculum* bisucaberin synthase. However, the gene organization is different from those previously reported displaying a major facilitator-family exporter-encoding gene and a duplication/fusion of the *tbsCD* gene (Supplementary Figure S2). A highly similar locus is observed in the genome of the *Tenacibaculum mesophilum* type strain (CIP 107215<sup>T</sup>; data not shown), likely responsible of the bisucaberin B siderophore biosynthesis as proposed by Fujita et al. (2013). Among the numerous TonB-dependent outer membrane receptors encoded in the *T. maritimum* genome, *MARIT\_0185*, located in the *tbs* locus neighborhood, likely encodes the bisucaberin siderophore-iron transporter.

In the human periodontal bacterium *Porphyromonas gingivalis*, the heme-binding lipoprotein HmuY, together with the outer-membrane receptor HmuR, are predicted to be virulence factors during bacterial infection (Olczak et al., 2008; Wójtowicz et al., 2009). In the *T. maritimum* genome, two genes (*MARIT\_1312-1313*), organized in tandem, encode HmuR and HmuY homologous proteins and are predicted to be involved in heme uptake.

In addition, *MARIT\_0141 - 0142* encoding FeoAB likely constitute a Fe<sup>2+</sup> uptake system (Lau et al., 2016) and two iron-regulated protein homologous genes (*MARIT\_1664* and *MARIT\_1661*), belonging to the imelysin family, might also be involved in iron acquisition, uptake or storage (Xu et al., 2011). The control of iron metabolism is likely carried out by the ferric uptake regulator Fur (*MARIT\_1835*; Fillat, 2014). Hence, such a variety of iron acquisition systems strongly suggests the

ability of this bacterium to survive under poor iron conditions (sea water) and/or to retrieve iron sequestered by host proteins (Avendaño-Herrera et al., 2005b).

## Motility, Adhesion, Quorum Sensing/Quenching, and Stress Response

Like most members of the family Flavobacteriaceae, *T. maritimum* moves over surfaces by gliding motility, an active process that does not involve pili or flagella. The genome of *T. maritimum* encodes all the proteins that form the gliding machinery, i.e., the 14 *gld* genes (*gldA* to *gldN*; McBride et al., 2009) and 10 *spr* genes (*sprA*, *sprB*, *sprC*, *sprD*, *sprE*, and five *sprF* paralogs; McBride and Zhu, 2013). *T. maritimum* NCIMB 2154<sup>T</sup> is extremely adherent to different surfaces including agar, plastic, and glass. Genes encoding (i) the biosynthesis of exopolysaccharides (*MARIT\_2522-2537*); (ii) the numerous adhesins ( $n = 17$ ); and (iii) the proteins displaying lectin or carbohydrate-binding motifs could be involved in these strong adhesive properties, in the biofilm-forming ability and in the hemagglutination properties of the bacterium (Pazos, 1997).

*Quorum sensing* is a bacterial communication process that controls a range of functions at the population level. In Gram-negative bacteria, the most studied *quorum sensing* system comprises the production and detection of acyl homoserine lactones (AHLs), diffusible compounds that act as signaling molecules between cells (Garg et al., 2014). Though AHL production was previously reported in *T. maritimum* (Romero et al., 2010), no homologous gene for AHL biosynthesis was detected in its genome. In contrast, *quorum quenching* refers to all processes involved in the inhibition of bacterial communication (Kalia, 2013). A *N*-acyl homoserine lactonase encoding gene (GenBank: KR232938.1) belonging to the metallo- $\beta$ -lactamase family has been proposed to be the quorum quencher of *T. maritimum* (Mayer et al., 2015). However, this gene is definitively absent from the *T. maritimum* genome and one must conclude that KR232938.1 does not belong to *T. maritimum* but rather to another fish pathogen, *T. discolor* (99.77 % nucleotide sequence identity).

Pathogenic bacteria have to adapt to the changing environments between their different lifestyles and to cope with various stresses including reactive oxygen species (ROS) produced by host macrophages. The genome of *T. maritimum* encodes three superoxide dismutases (SodA, SodB, and SodC). Most bacteria possess either a manganese-dependent (SodA) or an iron-dependent (SodB) superoxide dismutase in their cytoplasm, while zinc-dependent superoxide dismutases (SodC) have been detected mostly in pathogenic bacteria (Sheng et al., 2014). These enzymes convert superoxide anions to molecular oxygen and hydrogen peroxide, to be further metabolized by catalases or peroxidases. The presence of the three types of superoxide dismutases and two catalase/peroxidase (KatA and KatG) suggests that *T. maritimum* uses a sophisticated mechanism to face up oxidative stress. In addition, three loci involved in bacterial resistance to heavy metals have been


identified: (i) *MARIT\_0364-0366*, similar to the drug efflux system AcrA–AcrB–TolC of *E. coli* (Lee et al., 2012); (ii) *MARIT\_1200* encoding a putative arsenate reductase; and (iii) *MARIT\_1768-1771* encoding a heavy metal efflux pump-type ATPase. These loci are likely involved in the removal of cationic heavy metals to limit the production of ROS by the Fenton reaction.

## Transport and Secretion Systems

Transport systems are of great significance for virulence by addressing toxins to the bacterial surface. ABC-type transport systems, the Sec-dependent transport system, and the twin-arginine transport system were identified. In the phylum Bacteroidetes, the type IX secretion system (T9SS) allows the delivery of proteins to the cell surface (McBride and Zhu, 2013). All previously characterized components of the T9SS were identified in the *T. maritimum* genome. In *P. gingivalis*, the T9SS-secreted proteins comprise many virulence factors, including the extracellular and cell-surface cysteine proteinases gingipains (Sato et al., 2010, 2013). These T9SS-secreted proteins possess a conserved C-terminal domain (CTD), involved in secretion and cell-surface anchoring. These 70–100 amino acids long CTDs belong to two different TIGRFAM protein domain families, TIGR04183 and TIGR04131 (McBride and Nakane, 2015). The *T. maritimum* genome encompasses eight genes encoding TIGR04131-containing proteins (Supplementary Table S3). Most, if not all, are predicted to be adhesins, including SprB (*MARIT\_1321*), which is also required for gliding motility (Nelson et al., 2008). In addition, several predicted toxins were identified among the 43 genes encoding TIGR04183-containing proteins.

## Toxins


As *T. maritimum* is a pathogenic bacterium, this species should possess sophisticated mechanisms to invade and colonize host tissues. Accordingly, the *T. maritimum* genome encodes a bunch of predicted toxins and virulence factors including membrane-damaging enzymes potentially involved in host cells lysis. A gene encoding a sphingomyelinase with a lipoprotein signal (*MARIT\_1748*) homologous (30.2% identity/50% similarity) to the one of *B. cereus* was identified (Supplementary Figure S3A). A gene encoding a ceramidase with a signal peptide and a TIGR04183 domain (*MARIT\_2033*) homologous (33.5% identity/59.8% similarity) to the one of *Pseudomonas aeruginosa* (Supplementary Figure S3B) is also present in the *T. maritimum* genome. Sphingomyelinase has been reported to be cytotoxic to host cells by acting as a potent hemolytic factor (Oda et al., 2010), while the bacterial ceramidase functions as an exotoxin or activator of exotoxin (Okino et al., 2010; Ito et al., 2014). Indeed, the outer layer of the plasma membrane of eukaryotic cells contains phospholipids, which are hydrolyzed to phosphocholine and ceramide by sphingomyelinase, the latter being subsequently hydrolyzed to sphingosine and fatty acids by a ceramidase. To formally demonstrate that *MARIT\_1748* encodes the sphingomyelinase, we cloned the corresponding nucleotide sequence in the pFO4 vector (Groisillier et al., 2010). The recombinant protein



**FIGURE 2 |** *Tenacibaculum maritimum* sphingomyelinase activity. Fluorescence measurement (arbitrary units) following incubation of 1.3 ng (blue line) of recombinant protein with Amplex Red Sphingomyelinase assay kit. The control conditions were as follows: negative control, 1.3 ng of boiled recombinant protein (green line); positive control, the purified sphingomyelinase from *Bacillus cereus* provided by the manufacturer (red line). Results correspond to the mean of triplicates and SDs are included.


was produced in a soluble form in *E. coli* BL21(*DE3*) and the enzymatic activity of the purified sphingomyelinase was assayed in triplicate using the Amplex Red Sphingomyelinase assay kit (Figure 2). Another predicted hemolysin is encoded by *MARIT\_0124* and belongs to the cholesterol-dependent cytolysin family (Gilbert, 2010). These pore-forming toxins were originally identified in Gram-positive bacteria and encompass well-known examples including listeriolysin, perfringolysin, streptolysin, and pneumolysin (Los et al., 2013).

Glycosaminoglycans (GAGs) are highly sulfated polymers composed of repeated disaccharide units (an amino sugar and an uronic sugar). They represent major components of animal cell surface and extracellular matrix, mostly in the form of proteoglycans. Among them, chondroitin sulfate is an important component of cartilage and fish connective tissue (Arima et al., 2013). It is composed of a chain of alternating *N*-acetylgalactosamine and glucuronic acid to which proteins attach. Chondroitin sulfate lyases have been suggested to be virulence factors, for instance in the other fish pathogen *Flavobacterium columnare* (Suomalainen et al., 2006). One might predict that the *csIA* gene (*MARIT\_2107*) encoding a PL8\_3 family chondroitin AC lyase highly similar to that of *F. columnare* plays a similar role. *In vitro* analyses demonstrated that *T. maritimum* is able to degrade chondroitin sulfate A and C on marine agar 2216, as showed by the formation of a degradation halo around the bacterial growth (Figure 3A). Although a chondroitin C-lyase activity was recently suggested for *T. maritimum* (Rahman et al., 2014), our results demonstrate the ability of this bacterium to also degrade chondroitin sulfate A. Other phylogenetically close *Tenacibaculum* species, such as *T. discolor*, *T. jejuense*, or *T. soleae* do not display this chondroitin AC lyase activity under the same conditions. To formally demonstrate that gene *csIA* encodes the chondroitin



AC lyase, we cloned the nucleotide sequence corresponding to the PL8\_3 catalytic module in the pFO4 vector (Groisillier et al., 2010). The recombinant protein, referred to as *TmCslA*<sub>PL8</sub>, was produced in a soluble form in *E. coli* BL21(DE3). The enzymatic activity of the purified *TmCslA*<sub>PL8</sub> was assayed in triplicate by measuring the increase in absorbance at 232 nm of the reaction products using chondroitin A and C sulfates as substrates. As seen in **Figure 3B**, *TmCslA*<sub>PL8</sub> is highly active on both substrates, confirming the functional annotation of *MARIT\_2107*. As no sulfatase could be identified in the genome, it is likely that GAGs such as chondroitin sulfate cannot be assimilated by *T. maritimum*. Therefore, *CslA* might be a *bona fide* virulence factor allowing the pathogen to invade fish tissues.

*Capnocytophaga canimorsus*, another member of the family Flavobacteriaceae, is a commensal of cat and dog mouth that can cause dramatic infections in bitten humans (Pers et al., 1996). *C. canimorsus* has the unusual property to feed directly on cultured mammalian cells by harvesting the glycan moiety of cellular glycoproteins, a property dependent on *SiaC* (Mally



et al., 2008). Sialic acids are predominantly found in cell-surface exposed and secreted eukaryotic glycoproteins, being involved in many physiological, biological, and immunological functions (Varki and Varki, 2007). Mucosal surfaces are especially sialoglycan-rich and bacterial sialidases play important roles during the colonization and damage of mammalian mucosal surfaces (Lewis and Lewis, 2012). In *T. maritimum*, *siaA* (*MARIT\_2686*) encodes a predicted sialidase, which activity was formally demonstrated using the fluorogenic substrate MUAN (**Figure 4**). In contrast, other *Tenacibaculum* species including the two fish pathogens *T. soleae* and *T. discolor*, for which no *siaA* homologous gene has been identified (data not shown), were unable to degrade MUAN under the same experimental conditions. Indeed, the *siaA* gene is encompassed in a [GI N° 24 inserted in a Gln-tRNA (position 2,965,888), Supplementary Table S1] and has a predicted foreign origin. The 3' part of this GI (2,916,321–2,965,888) is mainly composed of pseudogenes including scars of transposases, Vgr family proteins and Rhs family proteins. On the other hand, the 5' part of this GI (2,916,321–2,915,448) contains nine *bona fide* genes, predicted to encode a PUL encompassing (i) a SusC/SusD outer-membrane importer system; (ii) a *N*-acetylneuraminatase lyase; (iii) a *N*-acyl-D-glucosamine 2-epimerase; and (iv) a *N*-acetylneuraminatase epimerase/sodium:sialic acid symporter-fusion inner-membrane protein. This PUL system is likely dedicated to the harvesting, import, and catabolism of sialic acids from host glycoproteins (Supplementary Figure S4). Intriguingly, the sialidase displays a very unique structure including (i) a signal peptide; (ii) a carbohydrate esterase family 6 domain; (iii) a family 40 carbohydrate-binding module; (iv) two adjacent, fully duplicated, family 33 glycoside hydrolase domains; and (v) a TIGR04183 domain for T9SS-mediated secretion and cell-surface anchoring. Moreover, the predicted mechanism of glycan harvesting by *T. maritimum* is likely different from the

**TABLE 1** | Summary of the predicted virulence-associated genes identified in this study in the genome of *T. maritimum* NCIMB 2154<sup>T</sup>.

Label	Predicted function	Activity present in the following species	Activity suspected in <i>T. maritimum</i>
<b>Iron uptake</b>			
MARIT_0169-0174	Siderophore biosynthesis system, Tbs	Deep sea metagenome (Fujita et al., 2012)	Avendaño-Herrera et al., 2005b
MARIT_1312-1313	Heme uptake mechanism, HmuYR	<i>Porphyromonas gingivalis</i> (Olczak et al., 2008; Wójtowicz et al., 2009)	Avendaño-Herrera et al., 2005b
MARIT_0141 – 0142	Fe <sup>2+</sup> uptake system, FeoAB	<i>Porphyromonas gingivalis</i> (Lau et al., 2016)	Avendaño-Herrera et al., 2005b
MARIT_1664	Iron acquisition, uptake or storage, imelysin family protein	<i>Synechococcus elongatus</i> (Xu et al., 2011)	Avendaño-Herrera et al., 2005b
MARIT_1661	Iron acquisition, uptake or storage, imelysin family protein	<i>Synechococcus elongatus</i> (Xu et al., 2011)	Avendaño-Herrera et al., 2005b
<b>Stress resistance</b>			
MARIT_3105	Superoxide dismutase [Mn/Fe], SodA	<i>Staphylococcus carnosus</i> (Barrière et al., 2001)	Suzuki et al., 2001
MARIT_1670	Superoxide dismutase [Fe], SodB	<i>Legionella pneumophila</i> (Amemura-Maekawa et al., 1996)	Suzuki et al., 2001
MARIT_1821	Superoxide dismutase [Cu-Zn], SodC	<i>Schistosoma mansoni</i> (da Silva et al., 1992)	Suzuki et al., 2001
MARIT_0946	KatG catalase	<i>Geobacillus stearothermophilus</i> (Singh et al., 2008)	Suzuki et al., 2001
MARIT_2408	KatA catalase	<i>Pseudomonas aeruginosa</i> (Brown et al., 1995)	Suzuki et al., 2001
<b>Toxins</b>			
MARIT_0124	Cholesterol-dependent cytolysin	<i>Capnocytophaga canimorsus</i> (Gilbert, 2010)	Baxa et al., 1988
MARIT_1085	Collagenase	<i>Cytophaga</i> sp. (Sasagawa et al., 1995)	Baxa et al., 1988
MARIT_1748	Sphingomyelinase	<i>Bacillus cereus</i> (Oda et al., 2010)	
MARIT_2033	Ceramidase	<i>Pseudomonas aeruginosa</i> (Ito et al., 2014)	
MARIT_2107	Chondroitin AC lyase	<i>Flavobacterium columnare</i> (Suomalainen et al., 2006)	Rahman et al., 2014
MARIT_2328	Streptopain family protease	<i>Streptococcus pyogenes</i> (Nelson et al., 2011)	Wakabayashi et al., 1986; Baxa et al., 1988
MARIT_2678 - 2687	Sialoglycan degradation and uptake	<i>Capnocytophaga canimorsus</i> (Mally et al., 2008)	

one identified in *C. canimorsus*. Indeed, the *T. maritimum* genome is devoid of the *gpdCDGEF* operon encoded by the *C. canimorsus* PUL5 and involved in deglycosylation and import of N-linked oligosaccharides (Renzi et al., 2011). In addition, the *C. canimorsus* sialidase is a periplasmic-exposed lipoprotein that processes oligosaccharides after SucC/SusD-mediated import, whereas the *T. maritimum* sialidase is predicted to be cell-surface exposed, therefore directly processing sialic acid from glycoproteins. Hence, one might predict that both species perform the same function (foraging host glycoproteins) using common strategies (sialidase, PUL) but in sequentially different ways.

Extracellular proteases may exhibit a wide range of virulence potentials when interacting with the host defense mechanisms and tissue components. Furthermore, they may promote the survival of pathogens under adverse environmental conditions encountered in the infected host (Dubin, 2002). Since early studies, *T. maritimum* was shown to be proteolytic (Wakabayashi et al., 1986) and its proteases were suspected to act synergistically with other virulence factors, leading to tissue destruction and mortality (Baxa et al., 1988). Eight cell-surface-exposed, TIGR04183 domain-containing proteases were predicted (Supplementary Table S3), two of which likely of importance. MARIT\_2328 encodes a multi-domain protein encompassing a C10 family peptidase, highly similar to streptopain (SpeB), an important streptococcal virulence factor likely playing a role in bacterial colonization, invasion, and inhibition of wound healing (Nelson et al., 2011). MARIT\_1085 encodes a collagenase

similar to that of *Cytophaga* sp. strain L43-1 (Sasagawa et al., 1995).

## CONCLUSIONS AND PERSPECTIVES

We report here the complete genome sequence of *T. maritimum*, a serious pathogen of marine fish in many geographical areas. *T. maritimum* shows a lack of host specificity, affecting a variety of wild and farmed fish species (Avendaño-Herrera et al., 2006b). Sequence analysis has revealed a combination of strategies that probably confers *T. maritimum* the ability to invade, colonize, and degrade fish tissues and to exploit some cellular compounds for growth. The central metabolism of *T. maritimum* is similar to that of the other flavobacteria sequenced to date (e.g., several *Flavobacterium*, *Gramella*, *Dokdonia*, and *Polaribacter* species). However, *T. maritimum* does not possess a proteorhodopsin-encoding gene as identified in close relatives such as *Polaribacter*, *Dokdonia*, or *Psychroflexus* species, suggesting the inability of this bacterium to use light to generate proton motive force. Comparison with the available genomes of the three other fish-pathogenic *Tenacibaculum* species *Tenacibaculum dicentrarchi*, *Tenacibaculum ovolyticum*, and *T. soleae* (Grothusen et al., 2016; Lujan et al., 2016; Teramoto et al., 2016), has revealed striking differences in virulence strategies as most, if not all, the aforementioned predicted toxins (Table 1) are absent from the genomes of the latter species. These elements point to very different paths in the evolution of virulence as suggested using

a subset of core-genome genes (Habib et al., 2014). The genome sequence of *T. maritimum* provides insights into the lifestyle of this poorly studied pathogen and may help in the development of efficient control strategies in fish farms. Indeed, the predicted virulence factors could lead to the development of attenuated *T. maritimum* variants for vaccine development. The genome of the type strain may also serve as a reference for future genomic comparisons for a better understanding of intraspecies and intragenus diversity and evolution as well as whole genome-based molecular epidemiology studies (Bayliss et al., 2017).

## AUTHOR CONTRIBUTIONS

DP-P performed genome annotation, phenotypic characterization, and drafting the manuscript; AL, AR, and CL-R performed DNA extraction, library construction, and sequencing; GhM performed the optical-mapping; ZR participated in genomic data analysis; RL, TB, AG, and GuM performed gene cloning, protein expression, and biochemical characterization with substantial intellectual contribution; J-FB substantial intellectual contribution throughout the study, data analysis, and manuscript preparation. ED substantial intellectual contribution throughout the study, gene mining, interpretation of data, manuscript preparation, and responsible for acquisition of funding. All authors read and approved the final manuscript.

## REFERENCES

- AbdEl-Galil, M. A. A., and Hashiem, M. (2011). Tenacibaculosis in Picasso tigger fish (*Rhinecanthus assasi*) and black damselfish (*Neoglyphidodon niger*) of Red sea at Hurghada, Egypt. *Life Sci. J.* 8, 1166–1171.
- Achaz, G., Boyer, F., Rocha, E. P., Viari, A., and Coissac, E. (2007). Repseek, a tool to retrieve approximate repeats from large DNA sequences. *Bioinformatics* 23, 119–121. doi: 10.1093/bioinformatics/btl519
- Amemura-Maekawa, J., Kura, F., and Watanabe, H. (1996). Cloning and nucleotide sequences of iron and copper-zinc superoxide dismutase genes of *Legionella pneumophila* and their distribution among *Legionella* species. *Jpn. J. Med. Sci. Biol.* 49, 167–186.
- Anderson, K. L., and Salyers, A. A. (1989). Genetic evidence that outer membrane binding of starch is required for starch utilization by *Bacteroides thetaiotaomicron*. *J. Bacteriol.* 171, 3199–3204. doi: 10.1128/jb.171.6.3199-3204.1989
- Arima, K., Fujita, H., Toita, R., Imazu-Okada, A., Tsutsumishita-Nakai, N., Takeda, N., et al. (2013). Amounts and compositional analysis of glycosaminoglycans in the tissue of fish. *Carbohydr. Res.* 366, 25–32. doi: 10.1016/j.carres.2012.11.010
- Avendaño-Herrera, R., Magariños, B., Morfiño, M., Romalde, J. L., and Toranzo, A. E. (2005a). A novel O-serotype in *Tenacibaculum maritimum* strains isolated from cultured sole (*Solea senegalensis*). *Bull. Eur. Assoc. Fish Pathol.* 25, 70–74.
- Avendaño-Herrera, R., Núñez, S., Barja, J. L., and Toranzo, A. E. (2008). Evolution of drug resistance and minimum inhibitory concentration to enrofloxacin in *Tenacibaculum maritimum* strains isolated in fish farms. *Aquac. Int.* 16, 1–11. doi: 10.1007/s10499-007-9117-y
- Avendaño-Herrera, R., Toranzo, A. E., and Magariños, B. (2006a). A challenge model for *Tenacibaculum maritimum* infection in turbot, *Scophthalmus maximus* (L.). *J. Fish Dis.* 29, 371–374. doi: 10.1111/j.1365-2761.2006.00712.x
- Avendaño-Herrera, R., Toranzo, A. E., and Magariños, B. (2006b). Tenacibaculosis infection in marine fish caused by *Tenacibaculum maritimum*: a review. *Dis. Aquat. Organ.* 71, 255–266.

## FUNDING

This work was supported by the Agence Nationale pour la Recherche (contract ANR-14-CE19-0020). It has benefited from the expertise of the High-throughput Sequencing platform of I2BC (<http://www.i2bc.paris-saclay.fr>) and GeT core facility platform (<http://get.genotoul.fr>) and was supported by the France Génomique national infrastructure, funded as part of “Investissement d’avenir” program managed by Agence Nationale pour la Recherche (contract ANR-10-INBS-09).

## ACKNOWLEDGMENTS

We are thankful to Institut Français de Bioinformatique (IFB), V. Barbe (CEA/Genoscope) for the optical mapping, V. Loux (Malage) from the INRA MIGALE bioinformatics platform (<http://migale.jouy.inra.fr>) for help with bioinformatics and C. Klopp (GenoToul Bioinfo) for his contribution to genome assembly.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2017.01542/full#supplementary-material>

- Avendaño-Herrera, R., Toranzo, A. E., Romalde, J. L., Lemos, M. L., and Magariños, B. (2005b). Iron uptake mechanisms in the fish pathogen *Tenacibaculum maritimum*. *Appl. Environ. Microbiol.* 71, 6947–6953.
- Barbeyron, T., Thomas, F., Barbe, V., Teeling, H., Schenowitz, C., Dossat, C., et al. (2016). Habitat and taxon as driving forces of carbohydrate catabolism in marine heterotrophic bacteria: example of the model algae-associated bacterium *Zobellia galactanivorans* Dsij(T). *Environ. Microbiol.* 18, 4610–4627. doi: 10.1111/1462-2920.13584
- Barrière, C., Leroy-Sétrin, S., and Talon, R. (2001). Characterization of catalase and superoxide dismutase in *Staphylococcus carnosus* 833 strain. *J. Appl. Microbiol.* 91, 514–519. doi: 10.1046/j.1365-2672.2001.01411.x
- Baxa, D. V., Kawai, K., and Kusuda, R. (1988). *In vitro* and *in vivo* activities of *Flexibacter maritimus* toxins. *Rep. United States Mar. Biol. Inst. Kochi Univ.* 10, 1–8.
- Bayliss, S. C., Verner-Jeffreys, D. W., Bartie, K. L., Aanensen, D. M., Sheppard, S. K., Adams, A., et al. (2017). The promise of whole genome pathogen sequencing for the molecular epidemiology of emerging aquaculture pathogens. *Front. Microbiol.* 8:121. doi: 10.3389/fmicb.2017.00121
- Bernardet, J.-F., Campbell, A. C., and Buswell, J. A. (1990). *Flexibacter maritimus* is the agent of “black patch necrosis” in Dover sole in Scotland. *Dis. Aquat. Org.* 8, 233–237. doi: 10.3354/dao008233
- Brown, S. M., Howell, M. L., Vasil, M. L., Anderson, A. J., and Hasset, D. J. (1995). Cloning and characterization of the *katB* gene of *Pseudomonas aeruginosa* encoding a hydrogen peroxide-inducible catalase: purification of KatB, cellular localization, and demonstration that it is essential for optimal resistance to hydrogen peroxide. *J. Bacteriol.* 177, 6536–6544.
- Burchard, R. P., Rittschof, D., and Bonaventura, J. (1990). Adhesion and motility of gliding bacteria on substrata with different surface free energies. *Appl. Environ. Microbiol.* 56, 2529–2534.
- da Silva, A. C., LePresle, T., Capron, A., and Pierce, R. J. (1992). Molecular cloning of a 16-kilodalton Cu/Zn superoxide dismutase from *Schistosoma mansoni*. *Mol. Biochem. Parasitol.* 52, 275–278. doi: 10.1016/0166-6851(92)90060-W
- Dubin, G. (2002). Extracellular proteases of *Staphylococcus* spp. *Biol. Chem.* 383, 1075–1086. doi: 10.1515/BC.2002.116

- Fillat, M. F. (2014). The FUR (ferric uptake regulator) superfamily: diversity and versatility of key transcriptional regulators. *Arch. Biochem. Biophys.* 546, 41–52. doi: 10.1016/j.abb.2014.01.029
- Florio, D., Gridelli, S., Fioravanti, M. L., and Zanoni, R. G. (2016). First isolation of *Tenacibaculum maritimum* in a captive sand tiger shark (*Carcharias taurus*). *J. Zoo Wildl. Med.* 47, 351–353. doi: 10.1638/2015-0064.1
- Fujita, M. J., Kimura, N., Yokose, H., and Otsuka, M. (2012). Heterologous production of bisucaberin using a biosynthetic gene cluster cloned from a deep sea metagenome. *Mol. Biosyst.* 8, 482–485. doi: 10.1039/c1mb05431g
- Fujita, M. J., Nakano, K., and Sakai, R. (2013). Bisucaberin B, a linear hydroxamate class siderophore from the marine bacterium *Tenacibaculum mesophilum*. *Molecules* 18, 3917–3926. doi: 10.3390/molecules18043917
- Garg, N., Manchanda, G., and Kumar, A. (2014). Bacterial quorum sensing: circuits and applications. *Antonie Van Leeuwenhoek* 105, 289–305. doi: 10.1007/s10482-013-0082-3
- Gil, R., Silva, F. J., Pereté, J., and Moya, A. (2004). Determination of the core of a minimal bacterial gene set. *Microbiol. Mol. Biol. Rev.* 68, 518–537. doi: 10.1128/MMBR.68.3.518-537.2004
- Gilbert, R. J. (2010). Cholesterol-dependent cytolysins. *Adv. Exp. Med. Biol.* 677, 56–66. doi: 10.1007/978-1-4419-6327-7\_5
- Groisillier, A., Hervé, C., Jeudy, A., Rebuffet, E., Pluchon, P. F., Chevolut, Y., et al. (2010). MARINE-EXPRESS: taking advantage of high throughput cloning and expression strategies for the post-genomic analysis of marine organisms. *Microb. Cell Fact.* 9:45. doi: 10.1186/1475-2859-9-45
- Grothusen, H., Castillo, A., Henríquez, P., Navas, E., Bohle, H., Araya, C., et al. (2016). First complete genome sequence of *Tenacibaculum dicentrarchi*, an emerging bacterial pathogen of salmonids. *Genome Announc.* 4:e01756–15. doi: 10.1128/genomeA.01756-15
- Habib, C., Houel, A., Lunazzi, A., Bernardet, J. F., Olsen, A. B., Nilsen, H., et al. (2014). Multilocus sequence analysis of the marine bacterial genus *Tenacibaculum* suggests parallel evolution of fish pathogenicity and endemic colonization of aquaculture systems. *Appl. Environ. Microbiol.* 80, 5503–5514. doi: 10.1128/AEM.01177-14
- Handler, J., Soltani, M., and Percival, S. (1997). The pathology of *Flexibacter maritimus* in aquaculture species in Tasmania, Australia. *J. Fish Dis.* 20, 159–168. doi: 10.1046/j.1365-2761.1997.00288.x
- Ito, M., Okino, N., and Tani, M. (2014). New insight into the structure, reaction mechanism, and biological functions of neutral ceramidase. *Biochim. Biophys. Acta* 1841, 682–691. doi: 10.1016/j.bbali.2013.09.008
- Kalia, V. C. (2013). Quorum sensing inhibitors: an overview. *Biotechnol. Adv.* 31, 224–245. doi: 10.1016/j.biotechadv.2012.10.004
- Lau, C. K., Krewulak, K. D., and Vogel, H. J. (2016). Bacterial ferrous iron transport: the Feo system. *FEMS Microbiol. Rev.* 40, 273–298. doi: 10.1093/femsre/fuv049
- Lee, M., Jun, S. Y., Yoon, B. Y., Song, S., Lee, K., and Ha, N. C. (2012). Membrane fusion proteins of type I secretion system and tripartite efflux pumps share a binding motif for TolC in gram-negative bacteria. *PLoS ONE* 7:e40460. doi: 10.1371/journal.pone.0040460
- Lewis, A. L., and Lewis, W. G. (2012). Host sialoglycans and bacterial sialidases: a mucosal perspective. *Cell Microbiol.* 14, 1174–1182. doi: 10.1111/j.1462-5822.2012.01807.x
- Li, N., Qin, T., Zhang, X. L., Huang, B., Liu, Z. X., Xie, H. X., et al. (2015). Gene deletion strategy to examine the involvement of the two chondroitin lyases in *Flavobacterium columnare* virulence. *Appl. Environ. Microbiol.* 81, 7394–7402. doi: 10.1128/AEM.01586-15
- López, J. R., Núñez, S., Magariños, B., Castro, N., Navas, J. I., de la Herran, R., et al. (2009). First isolation of *Tenacibaculum maritimum* from wedge sole, *Dicologlossa cuneata* (Moreau). *J. Fish Dis.* 32, 603–610. doi: 10.1111/j.1365-2761.2009.01029.x
- Los, F. C., Randis, T. M., Aroian, R. V., and Ratner, A. J. (2013). Role of pore-forming toxins in bacterial infectious diseases. *Microbiol. Mol. Biol. Rev.* 77, 173–207. doi: 10.1128/MMBR.00052-12
- Lujan, K. M., Eisen, J. A., and Coil, D. A. (2016). Draft genome sequence of *Tenacibaculum soleae* UCD-KL19. *Genome Announc.* 4:e01120–16. doi: 10.1128/genomeA.01120-16
- Magariños, B., Pazos, F., Santos, Y., Romalde, J. L., and Toranzo, A. E. (1995). Response of *Pasteurella piscicida* and *Flexibacter maritimus* to skin mucus of marine fish. *Dis. Aquat. Organ.* 21, 103–108. doi: 10.3354/dao021103
- Mally, M., Shin, H., Paroz, C., Landmann, R., and Cornelis, G. R. (2008). *Capnocytophaga canimorsus*: a human pathogen feeding at the surface of epithelial cells and phagocytes. *PLoS Pathog.* 4:e1000164. doi: 10.1371/journal.ppat.1000164
- Mayer, C., Romero, M., Muras, A., and Otero, A. (2015). Aii20J, a wide-spectrum thermostable N-acylhomoserine lactonase from the marine bacterium *Tenacibaculum* sp. 20J, can quench AHL-mediated acid resistance in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 99, 9523–9539. doi: 10.1007/s00253-015-6741-8
- McBride, M. J., and Nakane, D. (2015). *Flavobacterium* gliding motility and the type IX secretion system. *Curr. Opin. Microbiol.* 28, 72–77. doi: 10.1016/j.mib.2015.07.016
- McBride, M. J., Xie, G., Martens, E. C., Lapidus, A., Henrissat, B., Rhodes, R. G., et al. (2009). Novel features of the polysaccharide-digesting gliding bacterium *Flavobacterium johnsoniae* as revealed by genome sequence analysis. *Appl. Environ. Microbiol.* 75, 6864–6875. doi: 10.1128/AEM.01495-09
- McBride, M. J., and Zhu, Y. (2013). Gliding motility and Por secretion system genes are widespread among members of the phylum Bacteroidetes. *J. Bacteriol.* 195, 270–278. doi: 10.1128/JB.01962-12
- Michel, G., Pojasek, K., Li, Y., Sulea, T., Linhardt, R. J., Raman, R., et al. (2004). The structure of chondroitin B lyase complexed with glycosaminoglycan oligosaccharides unravels a calcium-dependent catalytic machinery. *J. Biol. Chem.* 279, 32882–32896. doi: 10.1074/jbc.M403421200
- Nelson, D. C., Garbe, J., and Collin, M. (2011). Cysteine proteinase SpeB from *Streptococcus pyogenes* - a potent modifier of immunologically important host and bacterial proteins. *Biol. Chem.* 392, 1077–1088. doi: 10.1515/BC.2011.208
- Nelson, S. S., Bollampalli, S., and McBride, M. J. (2008). SprB is a cell surface component of the *Flavobacterium johnsoniae* gliding motility machinery. *J. Bacteriol.* 190, 2851–2857. doi: 10.1128/JB.01904-07
- Oda, M., Takahashi, M., Matsuno, T., Uoo, K., Nagahama, M., and Sakurai, J. (2010). Hemolysis induced by *Bacillus cereus* sphingomyelinase. *Biochim. Biophys. Acta* 1798, 1073–1080. doi: 10.1016/j.bbame.2010.03.004
- Okino, N., Ikeda, R., and Ito, M. (2010). Expression, purification, and characterization of a recombinant neutral ceramidase from *Mycobacterium tuberculosis*. *Biosci. Biotechnol. Biochem.* 74, 316–321. doi: 10.1271/bbb.90645
- Olczak, T., Sroka, A., Potempa, J., and Olczak, M. (2008). *Porphyromonas gingivalis* HmuY and HmuR: further characterization of a novel mechanism of heme utilization. *Arch. Microbiol.* 189, 197–210. doi: 10.1007/s00203-007-0309-7
- Pazos, F. (1997). *Flexibacter Maritimus: Estudio Fenotípico, Inmunológico y Molecular*. Ph.D. thesis, University of Santiago de Compostela, Santiago.
- Pers, C., Gahrn-Hansen, B., and Frederiksen, W. (1996). *Capnocytophaga canimorsus* septicemia in Denmark, 1982–1995: review of 39 cases. *Clin. Infect. Dis.* 23, 71–75. doi: 10.1093/clinids/23.1.71
- Rahman, T., Suga, K., Kinya, K., and Yukitaka, S. (2014). Biological and serological characterization of a non-gliding strain of *Tenacibaculum maritimum* isolated from a diseased puffer fish *Takifugu rubripes*. *Fish Pathol.* 49, 121–129. doi: 10.3147/jfsfp.49.121
- Ratledge, C., and Dover, L. G. (2000). Iron metabolism in pathogenic bacteria. *Annu. Rev. Microbiol.* 54, 881–941. doi: 10.1146/annurev.micro.54.1.881
- Renzi, F., Manfredi, P., Mally, M., Moes, S., Jenö, P., and Cornelis, G. R. (2011). The N-glycan glycoprotein deglycosylation complex (Gpd) from *Capnocytophaga canimorsus* deglycosylates human IgG. *PLoS Pathog.* 7:e1002118. doi: 10.1371/journal.ppat.1002118
- Romero, M., Avendaño-Herrera, R., Magariños, B., Cámara, M., and Otero, A. (2010). Acylhomoserine lactone production and degradation by the fish pathogen *Tenacibaculum maritimum*, a member of the *Cytophaga-Flavobacterium-Bacteroides* (CFB) group. *FEMS Microbiol. Lett.* 304, 131–139. doi: 10.1111/j.1574-6968.2009.01889.x
- Sasagawa, Y., Izaki, K., Matsubara, Y., Suzuki, K., Kojima, H., and Kamio, Y. (1995). Molecular cloning and sequence analysis of the gene encoding the collagenase from *Cytophaga* sp. L43-1 strain. *Biosci. Biotechnol. Biochem.* 59, 2068–2073. doi: 10.1271/bbb.59.2068
- Sato, K., Naito, M., Yukitake, H., Hirakawa, H., Shoji, M., McBride, M. J., et al. (2010). A protein secretion system linked to bacteroidete gliding motility and pathogenesis. *Proc. Natl. Acad. Sci. U.S.A.* 107, 276–281. doi: 10.1073/pnas.0912010107
- Sato, K., Yukitake, H., Narita, Y., Shoji, M., Naito, M., and Nakayama, K. (2013). Identification of *Porphyromonas gingivalis* proteins secreted by the Por

- secretion system. *FEMS Microbiol. Lett.* 338, 68–76. doi: 10.1111/1574-6968.12028
- Sheng, Y., Abreu, I. A., Cabelli, D. E., Maroney, M. J., Miller, A. F., Teixeira, M., et al. (2014). Superoxide dismutases and superoxide reductases. *Chem. Rev.* 114, 3854–3918. doi: 10.1021/cr4005296
- Singh, R., Wiseman, B., Deemagarn, T., Jha, V., Switala, J., and Loewen, P. C. (2008). Comparative study of catalase-peroxidases (KatGs). *Arch. Biochem. Biophys.* 471, 207–214. doi: 10.1016/j.abb.2007.12.008
- Småge, S. B., Frisch, K., Brevik, ØJ., Kuninori, W., and Nylunda, A. (2016). First isolation, identification and characterisation of *Tenacibaculum maritimum* in Norway, isolated from diseased farmed sea lice cleaner fish *Cyclopterus lumpus* L. *Aquaculture* 464, 178–184. doi: 10.1016/j.aquaculture.2016.06.030
- Studier, F. W. (2005). Protein production by auto-induction in high density shaking cultures. *Protein Expr. Purif.* 41, 207–234. doi: 10.1016/j.pep.2005.01.016
- Suomalainen, L. R., Tirola, M., and Valtonen, E. T. (2006). Chondroitin AC lyase activity is related to virulence of fish pathogenic *Flavobacterium columnare*. *J. Fish Dis.* 29, 757–763. doi: 10.1111/j.1365-2761.2006.00771.x
- Suzuki, M., Nakagawa, Y., Harayama, S., and Yamamoto, S. (2001). Phylogenetic analysis and taxonomic study of marine *Cytophaga*-like bacteria: proposal for *Tenacibaculum* gen. nov. with *Tenacibaculum maritimum* comb. nov. and *Tenacibaculum ovolyticum* comb. nov., and description of *Tenacibaculum mesophilum* sp. nov. and *Tenacibaculum amylolyticum* sp. nov. *Int. J. Syst. Evol. Microbiol.* 51, 1639–1652. doi: 10.1099/00207713-51-5-1639
- Teramoto, M., Zhai, Z., Komatsu, A., Shibayama, K., and Suzuki, M. (2016). Genome sequence of the psychrophilic bacterium *Tenacibaculum ovolyticum* strain da5A-8 isolated from deep seawater. *Genome Announc.* 4:e00644–16. doi: 10.1128/genomeA.00644-16
- Terrapon, N., Lombard, V., Gilbert, H. J., and Henrissat, B. (2015). Automatic prediction of polysaccharide utilization loci in *Bacteroidetes* species. *Bioinformatics* 31, 647–655. doi: 10.1093/bioinformatics/btu716
- Vallenet, D., Belda, E., Calteau, A., Cruveiller, S., Engelen, S., Lajus, A., et al. (2013). MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res.* 41, D636–D647. doi: 10.1093/nar/gks1194
- van Gelderen, R., Carson, J., and Nowak, B. (2009). Effect of extracellular products of *Tenacibaculum maritimum* in Atlantic salmon, *Salmo salar* L. *J. Fish Dis.* 32, 727–731. doi: 10.1111/j.1365-2761.2009.01032.x
- Varki, N. M., and Varki, A. (2007). Diversity in cell surface sialic acid presentations: implications for biology and disease. *Lab. Invest.* 87, 851–857. doi: 10.1038/labinvest.3700656
- Vernikos, G. S., and Parkhill, J. (2006). Interpolated variable order motifs for identification of horizontally acquired DNA: revisiting the *Salmonella* pathogenicity islands. *Bioinformatics* 22, 2196–2203. doi: 10.1093/bioinformatics/btl369
- Waack, S., Keller, O., Asper, R., Brodag, T., Damm, C., Fricke, W. F., et al. (2006). Score-based prediction of genomic islands in prokaryotic genomes using hidden Markov models. *BMC Bioinformatics* 7:142. doi: 10.1186/1471-2105-7-142
- Wakabayashi, H., Hikida, M., and Masumura, K. (1986). *Flexibacter maritimum* sp. nov., a pathogen of marine fishes. *Int. J. Syst. Evol. Microbiol.* 36, 396–398. doi: 10.1099/00207713-36-3-396
- Wójtowicz, H., Wojaczyński, J., Olczak, M., Króliczewski, J., Latos-Grażyński, L., and Olczak, T. (2009). Heme environment in HmuY, the heme-binding protein of *Porphyromonas gingivalis*. *Biochem. Biophys. Res. Commun.* 383, 178–182. doi: 10.1016/j.bbrc.2009.03.148
- Xu, Q., Rawlings, N. D., Farr, C. L., Chiu, H. J., Grant, J. C., Jaroszewski, L., et al. (2011). Structural and sequence analysis of imelysin-like proteins implicated in bacterial iron uptake. *PLoS ONE* 6:e21875. doi: 10.1371/journal.pone.0021875
- Yin, Y., Mao, X., Yang, J., Chen, X., Mao, F., and Xu, Y. (2012). dbCAN: a web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res.* 40, W445–W451. doi: 10.1093/nar/gks479

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Pérez-Pascual, Lunazzi, Magdelenat, Rouy, Roulet, Lopez-Roques, Larocque, Barbeyron, Gobet, Michel, Bernardet and Duchaud. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.