



# Sperm-Mediated Transgenerational Inheritance

Corrado Spadafora\*

*Institute of Translational Pharmacology, National Research Council of Italy, Rome, Italy*

Spermatozoa of virtually all species can spontaneously take up exogenous DNA or RNA molecules and internalize them into nuclei. In this article I review evidence for a key role of a reverse transcriptase (RT) activity, encoded by LINE-1 retrotransposons, in the fate of the internalized nucleic acid molecules and their implication in transgenerational inheritance. LINE-1-derived RT, present in sperm heads, can reverse-transcribe the internalized molecules in cDNA copies: exogenous RNA is reverse-transcribed in a one-step reaction, whereas DNA is first transcribed into RNA and subsequently reverse-transcribed. Both RNA and cDNA molecules can be delivered from sperm cells to oocytes at fertilization, further propagated throughout embryogenesis and inherited in a non-Mendelian fashion in tissues of adult animals. The reverse-transcribed sequences are extrachromosomal, low-abundance, and mosaic distributed in tissues of adult individuals, where they are variably expressed. These “retrogenes” are transcriptionally competent and induce novel phenotypic traits in animals. Growing evidence indicate that cancer tissues produce DNA- and RNA-containing exosomes. We recently found that these exosomes are released in the bloodstream and eventually taken up into epididymal spermatozoa, consistent with the emerging view that a transgenerational flow of extrachromosomal RNA connects soma to germline and, further, to next generation embryos. Spermatozoa play a crucial bridging role in this process: they act as collectors of somatic information and as delivering vectors to the next generation. On the whole, this phenomenon is compatible with a Lamarckian-type view and closely resembles Darwinian pangenesis.

## OPEN ACCESS

### Edited by:

Guenther Witzany,  
Telos-Philosophische Praxis, Austria

### Reviewed by:

Masato Tsurudome,  
Mie University, Japan  
Carlos M. Guerrero-Bosagna,  
Linköping University, Sweden

### \*Correspondence:

Corrado Spadafora  
corrado.spadafora@gmail.com

### Specialty section:

This article was submitted to  
Virology,  
a section of the journal  
Frontiers in Microbiology

**Received:** 15 August 2017

**Accepted:** 20 November 2017

**Published:** 04 December 2017

### Citation:

Spadafora C (2017) Sperm-Mediated  
Transgenerational Inheritance.  
*Front. Microbiol.* 8:2401.  
doi: 10.3389/fmicb.2017.02401

**Keywords:** spermatozoa, LINE-1 retrotransposons, reverse transcriptase, exosomes, transgenerational inheritance, evolution

## SPERMATOZOA AS A SOURCE OF REVERSE TRANSCRIPTASE-MEDIATED EXTRACHROMOSOMAL INFORMATION: A LOOK TO THE PAST

It is a well-established notion that mature spermatozoa have the spontaneous ability to take up exogenous DNA molecules and to internalize them in their nuclei (reviewed by Spadafora, 1998). This permeability is a distinctive feature of spermatozoa, both epididymal and ejaculated (after wash-off of seminal fluid), from virtually all animal species including humans

(Smith and Spadafora, 2005). Thus, in spite of the highly compact and impenetrable structure of their nuclei, sperm cells are in fact highly permeable to foreign molecule intrusion. Intense investigations of this phenomenon revealed that the interaction of exogenous DNA molecules with sperm cells, as well as their subsequent nuclear internalization, are well-regulated processes mediated by a network of specific factors (Spadafora, 1998). Parallel studies have revealed that spermatozoa can also take up RNA molecules and internalize them in their nuclei. Somewhat unexpectedly, these RNAs are reverse-transcribed into cDNA copies by a biologically active reverse transcriptase (RT) activity encoded by LINE-1 retrotransposons and present in sperm nuclei (Giordano et al., 2000; Spadafora, 2008). The LINE-1-derived RT interplays with a DNA-dependent RNA polymerase, also present in spermatozoa (Fuster et al., 1977), which together amplify the cDNAs copy number, mimicking a “natural” PCR/RT-PCR process. Most newly generated cDNA copies are released from spermatozoa into the medium and can be taken up again by further spermatozoa and internalized in their nuclei. Through this continuously cycling process, cDNA copies are evenly distributed among the vast majority of sperm cells suspension. Work with murine models showed that the RT-generated cDNAs are delivered to oocytes at fertilization (Giordano et al., 2000; Pittoggi et al., 2006), maintained as low-copy number (below one copy/genome) non-integrated extrachromosomal sequences throughout development, mosaic propagated in the tissues of founder individuals, eventually transmitted in a non-Mendelian fashion to the next generation, transcriptionally competent and able to generate phenotypic variations in animals of both generations (Sciamanna et al., 2003; Pittoggi et al., 2006). These results suggest that spermatozoa provide a previously unrecognized source of RT-mediated information, not linked to chromosomal genes, and, at the same time, act as propagating vectors throughout generations.

These findings raise several puzzling questions. First, does the ability of sperm cells to take up foreign nucleic acid molecules reflects an enforced behavior when they come in contact with RNA under conditions of *in vitro* assays, or else do spermatozoa naturally collect and carry foreign molecules under physiological conditions *in vivo*? Second, does the RT activity stored in spermatozoa represent a functionless remnant of ancestral retrotransposon activity, brought to new life in response to occasional intrusions of foreign molecules, or does it exert an extant physiological role in development? These two issues, i.e., the sperm permeability to exogenous RNA, and the sperm RT that uses the latter as a substrate for retrotranscription, raise the third key question of whether these phenomena are physiologically relevant or, in other words, whether they occur in nature to generate a source of novel information. To begin to address these issues, it was imperative to characterize the RNA population stored in spermatozoa and possibly identify its origin. In recent years, high-throughput technologies and next generation sequence analysis have revealed a highly complex composition of spermatozoal RNA, whose components are increasingly emerging as key players in epigenetic inheritance

processes, as will be seen in more depth in the following paragraphs.

## THE COMPLEX TRANSCRIPTIONAL LANDSCAPE OF MATURE SPERMATOZOA

Traditional views considered spermatozoa as transcriptionally silent cells (Grunewald et al., 2005) and sperm RNAs as negligible remnants produced during spermatogenesis. More recent data however contrast with these views, showing that mature spermatozoa in fact contain a complex population of coding RNAs, small non-coding RNA classes, and, finally, LINE-1, SINE/Alu, and LTR repeat-associated transcripts (Jodar et al., 2013; Sendler et al., 2013; Miller, 2014). Small non-coding RNAs account for a considerable proportion of spermatozoal RNA (Krawetz et al., 2011; Kawano et al., 2012), mainly represented by piRNAs produced during spermatogenesis, tsRNA (tRNA-derived), and to a lesser extent, microRNAs (miRNAs) are instead predominant in epididymal spermatozoa (Chen et al., 2016b). Importantly, the composition of the spermatozoal RNA population varies in response to paternal exposure to a variety of stressing conditions (Rodgers et al., 2013; Brieno-Enriquez et al., 2015), a circumstance that can have crucial consequences for the fate and health of the progeny. Most importantly, growing data are revealing that RNAs of somatic origin also contribute to the composition of the sperm RNA cargo in the form of selectively retained RNAs derived from soma-to-spermatozoa intercellular communication. This flow is mediated by a special class of epididymis-derived nanovesicles, called epididymosomes, which shuttle miRNAs and tRNA fragments from the epididymal tissue to mature sperm cells (Belleannée et al., 2013; Vojtech et al., 2014; Sharma et al., 2016). The shuttled sperm RNA, containing several 100s of developmentally relevant small RNAs, are the product of a “sieving” process, as their profiles are distinct from those of the surrounding soma (Reilly et al., 2016). The modulation of the sperm RNA content occurs during maturation of spermatozoa between the proximal and distal epididymal segments, and identifies the epididymis as a key site for the establishment of the sperm epigenome (Nixon et al., 2015).

We recently reported that epididymal spermatozoa can incorporate RNA from somatic cell-released exosomes: indeed, we found that human melanoma cells, engineered to express EGFP and inoculated in nude mice, release EGFP RNA-containing nanovesicles in the bloodstream of the animals; a proportion of that RNA reaches the epididymis and becomes internalized in sperm heads (Cossetti et al., 2014). This finding shows that the flow of RNA delivered to spermatozoa originates not only from the surrounding epididymal soma, but also from distant, unrelated districts of the body. Nanovesicles act as the ideal vectors of such delivery. Sperm heads are the final recipients of this extrachromosomal information due to their ability to spontaneously take up exogenous molecules, as mentioned above. On the whole, these data indicate that the

impenetrable Weismann barrier, considered for a long time as a cornerstone of modern genetics, can in fact be breached by nanovesicle-mediated flows of extrachromosomal RNA (Eaton et al., 2015).

## **BREAKING THE WEISMANN BARRIER: A SPERM-MEDIATED RNA-BASED FLOW CONNECTS SOMA TO THE NEXT GENERATION EMBRYOS**

In a seminal article, Krawetz and collaborators (Ostermeier et al., 2004) first reported that the sperm-specific RNA cargo is delivered to oocytes at fertilization. That finding proved that not only the male genome, but also extrachromosomal RNA carried by sperms, contribute to the zygote formation. However, the sperm RNA *per se* is not strictly required for embryonic development, as parthenogenetic mice can be successfully generated by microinjecting haploid, or bimaternal embryonic stem cells in murine oocytes (Li et al., 2016; Zhong et al., 2016). The latter finding indicates that all the fundamental information to support the developmental program, from fertilization to adulthood, is linked to chromosomal genes.

A novel turn to the field is being provided by recent data indicating that the composition of sperm RNA reflects the lifestyle habits and carry the “memory” of paternal experiences; that RNA-based memory is transmissible to the offspring as paternally acquired characteristics, with the potential to affect the health and overall biological fate of the progeny (reviewed by Liebers et al., 2014; Klosin and Lehner, 2016). Of remarkable interest are recent experiments that have assessed the potential of sperm RNAs as transgenerational modifiers in response to parental environmental or stressing conditions (Carone et al., 2010; Rodgers et al., 2013, 2015), including diet (Fullston et al., 2013; Chen et al., 2016a; Huypens et al., 2016), cigarette smoke (Marczylo et al., 2012), odor sensitivity (Dias and Ressler, 2014), and cognitive and behavioral conditioning (Rodgers et al., 2013; Gapp et al., 2014). RNA was unambiguously identified as the transgenerational modifier in a large set of compelling experimental data, showing that offspring generated from normal zygotes microinjected with sperm RNA recapitulate the phenotypical traits of the RNA donor animals (Rassoulzadegan et al., 2006; Gapp et al., 2014; Grandjean et al., 2015; Chen et al., 2016a).

Together, these data show that inheritance is not exclusively linked to chromosomal genes. Indeed, a subtle yet effective flow of RNA is established between somatic tissues and the next generation embryos. Spermatozoa are the pivots, playing a dual role both as collectors of paternal extrachromosomal RNA and as their vectors to the offspring. The emerging evidence that RNA-based information can travel from soma to germline subvert the Weismann’s theory and provide a foundation for the inheritance of acquired traits with far reaching implications for evolutionary processes.

## **RT ENCODED BY LINE-1 RETROTRANSPOSONS AS MODULATOR OF EARLY EMBRYONIC DEVELOPMENT**


In addition to being stored in mature spermatozoa, LINE-1-encoded RT is also abundantly expressed in early embryos and is implicated in the genesis and propagation of extrachromosomal information. We have found that LINE-1 retrotransposon-encoded RT is triggered soon after fertilization in both zygotic pronuclei, predominantly in the paternal pronucleus (Vitullo et al., 2012), and remains active in early preimplantation embryos (Pittoggi et al., 2003). RT plays a crucial role in early development: indeed, RT inhibition, induced either by pharmacological RT inhibitors (Pittoggi et al., 2003), or by downregulating LINE-1 expression by microinjecting antisense oligonucleotides in zygotic pronuclei (Beraldi et al., 2006), causes a drastic arrest of embryo development at the 2- or 4-cell stages. These results suggest that RT is strictly necessary for the unfolding of the developmental program from the second cell division, as the first cleavage exploits the maternal RNA stored in oocytes (Tang et al., 2007).

Although neither the specific role(s) nor the mechanism of action of embryonic RT are yet fully clarified, emerging data suggest that RT controls the biogenesis of miRNAs, a class of RNA that is globally, yet transiently, suppressed in early embryogenesis (Suh et al., 2010), concomitant with the up-regulation of RT expression (Pittoggi et al., 2003; Vitullo et al., 2012).

The link between LINE-1-encoded RT and miRNA biogenesis has been investigated in some depth in cancer cells. In striking analogy with early embryos, RT is also highly expressed in most cancer types from very early stages (reviewed by Sinibaldi-Vallebona et al., 2011; De Luca et al., 2016). In parallel with high RT activity, the biogenesis of LINE-1-derived miRNAs (Lu et al., 2005) and siRNAs (Chen et al., 2012) is globally reduced in cancer compared to normal cells, with ensuing alterations of the gene expression regulatory network. Exposure of cancer cells to RT inhibitors restores the normal expression profile of miRNAs, with a direct impact on global gene expression (Sciamanna et al., 2013, 2014). These lines of evidence suggest therefore that high RT expression exerts: (i) a physiological control on the biogenesis of miRNA in early embryogenesis, and (ii) a pathological role in cells conveyed toward tumorigenesis, by impairing the production of miRNAs, with the ensuing dysregulation of downstream targets and the increase of transcriptional fluctuations.

## **THE REMODELING OF THE EMBRYONIC EPIGENETIC LANDSCAPE AND ITS IMPLICATION IN EVOLUTION. A MODEL**

In recapitulating the aspects discussed so far a framework is beginning to emerge: (1) RNA-containing nanovesicles are released from somatic tissues into the bloodstream; (2)



Epididymal spermatozoa take up nanovesicles and internalize them in their nuclei; (3) The internalized RNA molecules are processed and their copy number is amplified, via the RT/DNA-dependent RNA polymerase interplay; (4) Somatic RNAs, or

their cDNA copies, are delivered from sperm to embryos at fertilization.

The first three steps continuously renew the RNA storage in sperm heads. The last one, i.e., the delivery of processed somatic RNA to oocytes, can recur at each round of fertilization. Through this process sperm RNA is transmitted from one generation to the next, which can contribute to the embryo fitness and in principle expand the adaptation of the newborn to diverse environmental conditions. It is reasonable to assume that a large proportion of the males living in a same ecological niche and exposed to the same stimuli, produce sperm RNA cargos of similar composition; under constant environmental conditions, these RNA cargo would be continuously delivered to the progeny via fertilization throughout generations. It is not unreasonable to hypothesize that, in the long run (i.e., after a "sufficient" number of generations), the sperm RNAs promote the "assimilation" of new trait(s), to use Waddington's concept (Waddington, 1959). In other words, the cumulative effects of RNA delivery through generations may promote the emergence of novel functional "canalized" pathways (Waddington, 1959), via remodeling of the embryonic chromatin architecture; in consequence, novel genetic circuits might be activated and/or pre-existing ones might be "rewired." Mechanistically, the cumulative effects of regulatory miRNAs and tsRNAs delivered by sperm upon fertilization would drive the emergence of novel canalized pathways through two sequential steps: (i) first, by "rewiring" the expression profile of genes constituting canalized genetic circuits, and (ii) second, via targeted retrotransposition events that provide new regulatory sequences, which brings to completion the newly canalized circuits.

The first (epigenetic) step builds on the established regulatory functions of miRNAs and tsRNAs, which can modulate the expression of relevant genes. It is reasonable to hypothesize that RNAs delivered by sperm cells at fertilization also exert these regulatory functions and remodelate the gene expression profile in early embryos. Consequently, new genetic circuits (canalized circuits) become functionally active, or/and pre-existing ones are rewired.

Canalized circuits would then reach their final state through targeted retrotransposition events (genetic step). New retrotranspositions can provide additional layers of control, by inserting protein-binding sites (e.g., for transcription factors, hormones, splicing factors), as well as enhancers, promoters, insulators, etc. in new sites within the genome.

Thus, a hybrid epigenetic/genetic process drives the remodeling of the embryonic chromatin architecture, as schematized in **Figure 1**.

The process is seen as progressive in nature, based on the assumption that the "quanta" of sperm-derived regulatory RNAs delivered to the embryo at fertilization (represented by the red ball rolling down the groove in **Figure 1**) constitute minimal contributions toward the epigenetic activation of novel canalization(s) by "deepening the canal" (**Figures 1a-c**, right branch), while non-active pathways become shallower (**Figures 1a-d**, left branch). When the cumulative effects of the

sperm-delivered RNA overcome the buffering capacity of the embryos, the novel canalized genomic circuit (**Figure 1d**) has the potential to redirect the embryonic ontogenesis and generate phenotypic novelties.

Targeted retrotransposition events constitute the genetic component of the model, contributing to the functional reshaping of the embryo regulatory circuits. Targeted insertions (symbolized by different colored symbols in **Figure 1**) contribute to establish novel regulatory circuits in at least three ways: (i) they provide new protein-binding and regulatory sites; (ii) they contribute new miRNA-coding sequences that expand the overall diversity in the RNA population, and (iii) they stabilize the newly remodeled landscape by fixing the chromatin architecture.

Three considerations suggest that these changes could be permanently assimilated. First, zygotes and early embryos are thought to provide permissive, change-prone environments, consistent with the finding that the early embryonic genome is largely unstructured before zygotic genome activation, showing a low level of chromatin organization over long genomic distances (Hug et al., 2017). Second, as mentioned, LINE-1-encoded RT activity is high in preimplantation embryos (Vitullo et al., 2012) and, in parallel, the miRNA-based control system is globally suppressed (Suh et al., 2010). This is relevant in the light of evidence that miRNA-mediated control reduces random fluctuations in differentiated cells and in development, hence conferring robustness to genetic pathways (Li et al., 2009; Ebert and Sharp, 2012); on the contrary, miRNA suppression increases instability and random fluctuations in the developmental program (Hornstein and Shomron, 2006; Li et al., 2009; Ebert and Sharp, 2012). Moreover, retrotransposon families (i.e., LINE-1, Alus, LTRs) are de-repressed in embryos concomitant with global genomic hypomethylation (Lee et al., 2014; Smith et al., 2014), and constitute a potential source of both genetic and epigenetic variations (Macia et al., 2011; Vitullo et al., 2012; Fadloun et al., 2013). Overall, no massive retrotransposition events are required, with the exception of some crucial insertions that provide regulatory sequences to the newly formed canalized circuits. These crucial events would be targeted to specific hypersensitive sites generated during embryonic chromatin remodeling. Third, the RNA population that spermatozoa deliver to oocytes contains regulatory miRNAs and tsRNAs (Chen et al., 2016b), which can reshape the embryonic expression landscape and reprogram the transcription profiles of 100s of embryonic genes. Indeed, even small amounts of delivered regulatory RNAs can generate an ample spectrum of epigenetic variations with a potential impact on the phenotype. Thus, inheritable variations may be driven by small regulatory RNAs, assimilated in the change-prone genome architecture of embryos and translated into new phenotypic variants, with no major adverse effect on the permissive embryonic context.

It is worth recalling that small RNAs are involved both in macroevolutionary processes – as their number increases over time in parallel with complexity, while their loss is associated

with morphological simplification (Wheeler et al., 2009; Erwin et al., 2011) – and with the canalization of genetic programs (Hornstein and Shomron, 2006; Li et al., 2009; Vidigal and Ventura, 2015).

## CONCLUSION

The present model of transgenerational inheritance attempts to integrate data from different sources in a biologically coherent framework. Most aspects implicated in the process are experimentally tested and are potentially able to generate transgenerationally relevant novelties. The mechanism is predominantly epigenetic and independent of genomic mutations. Importantly, the “sieving force” of natural selection is not necessary in conventional terms, because the canalization process driven by sperm RNA would generate specific pathways, leaving little space for random variations, and favoring the ultimate emergence of one or few new phenotype(s). In analogy with Lamarckism, this hypothesis is based on the assumption that extrachromosomal transgenerational inheritance can affect ontogenesis and generate evolutionarily significant, stably acquired variations. Darwinian pangenesis (Holterhoff, 2014) is the other theory with which the model has significant overlap. The hypothesis that “gemmules” containing parental characters are released from tissues and transferred to the next generation via the germline can now be reinterpreted in the light of our current knowledge of circulating nanovesicles and exosomes, carrying nucleic acids and released from somatic tissues, which can be taken up by sperm cells, thus providing a foundation for spermatozoa-mediated transgenerational inheritance. It is amazing that so-called obsolete concepts, developed in the context of two historically rejected theories, are re-emerging from modern experimental data based on next generation genomic methodologies, thus confirming that history sometime repeats itself.

## AUTHOR CONTRIBUTIONS

CS has conceived the model and wrote the manuscript.

## FUNDING

This work was supported by a grant from Fondazione Roma to the Project “Investigating the cellular endogenous Reverse Transcriptase as a novel therapeutic target and an early tumor marker” to CS.

## ACKNOWLEDGMENT

The author grateful to Michele Spadafora and Graziano Bonelli for skilful assistance with drawing preparation.

## REFERENCES

- Belleannée, C., Calvo, E., Caballero, J., and Sullivan, R. (2013). Epididymosomes convey different repertoires of microRNAs throughout the bovine epididymis. *Biol. Reprod.* 89:30. doi: 10.1095/biolreprod.113.110486
- Beraldi, R., Pittoggi, C., Sciamanna, I., Mattei, E., and Spadafora, C. (2006). Expression of LINE-1 retroposons is essential for murine preimplantation development. *Mol. Reprod. Dev.* 3, 279–287. doi: 10.1002/mrd.20423
- Brieno-Enriquez, M. A., Garcia-Lopez, J., Cardenas, D. B., Guibert, S., Cleroux, E., Ded, L., et al. (2015). Exposure to endocrine disruptor induces transgenerational epigenetic deregulation of microRNAs in primordial germ cells. *PLOS ONE* 10:e0124296. doi: 10.1371/journal.pone.0124296
- Carone, B. R., Fauquier, L., Habib, N., Shea, J. M., Hart, C. E., Li, R., et al. (2010). Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals. *Cell* 143, 1084–1096. doi: 10.1016/j.cell.2010.12.008
- Chen, L., Dahlstrom, J. E., Lee, S. H., and Rangasamy, D. (2012). Naturally occurring endo-siRNA silences LINE-1 retrotransposons in human cells through DNA methylation. *Epigenetics* 7, 758–771. doi: 10.4161/epi.20706
- Chen, Q., Yan, M., Cao, Z., Li, X., Zhang, Y., Shi, J., et al. (2016a). Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* 351, 397–400. doi: 10.1126/science.aad7977
- Chen, Q., Yan, W., and Duan, E. (2016b). Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. *Nat. Rev. Genet.* 17, 733–743. doi: 10.1038/nrg.2016.106
- Cossetti, C., Lugini, L., Astrologo, L., Saggio, I., Fais, S., and Spadafora, C. (2014). Soma-to-germline transmission of RNA in mice xenografted with human tumour cells: possible transport by exosomes. *PLOS ONE* 9:e101629. doi: 10.1371/journal.pone.0101629
- De Luca, C., Guadagni, F., Sinibaldi-Vallebona, P., Sentinelli, S., Gallucci, M., Hoffmann, A., et al. (2016). Enhanced expression of LINE-1-encoded ORF2 protein in early stages of colon and prostate transformation. *Oncotarget* 7, 4048–4061. doi: 10.18632/oncotarget.6767
- Dias, B. G., and Ressler, K. J. (2014). Parental olfactory experience influences behavior and neural structure in subsequent generations. *Nat. Neurosci.* 17, 89–96. doi: 10.1038/nn.3594
- Eaton, S. A., Jayasooriah, N., Buckland, M. E., Martin, D. I. K., Cropley, J. E., and Suter, C. M. (2015). Roll over weismann: extracellular vesicles in the transgenerational transmission of environmental effects. *Epigenomics* 7, 1165–1171. doi: 10.2217/epi.15.58
- Ebert, M. S., and Sharp, P. A. (2012). Roles for MicroRNAs in conferring robustness to biological processes. *Cell* 149, 515–524. doi: 10.1016/j.cell.2012.04.005
- Erwin, D. H., Laflamme, M., Tweed, S. M., Sperling, E. A., Pisani, D., and Peterson, K. J. (2011). The cambrian conundrum: early divergence and later ecological success in the early history of animals. *Science* 334, 1091–1097. doi: 10.1126/science.1206375
- Fadloun, A., Le Gras, S., Jost, B., Ziegler-Birling, C., Takahashi, H., Gorab, E., et al. (2013). Chromatin signatures and retrotransposon profiling in mouse embryos reveal regulation of LINE-1 by RNA. *Nat. Struct. Mol. Biol.* 20, 332–338. doi: 10.1038/nsmb.2495
- Fullston, T., Ohlsson Teague, E. M., Palmer, N. O., DeBlasio, M. J., Mitchell, M., Corbett, M., et al. (2013). Paternal obesity initiates metabolic disturbances in two generations of mice with incomplete penetrance to the F2 generation and alters the transcriptional profile of testis and sperm microRNA content. *FASEB J.* 27, 4226–4243. doi: 10.1096/fj.12-224048
- Fuster, C. D., Farrell, D., Stern, F. A., and Hecht, N. B. (1977). RNA polymerase activity in bovine spermatozoa. *J. Cell Biol.* 74, 698–706. doi: 10.1083/jcb.74.3.698
- Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., et al. (2014). Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nat. Neurosci.* 17, 667–669. doi: 10.1038/nn.3695
- Giordano, R., Magnano, A. R., Zaccagnini, G., Pittoggi, C., Moscufo, N., Lorenzini, R., et al. (2000). Reverse transcriptase activity in mature spermatozoa of mouse. *J. Cell Biol.* 148, 1107–1113. doi: 10.1083/jcb.148.6.1107
- Grandjean, V., Fourré, S., De Abreu, D. A. F., Derieppe, M.-A., Remy, J.-J., and Rassoulzadegan, M. (2015). RNA-mediated paternal heredity of diet-induced obesity and metabolic disorders. *Sci. Rep.* 5:18193. doi: 10.1038/srep18193
- Grunewald, S., Paasch, U., Glander, H. J., and Andereg, U. (2005). Mature human spermatozoa do not transcribe novel RNA. *Andrologia* 37, 69–71. doi: 10.1111/j.1439-0272.2005.00656.x
- Holterhoff, K. (2014). The history and reception of charles darwin's hypothesis of pangenesis. *J. Hist. Biol.* 47, 661–695. doi: 10.1007/s10739-014-9377-9377-0
- Hornstein, E., and Shomron, N. (2006). Canalization of development by microRNAs. *Nat. Genet.* 38(Suppl.), S20–S24. doi: 10.1038/ng1803
- Hug, C. B., Grimaldi, A. G., Kruse, K., and Vaquerizas, J. M. (2017). Chromatin architecture emerges during zygotic genome activation independent of transcription. *Cell* 169, 216–228. doi: 10.1016/j.cell.2017.03.024
- Huypens, P., Sass, S., Wu, M., Dyckhoff, D., Tschopp, M., Theis, F., et al. (2016). Epigenetic germline inheritance of diet-induced obesity and insulin resistance. *Nat. Genet.* 48, 497–499. doi: 10.1038/ng.3527
- Jodar, M., Selvaraju, S., Sendler, E., Diamond, M. P., Krawetz, S. A., and Reproductive Medicine Network (2013). The presence, role and clinical use of spermatozoal RNAs. *Hum. Reprod. Update* 19, 604–624. doi: 10.1093/humupd/dmt031
- Kawano, M., Kawaji, H., Grandjean, V., Kiani, J., and Rassoulzadegan, M. (2012). Novel small noncoding RNAs in mouse spermatozoa, zygotes and early embryos. *PLOS ONE* 7:e44542. doi: 10.1371/journal.pone.0044542
- Klosin, A., and Lehner, B. (2016). Mechanisms, timescales and principles of transgenerational epigenetic inheritance in animals. *Curr. Opin. Genet. Dev.* 36, 41–49. doi: 10.1016/j.gde.2016.04.001
- Krawetz, S. A., Kruger, A., Lalancette, C., Tagett, R., Anton, E., Draghici, S., et al. (2011). A survey of small RNAs in human sperm. *Hum. Reprod.* 26, 3401–3412. doi: 10.1093/humrep/der329
- Lee, H. J., Hore, T. A., and Reik, W. (2014). Reprogramming the methylome: erasing memory and creating diversity. *Cell Stem Cell* 14, 710–719. doi: 10.1016/j.stem.2014.05
- Li, X., Cassidy, J. J., Reinke, C. A., Fischboeck, S., and Carthew, R. W. (2009). A microRNA imparts robustness against environmental fluctuation during development. *Cell* 137, 273–282. doi: 10.1016/j.cell.2009.01.058
- Li, Z., Wan, H., Feng, G., Wang, L., He, Z., Wang, Y., et al. (2016). Birth of fertile bimaterial offspring following intracytoplasmic injection of parthenogenetic haploid embryonic stem cells. *Cell Res.* 26, 135–138. doi: 10.1038/cr.2015.151
- Liebers, R., Rassoulzadegan, M., and Lyko, F. (2014). Epigenetic regulation by heritable RNA. *PLOS Genet.* 10:e1004296. doi: 10.1371/journal.pgen.1004296
- Lu, J., Getz, G., Miska, E. A., Alvarez-Saavedra, E., Lamb, J., Peck, D., et al. (2005). MicroRNA expression profiles classify human cancers. *Nature* 435, 834–838. doi: 10.1038/nature03702
- Macía, A., Muñoz-Lopez, M., Cortes, J. L., Hastings, R. K., Morell, S., Lucena-Aguilar, G., et al. (2011). Epigenetic control of retrotransposon expression in human embryonic stem cells. *Mol. Cell. Biol.* 31, 300–316. doi: 10.1128/MCB.00561-10
- Marczylo, E. L., Amoako, A. A., Konje, J. C., Gant, T. W., and Marczylo, T. H. (2012). Smoking induces differential miRNA expression in human spermatozoa: a potential transgenerational epigenetic concern? *Epigenetics* 7, 432–439. doi: 10.4161/epi.19794
- Miller, D. (2014). Sperm RNA as a mediator of genomic plasticity. *Adv. Biol.* 2014:179701. doi: 10.1155/2014/179701
- Nixon, B., Stanger, S. J., Mihalas, B. P., Reilly, J. N., Anderson, A. L., Tyagi, S., et al. (2015). The microRNA signature of mouse spermatozoa is substantially modified during epididymal maturation. *Biol. Reprod.* 93, 91. doi: 10.1095/biolreprod.115.132209
- Ostermeier, G. C., Miller, D., Huntriss, J. D., Diamond, M. P., and Krawetz, S. A. (2004). Reproductive biology: delivering spermatozoan RNA to the oocyte. *Nature* 429:154. doi: 10.1038/429154a
- Pittoggi, C., Beraldi, R., Sciamanna, I., Barberi, L., Giordano, R., Magnano, A. R., et al. (2006). Generation of biologically active retro-genes upon interaction of mouse spermatozoa with exogenous DNA. *Mol. Reprod. Dev.* 73, 1239–1246. doi: 10.1002/mrd.20550
- Pittoggi, C., Sciamanna, I., Mattei, E., Beraldi, R., Lobascio, A. M., Mai, A., et al. (2003). Role of endogenous reverse transcriptase in murine early embryo development. *Mol. Reprod. Dev.* 66, 225–236. doi: 10.1002/mrd.10349
- Rassoulzadegan, M., Grandjean, V., Gounon, P., Vincent, S., Gillot, I., and Cuzin, F. (2006). RNA mediated non-mendelian inheritance of an epigenetic change in the mouse. *Nature* 441, 469–474. doi: 10.1038/nature04674

- Reilly, J. N., McLaughlin, E. A., Stanger, S. J., Anderson, A. L., Hutcheon, K., Church, K., et al. (2016). Characterisation of mouse epididymosomes reveals a complex profile of microRNAs and a potential mechanism for modification of the sperm epigenome. *Sci. Rep.* 6:31794. doi: 10.1038/srep31794
- Rodgers, A. B., Morgan, C. P., Bronson, S. L., Revello, S., and Bale, T. L. (2013). Paternal stress exposure alters sperm microRNA content and reprograms offspring HPA stress axis regulation. *J. Neurosci.* 33, 9003–9012. doi: 10.1523/JNEUROSCI.0914-13.2013
- Rodgers, A. B., Morgan, C. P., Leu, N. A., and Bale, T. L. (2015). Transgenerational epigenetic programming via sperm microRNA recapitulates effects of paternal stress. *Proc. Natl. Acad. Sci. U.S.A.* 112, 13699–13704. doi: 10.1073/pnas.1508347112
- Sciamanna, I., Barberi, L., Martire, A., Pittoggi, C., Beraldi, R., Giordano, R., et al. (2003). Sperm endogenous reverse transcriptase as mediator of new genetic information. *Biochem. Biophys. Res. Commun.* 312, 1039–1046. doi: 10.1016/j.bbrc.2003.11.024
- Sciamanna, I., Gualtieri, A., Cossetti, C., Osimo, E. F., Ferracin, M., Macchia, G., et al. (2013). A tumor-promoting mechanism mediated by retrotransposon-encoded reverse transcriptase is active in human transformed cell lines. *Oncotarget* 4, 2271–2287. doi: 10.18632/oncotarget.1403
- Sciamanna, I., Gualtieri, A., Piazza, P. V., and Spadafora, C. (2014). Regulatory roles of LINE-1-encoded reverse transcriptase in cancer onset and progression. *Oncotarget* 5, 8039–8051. doi: 10.18632/oncotarget.2504
- Sendler, E., Johnson, G. D., Mao, S., Goodrich, R. J., Diamond, M. P., Hauser, R., et al. (2013). Stability, delivery and functions of human sperm RNAs at fertilization. *Nucleic Acids Res.* 41, 4104–4117. doi: 10.1093/nar/gkt132
- Sharma, U., Conine, C. C., Shea, J. M., Boskovic, A., Derr, A. G., Bing, X. Y., et al. (2016). Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science* 351, 391–396. doi: 10.1126/science.aad6780
- Sinibaldi-Vallebona, P., Matteucci, C., and Spadafora, C. (2011). Retrotransposon-encoded reverse transcriptase in the genesis, progression and cellular plasticity of human cancer. *Cancers* 3, 1141–1157. doi: 10.3390/cancers3011141
- Smith, K., and Spadafora, C. (2005). Sperm-mediated gene transfer: applications and implications. *Bioessays* 27, 551–562. doi: 10.1002/bies.20211
- Smith, Z. D., Chan, M. M., Humm, K. C., Karnik, R., Mekhoubad, S., Regev, A., et al. (2014). DNA methylation dynamics of the human preimplantation embryo. *Nature* 511, 611–615. doi: 10.1038/nature13581
- Spadafora, C. (1998). Sperm cells and foreign DNA: a controversial relation. *Bioessays* 20, 955–964. doi: 10.1002/(SICI)1521-1878(199811)20:11<955::AID-BIES11>3.0.CO;2-8
- Spadafora, C. (2008). Sperm-mediated 'reverse' gene transfer: a role of reverse transcriptase in the generation of new genetic information. *Hum. Reprod.* 23, 735–740. doi: 10.1093/humrep/dem425
- Suh, N., Baehner, L., Moltzahn, F., Melton, C., Shenoy, A., Chen, J., et al. (2010). MicroRNA function is globally suppressed in mouse oocytes and early embryos. *Curr. Biol.* 20, 271–277. doi: 10.1016/j.cub.2009.12.044
- Tang, F., Kaneda, M., O'Carroll, D., Hajkova, P., Barton, S. C., Sun, Y. A., et al. (2007). Maternal microRNAs are essential for mouse zygotic development. *Genes Dev.* 21, 644–648. doi: 10.1101/gad.418707
- Vidigal, J. A., and Ventura, A. (2015). The biological functions of miRNAs: lessons from *in vivo* studies. *Trends Cell Biol.* 25, 137–147. doi: 10.1016/j.tcb.2014.11.004
- Vitulo, P., Sciamanna, I., Baiocchi, M., Sinibaldi-Vallebona, P., and Spadafora, C. (2012). LINE-1 retrotransposon copies are amplified during murine early embryo development. *Mol. Reprod. Dev.* 79, 118–127. doi: 10.1002/mrd.22003
- Vojtech, L., Woo, S., Hughes, S., Levy, C., Ballweber, L., Sauteraud, L. P., et al. (2014). Exosomes in human semen carry a distinctive repertoire of small non-coding RNAs with potential regulatory functions. *Nucleic Acids Res.* 42, 7290–7304. doi: 10.1093/nar/gku347
- Waddington, C. H. (1959). Canalization of development and genetic assimilation of acquired characters. *Nature* 183, 1654–1655. doi: 10.1038/1831654a0
- Wheeler, B. M., Heimberg, A. M., Moy, V. N., Sperling, E. A., Holstein, T. W., Heber, S., et al. (2009). The deep evolution of metazoan microRNAs. *Evol. Dev.* 11, 50–68. doi: 10.1111/j.1525-142X.2008.00302.x
- Zhong, C., Xie, Z., Yin, Q., Dong, R., Yang, S., Wu, Y., et al. (2016). Parthenogenetic haploid embryonic stem cells efficiently support mouse generation by oocyte injection. *Cell Res.* 26, 131–134. doi: 10.1038/cr.2015.132

**Conflict of Interest Statement:** The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Spadafora. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.