



Overview of the Antimicrobial Compounds Produced by Members of the *Bacillus subtilis* Group

Simon Caulier^{1,2†}, Catherine Nannan^{1†}, Annika Gillis¹, Florent Licciardi¹, Claude Bragard^{2*} and Jacques Mahillon^{1*}

¹ Laboratory of Food and Environmental Microbiology, Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, Belgium, ² Laboratory of Phytopathology-Applied Microbiology, Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, Belgium

OPEN ACCESS

Edited by:

José E. Barboza-Corona,
Universidad de Guanajuato, Mexico

Reviewed by:

Stefan Junne,
Technische Universität Berlin,
Germany
Jeongdae Im,
Kansas State University, United States

*Correspondence:

Claude Bragard
claude.bragard@uclouvain.be
Jacques Mahillon
jacques.mahillon@uclouvain.be

† These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Microbiotechnology, Ecotoxicology
and Bioremediation,
a section of the journal
Frontiers in Microbiology

Received: 05 November 2018

Accepted: 05 February 2019

Published: 26 February 2019

Citation:

Caulier S, Nannan C, Gillis A,
Licciardi F, Bragard C and Mahillon J
(2019) Overview of the Antimicrobial
Compounds Produced by Members
of the *Bacillus subtilis* Group.
Front. Microbiol. 10:302.
doi: 10.3389/fmicb.2019.00302

Over the last seven decades, applications using members of the *Bacillus subtilis* group have emerged in both food processes and crop protection industries. Their ability to form survival endospores and the plethora of antimicrobial compounds they produce has generated an increased industrial interest as food preservatives, therapeutic agents and biopesticides. In the growing context of food biopreservation and biological crop protection, this review suggests a comprehensive way to visualize the antimicrobial spectrum described within the *B. subtilis* group, including volatile compounds. This classification distinguishes the bioactive metabolites based on their biosynthetic pathways and chemical nature: *i.e.*, ribosomal peptides (RPs), volatile compounds, polyketides (PKs), non-ribosomal peptides (NRPs), and hybrids between PKs and NRPs. For each clade, the chemical structure, biosynthesis and antimicrobial activity are described and exemplified. This review aims at constituting a convenient and updated classification of antimicrobial metabolites from the *B. subtilis* group, whose complex phylogeny is prone to further development.

Keywords: *Bacillus subtilis* group, bacteriocins, biocontrol, biosynthetic pathways, lipopeptides, polyketides, siderophores, volatile

INTRODUCTION

The genus *Bacillus* comprises 377 species¹ (last update in January 2019) of Gram-positive, rod-shaped bacteria (Gordon et al., 1973). Their ability to form endospores, their diversity in physiological properties, as well as their capacity to produce numerous antimicrobial compounds (AMCs) favor their ubiquitous distribution in soil, aquatic environments, food and gut microbiota of arthropods and mammals (Nicholson, 2002).

Bacteria from the *Bacillus subtilis* group consist of small vegetative cells (<1 μm -wide) for which the strain *B. subtilis* subsp. *subtilis* 168 is considered as model organism (Barbe et al., 2009). They are usually mesophilic and neutrophilic, although some can tolerate high pH. The four original species of the group (*B. subtilis*, *Bacillus licheniformis*, *Bacillus pumilus*, and *Bacillus amyloliquefaciens*) were discovered more than 40 years ago (Gordon et al., 1973; Priest et al., 1987). Since then, the evolution of their molecular, chemotaxonomic and physiological characterizations led to regular re-evaluations and (re-)description of numerous novel species and subspecies (see current taxonomy of the group in **Figure 1**) (Fan et al., 2017).

¹<http://www.bacterio.net/bacillus.html>

The potential of *B. subtilis* group strains to produce a wide diversity of secondary metabolites mediating antibiosis was recognized for decades. For any given strain of the *B. subtilis* group, it is now estimated that at least 4–5% of its genome is devoted to antimicrobial compounds (AMCs) production (Stein, 2005). These molecules are mainly antimicrobial peptides (AMPs). Their structures are usually cyclic, hydrophobic and contain peculiar moieties such as D-amino acids (AA) or intramolecular thioether bonds. In addition to AMPs, volatile metabolites also constitute a large family of antimicrobials exhibiting numerous metabolic and functional roles.

Due to the wide diversity of these molecules, their classification is rather complex and can be based on several criteria such as their biosynthetic machinery, sources, biological functions, properties, three-dimensional structure, covalent bonding pattern or molecular targets (Tagg et al., 1976; Wang et al., 2015). Here a classification of the *B. subtilis* group antimicrobial molecules is proposed, based on their biosynthetic pathways and their chemical nature as shown in **Figure 2**. This review will emphasize the biosynthesis pathway and the bioactivity of the main clades of AMCs within the *B. subtilis* group: *i.e.*, the ribosomal peptides (RPs) (bacteriocins and enzymes), the polyketides (PKs), the non-ribosomal peptides (NRPs) and the volatiles. A full overview of this chart is provided as **Supplementary Material (Supplementary Figure S1)**.

RIBOSOMAL PEPTIDES

Ribosomally synthesized peptides (RPs) are usually derived from short precursors (*ca.* 100 AA) and are processed to mature compounds through post-translational modifications (Oman and van der Donk, 2009). Various enzymes mediate these modifications and therefore generate a wide diversity of chemical structures. Most of these peptides were originally referred to as “bacteriocins,” characterized as low molecular weight molecules that exhibit inhibiting growth activities against bacteria closely related to the producing strain (Klaenhammer, 1988; Chopra et al., 2015). In addition to bacteriocins, other types of enzymes exhibiting antagonistic activities are also ribosomally synthesized. However, those compounds display diverse metabolic activities such as quorum sensing (QS) mediation, cell lysis or induction of genetic competence (Schmidt, 2010; Shafi et al., 2017). It should also be noted that molecules referred to as BLIS (bacteriocin-like inhibitory substances) include AMPs for which the ribosomal synthesis has not been confirmed yet (Abriouel et al., 2011).

B. subtilis Group Bacteriocins

It is estimated that 99% of the bacteria and archaea are able to produce at least one bacteriocin. Historically, lactic acid bacteria (LAB) were studied as main bacteriocin producers, mostly because of their long history of safe use in food fermentation (O’Sullivan et al., 2002). Nisin (**Figure 3C**), produced by *Lactobacillus lactis* subsp. *lactis*, was approved as a food additive in the 1960s and has since then been used in over 50 countries for its antimicrobial activity against Gram-positive pathogens such as *Clostridium* spp. and *Bacillus* spp. (Klaenhammer, 1988;

Delves-Broughton, 1990). However, the search for new bioactive molecules has rapidly expanded to other bacteriocin-producing genera, with a particular attention, in the late 1990s, to the GRAS (generally recognized as safe) *Bacillus* species whose bacteriocin antimicrobial spectra were broader than those of LAB (Pedersen et al., 2002; Riley and Wertz, 2002; Sumi et al., 2015).



The generic biosynthetic pathway of *Bacillus* species bacteriocins includes several post-translational modifications, including the proteolytic cleavage of the leader peptide at the N-terminal end (McIntosh et al., 2009). The modifications of active peptides, its secretion and the immunity to the bacteriocin (as described below) vary depending on the bacteriocin class.

While many classifications have been suggested over the years, one reasonable way to cope with the diversity of the *Bacillus* bacteriocins is to sort them on the basis of their biosynthetic pathway as previously reported for *Streptococcus* spp. and *Enterococcus* spp. bacteriocins (Nes et al., 2007) and reviewed in Abriouel et al. (2011). Accordingly, three main classes subdivided into several subclasses can be distinguished for the *B. subtilis* group. As detailed in **Table 1**, Class I includes the post-translationally modified peptides such as the lantibiotics whereas the non-modified peptides are grouped in Class II; Class III involved bacteriocins larger than 10 kDa (Abriouel et al., 2011). **Supplementary Table S1** summarizes the different RPs produced by the strains belonging to the *B. subtilis* group, as well as their reported antimicrobial activities.

Class I includes small AMPs (19–38 AA) with extensive post-translational modifications. Subclasses I.1, I.2, and I.3 have in common their lantibiotic structure, which refers to inter-residual thioester bonds made of modified AA residues. As illustrated in **Figure 3**, lantibiotics involve 2,3-didehydroalanine (Dha) and (Z)-2,3-didehydrobutyrine (Dhb), resulting from the dehydration of serine and threonine residues, respectively. The intra-molecular addition of Dha or Dhb on a cysteine residue leads to the respective formation of lanthionine and methyllanthionine bridges (Willey and Donk, 2007). Subtilin (**Figure 3B**), from subclass I.1, is one of the most studied bacteriocins from the *B. subtilis* group. Its structure shares several similarities with nisin A lantibiotics, shown in **Figure 3C** (Guder et al., 2000; Abriouel et al., 2011). Peptides from subclass I.4 undergo other types of modifications. For instance, subtilosin A is a head-to-tail cyclic peptide with unusual inter-residue linkages (*i.e.*, Cys-Phe bond) (Marx et al., 2001; Kawulka et al., 2004).


Class II bacteriocins include small (<10 kDa), linear and non-modified peptides, resistant to heat and acido-basic treatments. They are divided in three subclasses based on a conserved AA motif near their N-terminus. The YGNGVXC (X is any AA) motif is associated to pediocin-like peptides from subclass II.1 whereas DWTXWSXL is specific to thuricin-like peptides from subclass II.2. Subclass II.3 comprises the small non-modified AMPs without any typical motif in their AA sequence (Abriouel et al., 2011). Finally, class III bacteriocins consist into large and heat labile molecules, generally characterized by a phospholipase activity (Cleveland et al., 2001).

Because of their wide diversity, bacteriocins display different modes of action such as protoplasm vesicularization, pore formation or cell disintegration (Sumi et al., 2015). They are



generally bactericidal with some exceptions that exhibit bacteriostatic activities (Gautam and Sharma, 2009). For most class I and II bacteriocins, the target of their activity is the bacterial envelope due to their amphiphilic or hydrophobic

properties. For instance, lantibiotics from subclass I.1 have a dual mode of action. On the one hand, they can inhibit the cell wall synthesis of the targeted bacteria through binding to lipid II, the major transporter of peptidoglycan subunits



across the inner cell membrane. On the other hand, lipid II can be used as a docking molecule to insert the lantibiotic in the membrane leading to pore formation and ultimately to cell death as well described in Chatterjee et al. (2005) and Cotter et al. (2005). This duality has been reported for subtilin, a class I bacteriocin which is active against a broad range of Gram-positive bacteria such as *Staphylococcus simulans*, *B. subtilis*, and *Bacillus stearothermophilus* (Linnett and Strominger, 1973; Parisot et al., 2008).

Many regulation systems mediate bacteriocin production, secretion and immunity. Bacteriocin production is usually linked

to particular cellular events such as stress responses. For instance, subtilin production depends on cell density and is increased under starvation conditions (Abriouel et al., 2011). Lantibiotic production is also mediated by QS. For subtilin, it has been demonstrated that the peptide itself acts as an auto-inducer of its own production (Kleerebezem, 2004). The export of bacteriocins is generally ensured by a dedicated membrane-associated ATP-Binding Cassette (ABC) transporter. For some lantibiotics, the cleavage of the leader peptide often occurs in a proteolytic domain present in the ABC transporter as described in McAuliffe et al. (2001) and Cotter et al. (2005). The immunity

TABLE 1 | Classification of the *B. subtilis* group bacteriocins.

Class	Class description	Subclass	Subclass description
I	Post-translationally modified peptides	I.1	Single-peptide, elongated lantibiotics
		I.2	Other single-peptide lantibiotics
		I.3	Two-peptide lantibiotics
		I.4	Other modified peptides
II	Non-modified peptides	II.1	Pediocin-like peptides
		II.2	Thuricin-like peptides
		II.3	Other linear peptides
III	Large peptides (> 10 kDa)		


Adapted from Abriouel et al. (2011).

of the producing strains to its own active bacteriocin(s) can be achieved by several mechanisms like the secretion of immunity proteins sequestering the peptide, the bacteriocin re-export through an ABC transporter system or the alteration of the targeted peptidoglycans bonds (e.g., modification of the cell wall or cytoplasmic membrane charge) (Cotter et al., 2005; Dubois et al., 2009).

B. subtilis Group AMP Enzymes

Among the *B. subtilis* group, two major types of enzymes exhibit antagonistic activities (**Supplementary Table S1**): the lytic enzymes and those involved in quorum quenching (QQ). Several strains from the *B. subtilis* group have indeed been identified as capable to produce lytic enzymes with biocontrol potential (Herrera-Estrella and Chet, 1999; Kumar et al., 2012; Shafi et al., 2017). They include cellulases, glucanases, proteases and chitinases and are generally referred to as cell wall degrading enzymes (CWDE) (Ariffin et al., 2006; Alamri, 2015; Caulier et al., 2018). They are particularly active against fungi since chitin and glucan are the major constituents of their cell wall where various glycoproteins are embedded (Bowman and Free, 2006; Geraldine et al., 2013; Gomaa, 2012).


Quorum quenching is able to silence or block QS which is generally defined as the cell-to-cell communication mechanism through the production of signal molecules (Czajkowski and Jafra, 2009). *N*-acyl-homoserine lactones (AHLs), composed of a fatty acid side chain and a homoserine lactone (**Figure 4**) are the most characterized signal autoinducers in Gram-negative bacteria. When a bacterial population proliferates, concentration of AHLs increases so that all the cells coordinate their metabolic activities (e.g., biofilm formation, sporulation, virulence factors or antibiotic production) (Dong et al., 2004). As the QS system brings ecological advantages to a coordinate population, QQ is able to counteract QS. Four types of enzymes (i.e., lactonase, decarboxylase, acylase, and deaminase) are able to inactivate AHLs, as illustrated in **Figure 4** (Czajkowski and Jafra, 2009). *B. subtilis* AHL-lactonases have for instance attracted interest for biocontrol since they affect the growth of deleterious microbial pest such as *Pectobacterium carotovorum* subsp. *carotovorum* causing potato soft rot (González and Keshavan, 2006).



POLYKETIDES

Among the bioactive compounds produced by microorganisms, PKs are well known from the human health sector for their broad spectrum of activity encompassing antibacterial, immunosuppressive, antitumor and many more antagonistic abilities. Typical PKs structures from the *B. subtilis* group are presented in **Figure 5**. They are synthesized from acyl CoA precursors such as malonate and methyl malonate. Their biosynthesis depends on multifunctional polyketide synthases (PKSs). Their structure was first extrapolated from fatty acid synthases (FASs) that share similarities in terms of chain extension mechanisms, precursors and overall architecture design (Smith and Tsai, 2007). As shown in **Figure 6A**, PKS are composed of a succession of elongation modules, flanked by initiation and termination modules. The reactive mechanism of these three PKS domains is illustrated in **Figure 7A** and is well summarized in Hertweck (2009). The initiation module is composed of two domains: an acyltransferase (AT) domain that recruits and catalyzes the binding of a monomer substrate to an acyl carrier protein (ACP) domain. The ACP then acts as an arm with a second catalytic domain located on the next elongation module. This domain, a β -ketoacyl synthase (KS), catalyzes the chain-elongation reaction that occurs through a decarboxylative Claisen thioester condensation (Cane and Walsh, 1999; Hertweck, 2009). In addition to the three core domains, auxiliary domains can also be present on elongation modules (gray domains in **Figure 6A**). These auxiliary domains mediate ketoreduction (KR), dehydration (DH), or enoylacyl reduction (ER) occurring before the chain-elongation reaction. These modifications considerably enrich the structural complexity and diversity of mature PKs (Hertweck, 2009). Finally, a termination module harboring an additional thioesterase (TE) domain catalyzes the macrolactonization and the release of the mature PK (Cane and Walsh, 1999).

Polyketide synthases have been classified in three canonical types based on the structural organization of their functional domains. Type I PKSs involve large multifunctional enzymes housing several domains linearly arranged and covalently bonded. Type II PKSs are multienzyme complexes composed of separate monofunctional enzymes combined during the PK



synthesis. Type III PKSs are chalcone synthase-like PKSs that operate the acid CoA thioesters directly without any ACP domain (Chen and Du, 2016). Beside these structural differences, PKSs are classified as iterative or non-iterative depending on how many KS domains are used in the biosynthetic process. Within prokaryotes, the non-iterative type I PKSs is the most represented. They produce PK compounds that harbor a one-to-one correspondence with the PKS modular architecture. This conservation of collinearity is used for PKS discovery via genome mining (Challis, 2008).

Due to the diversity of PKSs, many exceptions and transition states between the three main types are observed. In some cases, mixed PKs pathways combine different types of PKSs or can even be associated with FASs or NRP synthetases (NRPSs) to form PK-peptide hybrid metabolites such as bacillaene, compactin, fusarin C or salinosporamide A (Moldenhauer et al., 2007; Hertweck, 2009; Fisch, 2013).












TABLE 2 | Major classes of polyketides.

Polyketide class	Structure description	Typical core unit or example structure	Reference
Acetogenins	Linear 32- or 34-carbon chains with oxygenated functional groups bearing a terminal γ -lactone ring		Li et al., 2008
Ansamycins	Bridge between an aromatic moiety and an aliphatic chain		Williams, 1975
Eneclines	Compounds characterized by a core structure formed by a double C-C bond conjugated to two acetylenic groups		Horsman et al., 2009
Macrolides	Large macrocyclic lactone ring with one or more deoxy sugars		Yuan et al., 2012a
Polyenes	Poly-unsaturated organic compounds containing at least three alternating single and double C-C bonds		Hertweck, 2009
Polyethers	Polymers containing more than one ether group		Hertweck, 2009
Tetracyclines	Compounds family characterized by a typical four-ring system		Rohr, 1992

To date, seven PKs families have been recognized based on their carbon skeletons and typical structures, as summarized in **Table 2** (Eustáquio et al., 2009). However, to our knowledge, only three antimicrobial PKs and their variants are produced within the *B. subtilis* group: bacillaene, difficidin, and macrolactin. These compounds exhibit antibacterial activities through selective inhibition of protein synthesis (**Table 3**). Bacillaene is a polyene

PK resulting from a hybrid synthesis by a type I PKS and a NRPS *bae* operon (*bae*, *baeL*, *baeM*, *baeN* and *baeR*) (Chen et al., 2006; Moldenhauer et al., 2007). Its exhibits antimicrobial activity against various bacteria (e.g., *Myxococcus xanthus* or *Staphylococcus aureus*) and fungi (e.g., *Trichoderma* spp. or *Fusarium* spp.) (Patel et al., 1995; Um et al., 2013; Müller et al., 2014). Difficidin, and its oxidized form oxydifficidin,

TABLE 3 | PKS and hybrids NRPS/PKS produced by strains of the *B. subtilis* group.

PKS or hybrids class*	Compound	Antimicrobial activity**		References
		Antibacterial activity	Antifungal activity	
Macrolides	7-O-malonyl-macrolactin A	<i>B. cepacia</i> ^c , <i>Enterococci faecalis</i> ^c , <i>R. solanacearum</i> ^c , <i>S. aureus</i> ^c	<i>F. oxysporum</i> f. sp. <i>cubense</i> ^c	Romero-Tabarez et al., 2006; Yuan et al., 2012a
Macrolides	7-O-succinyl-macrolactin F	<i>B. subtilis</i> ^c , <i>S. aureus</i> ^c	–	Jaruchoktaweetchai et al., 2000; Nagao et al., 2001
Macrolides	7-O-succinyl-macrolactin A	<i>B. subtilis</i> ^c , <i>R. solanacearum</i> ^c , <i>S. aureus</i> ^c	<i>F. oxysporum</i> f. sp. <i>cubense</i> ^c	Jaruchoktaweetchai et al., 2000; Yuan et al., 2012a
Macrolides	Macrolactin A	<i>R. solanacearum</i> ^c	<i>F. oxysporum</i> f. sp. <i>cubense</i> ^c	Yuan et al., 2012a
Macrolides	Macrolactin D	<i>S. aureus</i> ^c	<i>A. solani</i> ^c , <i>Pyricularia oryzae</i> ^c	Xue et al., 2008
Macrolides	Macrolactin F, G, H, I, J, K, L, M	<i>B. subtilis</i> ^c , <i>S. aureus</i> ^c	–	Jaruchoktaweetchai et al., 2000; Nagao et al., 2001
Macrolides	Macrolactin N	<i>E. coli</i> ^c , <i>S. aureus</i> ^c	–	Yoo et al., 2006
Macrolides	Macrolactin Q	<i>B. subtilis</i> ^c , <i>E. coli</i> ^c , <i>P. aeruginosa</i> ^c , <i>S. aureus</i> ^c	–	Mojid Mondol et al., 2011
Macrolides	Macrolactin S	<i>B. subtilis</i> ^c , <i>E. coli</i> ^c , <i>S. aureus</i> ^c	<i>P. oryzae</i> ^c	Lu et al., 2008
Macrolides	Macrolactin T	<i>S. aureus</i> ^c	<i>A. solani</i> ^c , <i>P. oryzae</i> ^c	Xue et al., 2008
Macrolides	Macrolactin W	<i>B. subtilis</i> ^c , <i>E. coli</i> ^c , <i>P. aeruginosa</i> ^c , <i>S. aureus</i> ^c	–	Mojid Mondol et al., 2011
Polyenes	Bacillaene A	<i>B. thuringiensis</i> ^c , <i>E. coli</i> ^c , <i>Klebsiella pneumoniae</i> ^c , <i>M. xanthus</i> ^c , <i>P. vulgaris</i> ^c , <i>Serratia marcescens</i> ^c , <i>S. aureus</i> ^c	<i>Corioliopsis</i> spp. ^c , <i>Fusarium</i> sp. ^c , <i>Pseudoxylaria</i> sp. ^c , <i>Trichoderma</i> sp. ^c , <i>Umbelopsis</i> sp. ^c	Patel et al., 1995; Um et al., 2013; Müller et al., 2014
Polyenes	Difficidin	<i>Actinomyces naeslundii</i> ^c , <i>Bacteroides distasonis</i> ^c , <i>C. perfringens</i> ^c , <i>E. amylovora</i> ^c , <i>E. coli</i> ^c , <i>Eubacterium limosum</i> ^c , <i>K. pneumoniae</i> ^c , <i>P. vulgaris</i> ^c , <i>P. aeruginosa</i> ^c , <i>S. marcescens</i> ^c , <i>S. aureus</i> ^c , <i>Streptococcus faecalis</i> ^c , <i>X. oryzae</i> ^c	–	Zimmerman et al., 1987; Chen et al., 2009; Wu et al., 2015b
Polyenes	Oxydifficidin	<i>A. naeslundii</i> ^c , <i>B. distasonis</i> ^c , <i>C. perfringens</i> ^c , <i>E. coli</i> ^c , <i>E. limosum</i> ^c , <i>K. pneumoniae</i> ^c , <i>P. vulgaris</i> ^c , <i>P. aeruginosa</i> ^c , <i>S. marcescens</i> ^c , <i>S. aureus</i> ^c , <i>S. faecalis</i> ^c	–	Zimmerman et al., 1987
Hybrids PKs/NRPs	Kanosamine	–	<i>C. albicans</i> ^p , <i>Saccharomyces cerevisiae</i> ^p	Janiak and Milewski, 2001; van Straaten et al., 2013

^c Activity of isolated compound confirmed by compound purification or mutant deletion, ^p putative activity of the compound contained in a broth mixture. * Two PKS classes are reported in this review (macrolides and polyenes) as well as the hybrids between PKs and NRPs. ** –, no activity known.

are polyenes synthesized by a type I PKS encoded in the *dif* operon. They both inhibit bacterial pathogens such as *Clostridium perfringens*, *Erwinia amylovora*, *Escherichia coli* or *Xanthomonas oryzae* (Zimmerman et al., 1987; Chen et al., 2009; Aleti et al., 2015; Wu et al., 2015b). Finally, macrolactins and their 7-O-succinyl- or 7-O-malonyl-derivatives are synthesized via a type I PKS. They show antibacterial and antifungal activities against *Burkholderia cepacia*, *Ralstonia solanacearum*, *S. aureus* or *Fusarium oxysporum* (Romero-Tabarez et al., 2006; Yoo et al., 2006; Yuan et al., 2012a). Some macrolactins, such as the macrolactin A, apparently also displays antiviral properties (e.g., against Herpes simplex viruses) (Gustafson et al., 1989).

NON-RIBOSOMAL PEPTIDES

Non-ribosomal peptides form a versatile family of secondary metabolites with growing interest in many industrial fields

as antibiotics, siderophores, surfactants, pigments, immunosuppressors or antitumor molecules (Wang et al., 2014). NRPs show a broad structural diversity, from linear to cyclic or branched structures (Kopp and Marahiel, 2007). As illustration, the Norine database counts almost 1.200 NRP molecules, including their structure, synthesis and evolution² (last update in January 2019) (Caboche et al., 2008).

Two categories of NRPs can be distinguished whether they are synthesized through a multi-enzyme thio-template mechanism or not (Sumi et al., 2015). The first ones usually result in structures with two to *ca.* 50 residues and other moieties such as fatty acid chains [*i.e.*, lipopeptides (LPs) and siderophores] whereas the second ones are generally smaller. **Figure 8** shows the chemical structures of typical NRPS from the *B. subtilis* group.

²<http://bioinfo.lifl.fr/norine>

Kim et al., 1997; Cosby et al., 1998). Warm temperature ($\geq 37^\circ\text{C}$) and anaerobic conditions increase the production of surfactins while lower temperatures ($25\text{--}37^\circ\text{C}$) and aerated bioreactors favor fengycins and iturins family metabolites (Jacques, 2011). The production of surfactins by *B. subtilis* is also QS-dependent and involves ComX and PhrC. These pheromones trigger complex cascades regulating cell density-dependent processes such as sporulation and competence (Hamoen et al., 2003; Ongena et al., 2005).

Iturins and fengycins are mainly known for their strong antifungal activity against several plant and human pathogenic fungi (**Supplementary Table S2**). In addition, iturin-like mycosubtilin, bacillomycin R, subtilene A and eumycin show antibacterial properties (Besson et al., 1976; Leclere et al., 2005; Thasana et al., 2010). Contrary to iturins and fengycins, surfactins mainly display antiviral and antibacterial activities (Ongena and Jacques, 2008). Their antiviral activity essentially targets enveloped viruses (e.g., herpes simplex or porcine epidemic diarrhea viruses). They also inhibit pathogenic bacteria such as *Legionella pneumophila*, *Listeria monocytogenes*, *R. solanacearum* or *X. oryzae* (Naruse et al., 1990; Yakimov et al., 1995; Sabaté and Audisio, 2013; Loiseau et al., 2015; Luo et al., 2015). However, some surfactins are able to control important fungal plant and human pathogens such as *Botrytis cinerea*, *Candida albicans*, *F. oxysporum* or *Rhizoctonia solani* (Jenny et al., 1991; Lee et al., 2007; Qi et al., 2010; Dimkić et al., 2013; Romano et al., 2013).

The mere composition of LPs, where a peptide moiety is bound to a lipid tail, gives them an amphiphilic property. This nature makes them excellent surfactants and plays a significant role in their biological functions and antimicrobial properties. Indeed, LPs are able to destabilize the plasma membrane via a pore forming activity leading to the cell death of the target microbes. Their antiviral activity is the result of a similar disintegration of the bi-lipid envelope of virions explaining the weak LPs activity against plant viruses among which very few are enveloped (Ongena and Jacques, 2008).

Bacillus spp. LPs have many other biological and ecological functions as fully documented by Raaijmakers et al. (2010). They are also known to impact other metabolic mechanisms such as biofilm formation, motility, virulence, plant root colonization, and plant defenses. Moreover, it has been suggested that their participation to the degradation of hydrophobic substrates could be used for polluted soils bioremediation (Mulligan et al., 2001). Although some lipopeptides have already been exploited as food biopreservatives or crop protection products, the industrial interest for LPs in specific applications is unsurprisingly continuously growing.

Thiotemplate NRPs – Siderophore

Itoic acid is a mono-peptide composed of a 2,3-dihydroxybenzoate (DHB) molecule bound to a glycine. It is used as a precursor by trimodular NRPS machinery to produce bacillibactin which is obtained after a condensation of three units of DHB-glycine-threonine (May et al., 2001). The synthesis of the final hexapeptide is catalyzed by a terminal thioesterase domain leading to the production of a methylated trilactone ring link to three catecholates moieties. It is this cyclic structure that enables

the sequestration of the metal atom (Dertz et al., 2006). Itoic acid and bacillibactin are both catecholic siderophores that chelates iron reducing its bioavailability. This is limited access to iron that allows *B. subtilis* to antagonize the growth of other surrounding microbes such as, for instance, *F. oxysporum* f. sp. *capsici* (Yu et al., 2011).

Non-thiotemplate NRPs

Bacteria from the *B. subtilis* group are also able to synthesize other antimicrobial NRPs through non-thiotemplate mechanism. Rhizocticins are di- and tri-phosphono-peptides. They are constituted of a L-2-amino-5-phosphono-3-*cis*-pentenoic acid (APPA) linked to an arginine (rhizocticin A). They can be supplemented with an additional valine (rhizocticin B), isoleucine (rhizocticine C) or leucine (rhizocticine D). After their integration into the target microbes, their cleavage by host cell peptidases releases the fungitoxic L-APPA moiety that interferes with threonine metabolism in fungal cells. Interestingly, rhizocticin A has also an antagonistic activity against nematodes such as *Caenorhabditis elegans* (Kugler et al., 1990).

In addition to rhizocticin compounds, two other dipeptide NRPs are produced by *B. subtilis*: bacilysin (also known as tetaïne) and its chlorinated derivative, chlorotetaïne. They contain L-alanine (or chlorine-L-alanine) bound to the non-proteinogenic L-anticapsin (Kenig and Abraham, 1976; Rapp et al., 1988). Despite their simple composition, these bioactive compounds display strong antibacterial activity mediated by the anticapsin moiety that inhibits the glucosamine-6-phosphate synthase. Its inhibition suppresses the biosynthesis of peptidoglycans that are the main constituents of bacterial cell wall (Steinborn et al., 2005; Mahlstedt and Walsh, 2010). For the fungi, it has been proposed that because anticapsin is able to inhibit the production of chitin and fungal membrane mannoproteins, bacilysin and chlorotetaïne exhibit antifungal activity against *Aspergillus fumigatus* or *C. albicans* (Milewski et al., 1986; Rapp et al., 1988).

Finally, bacitracin and mycobacillin are two cyclic polypeptides produced by *B. subtilis*. Bacitracins are dodecapeptides containing a cyclic heptapeptide linked to a thiazoline ring (Johnson et al., 1945). They are mostly active against Gram-positive bacteria where they inhibit the bacterial cell-wall biosynthesis by preventing the lipid carrier from re-entering in the reaction cycle of peptidoglycan synthesis (Siewert and Strominger, 1967). Besides this primary mode of action, bacitracin might also act through other mechanisms affecting membrane functions, hydrolytic enzymes and/or the biosynthesis of ubiquinone precursors (Konz et al., 1997). Mycobacillin is an antifungal cyclic tridecapeptide altering the membrane of fungi like *Aspergillus niger* (Majumdar and Bose, 1958). Interestingly, its biosynthesis is rather peculiar. Although it is catalyzed by a large NRPS complex, it is divided in three fractions (A, B, and C) and does not use a thio-template mechanism (Zuber et al., 1993). Each fraction of the enzymatic complex contains a single enzyme polypeptide that catalyzes the polymerization of a first pentapeptide (A), a second nonapeptide (B) and the final tridecapeptide.

VOLATILES

Besides RPs, NRPs and PKs, strains from the *B. subtilis* group are able to produce a wide diversity of volatile compounds encompassing important roles especially in soil, one of the major habitats of this group (**Supplementary Figure S1**). Volatiles are notably involved in the bioconversion of the food chain, in the biogeochemical cycles of essential elements, in many physiological and metabolic reactions (e.g., nitrification, nitrogen mineralization, electron acceptor or donor reactions) as well as in communication signals triggering QS/QQ or defense mechanisms well reviewed in Effmert et al. (2012). Volatile compounds are generally classified into inorganic (VICs) and organic (VOCs) categories.

Volatile Inorganic Compounds (VICs)

Volatile inorganic compounds synthesized by microorganisms are mainly by-products of primary metabolism. They are carbonated, hydrogenated, sulfur or nitrogen-containing compounds such as CO₂, CO, H₂, HCN, H₂S, N₂, NH₃ and NO. Nitrogen-containing compounds are mostly released in aerated upper sediments layers by denitrifying bacteria. In this process, nitric oxide is enzymatically produced by the nitric-oxide reductase or the nitric-oxide synthase (Adak et al., 2002). The range of antimicrobial activities exhibited by VIC nitrogen-containing compounds from the *B. subtilis* group is wide. For instance, NO is able to induce systemic acquired resistance (SAR) in plants against bacterial pathogens such as *R. solanacearum* (Wang et al., 2005). *A contrario*, ammonia, a secondary metabolite from the catabolism of the amino acids L-aspartate, is known to be active against soil-borne Oomycetes such as *Pythium* spp. (Howell et al., 1988). Hydrogen cyanide, derived from the glycine catabolism, shows a direct antagonistic activity against aerobic microorganisms by inhibiting metal-containing enzymes such as the cytochrome c oxidase active in the respiration chain (Cherif-Silini et al., 2016).

Deeper in the soil, under low oxygen concentration, bacteria tend to produce different VICs such as H₂ or H₂S. Those compounds can serve as electron acceptors, AA precursors or antimicrobial metabolites. Hydrogen sulfide could be produced by *B. subtilis* from sulfate reduction or as a by-product of L-methionine and L-cysteine catabolism via a direct cleavage of L-methionine or a transamination followed by reductive demethylations (Even et al., 2006; Schulz and Dickschat, 2007). It is known to exhibit antifungal activity against several plant pathogens such as *A. niger* or *Penicillium italicum* but also against some food-borne bacteria or human pathogens (Fu et al., 2014). Curiously, it is also known to act as a bacterial defense mechanism against antibiotics (Shatalin et al., 2011). Interestingly, ammonia increases the resistance of several Gram-negative and Gram-positive bacteria to antibiotics too (Bernier et al., 2011).

Volatile Organic Compounds (VOCs)

Volatile organic compounds are small compounds with fewer than 20 carbon atoms and are characterized by low molecular mass (100–500 Da), high vapor pressure, low boiling point and a lipophilic moiety. These features ensure an easy evaporation

and a long distance distribution which is convenient in a complex matrix like soil (Schmidt et al., 2015). Their diffusion and production by soil-borne microbes are strongly dependent on various factors such as nutrient and oxygen availability, temperature, pH, physiological state of microorganisms, soil moisture, texture and architecture (McNeal and Herbert, 2009; Insam and Seewald, 2010; Effmert et al., 2012). The majority of VOCs derives from glucose oxidation involving glycolysis and the subsequent cycles such as the tricarboxylic acid cycle (TCA) as it has been well summarized in Korpi et al. (2009) and Schmidt et al. (2015). However, their production can also result from various other pathways such as aerobic heterotrophic carbon metabolism, fermentations, AA degradation, terpenes synthesis or sulfur reduction (Peñuelas et al., 2014). Based on previous reviews presented in Schulz and Dickschat (2007); Peñuelas et al. (2014) and Audrain et al. (2015), five categories of VOCs can be distinguished: (1) fatty acids and derivatives, (2) terpenoids, (3) nitrogen-containing VOCs, (4) sulfur-containing VOCs, and (5) metallo- or halogenated-containing VOCs. To date, about 2,000 compounds produced by almost 1,000 species of microorganisms have been listed in the mVOC 2.0 database (Lemfack et al., 2018). According to this database, almost 70% of recorded *Bacillus* VOCs are fatty acids derivatives (alcohols, ketones, alkanes, aldehydes, alkenes, and acids) followed by sulfur- and nitrogen-containing compounds. **Supplementary Table S3** displays the VOCs produced within the *B. subtilis* group and their antimicrobial activity.

Since many volatile fatty acids and their derivatives result from the glucose metabolism, their precursors mostly derive from the Embden-Meyerhof (glycolysis), Entner-Doudoroff, heterolactic and homolactic fermentation pathways (Peñuelas et al., 2014). *B. subtilis* bacteria, for instance, ferment pyruvate to produce ketone compounds such as acetoin (3-hydroxy-2-butanone) or 2,3-butanedione under anaerobic conditions (Ryu et al., 2003). Other intermediates coming from fatty acid biosyntheses or their β -oxydations are also used as precursors by microbes and transformed into VOCs through a decarboxylation reaction or a reduction of their carboxyl group (Schulz and Dickschat, 2007). They provide essential hydrocarbons but also other fatty acid derivatives. An oxidative deamination of several amino acids can lead to the production of aldehyde, ketone or alcohol volatile too. For instance, the degradation of L-phenylalanine or L-tyrosine can be the first step of the aromatic volatile compounds synthesis such as benzene or its carbohydrate derivatives. Finally, benzenoid volatiles can also be synthesized by microbes through the shikimate pathway that leads to the formation of chorismate, a natural precursor of aromatic amino acids (Bentley and Haslam, 1990). Degradation of intermediates from the shikimate pathway or aromatic amino acids can also lead to the production of benzenoid volatiles (Dickschat et al., 2005).

This wide variety of volatile fatty acids and their derivatives make them the most important group of VOCs produce by microbes and represent up to 87% of known antimicrobial VOCs produced by *B. subtilis* bacteria (**Supplementary Table S3**). They can be divided in two main categories: hydrocarbons (alkanes, alkenes, alkynes) or carbohydrates (acids, alcohols, aldehydes, esters, furans, ketones, lactones, benzenoids). Among them,

benzenoids is the most represented sub-category followed by alkanes, aldehydes, ketones, acids, and alcohols. Even though benzenoids could be considered as an individual category, they can also be seen as fatty acids derivatives because a large majority of antimicrobial benzenoid volatile produced by *B. subtilis* harbor a benzene core linked to a fatty acid derivatives.

There is an important diversity of benzenoids, sometimes linked with carbohydrate chains containing nitrogen, sulfur or both. Most of these antimicrobial volatile exert fungicidal activities but some have been characterized for their antibacterial or nematocidal abilities, too. Their mode of action is rarely fully characterized. For instance, morphological abnormalities on fungal and bacterial cells have been documented after an exposition to *B. subtilis* VOCs (Tahir et al., 2017). Volatile such as 1,3-butadiene or 2,3-butanediol are also known to induce modifications in the expression of genes linked to the pathogenicity of *R. solanacearum* and *Pectobacterium carotovorum* (Marquez-Villavicencio et al., 2011; Tahir et al., 2017). In addition to direct antimicrobial activities, fatty acids volatile have also several other biological functions. For instance, acetoin and 2-butanone have the ability to stimulate plant defenses or to induce plant stress tolerance which then promote plant growth (Ryu et al., 2003; Ryu et al., 2004; Ryu, 2015). They are essentially produced by strains of *B. amyloliquefaciens*, *B. velezensis* or *B. subtilis* (Audrain et al., 2015).

Terpenes and their derivatives (also known as terpenoids or isoprenoids) are among the most abundant secondary metabolites found in living systems (Fisher et al., 2001; Gershenzon and Dudareva, 2007). They originate from two main precursors: isopentenyl pyrophosphate (IPP) and its allylic isomer the dimethylallyl pyrophosphate (DMAPP) (Schulz and Dickschat, 2007). IPP and DMAPP are also the end-products of the deoxy-xylulose phosphate pathway (DOXP) starting with pyruvate and glyceraldehyde-3-phosphate originating from the glucose metabolism (Fisher et al., 2001). Terpenoids can be synthesized from isoprene molecules too. Julsing et al. (2007) showed that, in *B. subtilis*, isoprene is not formed by the MVA or DOXP pathways but, as in plant systems, might be a product of the methylerythritol phosphate (MEP) pathway (Guan et al., 2015).

Isoprenoid compounds are produced by all living organisms for essential physiological functions such as electron transport, membrane fluidity, light harvesting, photoprotection, anchoring of molecules to specific membranes and signaling (Fisher et al., 2001). The signaling ability is particularly important and is associated with several antagonistic, mutualistic or multi-trophic interactions (Shrivastava et al., 2015). More than 25,000 terpenic compounds have been listed and, for the vast majority, their biological functions and roles remain unknown (Buckingham, 1997). Volatile terpenes are generally recognized for their ability to inhibit bacteria (Scortichini and Rossi, 1991), fungi (Hammer et al., 2003; Dambolena et al., 2008), nematodes (Gu et al., 2007) or insects (Lee et al., 2003; Justicia et al., 2005). They can be classified in three categories: isoprene, monoterpenes (C₁₀) and sesquiterpenes (C₁₅) (Schmidt et al., 2015).

The mode of action of these compounds might be linked to their lipophilic nature allowing them to destabilize the cell

membrane integrity (Cox et al., 2000; Inoue et al., 2004). To our knowledge, only two terpenes produced by *B. subtilis* show antimicrobial abilities: isoprene and monoterpene α -terpineol exhibit antagonistic activities against cyanobacteria and nematodes (Wright and Thompson, 1985; Gu et al., 2007).

Little is known about the biosynthetic pathways of nitrogen-containing VOCs. Nevertheless, it is accepted that two main routes can be used: a non-enzymatic amination of acylolins, that can lead to the formation of pyrazines (Schulz and Dickschat, 2007) or derived from α -aminoketone intermediates resulting from AA catabolism (Owens et al., 1997; Zhu et al., 2010).

Nitrogen-containing VOCs can be distinguished based on their cyclization rate. Within non-cyclic compounds, three groups are identified (amides, amines and imines) while there are five categories of cyclic compounds (azoles, pyrazines, pyridines, pyridazines, and pyrimidines). Pyrazines are strongly represented among microbial volatile and are separated in two classes: lower-alkylated and higher-alkylated pyrazines (Schulz and Dickschat, 2007). These compounds are characterized by a strong odor and several *B. subtilis* coming from the rhizosphere or from food fermentations have already been recognized as pyrazines producers (Sugawara et al., 1985; Kosuge and Kamiya, 1962; Larroche et al., 1999; Leejeerajumnean et al., 2001). Pyrazines from *B. subtilis* strains are known to exhibit antifungal and nematocidal activities (Gu et al., 2007; Chen et al., 2008; Chaves-López et al., 2015; Haidar et al., 2016). For instance, tetramethylpyrazine inhibits the growth of *Moniliophthora perniciosa* and *F. oxysporum* f. sp. *lactucae*. Additionally, it acts on sporulation and elongation of the germ-tube of *B. cinerea* (Chen et al., 2008; Chaves-López et al., 2015). It is interesting to note that *B. subtilis* pyrazines can also exhibit antibacterial activities such as pulcherrimic acid which inhibits the growth of *S. aureus*, *E. coli* and *Proteus vulgaris* (Coutts et al., 1965). Beside pyrazines, strains from the *B. subtilis* group are able to produce other nitrogen VOCs such as 1H-imidazole, 1-ethyl showing antifungal activities against numerous soil-borne phytopathogens (Lupetti et al., 2002; Liu et al., 2008; Snelders et al., 2009; Schmidt et al., 2015).

Microbial VOCs containing sulfur (VSCs) derive from two main pathways originated from inorganic or organic sources (Schulz and Dickschat, 2007): inorganic sulfate reduction in methylated inorganic sulfides compounds or, for some microbial VSCs, originate from catabolism of AA such as L-methionine or more rarely, L-cysteine (Schulz and Dickschat, 2007). Some VSCs are produced as secondary volatiles via the production of hydrogen sulfide or methanethiol. Indeed, these two compounds are important precursors for subsequent VSCs synthesis (Schulz and Dickschat, 2007; Sourabié et al., 2012). Within the *B. subtilis* group, multiple VSCs such as dimethyl disulfide (DMDS), dimethyl trisulfide (DMTS), S-methyl thioacetate or S-methyl butanethioate have been characterized for their antifungal and nematocidal activities (Coosemans, 2005; Gerik, 2005; Gu et al., 2007; Kai et al., 2009; Wang et al., 2009; de Vrieze et al., 2015; Schmidt et al., 2015; Velivelli et al., 2015; Gotor-Vila et al., 2017). A putative antibacterial effect of DMDS is not to exclude. Indeed, DMDS is known to affect the bacterial cell-to-cell communications through a decrease in the

amount of *N*-acyl homoserine lactone (AHL) mediating QS (Chernin et al., 2011).

Other volatile organic compounds such as halogenated, metalloids, tellurium or selenium compounds have also been described. However, at the time of writing, no *B. subtilis* strains have been proved to produce these type of VOCs (Schulz and Dickschat, 2007), although related bacteria, like *Bacillus arsenicoselenatis*, have been shown to generate them (Switzer Blum et al., 1998).

CONCLUSION AND PERSPECTIVES

The *B. subtilis* group offers a plethora of antagonistic compounds displaying a broad range of biological functions. This huge versatility increases the industrial and environmental interest of *B. subtilis* strains, especially when considering their range of action against foodborne or phytopathogenic flora as well as their history of safe use in food. The present review on known AMCs from the *B. subtilis* group proposes a consistent classification frame based on their biosynthetic pathways (*i.e.*, RPs, PKs, NRPs, volatiles) and chemical nature.

The present classification suggests to establish systematic approaches for novel molecules discoveries and characterizations (biosynthesis, chemical nature and activity). Indeed, most current publications report antimicrobial activity of partially purified fractions which can involve mixtures of bioactive compounds. To assess the activity of an unique compound, implementations of genetic confirmation such as knockout strategy are needed. Besides, very few studies have focused on the putative synergistic effects within these bio-active mixtures. Also, the concentration of purified or semi-purified compound(s) often remains uncharacterized or biologically irrelevant. Finally, there is no doubt that novel AMCs originating from *B. subtilis* bacteria remain to be identified, characterized and properly classified.

REFERENCES

- Abriouel, H., Franz, C. M., Omar, N. B., and Gálvez, A. (2011). Diversity and applications of *Bacillus* bacteriocins. *FEMS Microbiol. Rev.* 35, 201–232. doi: 10.1111/j.1574-6976.2010.00244.x
- Adak, S., Aulak, K. S., and Stuehr, D. J. (2002). Direct evidence for nitric oxide production by a nitric-oxide synthase-like protein from *Bacillus subtilis*. *J. Biol. Chem.* 277, 16167–16171. doi: 10.1074/jbc.M201136200
- Agrios, G. N. (1988). “1 - INTRODUCTION TO PLANT PATHOLOGY,” in *Plant Pathology*, 3rd Edition Edn, ed. G. N. Agrios (New York, NY: Academic Press), 3–39. doi: 10.1016/B978-0-12-044563-9.50005-0
- Alamri, S. A. (2015). Enhancing the efficiency of the bioagent *Bacillus subtilis* JF419701 against soil-borne phytopathogens by increasing the productivity of fungal cell wall degrading enzymes. *Arch. Phytopathol. Plant Prot.* 48, 159–170. doi: 10.1080/03235408.2014.884671
- Aleti, G., Sessitsch, A., and Brader, G. (2015). Genome mining: prediction of lipopeptides and polyketides from *Bacillus* and related Firmicutes. *Comput. Struct. Biotechnol. J.* 13, 192–203. doi: 10.1016/j.csbj.2015.03.003
- An, J., Zhu, W., Liu, Y., Zhang, X., Sun, L., Hong, P., et al. (2015). Purification and characterization of a novel bacteriocin CAMT2 produced by *Bacillus amyloquelificans* isolated from marine fish *Epinephelus areolatus*. *Food Control* 51, 278–282. doi: 10.1016/j.foodcont.2014.11.038

AUTHOR CONTRIBUTIONS

SC, CN, and FL conducted the bibliographic search. SC and CN wrote the manuscript. AG, CB, and JM edited and reviewed the manuscript. All authors have read and approved the final version.

FUNDING

This work was supported by the National Fund for Scientific Research (FNRS), the Université catholique de Louvain (UCLouvain), and the Brussels Institute for Research and Innovation (Innoviris, Doctiris programme to CN). SC was supported by the Foundation for Training in Industrial and Agricultural Research (FRIA, FNRS), AG holds a Chargé de Recherche fellowship from the FNRS (Grant 1.B.208.16F).

ACKNOWLEDGMENTS

We gratefully acknowledge members of SC's Ph.D. committee Prof A. Legrève, UCLouvain, Prof. M. Ongena, ULiège and Dr. J.-P. Goffart, CRAw for their valuable comments on this manuscript. We also acknowledge the Walloon Region for the long-term financial support through the WACOBI and ANTAGONIST projects (Conventions N° DGO3-D31-1330 and N° DGO3-D31-1383/S1, respectively).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2019.00302/full#supplementary-material>

- Arguelles Arias, A., Ongena, M., Devreese, B., Terrak, M., Joris, B., and Fickers, P. (2013). Characterization of amylolysin, a novel lantibiotic from *Bacillus amyloquelificans* GA1. *PLoS One* 8:e83037. doi: 10.1371/journal.pone.0083037
- Ariffin, H., Abdullah, N., Md Shah, U. K., Shirai, Y., and Hassan, M. A. (2006). Production and characterization of cellulases by *Bacillus pumilus* EB3. *Int. J. Eng. Technol.* 3, 47–53.
- Arrebola, E., Sivakumar, D., and Korsten, L. (2010). Effect of volatile compounds produced by *Bacillus* strains on postharvest decay in citrus. *Biol. Control* 53, 122–128. doi: 10.1016/j.biocontrol.2009.11.010
- Audrain, B., Farag, M. A., Ryu, C.-M., and Ghigo, J.-M. (2015). Role of bacterial volatile compounds in bacterial biology. *FEMS Microbiol. Rev.* 39, 222–233. doi: 10.1093/femsre/fuu013
- Bais, H. P., Fall, R., and Vivanco, J. M. (2004). Biocontrol of *Bacillus subtilis* against Infection of *Arabidopsis* Roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin Production. *Plant Physiol.* 134, 307–319. doi: 10.1104/pp.103.028712
- Barbe, V., Cruveiller, S., Kunst, F., Lenoble, P., Meurice, G., Sekowska, A., et al. (2009). From a consortium sequence to a unified sequence: the *Bacillus subtilis* 168 reference genome a decade later. *Microbiology* 155, 1758–1775. doi: 10.1099/mic.0.027839-0
- Béchet, M., Caradec, T., Hussein, W., Abderrahmani, A., Chollet, M., Leclère, V., et al. (2012). Structure, biosynthesis, and properties of kurstakins,

- nonribosomal lipopeptides from *Bacillus* spp. *Appl. Microbiol. Biotechnol.* 95, 593–600. doi: 10.1007/s00253-012-4181-2
- Begley, M., Cotter, P. D., Hill, C., and Ross, R. P. (2009). Identification of a novel two-peptide lantibiotic, lichenicidin, following rational genome mining for LanM proteins. *Appl. Environ. Microbiol.* 75, 5451–5460. doi: 10.1128/aem.00730-09
- Bentley, R., and Haslam, E. (1990). The shikimate pathway — a metabolic tree with many branches. *Crit. Rev. Biochem. Mol. Biol.* 25, 307–384. doi: 10.3109/10409239009090615
- Bernier, S. P., Létoffé, S., Delepierre, M., and Ghigo, J.-M. (2011). Biogenic ammonia modifies antibiotic resistance at a distance in physically separated bacteria. *Mol. Microbiol.* 81, 705–716. doi: 10.1111/j.1365-2958.2011.07724.x
- Besson, F., Peypoux, F., Michel, G., and Delcambe, L. (1976). Characterization of iturin A in antibiotics from various strains of *Bacillus subtilis*. *J. Antibiot.* 29, 1043–1049. doi: 10.7164/antibiotics.29.1043
- Besson, F., Peypoux, F., Michel, G., and Delcambe, L. (1979). Antifungal activity upon *Saccharomyces cerevisiae* of iturin A, mycosubtilin, bacillomycin L and of their derivatives; inhibition of this antifungal activity by lipid antagonists. *J. Antibiot.* 32, 828–833. doi: 10.7164/antibiotics.32.828
- Bowman, S. M., and Free, S. J. (2006). The structure and synthesis of the fungal cell wall. *Bioessays* 28, 799–808. doi: 10.1002/bies.20441
- Brötz, H., Bierbaum, G., Markus, A., Molitor, E., and Sahl, H. G. (1995). Mode of action of the lantibiotic mercadin: inhibition of peptidoglycan biosynthesis via a novel mechanism? *Antimicrob. Agents Chemother.* 39, 714–719. doi: 10.1128/aac.39.3.714
- Buckingham, J. (1997). *Dictionary of Natural Products*. London: CRC press. doi: 10.1007/978-1-4899-6850-0
- Caboche, S., Pupin, M., Leclère, V., Fontaine, A., Jacques, P., and Kucherov, G. (2008). NORINE: a database of nonribosomal peptides. *Nucleic Acids Res.* 36, D326–D331. doi: 10.1093/nar/gkm792
- Cane, D. E., and Walsh, C. T. (1999). The parallel and convergent universes of polyketide synthases and nonribosomal peptide synthetases. *Chem. Biol.* 6, R319–R325. doi: 10.1016/S1074-5521(00)80001-0
- Caulier, S., Gillis, A., Colau, G., Licciardi, F., Liépin, M., Desoignies, N., et al. (2018). Versatile antagonistic activities of soil-borne *Bacillus* spp. and *Pseudomonas* spp. against *Phytophthora infestans* and other potato pathogens. *Front. Microbiol.* 9:143. doi: 10.3389/fmicb.2018.00143
- Cawoy, H., Debois, D., Franzil, L., De Pauw, E., Thonart, P., and Ongena, M. (2015). Lipopeptides as main ingredients for inhibition of fungal phytopathogens by *Bacillus subtilis/amyloliquefaciens*. *Microb. Biotechnol.* 8, 281–295. doi: 10.1111/1751-7915.12238
- Challis, G. L. (2008). Genome mining for novel natural product discovery. *J. Med. Chem.* 51, 2618–2628. doi: 10.1021/jm700948z
- Challis, G. L., and Naismith, J. H. (2004). Structural aspects of non-ribosomal peptide biosynthesis. *Curr. Opin. Struct. Biol.* 14, 748–756. doi: 10.1016/j.sbi.2004.10.005
- Chatterjee, C., Paul, M., Xie, L., and van der Donk, W. A. (2005). Biosynthesis and mode of action of lantibiotics. *Chem. Rev.* 105, 633–684. doi: 10.1021/cr030105v
- Chaves-López, C., Serio, A., Gianotti, A., Sacchetti, G., Ndagijimana, M., Ciccarone, C., et al. (2015). Diversity of food-borne *Bacillus* volatile compounds and influence on fungal growth. *J. Appl. Microbiol.* 119, 487–499. doi: 10.1111/jam.12847
- Chen, H., and Du, L. (2016). Iterative polyketide biosynthesis by modular polyketide synthases in bacteria. *Appl. Microbiol. Biotechnol.* 100, 541–557. doi: 10.1007/s00253-015-7093-0
- Chen, H., Xiao, X., Wang, J., Wu, L., Zheng, Z., and Yu, Z. (2008). Antagonistic effects of volatiles generated by *Bacillus subtilis* on spore germination and hyphal growth of the plant pathogen, *Botrytis cinerea*. *Biotechnol. Lett.* 30, 919–923. doi: 10.1007/s10529-007-9626-9
- Chen, X., Scholz, R., Borriss, M., Junge, H., Mogel, G., Kunz, S., et al. (2009). Difficidin and bacilysin produced by plant-associated *Bacillus amyloliquefaciens* Dare efficient in controlling fire blight disease. *J. Biotechnol.* 140, 38–44. doi: 10.1016/j.jbiotec.2008.10.015
- Chen, X., Vater, J., Piel, J., Franke, P., Scholz, R., Schneider, K., et al. (2006). Structural and functional characterization of three polyketide synthase gene clusters in *Bacillus amyloliquefaciens* FZB42. *J. Bacteriol.* 188, 4024–4036. doi: 10.1128/jb.00052-06
- Cherif-Silini, H., Silini, A., Yahiaoui, B., Ouzari, I., and Boudabous, A. (2016). Phylogenetic and plant-growth-promoting characteristics of *Bacillus* isolated from the wheat rhizosphere. *Ann. Microbiol.* 66, 1087–1097. doi: 10.1007/s13213-016-1194-6
- Chernin, L., Toklikishvili, N., Ovadis, M., Kim, S., Ben-Ari, J., Khmel, I., et al. (2011). Quorum-sensing quenching by rhizobacterial volatiles. *Environ. Microbiol. Rep.* 3, 698–704. doi: 10.1111/j.1758-2229.2011.00284.x
- Chopra, L., Singh, G., Choudhary, V., and Sahoo, D. K. (2014). Sonorensin: an antimicrobial peptide, belonging to the heterocycloanthracin subfamily of bacteriocins, from a new marine isolate, *Bacillus sonorensis* MT93. *Appl. Environ. Microbiol.* 80, 2981–2990. doi: 10.1128/aem.04259-13
- Chopra, L., Singh, G., Jena, K. K., Verma, H., and Sahoo, D. K. (2015). Bioprocess development for the production of sonorensin by *Bacillus sonorensis* MT93 and its application as a food preservative. *Bioresour. Technol.* 175, 358–366. doi: 10.1016/j.biortech.2014.10.105
- Cleveland, J., Montville, T. J., Nes, I. F., and Chikindas, M. L. (2001). Bacteriocins: safe, natural antimicrobials for food preservation. *Int. J. Food Microbiol.* 71, 1–20. doi: 10.1016/S0168-1605(01)00560-8
- Coosemans, J. (2005). Dimethyl disulphide (DMS): a potential novel nematicide and soil disinfectant. *Acta Hort.* 698, 57–64. doi: 10.17660/ActaHortic.2005.698.6
- Cosby, W. M., Vollenbroich, D., Lee, O. H., and Zuber, P. (1998). Altered *srf* expression in *Bacillus subtilis* resulting from changes in culture pH is dependent on the Spo0K oligopeptide permease and the ComQX system of extracellular control. *J. Bacteriol.* 180, 1438–1445.
- Cotter, P. D., Hill, C., and Ross, R. P. (2005). Bacteriocins: developing innate immunity for food. *Nat. Rev. Microbiol.* 3, 777–788. doi: 10.1038/nrmicro1273
- Coutts, R. T., Pitkethly, W. N., and Wibberley, D. G. (1965). Antibacterial activity of some quinolines containing a cyclic hydroxamic acid group. *J. Pharm. Sci.* 54, 792–795. doi: 10.1002/jps.2600540530
- Cox, S. D., Mann, C. M., Markham, J. L., Bell, H. C., Gustafson, J. E., Warmington, J. R., et al. (2000). The mode of antimicrobial action of the essential oil of *Melaleuca alternifolia* (tea tree oil). *J. Appl. Microbiol.* 88, 170–175. doi: 10.1046/j.1365-2672.2000.00943.x
- Crane, J. M., Gibson, D. M., Vaughan, R. H., and Bergstrom, G. C. (2012). Iturin levels on wheat spikes linked to biological control of fusarium head blight by *Bacillus amyloliquefaciens*. *Phytopathology* 103, 146–155. doi: 10.1094/PHYTO-07-12-0154-R
- Czajkowski, R., and Jafra, S. (2009). Quenching of acyl-homoserine lactone-dependent quorum sensing by enzymatic disruption of signal molecules. *Acta Biochim. Pol.* 56, 1–16.
- Dambolena, J. S., López, A. G., Cánepa, M. C., Theumer, M. G., Zygodlo, J. A., and Rubinstein, H. R. (2008). Inhibitory effect of cyclic terpenes (limonene, menthol, menthone and thymol) on *Fusarium verticillioides* MRC 826 growth and fumonisin B1 biosynthesis. *Toxicol.* 51, 37–44. doi: 10.1016/j.toxicol.2007.07.005
- de Vrieze, M., Pandey, P., Bucheli, T. D., Varadarajan, A. R., Ahrens, C. H., Weisskopf, L., et al. (2015). Volatile organic compounds from native potato-associated *Pseudomonas* as potential anti-oomycete agents. *Front. Microbiol.* 6:1295. doi: 10.3389/fmicb.2015.01295
- Delves-Broughton, J. (1990). Nisin and its application as a food preservative. *Int. J. Dairy Technol.* 43, 73–76. doi: 10.1111/j.1471-0307.1990.tb02449.x
- Dertz, E. A., Xu, J., Stintzi, A., and Raymond, K. N. (2006). Bacillibactin-mediated iron transport in *Bacillus subtilis*. *J. Am. Chem. Soc.* 128, 22–23. doi: 10.1021/ja055898c
- Dickschat, J. S., Bode, H. B., Wenzel, S. C., Müller, R., and Schulz, S. (2005). Biosynthesis and identification of volatiles released by the myxobacterium *Stigmatella aurantiaca*. *Chembiochem* 6, 2023–2033. doi: 10.1002/cbic.20050174
- Dimkić, I., Živković, S., Berić, T., Ivanović, Ž, Gavrilović, V., Stanković, S., et al. (2013). Characterization and evaluation of two *Bacillus* strains, SS-12.6 and SS-13.1, as potential agents for the control of phytopathogenic bacteria and fungi. *Biol. Control* 65, 312–321. doi: 10.1016/j.biocontrol.2013.03.012
- Dong, Y.-H., Zhang, X.-F., Xu, J.-L., and Zhang, L.-H. (2004). Insecticidal *Bacillus thuringiensis* silences *Erwinia carotovora* virulence by a new form of microbial antagonism, signal interference. *Appl. Environ. Microbiol.* 70, 954–960. doi: 10.1128/aem.70.2.954-960.2004

- Dubois, J. Y. F., Kouwen, T. R., Schurich, A. K. C., Reis, C. R., Ensing, H. T., Trip, E. N., et al. (2009). Immunity to the bacteriocin sublancin 168 is determined by the SunI (YolF) protein of *Bacillus subtilis*. *Antimicrob. Agents Chemother.* 53, 651–661. doi: 10.1128/aac.01189-08
- Dunlap, C. A., Schisler, D. A., Price, N. P., and Vaughn, S. F. (2011). Cyclic lipopeptide profile of three *Bacillus subtilis* strains; antagonists of *Fusarium* head blight. *J. Microbiol.* 49, 603–609. doi: 10.1007/s12275-011-1044-y
- Effmert, U., Kalderás, J., Warnke, R., and Piechulla, B. (2012). Volatile mediated interactions between bacteria and fungi in the soil. *J. Chem. Ecol.* 38, 665–703. doi: 10.1007/s10886-012-0135-5
- Eshita, S. M., Roberto, N. H., Beale, J. M., Mamiya, B. M., and Workman, R. F. (1995). Bacillomycin Lc, a new antibiotic of the iturin group: isolations, structures, and antifungal activities of the congeners. *J. Antibiot.* 48, 1240–1247. doi: 10.7164/antibiotics.48.1240
- Eustáquio, A. S., McGlinchey, R. P., Liu, Y., Hazzard, C., Beer, L. L., Florova, G., et al. (2009). Biosynthesis of the salinosporamide A polyketide synthase substrate chloroethylmalonyl-coenzyme A from S-adenosyl-L-methionine. *Proc. Natl. Acad. Sci. U.S.A.* 106, 12295–12300. doi: 10.1073/pnas.0901237106
- Even, S., Burguière, P., Auger, S., Soutourina, O., Danchin, A., and Martin-Verstraete, I. (2006). Global control of cysteine metabolism by CymR in *Bacillus subtilis*. *J. Bacteriol.* 188, 2184–2197. doi: 10.1128/jb.188.6.2184-2197.2006
- Fan, B., Blom, J., Klenk, H.-P., and Borriss, R. (2017). *Bacillus amyloliquefaciens*, *Bacillus velezensis*, and *Bacillus siamensis* form an “operational group *B. amyloliquefaciens*” within the *B. subtilis* species complex. *Front. Microbiol.* 8:22. doi: 10.3389/fmicb.2017.00022
- Fickers, P., Guez, J.-S., Damblon, C., Leclère, V., Béchet, M., Jacques, P., et al. (2009). High-level biosynthesis of the anteiso-C17 isoform of the antibiotic mycosubtilin in *Bacillus subtilis* and characterization of its candidacidal activity. *Appl. Environ. Microbiol.* 75, 4636–4640. doi: 10.1128/aem.00548-09
- Fisch, K. M. (2013). Biosynthesis of natural products by microbial iterative hybrid PKS-NRPS. *RSC Adv.* 3, 18228–18247. doi: 10.1039/C3RA42661K
- Fisher, A. J., Rosenstiel, T. N., Shirk, M. C., and Fall, R. (2001). Nonradioactive assay for cellular dimethylallyl diphosphate. *Anal. Biochem.* 292, 272–279. doi: 10.1006/abio.2001.5079
- Fu, L.-H., Hu, K.-D., Hu, L.-Y., Li, Y.-H., Hu, L.-B., Yan, H., et al. (2014). An antifungal role of hydrogen sulfide on the postharvest pathogens *Aspergillus niger* and *Penicillium italicum*. *PLoS One* 9:e104206. doi: 10.1371/journal.pone.0104206
- Fuchs, S. W., Jaskolla, T. W., Bochmann, S., Kötter, P., Wichelhaus, T., Karas, M., et al. (2011). Entianin, a novel subtilin-like lantibiotic from *Bacillus subtilis* DSM 15029T with high antimicrobial activity. *Appl. Environ. Microbiol.* 77, 1698–1707. doi: 10.1128/aem.01962-10
- Furuta, Y., Harima, H., Ito, E., Maruyama, F., Ohnishi, N., Osaki, K., et al. (2018). Loss of bacitracin resistance due to a large genomic deletion among *Bacillus anthracis* strains. *mSystems* 3:e00182-18. doi: 10.1128/mSystems.00182-18
- Gao, L., Han, J., Liu, H., Qu, X., Lu, Z., and Bie, X. (2017). Plipastatin and surfactin coproduction by *Bacillus subtilis* pB2-L and their effects on microorganisms. *Antonie Van Leeuwenhoek* 110, 1007–1018. doi: 10.1007/s10482-017-0874-y
- Gao, Z., Zhang, B., Liu, H., Han, J., and Zhang, Y. (2017). Identification of endophytic *Bacillus velezensis* ZSY-1 strain and antifungal activity of its volatile compounds against *Alternaria solani* and *Botrytis cinerea*. *Biol. Control* 105, 27–39. doi: 10.1016/j.biocontrol.2016.11.007
- Gautam, N., and Sharma, N. (2009). Bacteriocin: safest approach to preserve food products. *Indian J. Microbiol.* 49, 204–211. doi: 10.1007/s12088-009-0048-3
- Geraldine, A. M., Lopes, F. A. C., Carvalho, D. D. C., Barbosa, E. T., Rodrigues, A. R., Brandão, R. S., et al. (2013). Cell wall-degrading enzymes and parasitism of sclerotia are key factors on field biocontrol of white mold by *Trichoderma* spp. *Biol. Control* 67, 308–316. doi: 10.1016/j.biocontrol.2013.09.013
- Gerik, J. S. (2005). Evaluation of soil fumigants applied by drip irrigation for liatris production. *Plant Dis.* 89, 883–887. doi: 10.1094/PD-89-0883
- Gershenson, J., and Dudareva, N. (2007). The function of terpene natural products in the natural world. *Nat. Chem. Biol.* 3, 408–414. doi: 10.1038/nchembio.2007.5
- Giorgio, A., De Stradis, A., Lo Cantore, P., and Iacobellis, N. S. (2015). Biocide effects of volatile organic compounds produced by potential biocontrol rhizobacteria on *Sclerotinia sclerotiorum*. *Front. Microbiol.* 6:1056. doi: 10.3389/fmicb.2015.01056
- Gomaa, E. Z. (2012). Chitinase production by *Bacillus thuringiensis* and *Bacillus licheniformis*: their potential in antifungal biocontrol. *J. Microbiol.* 50, 103–111. doi: 10.1007/s12275-012-1343-y
- Gong, A.-D., Li, H.-P., Yuan, Q.-S., Song, X.-S., Yao, W., He, W.-J., et al. (2015). Antagonistic mechanism of iturin A and plipastatin A from *Bacillus amyloliquefaciens* S76-3 from wheat spikes against *Fusarium graminearum*. *PLoS One* 10:e0116871. doi: 10.1371/journal.pone.0116871
- Gong, Q., Zhang, C., Lu, F., Zhao, H., Bie, X., and Lu, Z. (2014). Identification of bacillomycin D from *Bacillus subtilis* fmbj and its inhibition effects against *Aspergillus flavus*. *Food Control* 36, 8–14. doi: 10.1016/j.foodcont.2013.07.034
- González, J. E., and Keshavan, N. D. (2006). Messing with bacterial quorum sensing. *Microbiol. Mol. Biol. Rev.* 70, 859–875. doi: 10.1128/mmbr.00002-06
- Gordon, R., Haynes, W., Pang, C., and Smith, N. (1973). *The Genus Bacillus*. Washington, DC: United States Department of Agriculture, 109–126.
- Gotor-Vila, A., Teixidó, N., Di Francesco, A., Usall, J., Ugolini, L., Torres, R., et al. (2017). Antifungal effect of volatile organic compounds produced by *Bacillus amyloliquefaciens* CPA-8 against fruit pathogen decays of cherry. *Food Microbiol.* 64, 219–225. doi: 10.1016/j.fm.2017.01.006
- Gu, Y.-Q., Mo, M.-H., Zhou, J.-P., Zou, C.-S., and Zhang, K.-Q. (2007). Evaluation and identification of potential organic nematicidal volatiles from soil bacteria. *Soil Biol. Biochem.* 39, 2567–2575. doi: 10.1016/j.soilbio.2007.05.011
- Guan, Z., Xue, D., Abdallah, I. I., Dijkshoorn, L., Setroikromo, R., Lv, G., et al. (2015). Metabolic engineering of *Bacillus subtilis* for terpenoid production. *Appl. Microbiol. Biotechnol.* 99, 9395–9406. doi: 10.1007/s00253-015-6950-1
- Guder, A., Wiedemann, I., and Sahl, H.-G. (2000). Posttranslationally modified bacteriocins—the lantibiotics. *Biopolymers* 55, 62–73. doi: 10.1002/1097-0282(2000)55:1<62::AID-BIP60>3.0.CO;2-Y
- Guo, Q., Dong, W., Li, S., Lu, X., Wang, P., Zhang, X., et al. (2014). Fengycin produced by *Bacillus subtilis* NCD-2 plays a major role in biocontrol of cotton seedling damping-off disease. *Microbiol. Res.* 169, 533–540. doi: 10.1016/j.micres.2013.12.001
- Gustafson, K., Roman, M., and Fenical, W. (1989). The macrolactin, a novel class of antiviral and cytotoxic macrolides from a deep-sea marine bacterium. *J. Am. Chem. Soc.* 111, 7519–7524. doi: 10.1021/ja00201a036
- Haidar, R., Roudet, J., Bonnard, O., Dufour, M. C., Corio-Costet, M. F., Fert, M., et al. (2016). Screening and modes of action of antagonistic bacteria to control the fungal pathogen *Phaeoemoniella chlamydospora* involved in grapevine trunk diseases. *Microbiol. Res.* 192, 172–184. doi: 10.1016/j.micres.2016.07.003
- Hammami, I., Jaouadi, B., Bacha, A. B., Rebai, A., Bejar, S., Nesme, X., et al. (2012). *Bacillus subtilis* bacteriocin Bac 14B with a broad inhibitory spectrum: purification, amino acid sequence analysis, and physicochemical characterization. *Biotechnol. Bioprocess Eng.* 17, 41–49. doi: 10.1007/s12257-010-0401-8
- Hammer, K. A., Carson, C. F., and Riley, T. V. (2003). Antifungal activity of the components of *Melaleuca alternifolia* (tea tree) oil. *J. Appl. Microbiol.* 95, 853–860. doi: 10.1046/j.1365-2672.2003.02059.x
- Hamoen, L. W., Venema, G., and Kuipers, O. P. (2003). Controlling competence in *Bacillus subtilis*: shared use of regulators. *Microbiology* 149, 9–17. doi: 10.1099/mic.0.26003-0
- Heinzmann, S., Entian, K.-D., and Stein, T. (2006). Engineering *Bacillus subtilis* ATCC 6633 for improved production of the lantibiotic subtilin. *Appl. Microbiol. Biotechnol.* 69, 532–536. doi: 10.1007/s00253-005-0023-9
- Herrera-Estrella, A., and Chet, I. (1999). Chitinases in biological control. *EXS* 87, 171–184. doi: 10.1007/978-3-0348-8757-1_12
- Hertweck, C. (2009). The biosynthetic logic of polyketide diversity. *Angew. Chem. Int. Ed. Engl.* 48, 4688–4716. doi: 10.1002/anie.200806121
- Horsman, G. P., Van Lanen, S. G., and Shen, B. (2009). Chapter 5 Iterative type I polyketide synthases for enediene core biosynthesis. *Methods Enzymol.* 459, 97–112. doi: 10.1016/S0076-6879(09)04605-9
- Howell, C., Beier, R., and Stipanovic, R. (1988). Production of ammonia by *Enterobacter cloacae* and its possible role in the biological control of *Pythium* preemergence damping-off by the bacterium. *Phytopathology* 78, 1075–1078. doi: 10.1094/Phyto-78-1075
- Hsieh, F.-C., Lin, T.-C., Meng, M., and Kao, S.-S. (2008). Comparing methods for identifying *Bacillus* strains capable of producing the antifungal lipopeptide iturin A. *Curr. Microbiol.* 56, 1–5. doi: 10.1007/s00284-007-9003-x
- Hussein, A., and Al-Janabi, S. (2006). Identification of bacitracin produced by local isolate of *Bacillus licheniformis*. *Afr. J. Biotechnol.* 5, 1600–1601.

- Inoue, Y., Shiraishi, A., Hada, T., Hirose, K., Hamashima, H., and Shimada, J. (2004). The antibacterial effects of terpene alcohols on *Staphylococcus aureus* and their mode of action. *FEMS Microbiol. Lett.* 237, 325–331. doi: 10.1111/j.1574-6968.2004.tb09714.x
- Insam, H., and Seewald, M. S. A. (2010). Volatile organic compounds (VOCs) in soils. *Biol. Fertil. Soils* 46, 199–213. doi: 10.1007/s00374-010-0442-3
- Jacques, P. (2011). “Surfactin and other lipopeptides from *Bacillus* spp.,” in *Biosurfactants: From Genes to Applications*, ed. G. Soberón-Chávez (Berlin: Springer), 57–91. doi: 10.1007/978-3-642-14490-5_3
- Janiak, A. M., and Milewski, S. (2001). Mechanism of antifungal action of kanosamine. *Med. Mycol.* 39, 401–408. doi: 10.1080/mmy.39.5.401.408
- Jaruchoktaweetchai, C., Suwanborirux, K., Tanasupawatt, S., Kittakoop, P., and Menasveta, P. (2000). New Macrolactins from a Marine *Bacillus* sp. Sc026. *J. Nat. Prod.* 63, 984–986. doi: 10.1021/np990605c
- Jenny, K., Käppeli, O., and Fiechter, A. (1991). Biosurfactants from *Bacillus licheniformis*: structural analysis and characterization. *Appl. Microbiol. Biotechnol.* 36, 5–13. doi: 10.1007/bf00164690
- Johnson, B. A., Anker, H., and Meloney, F. L. (1945). Bacitracin: a new antibiotic produced by a member of the *B. subtilis* group. *Science* 102, 376–377. doi: 10.1126/science.102.2650.376
- Julsing, M. K., Rijpkema, M., Woerdenbag, H. J., Quax, W. J., and Kayser, O. (2007). Functional analysis of genes involved in the biosynthesis of isoprene in *Bacillus subtilis*. *Appl. Microbiol. Biotechnol.* 75, 1377–1384. doi: 10.1007/s00253-007-0953-5
- Justicia, J., Oltra, J. E., Barrero, A. F., Guadaño, A., González-Coloma, A., and Cuerva, J. M. (2005). Total synthesis of 3-hydroxydrimanes mediated by titanocene(III) – evaluation of their antifeedant activity. *Eur. J. Org. Chem.* 2005, 712–718. doi: 10.1002/ejoc.200400634
- Kai, M., Hausteiner, M., Molina, F., Petri, A., Scholz, B., and Piechulla, B. (2009). Bacterial volatiles and their action potential. *Appl. Microbiol. Biotechnol.* 81, 1001–1012. doi: 10.1007/s00253-008-1760-3
- Kawulka, K. E., Sprules, T., Diaper, C. M., Whittall, R. M., McKay, R. T., Mercier, P., et al. (2004). Structure of subtilosin A, a cyclic antimicrobial peptide from *Bacillus subtilis* with unusual sulfur to (-carbon cross-links: formation and reduction of (-Thio(-Amino Acid derivatives. *Biochemistry* 43, 3385–3395. doi: 10.1021/bi0359527
- Kenig, M., and Abraham, E. P. (1976). Antimicrobial activities and antagonists of bacilysin and anticapsin. *Microbiology* 94, 37–45. doi: 10.1099/00221287-94-1-37
- Kenig, M., Vandamme, E., and Abraham, E. P. (1976). The mode of action of bacilysin and anticapsin and biochemical properties of bacilysin-resistant mutants. *Microbiology* 94, 46–54. doi: 10.1099/00221287-94-1-46
- Kim, H.-S., Yoon, B.-D., Lee, C.-H., Suh, H.-H., Oh, H.-M., Katsuragi, T., et al. (1997). Production and properties of a lipopeptide biosurfactant from *Bacillus subtilis* C9. *J. Ferment. Bioeng.* 84, 41–46. doi: 10.1016/S0922-338X(97)82784-5
- Kim, P. I., Ryu, J., Kim, Y. H., and Chi, Y.-T. (2010). Production of biosurfactant lipopeptides iturin A, fengycin and surfactin A from *Bacillus subtilis* CMB32 for control of *Colletotrichum gloeosporioides*. *J. Microbiol. Biotechnol.* 20, 138–145. doi: 10.4014/jmb.0905.05007
- Klaenhammer, T. R. (1988). Bacteriocins of lactic acid bacteria. *Biochimie* 70, 337–349. doi: 10.1016/0300-9084(88)90206-4
- Kleerebezem, M. (2004). Quorum sensing control of lantibiotic production; nisin and subtilin autoregulate their own biosynthesis. *Peptides* 25, 1405–1414. doi: 10.1016/j.peptides.2003.10.021
- Konz, D., Klens, A., Schörgendorfer, K., and Marahiel, M. A. (1997). The bacitracin biosynthesis operon of *Bacillus licheniformis* ATCC 10716: molecular characterization of three multi-modular peptide synthetases. *Chem. Biol.* 4, 927–937. doi: 10.1016/S1074-5521(97)90301-X
- Kopp, F., and Marahiel, M. A. (2007). Macrocyclization strategies in polyketide and nonribosomal peptide biosynthesis. *Nat. Prod. Rep.* 24, 735–749. doi: 10.1039/B613652B
- Korpi, A., Järnberg, J., and Pasanen, A.-L. (2009). Microbial volatile organic compounds. *Crit. Rev. Toxicol.* 39, 139–193. doi: 10.1080/10408440802291497
- Kosuge, T., and Kamiya, H. (1962). Discovery of a pyrazine in a natural product: tetramethylpyrazine from cultures of a strain of *Bacillus subtilis*. *Nature* 193:776. doi: 10.1038/193776a0
- Kugler, M., Loeffler, W., Rapp, C., Kern, A., and Jung, G. (1990). Rhizoctin A, an antifungal phosphono-oligopeptide of *Bacillus subtilis* ATCC 6633: biological properties. *Arch. Microbiol.* 153, 276–281. doi: 10.1007/bf00249082
- Kumar, P., Dubey, R. C., and Maheshwari, D. K. (2012). *Bacillus* strains isolated from rhizosphere showed plant growth promoting and antagonistic activity against phytopathogens. *Microbiol. Res.* 167, 493–499. doi: 10.1016/j.micres.2012.05.002
- Kwon, J. W., and Kim, S. D. (2014). Characterization of an antibiotic produced by *Bacillus subtilis* JW-1 that suppresses *Ralstonia solanacearum*. *J. Microbiol. Biotechnol.* 24, 13–18. doi: 10.4014/jmb.1308.08060
- Landy, M., Warren, G. H., Rosenman, S. B., and Colio, L. G. (1948). Bacillomycin: an antibiotic from *Bacillus subtilis* active against pathogenic fungi. *Proc. Soc. Exp. Biol. Med.* 67, 539–541. doi: 10.3181/00379727-67-16367
- Larroche, C., Besson, I., and Gros, J.-B. (1999). High pyrazine production by *Bacillus subtilis* in solid substrate fermentation on ground soybeans. *Process Biochem.* 34, 667–674. doi: 10.1016/S0032-9592(98)00141-1
- Leclere, V., Bechet, M., Adam, A., Guez, J. S., Wathélet, B., Ongena, M., et al. (2005). Mycosubtilin overproduction by *Bacillus subtilis* BBG100 enhances the organism's antagonistic and biocontrol activities. *Appl. Environ. Microbiol.* 71, 4577–4584. doi: 10.1128/aem.71.8.4577-4584.2005
- Lee, D. W., and Kim, B. S. (2015). Antimicrobial cyclic peptides for plant disease control. *Plant Pathol. J.* 31, 1–11. doi: 10.5423/PPJ.RW.08.2014.0074
- Lee, H., Churey, J. J., and Worobo, R. W. (2008). Purification and structural characterization of bacillomycin F produced by a bacterial honey isolate active against *Byssoschlamys fulva* H25. *J. Appl. Microbiol.* 105, 663–673. doi: 10.1111/j.1365-2672.2008.03797.x
- Lee, S., Peterson, C. J., and Coats, J. R. (2003). Fumigation toxicity of monoterpenoids to several stored product insects. *J. Stored Prod. Res.* 39, 77–85. doi: 10.1016/S0022-474X(02)00020-6
- Lee, S.-C., Kim, S.-H., Park, I.-H., Chung, S.-Y., and Choi, Y.-L. (2007). Isolation and structural analysis of bamylocin A, novel lipopeptide from *Bacillus amyloliquefaciens* LP03 having antagonistic and crude oil-emulsifying activity. *Arch. Microbiol.* 188, 307–312. doi: 10.1007/s00203-007-0250-9
- Leejeerajumnean, A., Duckham, S. C., Owens, J. D., and Ames, J. M. (2001). Volatile compounds in *Bacillus*-fermented soybeans. *J. Sci. Food Agric.* 81, 525–529. doi: 10.1002/jsfa.843
- Lemfack, M. C., Gohlke, B.-O., Toguem, S. M. T., Preissner, S., Piechulla, B., and Preissner, R. (2018). mVOC 2.0: a database of microbial volatiles. *Nucleic Acids Res.* 46, D1261–D1265. doi: 10.1093/nar/gkx1016
- Li, L., Ma, M., Huang, R., Qu, Q., Li, G., Zhou, J., et al. (2012). Induction of chlamydospore formation in *Fusarium* by cyclic lipopeptide antibiotics from *Bacillus subtilis* C2. *J. Chem. Ecol.* 38, 966–974. doi: 10.1007/s10886-012-0171-1
- Li, N., Shi, Z., Tang, Y., Chen, J., and Li, X. (2008). Recent progress on the total synthesis of acetogenins from Annonaceae. *Beilstein J. Org. Chem.* 4:48. doi: 10.3762/bjoc.4.48
- Linnett, P. E., and Strominger, J. L. (1973). Additional antibiotic inhibitors of peptidoglycan synthesis. *Antimicrob. Agents Chemother.* 4, 231–236. doi: 10.1128/aac.4.3.231
- Liu, W. W., Mu, W., Zhu, B. Y., Du, Y. C., and Liu, F. (2008). Antagonistic activities of volatiles from four strains of *Bacillus* spp. and *Paenibacillus* spp. against soil-borne plant pathogens. *Agric. Sci. China* 7, 1104–1114. doi: 10.1016/S1671-2927(08)60153-4
- Liu, Y., Chen, Z., Ng, T. B., Zhang, J., Zhou, M., Song, F., et al. (2007). Bacisubin, an antifungal protein with ribonuclease and hemagglutinating activities from *Bacillus subtilis* strain B-916. *Peptides* 28, 553–559. doi: 10.1016/j.peptides.2006.10.009
- Loeffler, W., Tschen, J. S.-M., Vaniitanakom, N., Kugler, M., Knorpp, E., Hsieh, T. F., et al. (1986). Antifungal effects of bacilysin and fengycin from *Bacillus subtilis* F-29-3 a comparison with activities of other *Bacillus* antibiotics. *J. Phytopathol.* 115, 204–213. doi: 10.1111/j.1439-0434.1986.tb00878.x
- Loiseau, C., Schlusshelhuber, M., Bigot, R., Bertaux, J., Berjeaud, J.-M., and Verdon, J. (2015). Surfactin from *Bacillus subtilis* displays an unexpected anti-*Legionella* activity. *Appl. Microbiol. Biotechnol.* 99, 5083–5093. doi: 10.1007/s00253-014-6317-z
- Lu, X. L., Xu, Q. Z., Shen, Y. H., Liu, X. Y., Jiao, B. H., Zhang, W. D. et al. (2008). Macrolactin S, a novel macrolactin antibiotic from marine *Bacillus* sp. *Nat. Prod. Res.* 22, 342–347. doi: 10.1080/14786410701768162
- Luo, C., Liu, X., Zhou, X., Guo, J., Truong, J., Wang, X., et al. (2015). Unusual biosynthesis and structure of locillomycins from *Bacillus*

- subtilis* 916. *Appl. Environ. Microbiol.* 81, 6601–6609. doi: 10.1128/aem.01639-15
- Lupetti, A., Danesi, R., Campa, M., Tacca, M. D., and Kelly, S. (2002). Molecular basis of resistance to azole antifungals. *Trends Mol. Med.* 8, 76–81. doi: 10.1016/S1471-4914(02)02280-3
- Ma, Z., Hu, J., Wang, X., and Wang, S. (2013). NMR spectroscopic and MS/MS spectrometric characterization of a new lipopeptide antibiotic bacillopeptin B1 produced by a marine sediment-derived *Bacillus amyloliquefaciens* SH-B74. *J. Antibiot.* 67, 175–178. doi: 10.1038/ja.2013.89
- Mahlstedt, S. A., and Walsh, C. T. (2010). Investigation of anticapsin biosynthesis reveals a four-enzyme pathway to tetrahydrotyrosine in *Bacillus subtilis*. *Biochemistry* 49, 912–923. doi: 10.1021/bi9021186
- Majumdar, S. K., and Bose, S. K. (1958). Mycobaccillin, a new antifungal antibiotic produced by *B. subtilis*. *Nature* 181, 134–135. doi: 10.1038/181134a0
- Malfanova, N., Franzil, L., Lugtenberg, B., Chebotar, V., and Ongena, M. (2012). Cyclic lipopeptide profile of the plant-beneficial endophytic bacterium *Bacillus subtilis* HC8. *Arch. Microbiol.* 194, 893–899. doi: 10.1007/s00203-012-0823-0
- Marquez-Villavicencio, M. D. P., Weber, B., Witherell, R. A., Willis, D. K., and Charkowski, A. O. (2011). The 3-Hydroxy-2-Butanone pathway is required for *Pectobacterium carotovorum* pathogenesis. *PLoS One* 6:e22974. doi: 10.1371/journal.pone.0022974
- Marx, R., Stein, T., Entian, K.-D., and Glaser, S. J. (2001). Structure of the *Bacillus subtilis* peptide antibiotic subtilisin A determined by 1H-NMR and matrix assisted laser desorption/ionization time-of-flight mass spectrometry. *J. Protein Chem.* 20, 501–506. doi: 10.1023/a:1012562631268
- May, J. J., Wendrich, T. M., and Marahiel, M. A. (2001). The *dhb* operon of *Bacillus subtilis* encodes the biosynthetic template for the catecholic siderophore 2,3-Dihydroxybenzoate-Glycine-Threonine trimeric ester bacillibactin. *J. Biol. Chem.* 276, 7209–7217. doi: 10.1074/jbc.M009140200
- McAuliffe, O., Ross, R. P., and Hill, C. (2001). Lantibiotics: structure, biosynthesis and mode of action. *FEMS Microbiol. Rev.* 25, 285–308. doi: 10.1111/j.1574-6976.2001.tb00579.x
- McIntosh, J. A., Donia, M. S., and Schmidt, E. W. (2009). Ribosomal peptide natural products: bridging the ribosomal and nonribosomal worlds. *Nat. Prod. Rep.* 26, 537–559. doi: 10.1039/B714132G
- McNeal, K. S., and Herbert, B. E. (2009). Volatile organic metabolites as indicators of soil microbial activity and community composition shifts. *Soil Sci. Soc. Am. J.* 73, 579–588. doi: 10.2136/sssaj2007.0245
- Mhammedi, A., Peypoux, F., Besson, F., and Michel, G. (1982). Bacillomycin F, a new antibiotic of iturin group: isolation and characterization. *J. Antibiot.* 35, 306–311. doi: 10.7164/antibiotics.35.306
- Milewski, S., Chmara, H., and Borowski, E. (1986). Antibiotic tetaine — a selective inhibitor of chitin and mannoprotein biosynthesis in *Candida albicans*. *Arch. Microbiol.* 145, 234–240. doi: 10.1007/bf00443651
- Mojid Mondol, M. A., Kim, J. H., Lee, H.-S., Lee, Y.-J., and Shin, H. J. (2011). Macrolactin W, a new antibacterial macrolide from a marine *Bacillus* sp. *Bioorg. Med. Chem. Lett.* 21, 3832–3835. doi: 10.1016/j.bmcl.2010.12.050
- Moldenhauer, J., Chen, X.-H., Borriss, R., and Piel, J. (2007). Biosynthesis of the antibiotic bacillaene, the product of a giant polyketide synthase complex of the trans-AT family. *Angew. Chem. Int. Ed. Engl.* 46, 8195–8197. doi: 10.1002/anie.200703386
- Moyno, A.-L., Shelby, R., Cleveland, T. E., and Tuzun, S. (2001). Bacillomycin D: an iturin with antifungal activity against *Aspergillus flavus*. *J. Appl. Microbiol.* 90, 622–629. doi: 10.1046/j.1365-2672.2001.01290.x
- Müller, S., Strack, S. N., Hoefler, B. C., Straight, P., Kearns, D. B., and Kirby, J. R. (2014). Bacillaene and sporulation protect *Bacillus subtilis* from predation by *Myxococcus xanthus*. *Appl. Environ. Microbiol.* 80, 5603–5610. doi: 10.1128/aem.01621-14
- Mulligan, C. N., Yong, R. N., and Gibbs, B. F. (2001). Heavy metal removal from sediments by biosurfactants. *J. Hazard. Mater.* 85, 111–125. doi: 10.1016/S0304-3894(01)00224-2
- Nagao, T., Adachi, K., Sakai, M., Nishijima, M., and Sano, H. (2001). Novel macrolactins as antibiotic lactones from a marine bacterium. *J. Antibiot.* 54, 333–339. doi: 10.7164/antibiotics.54.333
- Naruse, N., Tenmyo, O., Kobaru, S., Kamei, H., Miyaki, T., Konishi, M., et al. (1990). Pumilacidin, a complex of new antiviral antibiotics. *J. Antibiot.* 43, 267–280. doi: 10.7164/antibiotics.43.267
- Nes, I. F., Diep, D. B., and Holo, H. (2007). Bacteriocin diversity in *Streptococcus* and *Enterococcus*. *J. Bacteriol.* 189, 1189–1198. doi: 10.1128/jb.01254-06
- Nicholson, W. L. (2002). Roles of *Bacillus* endospores in the environment. *Cell. Mol. Life Sci.* 59, 410–416. doi: 10.1007/s00018-002-8433-7
- Oman, T. J., and van der Donk, W. A. (2009). Follow the leader: the use of leader peptides to guide natural product biosynthesis. *Nat. Chem. Biol.* 6, 9–18. doi: 10.1038/nchembio.286
- Ongena, M., Duby, F., Jourdan, E., Beaudry, T., Jadin, V., Dommes, J., et al. (2005). *Bacillus subtilis* M4 decreases plant susceptibility towards fungal pathogens by increasing host resistance associated with differential gene expression. *Appl. Microbiol. Biotechnol.* 67, 692–698. doi: 10.1007/s00253-004-1741-0
- Ongena, M., and Jacques, P. (2008). *Bacillus* lipopeptides: versatile weapons for plant disease biocontrol. *Trends Microbiol.* 16, 115–125. doi: 10.1016/j.tim.2007.12.009
- O'Sullivan, L., Ross, R. P., and Hill, C. (2002). Potential of bacteriocin-producing lactic acid bacteria for improvements in food safety and quality. *Biochimie* 84, 593–604. doi: 10.1016/S0300-9084(02)01457-8
- Owens, J. D., Allagheny, N., Kipping, G., and Ames, J. M. (1997). Formation of volatile compounds during *Bacillus subtilis* fermentation of soya beans. *J. Sci. Food Agric.* 74, 132–140. doi: 10.1002/(SICI)1097-0010(199705)74:1<132::AID-JSFA779>3.0.CO;2-8
- Paik, S. H., Chakicherla, A., and Hansen, J. N. (1998). Identification and characterization of the structural and transporter genes for, and the chemical and biological properties of, sublancin 168, a novel lantibiotic produced by *Bacillus subtilis* 168. *J. Biol. Chem.* 273, 23134–23142. doi: 10.1074/jbc.273.36.23134
- Parisot, J., Carey, S., Breukink, E., Chan, W. C., Narbad, A., and Bonev, B. (2008). Molecular mechanism of target recognition by subtilin, a class I lanthionine antibiotic. *Antimicrob. Agents Chemother.* 52, 612–618. doi: 10.1128/aac.00836-07
- Patel, P. S., Huang, S., Fisher, S., Pirnik, D., Aklonis, C., Dean, L., et al. (1995). Bacillaene, a novel inhibitor of prokaryotic protein synthesis produced by *Bacillus subtilis*: production, taxonomy, isolation, physico-chemical characterization and biological activity. *J. Antibiot.* 48, 997–1003. doi: 10.7164/antibiotics.48.997
- Pattnaik, P., Kaushik, J. K., Grover, S., and Batish, V. K. (2001). Purification and characterization of a bacteriocin-like compound (Lichenin) produced anaerobically by *Bacillus licheniformis* isolated from water buffalo. *J. Appl. Microbiol.* 91, 636–645. doi: 10.1046/j.1365-2672.2001.01429.x
- Pedersen, P. B., Bjornvad, M. E., Rasmussen, M. D., and Petersen, J. N. (2002). Cytotoxic potential of industrial strains of *Bacillus* sp. *Regul. Toxicol. Pharmacol.* 36, 155–161. doi: 10.1006/rtp.2002.1574
- Peñuelas, J., Asensio, D., Tholl, D., Wenke, K., Rosenkranz, M., Piechulla, B., et al. (2014). Biogenic volatile emissions from the soil. *Plant Cell Environ.* 37, 1866–1891. doi: 10.1111/pce.12340
- Podile, A. R., and Prakash, A. P. (1996). Lysis and biological control of *Aspergillus niger* by *Bacillus subtilis* AF 1. *Can. J. Microbiol.* 42, 533–538. doi: 10.1139/m96-072
- Preecha, C., Sadowsky, M. J., and Prathuangwong, S. (2010). Lipopeptide surfactin produced by *Bacillus amyloliquefaciens* KPS46 is required for biocontrol efficacy against *Xanthomonas axonopodis* pv. *Glycines*. *Kasetsart J. Nat. Sci.* 44, 84–99.
- Priest, F. G., Goodfellow, M., Shute, L. A., and Berkeley, R. C. W. (1987). *Bacillus amyloliquefaciens* sp. nov., nom. rev. *Int. J. Syst. Evol. Microbiol.* 37, 69–71. doi: 10.1099/00207713-37-1-69
- Qi, G., Zhu, F., Du, P., Yang, X., Qiu, D., Yu, Z., et al. (2010). Lipopeptide induces apoptosis in fungal cells by a mitochondria-dependent pathway. *Peptides* 31, 1978–1986. doi: 10.1016/j.peptides.2010.08.003
- Raaijmakers, J. M., De Bruijn, I., Nybroe, O., and Ongena, M. (2010). Natural functions of lipopeptides from *Bacillus* and *Pseudomonas*: more than surfactants and antibiotics. *FEMS Microbiol. Rev.* 34, 1037–1062. doi: 10.1111/j.1574-6976.2010.00221.x
- Ramarathnam, R., Bo, S., Chen, Y., Dilantha Fernando, W. G., Xuewen, G., and Kievit, T. (2007). Molecular and biochemical detection of fengycin and bacillomycin D-producing *Bacillus* spp., antagonistic to fungal pathogens of canola and wheat. *Can. J. Microbiol.* 53, 901–911. doi: 10.1139/w07-049

- Rapp, C., Jung, G., Katzer, W., and Loeffler, W. (1988). Chlorotetain from *Bacillus subtilis*, an antifungal dipeptide with an unusual chlorine-containing amino acid. *Angew. Chem. Int. Ed. Engl.* 27, 1733–1734. doi: 10.1002/anie.198817331
- Raubitschek, F., and Dostrovsky, A. (1950). An antibiotic active against dermatophytes, derived from *Bacillus Subtilis*. *Dermatology* 100, 45–49. doi: 10.1159/000257151
- Raza, W., Ling, N., Yang, L., Huang, Q., and Shen, Q. (2016). Response of tomato wilt pathogen *Ralstonia solanacearum* to the volatile organic compounds produced by a biocontrol strain *Bacillus amyloliquefaciens* SQR-9. *Sci. Rep.* 6:24856. doi: 10.1038/srep24856
- Riley, M. A., and Wertz, J. E. (2002). Bacteriocins: evolution, ecology, and application. *Annu. Rev. Microbiol.* 56, 117–137. doi: 10.1146/annurev.micro.56.012302.161024
- Rohr, J. (1992). Comparison of multicyclic polyketides by folding analysis: a novel approach to recognize biosynthetic and/or evolutionary interrelationships of the natural products or intermediates and its exemplification on hepta-, octa-, and decaketides. *J. Org. Chem.* 57, 5217–5223. doi: 10.1021/jo00045a040
- Romano, A., Vitullo, D., Senatore, M., Lima, G., and Lanzotti, V. (2013). Antifungal cyclic lipopeptides from *Bacillus amyloliquefaciens* strain BO5A. *J. Nat. Prod.* 76, 2019–2025. doi: 10.1021/np400119n
- Romero, D., de Vicente, A., Rakotoaly, R. H., Dufour, S. E., Veening, J.-W., Arrebola, E., et al. (2007). The iturin and fengycin families of lipopeptides are key factors in antagonism of *Bacillus subtilis* Toward *Podosphaera fusca*. *Mol. Plant Microbe Interact.* 20, 430–440. doi: 10.1094/MPMI-20-4-0430
- Romero-Tabarez, M., Jansen, R., Sylla, M., Lünsdorf, H., Häußler, S., Santosa, D. A., et al. (2006). 7-O-Malonyl macrolactin A, a new macrolactin antibiotic from *Bacillus subtilis* active against methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant *Enterococci*, and a small-colony variant of *Burkholderia cepacia*. *Antimicrob. Agents Chemother.* 50, 1701–1709. doi: 10.1128/aac.50.5.1701-1709.2006
- Ryu, C.-M. (2015). “Bacterial volatiles as airborne signals for plants and bacteria,” in *Principles of Plant-Microbe Interactions: Microbes for Sustainable Agriculture*, ed. B. Lugtenberg (Cham: Springer International Publishing), 53–61.
- Ryu, C.-M., Farag, M. A., Hu, C.-H., Reddy, M. S., Kloepper, J. W., and Paré, P. W. (2004). Bacterial volatiles induce systemic resistance in *Arabidopsis*. *Plant Physiol.* 134, 1017–1026. doi: 10.1104/pp.103.026583
- Ryu, C.-M., Farag, M. A., Hu, C.-H., Reddy, M. S., Wei, H.-X., Paré, P. W., et al. (2003). Bacterial volatiles promote growth in *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 100, 4927–4932. doi: 10.1073/pnas.0730845100
- Sabaté, D. C., and Audisio, M. C. (2013). Inhibitory activity of surfactin, produced by different *Bacillus subtilis* subsp. *subtilis* strains, against *Listeria monocytogenes* sensitive and bacteriocin-resistant strains. *Microbiol. Res.* 168, 125–129. doi: 10.1016/j.micres.2012.11.004
- Saggese, A., Culurciello, R., Casillo, A., Corsaro, M., Ricca, E., and Baccigalupi, L. (2018). A marine isolate of *Bacillus pumilus* secretes a pumilacinin active against *Staphylococcus aureus*. *Mar. Drugs* 16:E180. doi: 10.3390/md16060180
- Schmidt, E. W. (2010). The hidden diversity of ribosomal peptide natural products. *BMC Biol.* 8:83. doi: 10.1186/1741-7007-8-83
- Schmidt, R., Cordovez, V., de Boer, W., Raaijmakers, J., and Garbeva, P. (2015). Volatile affairs in microbial interactions. *ISME J.* 9, 2329–2335. doi: 10.1038/ismej.2015.42
- Scholz, R., Molohon, K. J., Nachtigall, J., Vater, J., Markley, A. L., Süßmuth, R. D., et al. (2011). Plantazolicin, a novel microcin B17/Streptolysin S-Like natural product from *Bacillus amyloliquefaciens* FZB42. *J. Bacteriol.* 193, 215–224. doi: 10.1128/jb.00784-10
- Scholz, R., Vater, J., Budiharjo, A., Wang, Z., He, Y., Dietel, K., et al. (2014). Amylocyclin, a novel circular bacteriocin produced by *Bacillus amyloliquefaciens* FZB42. *J. Bacteriol.* 196, 1842–1852. doi: 10.1128/jb.01474-14
- Schulz, S., and Dickschat, J. S. (2007). Bacterial volatiles: the smell of small organisms. *Nat. Prod. Rep.* 24, 814–842. doi: 10.1039/b5507392h
- Scortichini, M., and Rossi, M. P. (1991). Preliminary *in vitro* evaluation of the antimicrobial activity of terpenes and terpenoids towards *Erwinia amylovora*. *J. Appl. Bacteriol.* 71, 109–112. doi: 10.1111/j.1365-2672.1991.tb02963.x
- Shafi, J., Tian, H., and Ji, M. (2017). *Bacillus* species as versatile weapons for plant pathogens: a review. *Biotechnol. Biotechnol. Equ.* 31, 446–459. doi: 10.1080/13102818.2017.1286950
- Shatalin, K., Shatalina, E., Mironov, A., and Nudler, E. (2011). H2S: a universal defense against antibiotics in bacteria. *Science* 334, 986–990. doi: 10.1126/science.1209855
- Shelburne, C. E., An, F. Y., Dholpe, V., Ramamoorthy, A., Lopatin, D. E., and Lantz, M. S. (2007). The spectrum of antimicrobial activity of the bacteriocin subtilisin A. *J. Antimicrob. Chemother.* 59, 297–300. doi: 10.1093/jac/dkl495
- Shrivastava, G., Ownley, B. H., Augé, R. M., Toler, H., Dee, M., Vu, A., et al. (2015). Colonization by arbuscular mycorrhizal and endophytic fungi enhanced terpene production in tomato plants and their defense against a herbivorous insect. *Symbiosis* 65, 65–74. doi: 10.1007/s13199-015-0319-1
- Siewert, G., and Strominger, J. L. (1967). Bacitracin: an inhibitor of the dephosphorylation of lipid pyrophosphate, an intermediate in the biosynthesis of the peptidoglycan of bacterial cell walls. *Proc. Natl. Acad. Sci. U.S.A.* 57, 767–773. doi: 10.1073/pnas.57.3.767
- Smith, S., and Tsai, S.-C. (2007). The type I fatty acid and polyketide synthases: a tale of two megasynthases. *Nat. Prod. Rep.* 24, 1041–1072. doi: 10.1039/B603600G
- Snelders, E., Huis in 't Veld, R. A., Rijs, A. J., Kema, G. H. J., Melchers, W. J. G., and Verweij, P. E. (2009). Possible environmental origin of resistance of *Aspergillus fumigatus* to medical triazoles. *Appl. Environ. Microbiol.* 75, 4053–4057. doi: 10.1128/aem.00231-09
- Sourabié, A. M., Spinnler, H.-E., Bourdat-Deschamps, M., Tallon, R., Landaud, S., and Bonnarme, P. (2012). S-methyl thioesters are produced from fatty acids and branched-chain amino acids by brevibacteria: focus on l-leucine catabolic pathway and identification of acyl-CoA intermediates. *Appl. Microbiol. Biotechnol.* 93, 1673–1683. doi: 10.1007/s00253-011-3500-3
- Spieß, T., Korn, S. M., Kötter, P., and Entian, K.-D. (2015). Autoinduction Specificities of the Lantibiotics Subtilin and Nisin. *Appl. Environ. Microbiol.* 81, 7914–7923. doi: 10.1128/aem.02392-15
- Stein, T. (2005). *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. *Mol. Microbiol.* 56, 845–857. doi: 10.1111/j.1365-2958.2005.04587.x
- Stein, T., Borchert, S., Conrad, B., Feesche, J., Hofemeister, B., Hofemeister, J., et al. (2002). Two different lantibiotic-like peptides originate from the ericin gene cluster of *Bacillus subtilis* A1/3. *J. Bacteriol.* 184, 1703–1711. doi: 10.1128/JB.184.6.1703-1711.2002
- Steinborn, G., Hajirezaei, M.-R., and Hofemeister, J. (2005). bac genes for recombinant bacilysin and anticapsin production in *Bacillus* host strains. *Arch. Microbiol.* 183, 71–79. doi: 10.1007/s00203-004-0743-8
- Sugawara, E., Ito, T., Odagiri, S., Kubota, K., and Kobayashi, A. (1985). Comparison of compositions of odor components of natto and cooked soybeans. *Agric. Biol. Chem.* 49, 311–317. doi: 10.1271/abb1961.49.311
- Sumi, C. D., Yang, B. W., Yeo, I.-C., and Hahm, Y. T. (2015). Antimicrobial peptides of the genus *Bacillus*: a new era for antibiotics. *Can. J. Microbiol.* 61, 93–103. doi: 10.1139/cjm-2014-0613
- Switzer Blum, J., Burns Bindi, A., Buzzelli, J., Stolz, J. F., and Oremland, R. S. (1998). *Bacillus arsenicoselenatis*, sp. nov., and *Bacillus selenitireducens*, sp. nov.: two haloalkaliphiles from Mono Lake, California that respire oxyanions of selenium and arsenic. *Arch. Microbiol.* 171, 19–30. doi: 10.1007/s002030050673
- Tagg, J. R., Dajani, A. S., and Wannamaker, L. W. (1976). Bacteriocins of gram-positive bacteria. *Bacteriol. Rev.* 40, 722–756.
- Tahir, H. A. S., Gu, Q., Wu, H., Niu, Y., Huo, R., and Gao, X. (2017). *Bacillus* volatiles adversely affect the physiology and ultra-structure of *Ralstonia solanacearum* and induce systemic resistance in tobacco against bacterial wilt. *Sci. Rep.* 7:40481. doi: 10.1038/srep40481
- Tanaka, K., Ishihara, A., and Nakajima, H. (2014). Isolation of anteiso-C17, iso-C17, iso-C16, and iso-C15 bacillomycin D from *Bacillus amyloliquefaciens* SD-32 and their antifungal activities against plant pathogens. *J. Agric. Food Chem.* 62, 1469–1476. doi: 10.1021/jf404531t
- Thasana, N., Prapagdee, B., Rangkadilok, N., Sallabhan, R., Aye, S. L., Ruchirawat, S., et al. (2010). *Bacillus subtilis* SSE4 produces subtilene A, a new lipopeptide antibiotic possessing an unusual C15 unsaturated (-amino acid. *FEBS Lett.* 584, 3209–3214. doi: 10.1016/j.febslet.2010.06.005
- Thimon, L., Peyoux, F., Maget-Dana, R., and Michel, G. (1992). Surface-active properties of antifungal lipopeptides produced by *Bacillus subtilis*. *J. Am. Oil Chem. Soc.* 69, 92–93. doi: 10.1007/bf02635884
- Um, S., Fraimout, A., Sapountzis, P., Oh, D.-C., and Poulsen, M. (2013). The fungus-growing termite *Macrotermes natalensis* harbors bacillaene-producing

- Bacillus* sp. that inhibit potentially antagonistic fungi. *Sci. Rep.* 3:3250. doi: 10.1038/srep03250
- van Straaten, K. E., Ko, J. B., Jagdhane, R., Anjum, S., Palmer, D. R. J., and Sanders, D. A. R. (2013). The structure of NtdA, a sugar aminotransferase involved in the kanosamine biosynthetic pathway in *Bacillus subtilis*, reveals a new sub-class of aminotransferases. *J. Biol. Chem.* 288, 34121–34130. doi: 10.1074/jbc.M113.500637
- Velivelli, S. L. S., Kromann, P., Lojan, P., Rojas, M., Franco, J., Suarez, J. P., et al. (2015). Identification of mVOCs from andean rhizobacteria and field evaluation of bacterial and mycorrhizal inoculants on growth of potato in its center of origin. *Microb. Ecol.* 69, 652–667. doi: 10.1007/s00248-014-0514-2
- Wang, B., Yuan, J., Zhang, J., Shen, Z., Zhang, M., Li, R., et al. (2013). Effects of novel bioorganic fertilizer produced by *Bacillus amyloliquefaciens* W19 on antagonism of Fusarium wilt of banana. *Biol. Fertil. Soils* 49, 435–446. doi: 10.1007/s00374-012-0739-5
- Wang, D., Rosen, C., Kinkel, L., Cao, A., Tharayil, N., and Gerik, J. (2009). Production of methyl sulfide and dimethyl disulfide from soil-incorporated plant materials and implications for controlling soilborne pathogens. *Plant Soil* 324, 185–197. doi: 10.1007/s11104-009-9943-y
- Wang, H., Fewer, D. P., Holm, L., Rouhiainen, L., and Sivonen, K. (2014). Atlas of nonribosomal peptide and polyketide biosynthetic pathways reveals common occurrence of nonmodular enzymes. *Proc. Natl. Acad. Sci. U.S.A.* 111, 9259–9264. doi: 10.1073/pnas.1401734111
- Wang, T., Liang, Y., Wu, M., Chen, Z., Lin, J., and Yang, L. (2015). Natural products from *Bacillus subtilis* with antimicrobial properties. *Chin. J. Chem. Eng.* 23, 744–754. doi: 10.1016/j.cjche.2014.05.020
- Wang, T., Wu, M.-B., Chen, Z.-J., Lin, J.-P., and Yang, L.-R. (2016). Separation, determination and antifungal activity test of the products from a new *Bacillus amyloliquefaciens*. *Nat. Prod. Res.* 30, 1215–1218. doi: 10.1080/14786419.2015.1048246
- Wang, Y., Yang, Q., Tosa, Y., Nakayashiki, H., and Mayama, S. (2005). Nitric oxide-overproducing transformants of *Pseudomonas fluorescens* with enhanced biocontrol of tomato bacterial wilt. *J. Gen. Plant Pathol.* 71, 33–38. doi: 10.1007/s10327-004-0157-0
- Willey, J. M., and Donk, W. A. V. D. (2007). Lantibiotics: peptides of diverse structure and function. *Annu. Rev. Microbiol.* 61, 477–501. doi: 10.1146/annurev.micro.61.080706.093501
- Williams, T. H. (1975). Naphthomycin, a novel ansa macrocyclic antimetabolite. Proton NMR spectra and structure elucidation using lanthanide shift reagent. *J. Antibiot.* 28, 85–86. doi: 10.7164/antibiotics.28.85
- Wise, C., Falardeau, J., Hagberg, I., and Avis, T. J. (2014). Cellular lipid composition affects sensitivity of plant pathogens to fengycin, an antifungal compound produced by *Bacillus subtilis* strain CU12. *Phytopathology* 104, 1036–1041. doi: 10.1094/PHYTO-12-13-0336-R
- Wong, J. H., Hao, J., Cao, Z., Qiao, M., Xu, H., Bai, Y., et al. (2008). An antifungal protein from *Bacillus amyloliquefaciens*. *J. Appl. Microbiol.* 105, 1888–1898. doi: 10.1111/j.1365-2672.2008.03917.x
- Wright, S. J. L., and Thompson, R. J. (1985). *Bacillus* volatiles antagonize cyanobacteria. *FEMS Microbiol. Lett.* 30, 263–267. doi: 10.1111/j.1574-6968.1985.tb01093.x
- Wu, L., Wu, H., Chen, L., Lin, L., Borriss, R., and Gao, X. (2015a). Bacilysin overproduction in *Bacillus amyloliquefaciens* FZB42 markerless derivative strains FZBREP and FZBSPA enhances antibacterial activity. *Appl. Microbiol. Biotechnol.* 99, 4255–4263. doi: 10.1007/s00253-014-6251-0
- Wu, L., Wu, H., Chen, L., Yu, X., Borriss, R., and Gao, X. (2015b). Difficidin and bacilysin from *Bacillus amyloliquefaciens* FZB42 have antibacterial activity against *Xanthomonas oryzae* rice pathogens. *Sci. Rep.* 5:12975. doi: 10.1038/srep12975
- Wu, L., Wu, H., Chen, L., Xie, S., Zang, H., Borriss, R., et al. (2014). Bacilysin from *Bacillus amyloliquefaciens* FZB42 has specific bactericidal activity against harmful algal bloom species. *Appl. Environ. Microbiol.* 80, 7512–7520. doi: 10.1128/aem.02605-14
- Wu, S., Jia, S., Sun, D., Chen, M., Chen, X., Zhong, J., et al. (2005). Purification and characterization of two novel antimicrobial peptides subpeptin JM4-A and subpeptin JM4-B produced by *Bacillus subtilis* JM4. *Curr. Microbiol.* 51, 292–296. doi: 10.1007/s00284-005-0004-3
- Xiu, P., Liu, R., Zhang, D., and Sun, C. (2017). Pumilacidin-like lipopeptides derived from marine bacterium *Bacillus* sp. strain 176 suppress the motility of *Vibrio alginolyticus*. *Appl. Environ. Microbiol.* 83:e00450-17. doi: 10.1128/aem.00450-17
- Xue, C., Tian, L., Xu, M., Deng, Z., and Lin, W. (2008). A new 24-membered lactone and a new polyene (-lactone from the marine bacterium *Bacillus marinus*. *J. Antibiot.* 61, 668–674. doi: 10.1038/ja.2008.94
- Yakimov, M. M., Timmis, K. N., Wray, V., and Fredrickson, H. L. (1995). Characterization of a new lipopeptide surfactant produced by thermotolerant and halotolerant subsurface *Bacillus licheniformis* BAS50. *Appl. Environ. Microbiol.* 61, 1706–1713.
- Yoo, J.-S., Zheng, C.-J., Lee, S., Kwak, J.-H., and Kim, W.-G. (2006). Macrolactin N, a new peptide deformylase inhibitor produced by *Bacillus subtilis*. *Bioorg. Med. Chem. Lett.* 16, 4889–4892. doi: 10.1016/j.bmcl.2006.06.058
- Yu, X., Ai, C., Xin, L., and Zhou, G. (2011). The siderophore-producing bacterium, *Bacillus subtilis* CAS15, has a biocontrol effect on *Fusarium* wilt and promotes the growth of pepper. *Eur. J. Soil Biol.* 47, 138–145. doi: 10.1016/j.ejsobi.2010.11.001
- Yuan, J., Li, B., Zhang, N., Waseem, R., Shen, Q., and Huang, Q. (2012a). Production of bacillomycin- and macrolactin-type antibiotics by *Bacillus amyloliquefaciens* NJN-6 for suppressing soilborne plant pathogens. *J. Agric. Food Chem.* 60, 2976–2981. doi: 10.1021/jf204868z
- Yuan, J., Raza, W., Shen, Q., and Huang, Q. (2012b). Antifungal activity of *Bacillus amyloliquefaciens* NJN-6 volatile compounds against *Fusarium oxysporum* f. sp. cubense. *Appl. Environ. Microbiol.* 78, 5942–5944. doi: 10.1128/aem.01357-12
- Zhang, B., Dong, C., Shang, Q., Han, Y., and Li, P. (2013). New insights into membrane-active action in plasma membrane of fungal hyphae by the lipopeptide antibiotic bacillomycin L. *Biochim. Biophys. Acta* 1828, 2230–2237. doi: 10.1016/j.bbamem.2013.05.033
- Zhao, P., Quan, C., Wang, Y., Wang, J., and Fan, S. (2014). *Bacillus amyloliquefaciens* Q-426 as a potential biocontrol agent against *Fusarium oxysporum* f. sp. spinaciae. *J. Basic Microbiol.* 54, 448–456. doi: 10.1002/jobm.201200414
- Zhao, Z., Wang, Q., Wang, K., Brian, K., Liu, C., and Gu, Y. (2010). Study of the antifungal activity of *Bacillus vallismortis* ZZ185 in vitro and identification of its antifungal components. *Bioresour. Technol.* 101, 292–297. doi: 10.1016/j.biortech.2009.07.071
- Zheng, M., Shi, J., Shi, J., Wang, Q., and Li, Y. (2013). Antimicrobial effects of volatiles produced by two antagonistic *Bacillus* strains on the anthracnose pathogen in postharvest mangos. *Biol. Control* 65, 200–206. doi: 10.1016/j.bioccontrol.2013.02.004
- Zhu, B.-F., Xu, Y., and Fan, W.-L. (2010). High-yield fermentative preparation of tetramethylpyrazine by *Bacillus* sp. using an endogenous precursor approach. *J. Ind. Microbiol. Biotechnol.* 37, 179–186. doi: 10.1007/s10295-009-0661-5
- Zimmerman, S., Schwartz, C., Monaghan, R., Pelak, B., Weissberger, B., Gilfillan, E., et al. (1987). Difficidin and oxydifficidin: novel broad spectrum antibacterial antibiotics produced by *Bacillus subtilis*. I. Production, taxonomy and antibacterial activity. *J. Antibiot.* 40, 1677–1681. doi: 10.7164/antibiotics.40.1677
- Zuber, P., Nakano, M. M., and Marahiel, M. A. (1993). “Peptide antibiotics,” in *Bacillus Subtilis and Other Gram-Positive Bacteria*, eds A. L. Sonenshein, J. A. Hoch, and R. Losick (Washington, DC: America Society for Microbiology), 897–916.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Caulier, Nannan, Gillis, Licciardi, Bragard and Mahillon. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.