



Moonlighting Proteins: The Case of the Hexokinases

Carolina Rodríguez-Saavedra, Luis Enrique Morgado-Martínez, Andrés Burgos-Palacios, Beatriz King-Díaz, Montserrat López-Coria and Sobeida Sánchez-Nieto*

Laboratorio de Transporte y Percepción de Azúcares en Plantas, Departamento de Bioquímica, Facultad de Química, Universidad Nacional Autónoma de México, Mexico City, Mexico

OPEN ACCESS

Edited by:

Mauro Salvi,
University of Padua, Italy

Reviewed by:

Shanshan Li,
Hubei University, China
Petr Heneberg,
Charles University, Czechia
Stefano Ugel,
University of Verona, Italy

*Correspondence:

Sobeida Sánchez-Nieto
sobeida@unam.mx

Specialty section:

This article was submitted to
Cellular Biochemistry,
a section of the journal
Frontiers in Molecular Biosciences

Received: 28 April 2021

Accepted: 24 May 2021

Published: 09 June 2021

Citation:

Rodríguez-Saavedra C,
Morgado-Martínez LE,
Burgos-Palacios A, King-Díaz B,
López-Coria M and Sánchez-Nieto S
(2021) Moonlighting Proteins: The
Case of the Hexokinases.
Front. Mol. Biosci. 8:701975.
doi: 10.3389/fmolb.2021.701975

Moonlighting proteins are defined as proteins with two or more functions that are unrelated and independent to each other, so that inactivation of one of them should not affect the second one and vice versa. Intriguingly, all the glycolytic enzymes are described as moonlighting proteins in some organisms. Hexokinase (HXK) is a critical enzyme in the glycolytic pathway and displays a wide range of functions in different organisms such as fungi, parasites, mammals, and plants. This review discusses HXKs moonlighting functions in depth since they have a profound impact on the responses to nutritional, environmental, and disease challenges. HXKs' activities can be as diverse as performing metabolic activities, as a gene repressor complexing with other proteins, as protein kinase, as immune receptor and regulating processes like autophagy, programmed cell death or immune system responses. However, most of those functions are particular for some organisms while the most common moonlighting HXK function in several kingdoms is being a glucose sensor. In this review, we also analyze how different regulation mechanisms cause HXK to change its subcellular localization, oligomeric or conformational state, the response to substrate and product concentration, and its interactions with membrane, proteins, or RNA, all of which might impact the HXK moonlighting functions.

Keywords: hexokinase, moonlighting function, sensor proteins, glycolytic moonlighting proteins, hexokinase regulation mechanisms

INTRODUCTION

Many proteins can perform different independent functions, which are known as “moonlighting proteins” (Jeffery, 1999). This term has been well-accepted to designate proteins with other “jobs” in the cells (Huberts and van der Klei, 2010). Although this characteristic may be seen as multifunctionality, moonlighting proteins are defined as proteins with two or more functions that are unrelated to each other. Proteins whose different functions are due to gene fusion, splice variants, or those that are distributed in separate subcellular compartments but perform the same role in each of them are excluded (Jeffery, 1999; Gancedo and Flores, 2008).

The first reports describing moonlighting proteins were published in the 1980s. Piatigorsky and Wistow (1989) noticed that some crystallins (structural proteins in the lens of vertebrate eyes) were metabolic enzymes: duck ϵ -crystallin was actually lactate dehydrogenase B4 (Hendriks et al., 1988), and τ -crystallin was α -enolase (Wistow et al., 1988). The enzymes' ability to function as crystallin proteins is based on their thermodynamic stability and their ability to concentrate at high levels without precipitating. Additionally, the enzymes' metabolic role in the lens tissue is unlikely, having a

structural role instead, indicating that both proteins' function depends on cell type (Wistow and Piatigorsky, 1988).

Up until 2015, around 300 moonlighting proteins had been characterized in many different organisms performing equally diverse functions (Mani et al., 2015), which can be found in databases such as MoonProt (<http://www.moonlightingproteins.org>; Mani et al., 2015) and MultitaskProtDB (<http://wallace.uab.es/multitask/>; Hernández et al., 2014). From the moonlighting proteins that have been experimentally characterized, there are several bioinformatic approaches to predict whether a protein can moonlight (Hernández et al., 2015).

A crucial feature to designate a protein as moonlighting is that its functions must be independent, so that inactivation of one of them should not affect the second one and vice versa (Huberts and van der Klei, 2010). Moonlighting activities can be two different enzymatic activities, combine a catalytic function with a non-catalytic activity, or even two non-catalytic functions (Rad et al., 2018). Moonlighting proteins are distributed in all living beings, and a wide range of examples have been described in yeasts (Gancedo and Flores, 2008), mammals (Sriram et al., 2005), protozoa (Collinridge et al., 2010), bacteria (Arnér and Holmgren 2000) and plants (Aguilera-Alvarado and Sánchez-Nieto, 2017). Most moonlighting proteins were initially described for a well-defined function in a biochemical pathway, like human aconitase (Kennedy et al., 1992) or *Trypanosoma cruzi* mevalonate kinase (Goldstein and Brown, 1990); or for participating in essential physiological processes, as reactive oxygen species (ROS)-defense mechanisms, like superoxide dismutase 1 (Sod1) in *Saccharomyces cerevisiae* (Tsang et al., 2014); or for assistance in protein folding or assembly of macromolecular complexes, like bacterial molecular chaperones (Hagemann et al., 2017). In general, proteins with the ability to moonlight can be a remarkably diverse group of proteins. Their moonlighting functions are different and very diverse depending on the organism. Considering the examples mentioned above, human aconitase moonlights as an intracellular sensor of iron levels (Artymiuk and Green, 2006); *T. cruzi* mevalonate kinase has an additional function as binding factor to cell host membranes (Bahia, 2017); *S. cerevisiae* Sod 1 acts as a transcription factor of genes related to oxidative stress response (Gancedo et al., 2016) and bacterial molecular chaperones moonlight as host cell adhesion factors (Henderson and Martin, 2011). The switching between one moonlighting activity to another can vary as a consequence of changes in subcellular localization (Sun et al., 2018), oligomeric or conformational state (Vega et al., 2016), ligands, substrates, cofactors or product concentration (Artymiuk and Green, 2006), interactions with DNA or RNA (Jeffery, 1999), membrane interactions (Jeffery, 2009), cell types (Wistow and Piatigorsky, 1988) and even in response to environmental factors (Copley, 2014).

This review addresses the diverse moonlighting functions ascribed to hexokinases (HXKs) in different kingdoms. HXK is a key enzyme in the metabolism and belongs to the most recognized moonlighting protein group: the glycolytic enzymes. This review also focuses on the consequences of the

regulation of the HXK by changes in its subcellular location, oligomeric conformation, RNA and membrane interaction.

MOONLIGHTING PROTEINS IN THE GLYCOLYTIC PATHWAY

According to Singh and Bhalla (Singh and Bhalla, 2020), proteins constitutively expressed, as those belonging to metabolic pathways, are better candidates to evolving moonlighting functions as they have a higher chance of interacting in a transient or a stable manner with different molecules, such as RNA, DNA or even other proteins, hence, leading to a benefit. This is supported by the REM (RNA-enzyme-metabolite) hypothesis, which postulates the existence of regulatory networks between gene expression and intermediary metabolism mediated by moonlighting RNA-binding metabolic enzymes (Hentze and Preiss, 2010). Examples of enzymes leading to the REM hypothesis are hexokinase (HXK), aldolase (ALDO), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), enolase (ENO) and pyruvate kinase (PK) in HeLa, HEK293 and mouse embryonic stem cells (Castello et al., 2015). Furthermore, HXK, Phosphofructokinase (PFK), ALDO, GAPDH, Phosphoglycerate Kinase (PGK), Phosphoglycerate Mutase (PGM), ENO and PK, have been found inside the nucleus of *Saccharomyces cerevisiae* and cancer and muscle cells from *Homo sapiens* (Kim and Dang, 2005; Boukouris et al., 2016). At this location, these proteins carry out a wide range of functions either in complex with DNA or in complex with other proteins, some of which are listed in **Table 1** (for further information see Boukouris et al., 2016 and references within). It is not clear if the main purpose of these proteins in the nucleus is providing ATP and NADH, yet it is thought that their unusual location may be the result of evolutionary pressure. Around the REM hypothesis, several questions arise: are glycolytic enzymes capable of binding directly to RNA, or do they have to interact with other partners to do so? Interestingly, many of them share moonlighting functions as adhesion or transcriptional factors in different organisms: how can certain moonlighting functions be conserved throughout species? And, regarding this, it is also intriguing that since glycolysis is an essential and conserved metabolic pathway, it is remarkable that every of its enzymes play very different moonlighting roles, so how is it that they were selected to carry out different functions in different cell locations, under different environmental conditions and/or different stages of development?

In bacteria, the fact that most of the glycolytic enzymes help the microorganisms colonize or establish on host's cells has led to hypothesize that the reason why glycolytic enzymes moonlight is because as both bacterial and human glycolytic enzymes present a high degree of conservation, the host immune system will not elicit antibodies against the pathogens' glycolytic enzymes (Franco-Serrano et al., 2018). According to Wang et al., 2015, autoimmune diseases affect from 3–5% of the world population, with autoimmune thyroid disease and type one diabetes being the two most common ones. Amela et al., 2007 (a previous paper published by the same group as Franco-Serrano, et al., 2018), showed that only 1% of 2,175 proteins analyzed contain a

TABLE 1 | Moonlighting function of nine glycolytic enzymes.

Enzyme	Canonical function	Moonlighting function	Organism	References
Glucose-6-phosphate isomerase	Glucose-6-phosphate \rightleftharpoons fructose-6-phosphate	Adhesion protein	<i>Lactobacillus crispatus</i>	Kainulainen et al. (2012)
		Autocrine motility factor Neuroleukin Virulence	<i>Homo sapiens</i> <i>Xanthomonas oryzae</i>	Wantanabe et al. (1996) Sun et al. (1999) Tsuge et al. (2004)
Phosphofructokinase	Fructose-6-phosphate \rightarrow fructose-1,6-bisphosphate	Autophagy regulator	<i>Pichia pastoris</i>	Yuan et al. (1997)
		Part of RNA degradosome complex Transcriptional regulator	<i>Bacillus subtilis</i> <i>Drosophila melanogaster</i>	Commichau et al. (2009) Enzo et al. (2015)
Aldolase	Fructose-1,6-bisphosphate \rightleftharpoons glyceraldehyde-3-phosphate + dihydroxyacetone phosphate	Actin dynamics controller	<i>Oryctolagus cuniculus</i>	Ritterson-Lew and Tolan (2013)
		GLUT4 regulator Involved in pathogen-host interaction	<i>Homo sapiens</i> <i>Candida albicans</i> <i>Plasmodium berghei</i> <i>Streptococcus pneumoniae</i> <i>Neisseria meningitidis</i> <i>Paracoccidioides</i> spp.	Kao et al. (1999) Crowe et al. (2003) Bosch et al. (2007) Blau et al. (2007) Tunio et al. (2010b) Marcos et al. (2014)
		Transcriptional regulator	<i>Mus musculus</i> <i>Francisella tularensis</i> <i>Mus musculus</i>	Kiri and Goldspink (2002) Ziveri et al. (2017) Lu et al. (2001)
		V-ATPase assembly and activity mediator Zona pellucida recognition	<i>Homo sapiens</i>	Petit et al. (2013)
		Endoglucanase	<i>Pyrococcus furiosus</i>	Sharma and Guptasarma (2017)
Triosephosphate isomerase	Dihydroxyacetone phosphate \rightleftharpoons glyceraldehyde-3-phosphate	Pathogen establishment in host	<i>Paracoccidioides brasiliensis</i> <i>Streptococcus anginosus</i> <i>Streptococcus oralis</i> <i>Lactobacillus plantarum</i> <i>Staphylococcus aureus</i>	Pereira et al. (2007) Kinnby et al. (2008) Kinnby et al. (2008) Ramiah et al. (2008) Ikeda and Ichikawa (2014)
		Zona pellucida recognition Acts as SSBP Apoptosis regulator	<i>Homo sapiens</i> <i>Bos taurus</i> <i>Mus musculus</i>	Petit et al. (2013) Grosse et al. (1986) Dastoor and Dreyer (2001)
		Autophagy regulator	<i>Rattus norvegicus</i> <i>Homo sapiens</i> <i>Arabidopsis thaliana</i> <i>Nicotiana benthamiana</i>	Hara et al. (2005) Yego and Mohr (2010) Henry et al. (2015) Han et al. (2015)
		Extracellular polysaccharide synthesis Flagellar protein Immunomodulator	<i>Xanthomonas campestris</i> <i>Trypanosoma brucei</i> <i>Streptococcus agalactiae</i> <i>Haemonchus contortus</i>	Lu et al. (2009) Brown et al. (2014) Madureira et al. (2007) Sahoo et al. (2013)
		Pathogen establishment in host	Group a streptococci <i>Lactobacillus crispatus</i> <i>Escherichia coli</i> <i>Schistosoma bovis</i> <i>Bacillus anthracis</i> <i>Neisseria meningitidis</i> <i>Paracoccidioides</i> spp <i>Erysipelothrix rhusiopathiae</i> Streptococci	Pancholi and Fischetti (1992) Hurmalaianen et al. (2007) Egea et al. (2007) Ramajo-Hernández et al. (2007) Matta et al. (2010) Tunio et al. (2010a) Marcos et al. (2014) Zhu et al. (2017) Modun et al. (2000)

(Continued on following page)

TABLE 1 | (Continued) Moonlighting function of nine glycolytic enzymes.

Enzyme	Canonical function	Moonlighting function	Organism	References
Phosphoglycerate kinase	1,3-bisphosphoglycerate \rightleftharpoons 3-phosphoglycerate	Transcriptional regulator	<i>Homo sapiens</i> <i>Mus musculus</i> <i>Arabidopsis thaliana</i>	Zheng et al. (2003) Ikeda et al. (2012) Kim et al. (2020)
		DNA glycosylase	<i>Homo sapiens</i>	Meyer-Siegler et al. (1991)
		Zona pellucida recognition	<i>Homo sapiens</i>	Petit et al. (2013)
		Binds to polymerase increasing DNA synthesis	<i>Homo sapiens</i>	Popanda et al. (1998)
		Flagellar protein	<i>Mus musculus</i> <i>Trypanosoma brucei</i>	Sawyer et al. (2008) Brown et al. (2014)
Phosphoglycerate mutase	3-phosphoglycerate \rightleftharpoons 2-Phosphoglycerate	Pathogen establishment in host	Group B streptococci	Boone et al. (2011)
		Plasmin reductase	<i>Homo sapiens</i>	Lay et al. (2000)
		Transcriptional regulator	<i>Homo sapiens</i>	Shetty et al. (2004)
Enolase	2-phosphoglycerate \rightleftharpoons phosphoenolpyruvate	Involved in bacterial adhesion to host	<i>Candida albicans</i> <i>Bifidobacterium lactis</i>	Crowe et al. (2003) Candela et al. (2007)
		Immunosuppressing protein	<i>Streptococcus sobrinus</i> <i>Steinemema glaseri</i> <i>Entamoeba invadens</i>	Veiga-Malta et al. (2004) Liu et al. (2012) Segovia-Gamboa et al. (2010)
Pyruvate kinase	Phosphoenolpyruvate \rightleftharpoons Pyruvate	Involved in encystation mechanism	<i>Naegleria fowleri</i>	Chávez-Munguía et al. (2011)
		Neurotrophic and neuroprotective factor	<i>Rattus norvegicus</i>	Hattori et al. (1995)
		Pathogen establishment in host	<i>Pneumocystis carinii</i> <i>Candida albicans</i> <i>Streptococcus mutans</i> <i>Leishmania mexicana</i> <i>Bacillus anthracis</i> <i>Aeromonas hydrophila</i> <i>Paracoccidioides brasiliensis</i>	Fox and Smulian (2001) Jong et al. (2003) Jones and Holt (2007) Vanegas et al. (2007) Agarwal et al. (2008) Sha et al. (2009) Nogueira et al. (2010)
		Tau lens crystalline	<i>Pseudemys scripta</i>	Wistow et al. (1988)
		Transcriptional regulator	<i>Homo sapiens</i>	Subramanian and Miller (2000)
		Bacterial adhesion	<i>Toxoplasma gondii</i> <i>Lactococcus lactis</i>	Mouveaux et al. (2014) Katakura et al. (2010)
		Hormone binding protein	<i>Rattus norvegicus</i>	Parkinson et al. (1991)
		Protein kinase	<i>Homo sapiens</i> <i>Saccharomyces cerevisiae</i>	Gupta and Bamezai (2010) Li et al. (2015)
		Transcriptional regulator	<i>Homo sapiens</i>	Gao et al. (2012)
		Zona pellucida recognition	<i>Homo sapiens</i>	Petit et al. (2013)

sequence that can be found in both pathogen and host, suggesting that this might be a mechanism for a pathogen to avoid the host's immune system. In spite of this, it has been found that it is possible to generate autoantibodies raised against glycolytic enzymes. Antibodies against TPI (Vermeulen et al., 2011), ALDO (Magrys et al., 2007; Goëb et al., 2009; Vermeulen et al., 2011), GAPDH (Adamus et al., 2011), PGK (Goëb et al., 2009), PGM (Kimura et al., 2010; Vermeulen et al., 2011) and ENO (Terrier et al., 2007; Goëb et al., 2009; Adamus et al., 2011; Vermeulen et al., 2011; Maccallini et al., 2018) are linked to the development of retinopathy, rheumatoid arthritis, autoimmune hepatitis, inflammatory bowel disease and Lyme disease. The high expression of the glycolytic proteins in the diseases mentioned

before, together with frequent infections and chronic inflammation, and the high extracellular exposure of some of these protein may promote the production of autoantibodies (Adamus, 2017), suggest that "mimicking" glycolytic enzymes in order to establish on a host, might not be the best mechanism.

HEXOKINASE

Hexokinase was first described by Otto Meyerhof who demonstrated the stimulatory effect of a baker's yeast extract on the fermentation of hexoses (Meyerhof, 1927). Almost a decade later, von Euler and Adler (1935) and Meyerhof (1935),




FIGURE 1 | Structural overview of hexokinases. **A, D and G** Human hexokinase IV (HXKIV), **B, E and H** *Kluyveromyces lactis* hexokinase 1 (KHXK1) and **C, F and I** *A. thaliana* hexokinase 1 (AtHXK1). **A, B and C** are the Apo-form of the HXKs, **D, E and F** are the glucose-bound form and **G, H and I** are the comparison between the small domain of the Apo-HXK (dark green) and the Glc-HXK (light green). **A, B and C** exhibit a similar tertiary structure; they display a large (blue) and a small (orange) domain. Between both domains, there is a cleft (white arrows) where the active site is located. In **D, E and F** the small domain approaches to the large one to interact with glucose and carry out the catalysis and although domain approaching occurs in all of the three cases mentioned, it does not occur to the same extent; **G, H and I** show a comparison between the small domain of the Apo-HXK (dark green) vs. the Glc-HXK (light green). The white dotted line represents the angle of rotation of the small domain toward the big one. Glucose is represented as white sticks in **D, E and F**. Images of 1V4T (Kamata et al., 2004), 3O80 (Kuettnner et al., 2010), 4QS8 (Feng et al., 2015), 3IDH (Petit et al., 2011), 3O8M (Kuettnner et al., 2010) and 4QS7 (Feng et al., 2015) were created using PyMOL. The PyMOL Molecular Graphics System. Version 2.3 Schrödinger, LLC.

TABLE 2 | Confirmed non-canonical activities of hexokinases.

Moonlighting function	Organism	References
Apoptosis regulator	<i>Homo sapiens</i>	Pastorino et al. (2002)
	<i>Nicotiana tabacum</i>	Godbole et al. (2013)
	<i>Nicotiana benthamiana</i>	Kim et al. (2006)
Autophagy regulator	<i>Homo sapiens</i>	Kundu (2014)
Intracellular glucose sensor	<i>Arabidopsis thaliana</i>	Moore et al. (2003)
	<i>Chromochloris zofingiensis</i>	Roth et al. (2019)
	<i>Klebsormodium nitens</i>	Ulfstedt et al. (2018)
	<i>Oryza sativa</i>	Cho et al. (2009), Kim et al. (2016)
	<i>Nicotiana tabacum</i>	Kim et al. (2013b)
	<i>Saccharomyces cerevisiae</i>	Johnston (1999)
	<i>Solanum tuberosum</i>	Veramendi et al. (2002)
	<i>Homo sapiens</i>	Adams et al. (1991)
	<i>Malus domestica</i>	Hu et al. (2016), Sun et al. (2018)
	<i>Saccharomyces cerevisiae</i>	Herrero et al. (1989)
Protein kinase	<i>Arabidopsis thaliana</i>	Cho et al. (2006b)
	<i>Candida albicans</i>	Laurian et al. (2019)
	<i>Homo sapiens</i>	Sheikh et al. (2018)
	<i>Kluyveromyces lactis</i>	Prior et al. (1993)
	<i>Oryza sativa</i>	Huang et al. (2015)
	<i>Saccharomyces cerevisiae</i>	Herrero et al. (1998)
	<i>Vitis vinifera</i>	Wang et al. (2017)

determined that this enzyme catalyzes the phosphorylation of hexoses using ATP as a phosphoryl donor. Later, Colowick and Kalckar (1943) showed that only the γ -phosphate group was transferred from ATP to the hexoses. By 1953, the first plant hexokinase (from *Triticum aestivum*) was described (Saltman, 1953), giving enough evidence to state that hexokinases can be found in microorganisms, animals, and plants.

The existence of different hexokinase isoenzymes was first found in *Saccharomyces cerevisiae* in 1961 (Tayser and Colowick, 1961). In the same decade (1960s) mammal hexokinase isoenzymes were reported in rat liver (González et al., 1964), human and dogs (Brown et al., 1967). At last, Jang et al. (1997) demonstrated that there are also hexokinase isozymes in plants.

Hexokinases have similar structures between different organisms (Figure 1); they have two domains, the large domain has an oval shape that may or may not contain a targeting location sequence on the N-terminus, and a small domain with a three-layer architecture (Kamata et al., 2004; Feng et al., 2015). It should be clarified that in this review and in previous papers related to hexokinases' structure, the term "domain" is used to refer to a subregion of a protein that is autonomous in the sense that it possesses all the characteristics of a complete globular protein, as stated by Schulz and Schirmer (1979). Not only do hexokinases have a similar tertiary structure, but also a similar number of secondary structures, i.e. human hexokinase IV, HXKIV, has 15 α -helices (Figure 1A), whilst *Kluyveromyces lactis* hexokinase 1, KHXX1 (Figure 1B), has 13 and *Arabidopsis thaliana* hexokinase 1, AtHXX1 (Figure 1C), has 14; AtHXX1 has 12 β -strands and HXKIV has 13 and KHXX1 has 18. Another characteristic shared by hexokinases is that upon glucose binding, their domains approach to each other to a different extent, adopting the catalytically active closed conformation (Figures 1D–F) (Bennett and Steitz, 1980; Kamata et al., 2004; Petit et al., 2011; He et al., 2019). Besides those structural similarities, AtHXX1 is not only a glucose

phosphorylating enzyme but also a glucose sensor protein, so deeper analysis needs to be done to find the features that are important for HXKs to moonlighting.

FUNCTION OF THE HEXOKINASES AS GLUCOSE-SENSOR IS CONSERVED

Out of the non-canonical functions of the hexokinase, probably the one that has been described in a larger number of organisms is the glucose-sensing ability (Table 2). Hexokinases able to act as glucose sensors have been described in mammals and several other vertebrates, including fishes (Matschinsky, 1993; Irwin and Tan, 2014; Heneberg, 2019), *Saccharomyces cerevisiae* (Rolland et al., 2002), *Arabidopsis thaliana* (Moore et al., 2003), *Solanum tuberosum* (Veramendi et al., 2002), *Oryza sativa* (Cho et al., 2009; Kim et al., 2016), *Nicotiana tabacum* (Kim et al., 2013b), *Chromochloris zofingiensis* (Roth et al., 2019), *Klebsormidium nitens* (Ulfstedt et al., 2018) and *Synechocystis sp* (Ryu et al., 2008). Despite being well-known glucose-sensing proteins, the precise mechanism responsible for this moonlighting activity is not clearly understood yet. Several recent advances in the single molecule enzymology and the development of new computational analysis are giving insight on the enzymes' behavior that could lead to explain its moonlighting ability, for instance, mammal HXKIV (GCK), it has been suggested that its glucose-sensor activity is due to its low glucose affinity ($K_m \approx 10$ mM, Bustos and Iglesias, 2000; Larion and Miller, 2012) together with its kinetic cooperativity. The discovery of different enzyme conformers after the first enzyme cycle, explains the cooperativity that enables an adequate response to low glucose concentration and avoid depleting blood glucose levels (Larion et al., 2015; Šimčíková et al., 2017). A similar explanation might explain why SchXX2 suffers a conformational change in response to cytoplasmic glucose levels that allows the

enzyme to form the repressor complex that leads to the repression of *SUC2* gene (Vega et al., 2016). In some cases, determining the three-dimensional structure has shed light on elucidating how a protein can have two different functions (Jeffery, 2004). Unfortunately, as far as hexokinases are concerned, this task is limited because of the few amount of crystallized glucose sensing hexokinases, combined with the absence of non-glucose sensing HXK structures to make a comparison. In fact, from the organisms listed previously, it has only been possible to crystallize hexokinase II from *S. cerevisiae* (Kuser et al., 2000), hexokinase IV from *Homo sapiens* (Kamata et al., 2004), hexokinase 1 from *A. thaliana* (AtHXK1; Feng et al., 2015) and hexokinase 6 from *O. sativa* (OsHXK6; He et al., 2019).

The crystal structure of ScHXK2, AtHXK1, and OsHXK6 show great resemblance with other non-moonlighting hexokinases: they have almost the same number of secondary structures forming two domains which are connected by hinges, and between these two domains the active site is located. When bound to glucose, one domain rotates toward the other about 20° (Anderson et al., 1979; Bennett and Steltz, 1980; Feng et al., 2015; He et al., 2019). Besides, amino acids involved in catalysis are conserved.

One of the characteristics of moonlighting proteins is that the inactivation of the canonical function does not hamper the non-canonical function or vice-versa. This was proved by crystallizing the catalytically inactive mutant AtHXK1S177A (which still retains its glucose-sensing ability according to Moore et al., 2003) complexed with glucose (Feng et al., 2015); the 3D structure revealed that both domains still approach and are still able to interact with glucose in a similar fashion to wild type hexokinase, with an overall root mean square deviation (r.m.s.d.) of 0.45 Å. Additionally, AtHXK1 forming a complex with VHA-B1 and RPT5B is known to regulate the transcription of photosynthesis-related genes when bound to glucose (Cho et al., 2006b). These two reports suggest that the AtHXK1-Glc complex could be the active conformation able to moonlight, although further work is still needed to find the contacts between the different HXK patterns.

HEXOKINASES MOONLIGHTING FUNCTIONS IN DIVERSE LIFE KINGDOMS

Protozoa

In *Trypanosoma brucei*, TbHK1 and TbHK2 are expressed in bloodstream-form and procyclic stage parasites in different developmental stages, and TbHK2 has been localized in the flagellum inside and outside of the glycosomes, specialized structures that contain glycolytic enzymes (Blattner et al., 1995; Michaels et al., 2006). The ability of TbHK2 to be on different compartments in the cell might help to have different functions. Additionally, recombinant TbHXK2 is inactive, but when mixed with TbHXK1 also as a recombinant protein, the heterooligomer is active with kinetic properties different from the TbHXK1 recombinant enzyme alone, suggesting a regulatory role of TbHXK1, likely a glucose sensor to regulate the glycolysis flux and ATP production (Joice et al., 2012). However, these


hypotheses remain to be demonstrated, as well as to elucidate the mechanism involved in the translocation of TbHK2 from glycosomes to the flagellum, the interaction partners involved, and whether the functions of both HXKs as glucose-phosphorylating enzymes and as Hb binding protein or sugar sensor are uncoupled or not.

Yeasts and Fungi

Yeasts comprise an excellent example of the glucose sensor moonlighting activity of HXKs. *S. cerevisiae* expresses two cytosolic hexokinases: ScHXK1 and ScHXK2; and one glucokinase: ScGLK (Randez-Gil et al., 1998). ScHXK2 has been extensively described for its moonlighting properties participating in both glucose catabolism and glucose catabolite repression, which has been demonstrated to be mutually uncoupled functions (Ahuatzi et al., 2004; Gancedo et al., 2014). In the glucose catabolite repression process, ScHXK2 functions as a regulator of gene transcription in the nucleus, so that under high glucose conditions ScHXK2 translocates from the cytosol to the nucleus and once inside, it interacts with C2H2 zinc-finger transcription factor Mig1 (Randez-Gil et al., 1998; Ahuatzi et al., 2004) to recruit other proteins in order to form a repressor complex at gene promoters encoding catabolic enzymes of non-fermentable sugars, such as *SUC2* gene, repressing their transcription (Figure 2A) (Herrero et al., 1998; Ahuatzi et al., 2004; Vega et al., 2016). Contrary, low glucose concentration abolish the interaction between ScHXK2 and Mig1, and the repressor complex is disassembled, thus allowing transcription of *SUC2* gene, which encodes cytoplasmic and extracellular invertases (Figure 2B) (Ahuatzi et al., 2007; Fernández-García et al.).

Hexokinase 1 from *Yarrowia lipolytica* (YlHXK1) might also be a moonlighting protein; according to Hapeta et al., 2021, this hexokinase presents a unusual loop of 37 amino acids long that is not present in other hexokinases from the *Yarrowia* genus. *Y. lipolytica* cells, expressing YlHXK1 lacking this loop (YlHXK1Δloop), are unable to phosphorylate neither glucose nor fructose, apparently due to the conformational changes triggered by the loop deletion. Furthermore, by overexpressing the YlHXK1Δloop, the expression of lipases 2 and 8 (LIP2 and LIP8) is diminished. Unluckily, research on the effect that point mutations of amino acids involved in catalysis have on the expression of LIP2 and LIP8 is still needed in order to prove whether the downregulation of these lipases is due to incapability to phosphorylate glucose rather than the conformational changes of YlHXK1Δloop.

Studies on other fungi HXKs have been made by analyzing the phenotype of HXKs null mutants. *Fusarium graminearum*, the cause of *Fusarium* head blight (FHB), produces a range of mycotoxins, like type B trichothecenes, potent inhibitors of eukaryotic protein synthesis, being deoxynivalenol (DON) the main trichothecene produced having a serious impact on crops, human and animal health (Alexander et al., 2009). Null mutants of the two HXKs in *F. graminearum* showed that FgHXK1, but no FgHXK2 deletion produced significantly low levels of DON, indicating that FgHXK1 might be involved in increased DON production, and may have a possible indirect role on virulence




and *F. graminearum* spread within spikes. Additionally, *fghxk1* null mutants also exhibited defects in growth, pathogenicity, conidial, and perithecia production, suggesting a role in such processes (Zhang et al., 2016). Null *hxx1* mutants in *F. verticilloides*, the causal agent of maize ear rot worldwide, were substantially less virulent and produced fivefold less fumonisin B1 (FB1) toxin than the wild-type, and the production of FB1 was restored in the complemented strain, indicating that HXX1 is required for FB1 biosynthesis. Though the exact mechanism through which HXX1 affects FB1 production is still unknown, authors suggest that carbon catabolite repressor protein, CreA, a C2H2 zinc-finger protein, may indirectly regulate FB1 biosynthesis through the regulation of sugar transporters and sugar kinases, as they found that HXX1 has four putative CreA binding sites within 500 bp upstream of the start codon (Desai et al., 2002; Kim et al., 2011). Taken together, these reports suggest that fungi HXXs may have a much broader regulatory role that remains to be explored.

Mammals

In mammals, hexokinases present a wide range of functions, but in which it is still questionable their moonlighting ability (Dubey et al., 2016; Affourtit et al., 2018). For example, the human

genome encodes five hexokinases that phosphorylate glucose using ATP as a phosphoryl donor: HXXI, HXXII, HXXIII, HXXIV, the latter also known as glucokinase (GCK), and HXX domain protein containing 1 or HXXDC1. HXXDC1 was recently found in a genome wide association study and it is implicated in glucose tolerance, especially during pregnancy. It has a high degree of similarity with HXXI and show hexokinase catalytic activity, however, it is possible that it also regulates the activity of other HXXs (Guo et al., 2015; Ludvik et al., 2016). Another glucose phosphorylating enzyme found in humans is ADPGK, this endoplasmic reticulum enzyme is quite different from the other HXXs; it uses ADP as a phosphoryl donor and shows optimal activity at high temperature (42°C) and acidic pH and seems to be implicated in the regulation of the energy metabolism and protein glycosylation due to its subcellular location (Imle et al., 2019). Each of the hexokinase isoforms expresses differentially through tissues, being HXXI the one that expresses ubiquitously, though it is mainly expressed in the brain, and therefore suggesting a mere catabolic role. HXXII is the predominant isoform in skeletal and cardiac muscle, as well as in adipocytes, though an increase in its expression occurs in cancer cells, even in tissues where its expression is negligible under health physiological conditions (Mathupala et al., 2006). HXXIII



is also ubiquitously expressed, but is not predominant in any tissue, while glucokinase is found in pancreatic α - and δ -cells, adrenal gland, glucose-sensitive neurons, enteroendocrine cells and anterior pituitary cells (Roberts and Miyamoyto, 2015; Matschinsky and Wilson, 2019). HXKDC1 is only absent in muscle and adipose tissue, with high expression in colon, kidney and liver (Ludvik et al., 2016). ADPGK is highly expressed in immune cells of myeloid and lymphoid lineages (Imle et al., 2019).

From all the human isoforms, HXKII has been extensively studied due to its implication in the progression and maintenance of tumor cells, hence other functions in which it is involved have been discovered. HXKI and HXKII have a mitochondrial anchorage sequence at their N-terminus that allows its association to the MOM and its interaction with the voltage-

dependent anion channel (VDAC), the most abundant porin in MOM (Arora and Pedersen, 1988; Wilson, 2003). This association was demonstrated to be a “cellular strategy” to glycolytic enzymes and metabolites to gain “preferential access” to mitochondrial ATP, thus intimately coupling oxidative phosphorylation with carbon metabolism through ATP channeling from VDAC to HXKII (Mathupala et al., 2006). But HXKII association to MOM and VDAC offers an additional advantage: inhibiting the formation of mitochondrial permeability transition pore (mPTP) in cancer cells that overexpress HXKII and consequently, inhibiting apoptosis programmed cell death which is elicited by apoptotic Bcl-2 protein family and the release of cytochrome c through mPTP (Figure 3A). The entire inhibition mechanism is still to be elucidated, but there is evidence involving HXKII regulation.

Furthermore, indirect HXKII interaction with cyclophilin-D, a mitochondrial matrix protein related to the opening of mPTP (Baines et al., 2005), seems to stabilize HXKII association to MOM, thus promoting a closed state of mPTP and providing protection to mitochondrial death pathways. In contrast, dissociation of HXKII from the mitochondria facilitates the formation of mPTP mediated by cyclophilin-D and promoting apoptosis (Machida et al., 2006; Roberts and Miyamoto, 2015). Additionally, several reports have demonstrated that heterologous expression of VDAC can induce cell death, but the process can be mitigated by concomitant overexpression of HXKII (Pastorino and Hoek, 2008). In relation to this, an investigation showed that VDAC1, which is synthesized in cytosolic ribosomes, is found in both, cytosol and mitochondria, and cell death only occurs when the protein migrates to mitochondria. Translocation of VDAC1 seems to be regulated by HXKII protein levels, which apparently requires its catalytic activity, its N-terminal region, and the Glu73 residue to retain VDAC in the cytosol (Dubey et al., 2016). There is no information about the factors or signals involved in mitochondrial-translocation or cytosol-retention of the VDAC-HXKII complex, but it is likely to tilt the survival/death balance of the cell.

Furthermore, HXKII has been confirmed to mediate the regulation of autophagy in the absence of glucose-6-phosphate (G6P) (Roberts et al., 2014). Autophagy is a catabolic self-digestion system initiated by limited energy availability to ensure cellular energy homeostasis and survival. Overexpression of HXKII has been reported to promote autophagy, while its deletion inhibits the process even in the absence of glucose. So far, the elucidated mechanism involves HXKII binding to TORC1 (Target of Rapamycin Complex one), a negative regulator of autophagy, by its TOS (TOR signaling) motif, inactivating it (Roberts et al., 2014). According to experimental data, the key factor responsible for activation/inactivation of autophagy is the glucose phosphorylating activity, but not glucose binding (Figure 3B). However, the uncoupling between catabolic activity and autophagy regulation functions of HXKII is ambiguous since the experimental data did not show that the sugar-kinase activity of HXKII is independent of its G6P sensing function. It would require additional experiments, as mutating the HXKII catalytic site and determining whether it is still able to the allosteric recognition of G6P and autophagy prevention.

HXKII has also a role in the activation of the NOD-like receptor pyrin domain-containing 3 (NLRP3) inflammasome (Wolf et al., 2016), a cytosolic component of the innate immune system that regulates caspase-1 activation and secretion of proinflammatory cytokines IL-1 β and IL-18 in response to cellular infections and cellular damage (Kelley et al., 2019). Wolf et al. (2016) identified that purified N-acetylglucosamine (NAG) or derived from degradation of bacterial peptidoglycan in phagosomes inactivate HXKII, leading to its dissociation from MOM and inducing inflammasome NLRP3 activation, indicating that HXKII is likely acting as a pattern recognition receptor. Interestingly detachment of HXKII from MOM can also be caused by the

addition of G6P and jasmonate, a plant hormone (Goldin et al., 2008).

As discussed above, interaction of HXKII with VDAC in MOM allows HXKII to have preferential access to ATP, so inactivation and dissociation of HXKII from MOM implies a reduction in glycolytic flux and leads to interaction of VDAC to Bcl-protein family, which promotes opening of mPTP and subsequent apoptosis, so chemicals that induce this process could be used to reduce cancer cell proliferation, just as jasmonate does (Goldin et al., 2008). Moreover, metabolic disorders, such as diabetes or obesity, can directly induce inflammation (Wen et al., 2012) but it is still unclear whether the activation of NPLR3 by HXK follows the same mechanism in diabetes and obesity, as it does in inflammation caused by bacterial infection. Recently in the Pacific oyster *Crassostrea gigas*, it was described an HXK capable of recognizing various pathogen associated molecular patterns to activate an immune response against bacteria (Chen et al., 2021a), pointing out that the HXK receptor activity is necessary in different kingdoms. More research needs to be addressed to fully understand the different steps of the innate immune response mediated by the HXK and also to explore whether other organisms have a similar response against pathogens.

There is abundant information ascribing the role of GCK as a glucose sensor and single regulator of glucose homeostasis (Matschinsky and Wilson, 2019); however, GCK activity is important to the proposed Glc sensor activity, so it is difficult to determine if it is a moonlighting activity. Mice deficient in GCK (heterozygotes and homozygotes) and transgenic mice expressing GCK only in pancreatic β -cells were used to demonstrate that mice with only one GCK allele had elevated blood glucose levels and reduced insulin secretion, while *gck*-null mutants were lethal, and expression of GCK only in pancreatic β -cells was sufficient for survival. These experiments along with the investigation, description and screening of hundreds of nonsynonymous substitutions of GCK gene that causes inactivation or activation of GCK in many populations (Šimčíková et al., 2017) have been crucial to provide strong support to the hypothesis that GCK is a glucose sensor and a master regulator of glucose homeostasis (Grupe et al., 1995). Other experiments in mice provided detailed GCK links to diabetes mellitus and hypoglycemia diseases (Velho et al., 1992; Froguel et al., 1993; Grupe et al., 1995). There is also compelling evidence supporting the glucose-sensing function of GCK, and its role in glucose homeostasis in cells that participate in glucose homeostasis such as pancreatic α - and δ -cells, adrenal gland, glucose-sensitive neurons, enteroendocrine cells and anterior pituitary cells (Matschinsky and Wilson, 2019). Thus, GCK sensor activity in different cells seems to be vital to control the glucose homeostasis at the whole organism level and its mechanism of action is interconnected with the hormone production and secretion, so it is important to have a deeper understanding on how the sensor activity of GCK changes in different cells to provide strategies that might lead to manage the disorders associated with the low or high glucose blood levels.

Finally, for HXKI, Adams et al. (1991) showed evidence indicating that this hexokinase can also function as a protein-




FIGURE 4 | Plant hexokinases moonlighting functions. **(A)** At high glucose levels, AtHXK1 detaches from mitochondria and translocates to the nucleus in order to form a trimeric repressor complex with VHA-B1 and RPT5B; in turn, trimeric complex binds to putative transcription factors to bind on the CAB2 promoter to repress transcription of CAB2 (Cho et al., 2006b). **(B)** It has been suggested that the association of plant HXKs to MOM occurs through interaction with VDAC. In turn, this interaction functions as an inhibitor of the formation of mPTP and subsequently of PCD (Kim et al., 2006; Godbole et al., 2013). **(C)** Apple MdHXK1 seems to be involved in the regulation of anthocyanins biosynthesis by interacting and phosphorylating an anthocyanin-associated bHLH transcription factor to stabilize it and enhance transcription of the anthocyanin biosynthesis genes, increasing anthocyanin production (Hu et al., 2016). It is not clear where the interaction between MdHXK1 and bHLH occurs: MdHXK1 is likely to interact and phosphorylate bHLH on cytoplasm and then phosphorylated-bHLH enters the nucleus to activate transcription of anthocyanin genes. However, it is also possible that MdHXK1 enters to the nucleus to interact with bHLH and phosphorylate it. None of the possibilities has yet been described. **(D)** MdHXK1 also contributes to glucose-mediated salinity tolerance by interacting and phosphorylating a vacuolar Na⁺/H⁺ exchanger (Sun et al., 2018). **(E)** *Oryza sativa* OsHXK6 helps restore the fertility of Honglian (HL) cytoplasmic male sterility (CMS). OsHXK6 and RF6 protein form a complex inside mitochondria which is capable of cleaving atp6-orfH79 transcript at nucleotide 1,238, avoiding orfH79 mRNA from its translation and therefore reestablishing pollen fertility (Huang et al., 2015). Similarly, OsHXK5 deficiency causes male sterility, however, the expression of a non-catalytic version of this protein, OsHXK5G113D, is not able to rescue the male-sterile phenotype (Lee et al., 2019).

kinase capable of autophosphorylation and phosphorylation of histone 2A, solely in the absence of glucose, since when present, ATP is preferentially used for glucose-6-P production. Although there is still no physiological explanation for this protein-kinase activity, mammal HXKI is not the only hexokinase reported with this function: *S. cerevisiae* HXK2 have an autophosphorylation activity, which is stimulated *in vivo* by D-xylose (Fernández et al., 1988), the structural implications are discussed later. *Malus domestica* HXK1 has also been reported with a protein-kinase activity that apparently can lead to its self-phosphorylation *in vitro* and the participation of the protein in response to salt stress (Hu et al., 2016; Sun et al., 2018).

Plants

In plants, hexokinases display a wide range of functions from glucose sensing to regulation of programmed cell death (PCD). *Arabidopsis thaliana*, for instance, expresses three catalytic hexokinases: mitochondrial hexokinases AtHXK1 and AtHXK2, and plastid AtHXK3 (Sarowar et al., 2008; Aguilera-Alvarado and Sánchez-Nieto, 2017). From the three of them, AtHXK1 has been widely studied for its additional function as a glucose sensor, in which it was demonstrated that at high glucose levels AtHXK1 is one of the key factors to repress photosynthetic gene transcription and arrest seedling growth (Moore et al., 2003). The mechanism involved has been partially elucidated:

at high glucose levels, AtHXX1 form a trimeric repressor complex with the vacuolar H⁺-ATPase B1 (VHA-B1) and the 26S regulatory particle of proteasome subunit 5B (RPT5B); in turn, trimeric complex binds to putative transcription factors to bind on the CAB2 promoter to repress transcription of CAB2 (**Figure 4A**) (Cho et al., 2006b). It is unknown how AtHXX1 detaches from mitochondrial outer membrane and how AtHXX1, H⁺-ATPase B1, and RPT5B are translocated to the nucleus. However, it has been made clear that AtHXX1 renders its glucose-sensing function without needing its catalytic function, as observed by AtHXX1 catalytic mutants used to restore glucose-sensing phenotype on Arabidopsis null HXX1 mutants, *gin2-1* (Moore et al., 2003). Remarkably, *vha-B1* and *rpt5b* mutants display a glucose insensitive phenotype, even in the presence of AtHXX1, indicating that all three proteins are essential for glucose sensing and repression of photosynthetic genes (Cho et al., 2006b).

Kim and collaborators (2006) found that when *Nicotiana benthamiana* was silenced on its hexokinase 1, NbHXX1, necrotic lesions on leaves due to PCD were induced, suggesting a role of the HXX in the PCD (**Figure 4B**). On the other hand, Godbole and collaborators (2013) showed that overexpression of VDAC in tobacco cells protoplasts carried out PCD, but concomitant co-expression of mitochondrial hexokinase 3 from *N. benthamiana*, NbHXX3, rendered healthy protoplasts, suggesting a likely HXX-VDAC association in plant cells like that reported for mammals. This association between mitochondrial hexokinases and VDAC in plant cells has been hypothesized due to results found on beetroot cells, in which channeling of mitochondrial ATP to mitochondrial bound HXX through VDAC was evidenced (Alcántar-Aguirre et al., 2013). However, direct interaction between plant HXXs and VDAC remains to be demonstrated, and the mechanism involved in HXX-dependent programmed cell death pathway to be confirmed and described.

Additional functions of mitochondrial HXXs in plants have been described. Apple (*Malus domestica*) mitochondrial hexokinase 1, MdHXX1, seems to be involved in both regulation of anthocyanins biosynthesis and glucose-mediated salt stress tolerance (Hu et al., 2016; Sun et al., 2018). Apparently, at high exogenous glucose levels, MdHXX1 is capable to interact and phosphorylate an anthocyanin-associated bHLH transcription factor at its Ser361 site, in order to stabilize it and thus enhancing transcription of the anthocyanin biosynthesis genes, thereby increasing anthocyanin production (**Figure 4C**) (Hu et al., 2016). Additionally, it was reported that the same MdHXX1 contributes to glucose-mediated salinity tolerance by interacting and phosphorylating a vacuolar Na⁺/H⁺ exchanger, MdHXX1, at its Ser275 site (**Figure 4D**). According to the authors, phosphorylation improved the stability of MdHXX1 and enhanced its Na⁺/H⁺ transport activity when MdHXX1 was overexpressed (Sun et al., 2018). Remarkably, assays utilizing catalytic mutants of MdHXX1 showed that both functions seem to be independent of its glucose-phosphorylation activity. No other plant HXXs have been reported to have protein phosphorylating activity, further investigation needs to be done to determine if other plant HXXs have that capacity.

In rice, besides the ability of OsHXX6 to sense glucose (Cho et al., 2009), this protein can process an mRNA transcript (Huang et al., 2015). To carry out this last function, OsHXX6 interacts with RF6 protein (restorer of fertility six); this interaction occurs between each protein C-terminus. Once inside the mitochondria, the complex OsHXX6/RF6 cleaves the *atp6-orfH79* transcript at nucleotide 1,238, stopping *orfH79* mRNA from being translated into protein (**Figure 4E**). *orfH79* codes the ORFH79 protein, which interacts with a subunit of the mitochondrial complex III, impairing this complex activity and leading to pollen sterility. As no ORFH79 protein is synthesized, pollen becomes fertile again. RF6 and OsHXX6 are essential for processing the *atp6-orfH79* transcript, but neither of them can bind directly to mRNA, and for this reason, additional proteins may be involved in this mRNA processing activity (Huang et al., 2015). Although OsHXX5 is also capable of rescuing the rice plant from pollen sterility, it may not follow the same mechanism that OsHXX6 does. The fact that pollen from plants lacking OshHXX5 show reduced hexokinase activity and the catalytic inactive protein, OsHXX5G113D, does not restore fertility, may indicate that this additional activity relies on the hexokinase activity rather than on the sugar sensing ability (Lee et al., 2019).

Finally, hexokinases 1, 2 and 3 from *Jatropha curcas* (JcHXX1, JcHXX2 and JcHXX3) seems to play a role in a different type of abiotic stress (Wang et al., 2019); qRT-PCR analysis showed that when 14-days old seedlings of *J. curcas* were exposed to low temperature during different periods of time, the levels of the three JcHXXs were upregulated. In this case, more studies need to be carried out to clearly determine the moonlighting role of these hexokinases.

REGULATION MECHANISMS OF HEXOKINASES' MOONLIGHTING FUNCTIONS

The glucose phosphorylating activity of the HXX has been widely studied. Reviews on structural and enzyme kinetic features of HXXs are found in Cardenas et al., 1998 (HXXs from various organisms), Wilson, 2003 (mammal HXXs), Claeysen and Rivoal, 2007 (plant HXXs) and Steitz et al., 1981 (yeast HXXs). It is known that mammal HXXI-III glucose phosphorylating activity are product-regulated: physiological concentrations of G6P can allosterically inhibit its activity in a non-competitive way with respect to glucose, but competitive to ATP (Van Schaftingen, 2013). In maize, mitochondrial attached HXXs are less inhibited by G6P and ADP than the cytosolic HXXs (Aguilera-Alvarado et al., 2019), suggesting that their subcellular localization is a key component to control its enzyme activity.

On the other hand, HXXs' moonlighting activities are regulated by a wide range of mechanisms. A recent review analyzes the effect of redox state to change the function of diverse proteins in cancer cells, some of them are transformed in moonlighting proteins (Jiang et al., 2021). Understanding the existence of different multigene families with catalytic proteins, non-catalytic activities and pseudoenzymes which might appear

to have different functions through evolution, can shed light on how a protein shows a different function depending on the cell context or the presence of some structural or motif differences (Jeffery, 2020). In this review, we collect different and varied examples that show how hexokinases are regulated in accord with different cellular localization, oligomeric state, phosphorylation/dephosphorylation state, concentration, and cell type to maintain cellular homeostasis and function.

Transcriptional Regulation

In all living beings, to reduce the wasteful consumption of resources and energy a tight regulation of gene expression is crucial. For instance, in *S. cerevisiae* a complex regulation takes place to limit and specifically localize ScHXK2. In response to high glucose availability, ScHXK2 abundance increases and its localization changes. On glucose-supplemented medium, Mig1 sequesters ScHXK2 in the nucleus and the complex function as a gene repressor (Ahuatzi et al., 2004). Besides, the protein encoded by the *RGT1* gene takes part in the glucose-induced expression of hexose transporter (HXT) genes (Ozcan et al., 1996), Rgt1 together with the Med8 protein represses the ScHXK2 gene expression when glucose is absent. This repression occurs when Rgt1 binds to a motif (CGGAAAA) located 395 bp upstream of the start codon in the ScHXK2 promoter (Palomino et al., 2005).

In mammals, as described earlier, HXKII has a role in tumorigenesis, since its overexpression provides a metabolic benefit and suppression of apoptosis, allowing cancer cells to grow indefinitely (Pastorino and Hoek, 2008). In cancer cells, expression of HXKII is regulated by several transcriptional regulators such as microRNAs, long non-coding RNA and some transcription factors (Ma et al., 2021). For example, Zheng et al. (2017) reviewed that HXKII expression is controlled by miR-155/miR-143 cascade in breast cancer, while in renal cell carcinoma, HXKII is co-regulated by miR-143/miR-451 and in hepatocellular carcinoma cells it is regulated by miR-199a. On the other hand, it has been reported that long non-coding RNA urothelial carcinoma associated 1 (UCA1) overexpression promotes glycolysis by sequestering miR-203, which has an inhibitory effect on HXKII. This way, inhibition of miR-203 increases HXKII levels and, subsequently glycolysis rate (Liu et al., 2020). Finally, several transcription factors have been implicated in HXKII regulation in a variety of cancers, like hypoxia inducible factor 1 α (HIF-1 α) (Ma et al., 2021), the signal transducer and activator of transcription 3 (STAT3) and p53 (Rodríguez-Enríquez et al.,).

All these findings regarding the role of HXKII in cancer establishment and progression have been crucial to propose new therapeutic molecules that target this enzyme, as well as its regulators, to inhibit cancer cells proliferation. For instance, 3-bromopyruvate has been proved to break HXKII interaction with Apoptosis Inducing Factor (AIF) due to a covalent modification of mitochondria-bound HXKII (Wanga et al., 2009). On the other hand, methyl-jasmonate, a phytohormone related to defense mechanisms in plants, has been found to have anti-cancer properties due, in part, to its ability to remove HXKII from VDAC in apparently a selective manner. Detachment of HXKII

from mitochondria triggers the reduction in glycolysis flux and mitochondrial membrane potential, which subsequently leads to the activation of apoptosis in cancer cells (Cohen and Flescher, 2009; Cesari et al., 2014).

Posttranscriptional Regulation

Another way to regulate a gene expression is through posttranscriptional regulation mediated by AU-rich elements (AREs) (Shaw and Kamen, 1986) and ARE-binding proteins (Shyu and Wilkinson, 2000). Tristetraprolin (TTP), is a well-known ARE-binding protein able to promote the degradation of ARE-containing transcripts (Carballo et al., 1998; Chen et al., 2001; Lykke-Andersen and Wagner, 2005). In mammals, TTP is critical for the downregulation of HXKII expression in cancer cells as it enhances the degradation of the HXKII transcript, thus, suppressing the glycolytic capacity of cancer cells and reducing both the extracellular acidification rate and the oxygen consumption rate of cancer cells (Kim et al., 2019).

Post-Translational Modifications

Post-translational modifications (PTMs) on moonlighting proteins allow the cells, to certain extent, to deal with environmental changes (Jeffery, 2016). It has been shown that several hexokinase functions are regulated by the phosphorylation-dephosphorylation process. As discussed earlier, in mammals, HXKII-mitochondria interaction provides protection to mitochondrial-mediated death pathways (Machida et al., 2006). HXKII, but not other isoforms contain a serine/threonine kinase Akt consensus sequence and a TOS motif, thus retaining two active kinase domains (Tsai and Wilson, 1996). Kinase Akt is required to inhibit apoptosis (Majewski et al., 2004) by phosphorylating HXKII at Thr473 (Roberts and Miyamoyto, 2015), leading to an increase of the association of HXKII to mitochondria and the reduction of its dissociation induced by G6P. Thr-473 phosphorylation does not seem to influence G6P binding to HXKII, but it does stabilize the HXKII-mitochondrial interaction (Roberts et al., 2013). It has also been reported that insulin treatment in adipose and skeletal muscle cell lines increases HXKII mRNA and protein levels through Akt pathways (Osawa et al., 1996; Culbert and Tavaré, 2002). The phosphorylation of hexokinase II by Akt is accompanied by increased binding of the enzyme to the mitochondria and contributes to the anti-apoptotic effects of Akt against ischemia/reperfusion injury in cardiomyocytes (Roberts et al., 2013). These observations illustrate the ability of Akt to positively regulate the binding of hexokinase II to VDAC (Pastorino and Hoek, 2008).

On the other hand, HXKII abundance can be regulated by degradation mediated by its ubiquitination in liver cancer cells. HXKII ubiquitinated on the Lys63 residue by the E3 ligase TRAF6 (TNF receptor-associated factor 6), thus promoting its recognition by the autophagy receptor sequestosome 1 (SQSTM1, a human protein ubiquitin-binding protein), leading to its selective degradation, and promoting the autophagy-mediated suppression of glycolysis (Figure 3C) (Jiao et al., 2018).

In *S. cerevisiae* phosphorylated ScHXX2 exists *in vitro* as two oligomeric states, a monomer, and a dimer (Furman and Neet, 1983). *In vivo*, the relative abundance of either the monomer or the dimer is affected by the phosphorylated state of ScHXX2. ScHXX2 has two sites for phosphorylation: one on serine-14, and this modification *in vivo* is inversely related to the extracellular glucose concentration (Kriegel et al., 1994), and another for serine-158 (Heidrich et al., 1997). Snf1 kinase is responsible for phosphorylating ScHXX2 at serine 14, while Glc7- Reg1 acts as a phosphatase (Ahuatzi et al., 2004; Vega et al., 2016). Glucose, and so does maltose and sucrose, are able to trigger the loss of the phosphate group from phosphorylated ScHXX2. Unlike these three sugars, raffinose, galactose, and ethanol trigger the appearance of both phosphorylated and dephosphorylated forms of ScHXX2. Even though the phosphorylated state of hexokinase is affected by the carbon source, it does not depend on glucose repression (Fernández-García et al.).

ScHXX2 phosphorylation leads to: 1) Higher quantities of its monomeric form (Behlke et al., 1998), 2) increase of substrate affinity and inhibition by ATP (Golbik et al., 2001), and 3) stimulation of nucleocytoplasmic translocation of the phosphoenzyme (Fernández-García et al.). Despite the findings made by Fernández-García et al. (2012) and collaborators regarding the phosphorylation site of ScHXX2, Kaps et al. (2015) found out that it is not serine-14, but serine 15 and that either Ymr291w or a Ymr291w-dependent protein are the responsible kinases. According to the authors, when yeast cells grow on a low glucose containing media, higher amounts of Ymr291w are detected, compared with a high-glucose one, therefore Ymr291w is likely to contribute to glucose signaling in a situation of limited external glucose availability. However, further investigation regarding the phosphorylation is needed to understand the structural and functional aspects of yeasts hexokinases *in vivo* phosphorylation.

When wild type *S. cerevisiae* cells are grown under repressing conditions, the Reg1-Glc7 complex dephosphorylates Mig1-ScHXX2, so that they can be imported into the nucleus, where ScHXX2 crosses the nuclear envelope by the α/β -importin (Kap60/Kap95) pathway (Peláez et al., 2012) and once inside forms a Mig1-ScHXX2 complex to recruit other proteins (Mig2, Snf1, Snf4, Gal83, and Reg1). Once all the proteins have been recruited, the expression of genes associated with growth on non-fermentable carbon sources, like *SUC2*, is repressed (Figure 2) (Moreno and Herrero, 2002; Ahuatzi et al., 2007). On the other hand, when *S. cerevisiae* cells are grown on low-glucose levels, the interaction between ScHXX2 and Mig1 is abolished by phosphorylation of ScHXX2 and Mig1 by Snf1 at serine 14 and serine 311, respectively (Ahuatzi et al., 2007; Fernández-García et al.). Phosphorylation of both ScHXX2 and Mig1 results in their export from the nucleus to the cytoplasm by Xpo1 carrier protein (Peláez et al., 2009) and the subsequent disassembly of the repressor complex, allowing transcription of genes encoding metabolism proteins for non-fermentable carbon sources (Figure 2).

Nevertheless, the complex formation at the *SUC2* gene induced by ScHXX2 does not always depend on the

phosphorylation state of ScHXX2, thus indicating that the phosphorylation state of ScHXX2 regulates only its location (nucleus or cytoplasm), but not its assembly and disassembly to the *SUC2* repressor complex (Vega et al., 2016). ScHXX2 assembly to the *SUC2* promoter may be regulated by conformational changes induced by glucose levels. As illustrated in Figures 1B,E, ScHXX2 is formed by the large and the small domains, separated by a deep cleft where the active site is located; when glucose is not present, ScHXX2 adopts an open, catalytically inactive conformation, but in the presence of glucose, the cleft closes due to a movement of both domains. Vega and collaborators (2016) found a correlation between ScHXX2 catalytically inactive conformation and its binding to the *SUC2* repressor complex, while an open conformation of ScHXX2 correlates with its dissociation from the *SUC2* promoter. With these findings, the authors suggest that the signaling activity of ScHXX2 is linked to conformational changes induced by glucose levels. However, it remains to be investigated if there are other factors or interactors in the process.

A distinct feature of some hexokinases that might contribute to its moonlighting activity is their subcellular location: some of them, like *S. cerevisiae* HXX2 (Fernández-García et al.) and maize HXX7 and HXX8 (Aguilera-Alvarado et al., 2019) are located at the cytoplasm; others, in glycosomes, like *T. brucei* HXX2 (Blattner et al., 1995; Michaels et al., 2006); many in the mitochondrial outer membrane or MOM, like human HXXII (Arora and Pedersen, 1988; Dubey et al., 2016), *A. thaliana* HXX1 (Cho et al., 2006a) and apple HXX1 (Sun et al., 2018); or in the chloroplast, like *P. patens* HXX1 (Olsson et al., 2003) and *Oryza sativa* HXX4 (Cho et al., 2006a). And even there are HXXs that are translocated from one subcellular compartment to the nucleus, like AtHXX1 (Cho et al., 2006b) and ScHXX2 (Vega et al., 2016). The subcellular location depends partially on the presence of membrane anchor sequences, mitochondrial-targeting sequence (MTS), nuclear localization signals (NLS), or dual targeting signals (Cho et al., 2006b; Aguilera-Alvarado and Sánchez-Nieto, 2017). However post-translational modifications (PTM) might take place to change the localization of the protein from one compartment to the nucleus (Hashiguchi and Komatsu, 2017; Chen et al., 2021b), as occurs with ScHXX2. However, there are so many questions that need to be addressed to know the exact mechanism of the translocation process of various hexokinases to the nucleus, such as the type of PTM that takes place to change the protein destination. Hexokinase translocation between subcellular compartments, mainly to the nucleus, suggests that moonlighting proteins act as signaling intermediaries between organelles to promote rapid response to changes in the environment, substrate, or product concentrations and direct link between metabolic activity to genome integrity and gene expression (Kotiadis et al., 2014; Monaghan and Whitmarsh, 2015).

Protein-Protein Interactions

As mentioned above, mammal HXXII attaches to MOM by interactions to VDAC and/or cyclophilin D, apparently as a strategy to couple oxidative phosphorylation with carbon

metabolism through ATP channeling and to inhibit the formation of mPTP and subsequently, apoptosis programmed cell death (**Figure 3A**) (Mathupala et al., 2006; Pastorino and Hoek, 2008). Cyclophilin D, an immunophilin with *cis-trans* isomerase activity, affects mitochondrial binding of HXKII (Machida et al., 2006). Earlier studies have shown that the anti-apoptotic effects of cyclophilin D may be exerted by the stabilization of HXKII binding to mitochondria (Broekemeier et al., 1989). Inactivation of cyclophilin D with cyclosporine A or knock-down of its expression utilizing siRNA caused a release of mitochondrially bound HXKII. Moreover, the anti-apoptotic effects of cyclophilin D were overridden by the forced detachment of hexokinase II from the mitochondria. Such observations agree with the concept that HXKII prevents access of proapoptotic proteins such as Bax to the mitochondria or inhibits its pore-forming abilities. However, it is presently unclear how cyclophilin D activity regulates the binding of HXKII to the mitochondria (Pastorino and Hoek, 2008).

Product Regulation

HXKII is required for the autophagy-mediated regulation of glycolysis, maintaining cellular homeostasis by modulating the level of glycolysis (Ros and Schulze, 2013), as mentioned earlier. HXKII and autophagy are linked through mTORC1 (Roberts et al., 2014); in cardiomyocytes, treated with 2-deoxy-D-glucose (2-DG), a non-metabolizable glucose analog, mTORC1 activity is decreased (Chantranupong and Sabatini, 2016), induction of autophagy is inhibited, and cell death induced by glucose deprivation is also increased (Tan and Miyamoto, 2016). Through its TOS motifs, HXKII binds to the Raptor domain of mTORC1 and this binding is largely increased by glucose absence (Schalm and Blenis, 2002; Nojima et al., 2003). Interestingly, HXK glycolytic and autophagic activities are regulated by G6P (Roberts et al., 2014); decreased levels of G6P promote HXKII-mTORC1 interaction and facilitate autophagy, while high levels of G6P inhibit HXKII-mTORC1 association and therefore cellular metabolism and growth can happen. This regulating function of HXKII occurs somewhere else but mitochondria, so it is likely that G6P changes intracellular localization of HXKII to regulate its association to mTORC1 (**Figure 3B**) (Roberts and Miyamoto, 2015).

WHAT REMAINS TO BE UNRAVELED?

The available evidence from moonlighting proteins indicates that these moonlighting functions are not conserved across life domains, and the same happens with HXKs. The reported moonlighting activities are broad, from glucose sensor to hemoglobin binding protein, protein phosphorylating enzyme, immune receptor, apoptosis inhibitor and autophagy control regulator, among others. However, many of these proposed functions remain to be confirmed or need a deeper understanding.

As mentioned before, the glucose sensor activity is the most common moonlighting function of HXKs, though many aspects

of the mechanism of sugar sensing in each kingdom need further investigation. Concerning ScHXK2, moonlighting functions are already known (Herrero et al., 1998; Randez-Gil et al., 1998; Ahuatzzi et al., 2004; Fernández-García et al.), but efforts are still to be made to demonstrate the signal perceived by ScHXK2 to assemble and disassemble the repressor complex in the *SUC2* promoter. If the conformational change in the ScHXK2 state is the signal needed to start the process (Vega et al., 2016) it will be a powerful tool to predict moonlighting proteins and their functions in the future.

It is important to emphasize that sugars are critical regulators of many processes in plants, such as germination, seedling development, photosynthesis, flowering, senescence, stress responses and carbon and nitrogen metabolism (Moore et al., 2003). It seems that moonlighting hexokinases may also be critical regulators in many of these processes. It is noticeable that it appears that moonlighting proteins are subservient to their “canonical” function so that they seem to take advantage of enzymatic activity or subcellular location to perform another function somehow related to the first one. In this way, plant mitochondrial HXKs seem to exploit their ability to phosphorylate glucose to sense it (or vice versa) and then generate a response to glucose levels that allow HXKs to translocate to the nucleus to repress or activate genes. However, it is unknown how AtHXK1 and its two partners are translocated from their original location to the nucleus to form the trimeric repressor complex (Cho et al., 2006b). It is possible that the complex is assembled inside the nucleus. If it is the case, it remains to know how the other two proteins in the complex perceive the glucose signal to start the translocation process to the nucleus.

Intriguingly, MdHXK1 upon high glucose levels activates anthocyanin biosynthesis genes (Hu et al., 2016) and increases the Na^+/H^+ transport activity at the vacuole to respond to saline stress (Sun et al., 2018). For these two additional functions described for MdHXK1, experiments need to be carried out to define the subcellular localization of MdHXK1-bHLH1 interaction to activate anthocyanin biosynthesis (Hu et al., 2016): does the interaction occurs in the cytoplasm and then both proteins are imported to the nucleus? Or does MdHXK1 translocates from the mitochondria to the nucleus and then interact and phosphorylates MdbHLH1? Furthermore, about MdHXK1 response to salt stress (Sun et al., 2018), what is the pathway involved in MdHXK1 migration to vacuole to interact with MdNHX1? What are the specific signals MdHXK1 perceives to perform one function or another specifically? Which cellular partners take part in each function so that the specific response the cell requires is generated?

Mitochondrial HXKs seem to display different functions during abiotic stress conditions, but in plants are likely to transduce also stress signals by pathogens, as overexpression of AtHXK1 and AtHXK2 induces PR genes expression (Sarowar et al., 2008).

Many reports regarding mitochondrial protein translocation to the nucleus, namely retrograde transport, exhibit nuclear localization upon cellular and environmental stimuli (Lionaki et al., 2016). Most of these dual-localization proteins contain a

mitochondrial targeting sequence and a nuclear localization signal. However, some moonlighting proteins lack peptide signals; Cytochrome C is an example of it (González-Arzola et al., 2019). Due to lack of NLS, maybe cytochrome C could be exported to cytosol under non-apoptotic conditions, without mitochondrial membrane permeabilization, and then use a nuclear import mechanism by binding to nucleoporins (Jagot-Lacoussiere et al., 2015). Regarding plant mitochondrial HXKs, for several of them were reported the presence of an NLS within their amino acid sequence, adjacent to their MTS. Nevertheless, among species of higher plants as *Arabidopsis*, *Zea mays*, *Oryza sativa*, or the moss *Physcomitrella patens*, the organelle targeting signals could be ambiguous (Xu et al., 2013). For example, four hexokinases of moss *Physcomitrella patens*: PpHXK2, PpHXK3, PpHXK7 and, PpHXK11 have an ambiguous targeting signal that leads to dual protein targeting to the outer membrane of mitochondria and the outer envelope of chloroplast (Nilsson et al., 2011; Xu et al., 2013). PpHXK4 contains a truncated N-terminus that differs from the plastid or mitochondrial hexokinases, and it is in the cytosol and nucleus. In *O. sativa*, only HXK7 and HXK8 are predicted with N-terminus truncated but do not resemble the PpHXK4 sequence (Nilsson et al., 2011).

Changes in the subcellular localization also can be due to post-transcriptional modifications related to the redox cellular state. Some proteins respond to the redox state, changing their subcellular localization. ROS and redox state affect the protein import machinery (Bölter et al., 2015; Ling and Jarvis, 2015), and also lead to changes in microtubule orientation in plant cells (Dang et al., 2018) and is well documented that in cancer cells low mitochondria activity promotes ROS production which affect several metabolic proteins including some glycolytic enzymes and moonlighting proteins, observations that are reviewed by Jiang et al. (2021). Redox post-transcriptional modifications seem to be site-specific by Cys residues and subject to temporal and spatial control (Foyer et al., 2020). Due to the perception of sugar signals or methyl-jasmonate related to mitochondrial ROS production, AtHXK1 could be translocated between the mitochondrion and nucleus (Claeyssen and Rivoal, 2007; Xiang et al., 2011), after being subject to redox modification, since there is a Cys-159 potentially oxidable near the HXK1 sugar-binding site (McConell et al., 2019). Finally, suggestions have been made around a glucose-induced conformational change, as suggested for ScHXK2 (Vega et al., 2016) or a variation on its oligomeric state (Moore, 2004).

Mammal HXKII is also a mitochondrial protein with non-canonical functions, like inhibition of apoptosis programmed cell death, demonstrated from research on cancer and neurodegeneration (Pastorino et al., 2002; Mathupala et al., 2006). Nevertheless, how HXKII and its different interaction partners modulate apoptosis in normal conditions is still a matter of investigation: which are the interaction partners and signals participating in the process? Where does HXKII-VDAC interaction occurs, and which signals modulate HXKII detachment from VDAC or MOM? On the other hand, regarding HXKII role in tumorigenesis, more investigation regarding the molecular signals and mechanisms that cause re-programming of HXKII to function aberrantly in cancer cells are

needed in order to find compelling cancer therapeutics. It is not known if HXKII moonlighting ability is somehow related to its susceptibility to be re-programmed during cancer establishment; thus, keep digging on HXK moonlighting functions may shed light on the understanding of this possibility. As to HXKII regulation of autophagy, it is still unclear where HXKII translocates and what is the mechanism involved for this protein to interact with mTORC1 to activate the process. Is this function, in fact, a moonlighting function? Finally, referring to glucose homeostasis regulation by glucokinase (HXKIV), more data and experiments need to be carried out, as it is still a matter of discussion (Matschinsky and Wilson, 2019). Are suggestions around GCK as a single glucose sensor for glucose homeostasis regulation in mammals genuinely reliable? Does the system require additional factors for glucose sensitivity, as already discussed in plants?

CONCLUDING REMARKS

HXKs have been extensively studied in mammals and yeasts. Nowadays, we have a significant amount of information to keep studying, analyzing, and investigating the impact of the moonlighting HXK functions in those species' physiology. However, in plants, there is still a lack of information regarding not only hexokinases but moonlighting proteins in general. As sessile organisms, plants have been found to have large multigene protein families. Besides that, they have a complex phytohormone regulation and robust control of carbon and nitrogen sources. Thus, the presence of moonlighting proteins in plants may be more abundant and interconnected with different signaling pathways.

Thus, research is still needed to elucidate the regulatory mechanisms for the translocation of mitochondrial hexokinases to their different target organelles and to identify the pathway components and possible interaction partners that facilitate the responses to which plant mitochondrial hexokinases are associated.

AUTHOR CONTRIBUTIONS

CR-S, LM-M, BK-D, and SS-N developed the initial concept. CR-S, LM-M, and SS-N wrote the abstract. CR-S wrote the introduction, LM-M wrote "How are moonlighting proteins studied?", "Moonlighting proteins in the glycolytic pathway," and developed HXK crystal structure details and figures. LMM and ML-C wrote "hexokinase." CR-S wrote "Moonlighting proteins in life domains." CR-S and LM-M wrote "Hexokinases moonlighting activity in diverse organisms." CR-S, BK-D, ML-C, and SS-N wrote "Regulation mechanisms of hexokinases' moonlighting functions." CR-S and ML-C wrote "What remains to be unraveled?" and "Concluding remarks." AB-P and ML-C prepared the figures. LM-M created **Table 1**. CR-S, LM-M, AB-P, BK-D, ML-C, and SSN reviewed and corrected the whole manuscript. CR-S formatted the manuscript for submission. All authors reviewed the final manuscript.

FUNDING

This work was supported by Consejo Nacional de Ciencia y Tecnología (No. A1-S-17269 and FOP02-2021-316424), Universidad Nacional Autónoma de México (PAPIIT IN225220) and Facultad de Química (Grant Number 50009125). CR-S, Grant 590108; LM-M, Grant 1002862; AB-P, Grant 957302; and ML-C, Grant 429315, received a

scholarship from Consejo Nacional de Ciencia y Tecnología (CONACyT).

ACKNOWLEDGMENTS

We are grateful to Dr. José Antonio Pedroza-García for his valuable comments and suggestions for the improvement of the manuscript.

REFERENCES

- Abolhassani Rad, S., Clayton, E. J., Cornelius, E. J., Howes, T. R., and Kohalmi, S. E. (2018). Moonlighting Proteins: Putting the Spotlight on Enzymes. *Plant Signal. Behav.* 13 (10), 1–6. doi:10.1080/15592324.2018.1517075
- Adams, V., Griffin, L. D., Gelb, B. D., and McCabe, E. R. B. (1991). Protein Kinase Activity of Rat Brain Hexokinase. *Biochem. Biophysical Res. Commun.* 177 (3), 1101–1106. doi:10.1016/0006-291x(91)90652-n
- Adamu, G., Brown, L., Schiffman, J., and Iannaccone, A. (2011). Diversity in Autoimmunity against Retinal, Neuronal, and Axonal Antigens in Acquired Neuro-Retinopathy. *J. Ophthalm. Inflamm. Infect.* 1 (3), 111–121. doi:10.1007/s12348-011-0028-8
- Adamus, G. (2017). Impact of Autoantibodies against Glycolytic Enzymes on Pathogenesis of Autoimmune Retinopathy and Other Autoimmune Disorders. *Front. Immunol.* 8, 505. doi:10.3389/fimmu.2017.00505
- Affourtit, C., Alberts, B., Barlow, J., Carré, J. E., and Wynne, A. G. (2018). Control of Pancreatic β -cell Bioenergetics. *Biochem. Soc. Trans.* 46, 555–564. doi:10.1042/BST20170505
- Agarwal, S., Kulshreshtha, P., Bambah Mukku, D., and Bhatnagar, R. (2008). α -Enolase Binds to Human Plasminogen on the Surface of Bacillus Anthracis. *Biochim. Biophys. Acta (Bba) - Proteins Proteomics* 1784, 986–994. doi:10.1016/j.bbapap.2008.03.017
- Aguilera-Alvarado, G. P., Guevara-García, Á. A., Estrada-Antolín, S. A., and Sánchez-Nieto, S. (2019). Biochemical Properties and Subcellular Localization of Six Members of the HXK Family in maize and its Metabolic Contribution to Embryo Germination. *BMC Plant Biol.* 19 (1), 27. doi:10.1186/s12870-018-1605-x
- Aguilera-Alvarado, G. P., and Sánchez-Nieto, S. (2017). Plant Hexokinases Are Multifaceted Proteins. *Plant Cell Physiol.* 58 (7), 1151–1160. doi:10.1093/pcp/pcx062
- Ahuatzi, D., Herrero, P., de la Cera, T., and Moreno, F. (2004). The Glucose-Regulated Nuclear Localization of Hexokinase 2 in *Saccharomyces cerevisiae* Is Mig1-dependent. *J. Biol. Chem.* 279 (14), 14440–14446. doi:10.1074/jbc.M313431200
- Ahuatzi, D., Riera, A., Pela'ez, R., Herrero, P., and Moreno, F. (2007). Hxk2 Regulates the Phosphorylation State of Mig1 and Therefore its Nucleocytoplasmic Distribution. *J. Biol. Chem.* 282, 4485–4493. doi:10.1074/jbc.M606854200
- Alcántar-Aguirre, F. C., Chagolla, A., Tiessen, A., Délano, J. P., and González de la Vara, L. E. (2013). ATP Produced by Oxidative Phosphorylation Is Channeled toward Hexokinase Bound to Mitochondrial Porin (VDAC) in Beetroots (Beta Vulgaris). *Planta* 237, 1571–1583. doi:10.1007/s00425-013-1866-4
- Alexander, N. J., Proctor, R. H., and McCormick, S. P. (2009). Genes, Gene Clusters, and Biosynthesis of Trichothecenes and Fumonisin in *Fusarium*. *Toxin Rev.* 28, 198–215. doi:10.3390/toxins1101003610.1080/15569540903092142
- Amela, I., Cedano, J., and Querol, E. (2007). Pathogen Proteins Eliciting Antibodies Do Not Share Epitopes with Host Proteins: A Bioinformatics Approach. *PLoS One* 2 (6), e512. doi:10.1371/journal.pone.0000512
- Anderson, C., Zucker, F., and Steitz, T. (1979). Space-filling Models of Kinase Clefs and Conformation Changes. *Science* 204 (4391), 375–380. doi:10.1126/science.220706
- Arnér, E. S. J., and Holmgren, A. (2000). Physiological Functions of Thioredoxin and Thioredoxin Reductase. *Eur. J. Biochem.* 267 (20), 6102–6109. doi:10.1046/j.1432-1327.2000.01701.x
- Arora, K. K., and Pedersen, P. L. (1988). Functional Significance of Mitochondrial Bound Hexokinase in Tumor Cell Metabolism. Evidence for Preferential phosphorylation of Glucose by Intramitochondrially Generated ATP. *J. Biol. Chem.* 263 (33), 17422–17428. doi:10.1016/S0021-9258(19)77853-3
- Artymiuk, P. J., and Green, J. (2006). The Double Life of Aconitase. *Structure* 14 (1), 2–4. doi:10.1016/j.str.2005.12.001
- Bahia, D. (2017). A New Trick for a Conserved Enzyme: Mevalonate Kinase, a Glycosomal Enzyme, Can Be Secreted by Trypanosoma Cruzi and Modulate Cell Invasion and Signaling. Is it Another Moonlighting Enzyme? *Front. Cell. Infect. Microbiol.* 7, 426. doi:10.3389/fcimb.2017.00426
- Baines, C. P., Kaiser, R. A., Purcell, N. H., Blair, N. S., Osinska, H., Hambleton, M. A., et al. (2005). Loss of Cyclophilin D Reveals a Critical Role for Mitochondrial Permeability Transition in Cell Death. *Nature* 434, 658–662. doi:10.1038/nature03434
- Behlke, J., Heidrich, K., Naumann, M., Müller, E.-C., Otto, A., Reuter, R., et al. (1998). Hexokinase 2 from *Saccharomyces Cerevisiae*: Regulation of Oligomeric Structure by *In Vivo* Phosphorylation at Serine-14†. *Biochemistry* 37 (34), 11989–11995. doi:10.1021/bi980914m
- Bennett, W. S., Jr, and Steitz, T. A. (1980). Structure of a Complex between Yeast Hexokinase A and Glucose. *J. Mol. Biol.* 140 (2), 183–209. doi:10.1016/0022-2836(80)90102-3
- Betz, C., and Hall, M. N. (2013). Where Is mTOR and what Is it Doing There? *J. Cell Biol.* 203 (4), 563–574. doi:10.1083/jcb.201306041
- Blattner, J., Dörsam, H., and Clayton, C. E. (1995). Function of N-Terminal Import Signals in Trypanosome Microbodies. *FEBS Lett.* 360 (3), 310–314. doi:10.1016/0014-5793(95)00128-v
- Blau, K., Portnoi, M., Shagan, M., Kaganovich, A., Rom, S., Kafka, D., et al. (2007). Flamingo Cadherin: A Putative Host Receptor for *Streptococcus Pneumoniae*. *J. Infect. Dis.* 195 (12), 1828–1837. doi:10.1086/518038
- Bölter, B., Soll, J., and Schwenkert, S. (2015). Redox Meets Protein Trafficking. *Biochim. Biophys. Acta (Bba) - Bioenerg.* 1847 (9), 949–956. doi:10.1016/j.bbabi.2015.01.010
- Boone, T. J., Burnham, C.-A. D., and Tyrrell, G. J. (2011). Binding of Group B Streptococcal Phosphoglycerate Kinase to Plasminogen and Actin. *Microb. Pathogenesis* 51 (4), 255–261. doi:10.1016/j.micpath.2011.06.005
- Bosch, J., Buscaglia, C. A., Krumm, B., Ingason, B. P., Lucas, R., Roach, C., et al. (2007). Aldolase Provides an Unusual Binding Site for Thrombospondin-Related Anonymous Protein in the Invasion Machinery of the Malaria Parasite. *Proc. Natl. Acad. Sci.* 104 (17), 7015–7020. doi:10.1073/pnas.0605301104
- Boukouris, A. E., Zervopoulos, S. D., and Michelakis, E. D. (2016). Metabolic Enzymes Moonlighting in the Nucleus: Metabolic Regulation of Gene Transcription. *Trends Biochem. Sci.* 41 (8), 712–730. doi:10.1016/j.tibs.2016.05.013
- Broekemeier, K. M., Dempsey, M. E., and Pfeiffer, D. R. (1989). Cyclosporin A Is a Potent Inhibitor of the Inner Membrane Permeability Transition in Liver Mitochondria. *J. Biol. Chem.* 264 (14), 7826–7830. doi:10.1016/S0021-9258(18)83116-7
- Brown, J., Miller, D. M., Holloway, M. T., and Leve, G. D. (1967). Hexokinase Isoenzymes in Liver and Adipose Tissue of Man and Dog. *Science* 155 (3759), 205–207. doi:10.1126/science.155.3759.205
- Brown, R. W. B., Collingridge, P. W., Gull, K., Rigden, D. J., and Ginger, M. L. (2014). Evidence for Loss of a Partial Flagellar Glycolytic Pathway during Trypanosomatid Evolution. *PLoS ONE* 9 (7), e103026. doi:10.1371/journal.pone.0103026
- Bustos, D. M., and Iglesias, A. A. (2000). The Kinetic Properties of Liver Glucokinase and its Function in Glucose Physiology as a Model for the Comprehensive Study of Enzymes' Kinetic Parameters and Reversible Inhibitors. *Biochem. Mol. Biol. Edu.* 28, 332–337. doi:10.1016/S1470-8175(00)00027-810.1111/j.1539-3429.2000.tb00186.x

- Candela, M., Bergmann, S., Vici, M., Vitali, B., Turrone, S., Eikmanns, B. J., et al. (2007). Binding of Human Plasminogen to Bifidobacterium. *J. Bacteriol.* 189, 5929–5936. doi:10.1128/JB.00159-07
- Carballo, E., Lai, W. S., and Blackshear, P. J. (1998). Feedback Inhibition of Macrophage Tumor Necrosis Factor- Production by Tristetraprolin. *Science* 281 (5379), 1001–1005. doi:10.1126/science.281.5379.1001
- Cárdenas, M. L., Cornish-Bowden, A., and Ureta, T. (1998). Evolution and Regulatory Role of the Hexokinases. *Biochim. Biophys. Acta (Bba) - Mol. Cell Res.* 1401 (3), 242–264. doi:10.1016/s0167-4889(97)00150-x
- Castello, A., Hentze, M. W., and Preiss, T. (2015). Metabolic Enzymes Enjoying New Partnerships as RNA-Binding Proteins. *Trends Endocrinol. Metab.* 26 (12), 746–757. doi:10.1016/j.tem.2015.09.012
- Cesari, I. M., Carvalho, E., Figueiredo Rodrigues, M., Mendonça, B. d. S., Amôedo, N. D., and Rumjanek, F. D. (2014). Methyl Jasmonate: Putative Mechanisms of Action on Cancer Cells Cycle, Metabolism, and Apoptosis. *Int. J. Cell Biol.* 2014, 1–25. doi:10.1155/2014/572097
- Chantranupong, L., and Sabatini, D. M. (2016). The TORC1 Pathway to Protein Destruction. *Nature* 536 (7615), 155–156. doi:10.1038/nature18919
- Chávez-munguía, B., Segovia-Gamboa, N., Salazar-Villatoro, L., Omaña-molina, M., Espinosa-Cantellano, M., and Martínez-palomo, A. (2011). Naegleria Fowleri: Enolase Is Expressed during Cyst Differentiation. *J. Eukaryot. Microbiol.* 58 (5), 463–468. doi:10.1111/j.1550-7408.2011.00574.x
- Chen, C.-Y., Gherzi, R., Ong, S.-E., Chan, E. L., Raijmakers, R., Puijijn, G. J. M., et al. (2001). AU Binding Proteins Recruit the Exosome to Degrade ARE-Containing mRNAs. *Cell* 107 (4), 451–464. doi:10.1016/s0092-8674(01)00578-5
- Chen, X., Liu, Z., Gu, Y., Zhang, Y., Liu, Y., Wang, L., et al. (2021b). A Hexokinase from the Oyster *Crassostrea gigas* Is Involved in Immune Recognition as a Pattern Recognition Receptor. *Developmental Comp. Immunol.* 122, 104083. doi:10.1016/j.dci.2021.104083
- Chen, Y., Wang, Y., Yang, J., Zhou, W., and Dai, S. (2021a). Exploring the Diversity of Plant Proteome. *J. Integr. Plant Biol.* doi:10.1111/jipb.13087
- Cho, J.-I., Ryoo, N., Eom, J.-S., Lee, D.-W., Kim, H.-B., Jeong, S.-W., et al. (2009). Role of the Rice Hexokinases OsHXK5 and OsHXK6 as Glucose Sensors. *Plant Physiol.* 149 (2), 745–759. doi:10.1104/pp.108.131227
- Cho, J.-I., Ryoo, N., KoLee, S. S. K., Lee, S.-K., Lee, J., Jung, K.-H., et al. (2006a). Structure, Expression, and Functional Analysis of the Hexokinase Gene Family in rice (*Oryza Sativa* L.). *Planta* 224 (3), 598–611. doi:10.1007/s00425-006-0251-y
- Cho, Y.-H., Yoo, S.-D., and Sheen, J. (2006b). Regulatory Functions of Nuclear Hexokinase1 Complex in Glucose Signaling. *Cell* 127 (3), 579–589. doi:10.1016/j.cell.2006.09.028
- Claeyssen, É., and Rivoal, J. (2007). Isozymes of Plant Hexokinase: Occurrence, Properties and Functions. *Phytochemistry* 68 (6), 709–731. doi:10.1016/j.phytochem.2006.12.001
- Cohen, S., and Fleischer, E. (2009). Methyl Jasmonate: A Plant Stress Hormone as an Anti-cancer Drug. *Phytochemistry* 70 (13-14), 1600–1609. doi:10.1016/j.phytochem.2009.06.007
- Collinridge, P. W., Brown, R. W. B., and Ginger, M. L. (2010). Moonlighting Enzymes in Parasitic Protozoa. *Parasitology* 137 (9), 1467–1475. doi:10.1017/S0031182010000259
- Colowick, S. P., and Kalckar, H. M. (1943). The Role of Myokinase in Transphosphorylations. *J. Biol. Chem.* 148, 117–126. doi:10.1016/S0021-9258(18)72324-7
- Commichau, F. M., Rothe, F. M., Herzberg, C., Wagner, E., Hellwig, D., Lehnik-Habrink, M., et al. (2009). Novel Activities of Glycolytic Enzymes in *Bacillus Subtilis*. *Mol. Cell Proteomics* 8 (6), 1350–1360. doi:10.1074/mcp.M800546-MCP200
- Copley, S. D. (2014). An Evolutionary Perspective on Protein Moonlighting. *Biochem. Soc. Trans.* 42 (6), 1684–1691. doi:10.1042/BST20140245
- Crowe, J. D., Siewright, I. K., Auld, G. C., Moore, N. R., Gow, N. A. R., and Booth, N. A. (2003). Candida Albicans Binds Human Plasminogen: Identification of Eight Plasminogen-Binding Proteins. *Mol. Microbiol.* 47 (6), 1637–1651. doi:10.1046/j.1365-2958.2003.03390.x
- Culbert, A. A., and Tavaré, J. M. (2002). Multiple Signalling Pathways Mediate Insulin-Stimulated Gene Expression in 3T3-L1 Adipocytes. *Biochim. Biophys. Acta (Bba) - Gene Struct. Expr.* 1578, 43–50. doi:10.1016/s0167-4781(02)00481-5
- Dang, X., Yu, P., Li, Y., Yang, Y., Zhang, Y., Ren, H., et al. (2018). Reactive Oxygen Species Mediate Conical Cell Shaping in *Arabidopsis thaliana* Petals. *Plos Genet.* 14 (10), e1007705. doi:10.1371/journal.pgen.1007705
- Dastoor, Z., and Dreyer, J. L. (2001). Potential Role of Nuclear Translocation of Glyceraldehyde-3-Phosphate Dehydrogenase in Apoptosis and Oxidative Stress. *J. Cel Sci.* 114 (9), 1643–1653. doi:10.1242/jcs.114.9.1643
- Desai, K., Sullards, M. C., Allegood, J., Wang, E., Schmelz, E. M., Hartl, M., et al. (2002). Fumonisin and Fumonisin Analogs as Inhibitors of Ceramide Synthase and Inducers of Apoptosis. *Biochim. Biophys. Acta (Bba) - Mol. Cell Biol. Lipids* 1585, 188–192. doi:10.1016/s1388-1981(02)00340-2
- Dubey, A. K., Godbole, A., and Mathew, M. K. (2016). Regulation of VDAC Trafficking Modulates Cell Death. *Cel Death Discov.* 2 (2), 16085. doi:10.1038/cddiscovery.2016.85
- Egea, L., Aguilera, L., Giménez, R., Sorolla, M. A., Aguilar, J., Badía, J., et al. (2007). Role of Secreted Glyceraldehyde-3-Phosphate Dehydrogenase in the Infection Mechanism of Enterohemorrhagic and Enteropathogenic *Escherichia coli*: Interaction of the Extracellular Enzyme with Human Plasminogen and Fibrinogen. *Int. J. Biochem. Cel Biol.* 39 (6), 1190–1203. doi:10.1016/j.biocel.2007.03.008
- Enzo, E., Santinon, G., Pocaterra, A., Aragona, M., Bresolin, S., Forcato, M., et al. (2015). Aerobic Glycolysis Tunes YAP/TAZ Transcriptional Activity. *EMBO J.* 34 (10), 1349–1370. doi:10.15252/embj.201490379
- Feng, J., Zhao, S., Chen, X., Wang, W., Dong, W., Chen, J., et al. (2015). Biochemical and Structural Study of Arabidopsis Hexokinase 1. *Acta Cryst. D Biol. Crystallogr.* 71 (2), 367–375. doi:10.1107/S1399004714026091
- Fernández, R., Herrero, P., Fernández, E., Fernández, T., López-Boado, Y. S., and Moreno, F. (1988). Autophosphorylation of Yeast Hexokinase PII. *Microbiology* 134, 2493–2498. doi:10.1099/00221287-134-9-2493
- Fernández-García, P., Peláez, R., Herrero, P., and Moreno, F. (2012). Phosphorylation of Yeast Hexokinase 2 Regulates its Nucleocytoplasmic Shuttling. *J. Biol. Chem.* 287 (50), 42151–42164. doi:10.1074/jbc.M112.401679
- Fox, D., and Smulian, A. G. (2001). Plasminogen-binding Activity of Enolase in the Opportunistic Pathogen *Pneumocystis Carinii*. *Med. Mycol.* 39 (6), 495–507. doi:10.1080/mmy.39.6.495.507
- Foyer, C. H., Baker, A., Wright, M., Sparkes, I. A., Mhamdi, A., Schippers, J. H. M., et al. (2020). On the Move: Redox-dependent Protein Relocation in Plants. *J. Exp. Bot.* 71 (2), 620–631. doi:10.1093/jxb/erz330
- Franco-Serrano, L., Cedano, J., Perez-Pons, J. A., Mozo-Villarias, A., Piñol, J., Amela, I., et al. (2018). A Hypothesis Explaining Why So many Pathogen Virulence Proteins Are Moonlighting Proteins. *Pathog. Dis.* 76 (5), fty046. doi:10.1093/femsdpd/fty046
- Froguel, P., Zouali, H., Vionnet, N., Velho, G., Vaxillaire, M., Sun, F., et al. (1993). Familial Hyperglycemia Due to Mutations in Glucokinase -- Definition of a Subtype of Diabetes Mellitus. *N. Engl. J. Med.* 328 (10), 697–702. doi:10.1056/NEJM199303113281005
- Furman, T. C., and Neet, K. E. (1983). Association Equilibria and Reacting Enzyme Gel Filtration of Yeast Hexokinase. *J. Biol. Chem.* 258 (8), 4930–4936. doi:10.1016/s0021-9258(18)32517-1
- Gancedo, C., Flores, C.-L., and Gancedo, J. M. (2014). Evolution of Moonlighting Proteins: Insight from Yeasts. *Biochem. Soc. Trans.* 42 (6), 1715–1719. doi:10.1042/BST20140199
- Gancedo, C., Flores, C.-L., and Gancedo, J. M. (2016). The Expanding Landscape of Moonlighting Proteins in Yeasts. *Microbiol. Mol. Biol. Rev.* 80 (3), 765–777. doi:10.1128/MMBR.00012-16
- Gancedo, C., and Flores, C.-L. (2008). Moonlighting Proteins in Yeasts. *Mmbr* 72 (1), 197–210. doi:10.1128/MMBR.00036-07
- Gao, X., Wang, H., Yang, J. J., Liu, X., and Liu, Z. R. (2012). Piruvate Kinase M2 Regulates Gene Transcription by Acting as a Protein Kinase. *Mol. Cell.* 45, 598–609. doi:10.1016/j.molcel.2012.01.001
- Godbole, A., Dubey, A. K., Reddy, P. S., Udayakumar, M., and Mathew, M. K. (2013). Mitochondrial VDAC and Hexokinase Together Modulate Plant Programmed Cell Death. *Protoplasma* 250 (4), 875–884. doi:10.1007/s00709-012-0470-y
- Goëb, V., Thomas-L'Otellier, M., Daveau, R., Charlionet, R., Fardellone, P., Le Loët, X., et al. (2009). Candidate Autoantigens Identified by Mass Spectrometry in Early Rheumatoid Arthritis Are Chaperones and Citrullinated Glycolytic Enzymes. *Arthritis Res. Ther.* 11 (2), R38. doi:10.1186/ar2644

- Golbik, R., Naumann, M., Otto, A., Müller, E.-C., Behlke, J., Reuter, R., et al. (2001). Regulation of Phosphotransferase Activity of Hexokinase 2 from *Saccharomyces cerevisiae* by Modification at Serine-14†. *Biochemistry* 40 (4), 1083–1090. doi:10.1021/bi001745k
- Goldin, N., Arzoine, L., Heyfets, A., Israelson, A., Zaslavsky, Z., Bravman, T., et al. (2008). Methyl Jasmonate Binds to and Detaches Mitochondria-Bound Hexokinase. *Oncogene* 27, 4636–4643. doi:10.1038/onc.2008.108
- Goldstein, J. L., and Brown, M. S. (1990). Regulation of the Mevalonate Pathway. *Nature* 343 (6257), 425–430. doi:10.1038/343425a0
- González, C., Ureta, T., Sánchez, R., and Niemeyer, H. (1964). Multiple Molecular Forms of ATP: Hexose 6-phosphotransferase from Rat Liver. *Biochem. Biophysical Res. Commun.* 16 (4), 347–352. doi:10.1016/0006-291x(64)90038-5
- González-Arzola, K., Velázquez-Cruz, A., Guerra-Castellano, A., Casado-Combreras, M., Pérez-Mejías, G., Díaz-Quintana, A., et al. (2019). New Moonlighting Functions of Mitochondrial Cytochrome C in the Cytoplasm and Nucleus. *FEBS Lett.* 593 (22), 3101–3119. doi:10.1002/1873-3468.13655
- Grosse, F., Nasheuer, H.-P., Scholtissek, S., and Schomburg, U. (1986). Lactate Dehydrogenase and Glyceraldehyde-Phosphate Dehydrogenase Are Single-Stranded DNA-Binding Proteins that Affect the DNA-Polymerase-Alpha-Primase Complex. *Eur. J. Biochem.* 160 (3), 459–467. doi:10.1111/j.1432-1033.1986.tb10062.x
- Grupe, A., Hultgren, B., Ryan, A., Ma, Y. H., Bauer, M., and Stewart, T. A. (1995). Transgenic Knockouts Reveal a Critical Requirement for Pancreatic β Cell Glucokinase in Maintaining Glucose Homeostasis. *Cell* 83 (1), 69–78. doi:10.1016/0092-8674(95)00235-x
- Guo, C., Ludvik, A. E., Arlotto, M. E., Hayes, M. G., Armstrong, L. L., Scholtens, D. M., et al. (2015). Coordinated Regulatory Variation Associated with Gestational Hyperglycaemia Regulates Expression of the Novel Hexokinase HKDC1. *Nat. Commun.* 6, 6069. doi:10.1038/ncomms7069
- Gupta, V., and Bamezai, R. N. K. (2010). Human Pyruvate Kinase M2: a Multifunctional Protein. *Protein Sci.* 19 (11), 2031–2044. doi:10.1002/pro.505
- Hagemann, L., Gründel, A., Jacobs, E., and Dumke, R. (2017). The Surface-Displayed Chaperones GroEL and DnaK of *Mycoplasma pneumoniae* Interact with Human Plasminogen and Components of the Extracellular Matrix. *FEMS J. Pathog. Dis.* 75 (3). doi:10.1093/femspd/ftx017
- Han, S., Wang, Y., Zheng, X., Jia, Q., Zhao, J., Bai, F., et al. (2015). Cytoplasmic Glyceraldehyde-3-Phosphate Dehydrogenases Interact with ATG3 to Negatively Regulate Autophagy and Immunity in *Nicotiana glauca*. *Plant Cell* 27 (4), 1316–1331. doi:10.1105/tpc.114.134692
- Hapeta, P., Szczepańska, P., Neuveglise, C., and Lazar, Z. (2021). A 37-amino Acid Loop in the *Yarrowia lipolytica* Hexokinase Impacts its Activity and Affinity and Modulates Gene Expression. *Sci. Rep.* 11 (1), 6412. doi:10.1038/s41598-021-85837-8
- Hara, M. R., Agrawal, N., Kim, S. F., Cascio, M. B., Fujimuro, M., Ozeki, Y., et al. (2005). S-nitrosylated GAPDH Initiates Apoptotic Cell Death by Nuclear Translocation Following Siah1 Binding. *Nat. Cell Biol.* 7 (7), 665–674. doi:10.1038/ncb1268
- Hashiguchi, A., and Komatsu, S. (2017). Posttranslational Modifications and Plant-Environment Interaction. *Methods Enzymol.* 586, 97–113. doi:10.1016/bs.mie.2016.09.030
- Hattori, T., Takei, N., Mizuno, Y., Kato, K., and Kohsaka, S. (1995). Neurotrophic and Neuroprotective Effects of Neuron-specific Enolase on Cultured Neurons from Embryonic Rat Brain. *Neurosci. Res.* 21 (3), 191–198. doi:10.1016/0168-0102(94)00849-b
- He, C., Chen, J., Wang, H., Wan, Y., Zhou, J., Dan, Z., et al. (2019). Crystal Structures of rice Hexokinase 6 with a Series of Substrates Shed Light on its Enzymatic Mechanism. *Biochem. Biophysical Res. Commun.* 515 (4), 614–620. doi:10.1016/j.bbrc.2019.05.139
- Heidrich, K., Otto, A., Behlke, J., Rush, J., Wenzel, K.-W., and Kriegel, T. (1997). Autophosphorylation–Inactivation Site of Hexokinase 2 in *Saccharomyces cerevisiae*†. *Biochemistry* 36, 1960–1964. doi:10.1021/bi9623643
- Henderson, B., and Martin, A. (2011). Bacterial Virulence in the Moonlight: Multitasking Bacterial Moonlighting Proteins Are Virulence Determinants in Infectious Disease. *Infect. Immun.* 79 (9), 3476–3491. doi:10.1128/IAI.00179-11
- Hendriks, W., Mulders, J. W., Bibby, M. A., Slingsby, C., Bloemendal, H., and de Jong, W. W. (1988). Duck Lens Epsilon-Crystallin and Lactate Dehydrogenase B4 Are Identical: a Single-Copy Gene Product with Two Distinct Functions. *Proc. Natl. Acad. Sci.* 85, 7114–7118. doi:10.1073/pnas.85.19.7114
- Heneberg, P. (2019). Redox Regulation of Hexokinases. *Antioxid. Redox Signaling* 30 (3), 415–442. doi:10.1089/ars.2017.72510.1089/ars.2017.7255
- Henry, E., Fung, N., Liu, J., Drakakaki, G., and Coaker, G. (2015). Beyond Glycolysis: GAPDHs Are Multi-Functional Enzymes Involved in Regulation of ROS, Autophagy, and Plant Immune Responses. *Plos Genet.* 11 (4), e1005199. doi:10.1371/journal.pgen.1005199
- Hentze, M. W., and Preiss, T. (2010). The REM Phase of Gene Regulation. *Trends Biochem. Sci.* 35 (8), 423–426. doi:10.1016/j.tibs.2010.05.009
- Hernández, S., Ferragut, G., Amela, I., Perez-Pons, J., Piñol, J., Mozo-Villarias, A., et al. (2014). MultitaskProtDB: a Database of Multitasking Proteins. *Nucl. Acids Res.* 42, D517–D520. doi:10.1093/nar/gkt1153
- Hernández, S., Franco, L., Calvo, A., Ferragut, G., Hermoso, A., Amela, I., et al. (2015). Bioinformatics and Moonlighting Proteins. *Front. Bioeng. Biotechnol.* 3 (90), 1–16. doi:10.3389/fbioe.2015.0009010.3389/fbioe.2015.00090
- Herrero, P., Fernández, R., and Moreno, F. (1989). The Hexokinase Isoenzyme PII of *Saccharomyces cerevisiae* Is a Protein Kinase. *Microbiology* 135 (5), 1209–1216. doi:10.1099/00221287-135-5-1209
- Herrero, P., Martínez-Campa, C., and Moreno, F. (1998). The Hexokinase 2 Protein Participates in Regulatory DNA-Protein Complexes Necessary for Glucose Repression of the SUC2 Gene in *Saccharomyces cerevisiae*. *FEBS Lett.* 434 (2), 71–76. doi:10.1016/s0014-5793(98)00872-2
- Hu, D.-G., Sun, C.-H., Zhang, Q.-Y., An, J.-P., You, C.-X., and Hao, Y.-J. (2016). Glucose Sensor MdHXK1 Phosphorylates and Stabilizes MdbHLH3 to Promote Anthocyanin Biosynthesis in Apple. *Plos Genet.* 12 (8), e1006273. doi:10.1371/journal.pgen.1006273
- Huang, W., Yu, C., Hu, J., Wang, L., Dan, Z., Zhou, W., et al. (2015). Pentatricopeptide-repeat Family Protein RF6 Functions with Hexokinase 6 to rescue rice Cytoplasmic Male Sterility. *Proc. Natl. Acad. Sci. U.S.A.* 112 (48), 14984–14989. doi:10.1073/pnas.1511748112
- Huberts, D. H. E. W., and van der Klei, I. J. (2010). Moonlighting Proteins: An Intriguing Mode of Multitasking. *Biochim. Biophys. Acta (Bba) - Mol. Cell Res.* 1803, 520–525. doi:10.1016/j.bbamcr.2010.01.022
- Hurmalainen, V., Edelman, S., Antikainen, J., Baumann, M., Lähteenmäki, K., and Korhonen, T. K. (2007). Extracellular Proteins of *Lactobacillus crispatus* Enhance Activation of Human Plasminogen. *Microbiology (Reading)* 153 (4), 1112–1122. doi:10.1099/mic.0.2006/000901-0
- Ikeda, R., and Ichikawa, T. (2014). Interaction of Surface Molecules on *Cryptococcus neoformans* with Plasminogen. *FEMS Yeast Res.* 14 (3), 445–450. doi:10.1111/1567-1364.12131
- Ikeda, Y., Yamaji, R., Irie, K., Kioka, N., and Murakami, A. (2012). Glyceraldehyde-3-phosphate Dehydrogenase Regulates Cyclooxygenase-2 Expression by Targeting mRNA Stability. *Arch. Biochem. Biophys.* 528 (2), 141–147. doi:10.1016/j.abb.2012.09.004
- Imle, R., Wang, B.-T., Stützenberger, N., Birkenhagen, J., Tandon, A., Carl, M., et al. (2019). ADP-dependent Glucokinase Regulates Energy Metabolism via ER-Localized Glucose Sensing. *Sci. Rep.* 9, 14248. doi:10.1038/s41598-019-50566-6
- Irwin, D. M., and Tan, H. (2014). Evolution of Glucose Utilization: Glucokinase and Glucokinase Regulator Protein. *Mol. Phylogenet. Evol.* 70, 195–203. doi:10.1016/j.ympev.2013.09.016
- Jagot-Lacoussiere, L., Faye, A., Bruzzoni-Giovanelli, H., Villoutreix, B. O., Rain, J.-C., and Poyet, J.-L. (2015). DNA Damage-Induced Nuclear Translocation of Apaf-1 Is Mediated by Nucleoporin Nup107. *Cell Cycle* 14 (8), 1242–1251. doi:10.1080/15384101.2015.1014148
- Jang, J. C., León, P., Zhou, L., and Sheen, J. (1997). Hexokinase as a Sugar Sensor in Higher Plants. *Plant Cell* 9, 5–19. doi:10.1105/tpc.9.1.5
- Jeffery, C. J. (2020). Enzymes, Pseudoenzymes, and Moonlighting Proteins: Diversity of Function in Protein Superfamilies. *FEBS J.* 287, 4141–4149. doi:10.1111/febs.15446
- Jeffery, C. J. (2004). Molecular Mechanisms for Multitasking: Recent crystal Structures of Moonlighting Proteins. *Curr. Opin. Struct. Biol.* 14 (6), 663–668. doi:10.1016/j.sbi.2004.10.001
- Jeffery, C. J. (1999). Moonlighting Proteins. *Trends Biochem. Sci.* 24 (1), 8–11. doi:10.1016/s0968-0004(98)01335-8
- Jeffery, C. J. (2009). Moonlighting Proteins—An Update. *Mol. Biosyst.* 5 (4), 345–350. doi:10.1039/b900658n

- Jeffery, C. J. (2016). Protein Species and Moonlighting Proteins: Very Small Changes in a Protein's Covalent Structure Can Change its Biochemical Function. *J. Proteomics* 134, 19–24. doi:10.1016/j.jprot.2015.10.003
- Jiang, J., Peng, L., Wang, K., and Huang, C. (2021). Moonlighting Metabolic Enzymes in Cancer: New Perspectives on the Redox Code. *Antioxid. Redox Signaling* 34 (13), 979–1003. doi:10.1089/ars.2020.8123
- Jiao, L., Zhang, H.-L., Li, D.-D., Yang, K.-L., Tang, J., Li, X., et al. (2018). Regulation of Glycolytic Metabolism by Autophagy in Liver Cancer Involves Selective Autophagic Degradation of HK2 (Hexokinase 2). *Autophagy* 14 (4), 671–684. doi:10.1080/15548627.2017.1381804
- Johnston, M. (1999). Feasting, Fasting and Fermenting: Glucose Sensing in Yeast and Other Cells. *Trends Genet.* 15 (1), 29–33. doi:10.1016/s0168-9525(98)01637-0
- Joice, A. C., Lyda, T. L., Sayce, A. C., Verplaetse, E., Morris, M. T., Michels, P. A. M., et al. (2012). Extra-glycosomal Localisation of Trypanosoma Brucei Hexokinase 2. *Int. J. Parasitol.* 42, 401–409. doi:10.1016/j.ijpara.2012.02.008
- Jones, M. N., and Holt, R. G. (2007). Cloning and Characterization of an α -enolase of the Oral Pathogen Streptococcus Mutans that Binds Human Plasminogen. *Biochem. Biophysical Res. Commun.* 364 (4), 924–929. doi:10.1016/j.bbrc.2007.10.098
- Jong, A. Y., Chen, S. H. M., Stins, M. F., Kim, K. S., Tuan, T.-L., and Huang, S.-H. (2003). Binding of Candida Albicans Enolase to Plasmin(ogen) Results in Enhanced Invasion of Human Brain Microvascular Endothelial Cells. *J. Med. Microbiol.* 52 (8), 615–622. doi:10.1099/jmm.0.05060-0
- Kainulainen, V., Loimaranta, V., Pekkala, A., Edelman, S., Antikainen, J., Kylväjä, R., et al. (2012). Glutamine Synthetase and Glucose-6-Phosphate Isomerase Are Adhesive Moonlighting Proteins of Lactobacillus Crispatus Released by Epithelial Cathelicidin LL-37. *J. Bacteriol.* 194 (10), 2509–2519. doi:10.1128/JB.06704-11
- Kamata, K., Mitsuya, M., Nishimura, T., Eiki, J.-i., and Nagata, Y. (2004). Structural Basis for Allosteric Regulation of the Monomeric Allosteric Enzyme Human Glucokinase. *Structure* 12 (3), 429–438. doi:10.1016/j.str.2004.02.005
- Kao, A. W., Noda, Y., Johnson, J. H., Pessin, J. E., and Saltiel, A. R. (1999). Aldolase Mediates the Association of F-Actin with the Insulin-Responsive Glucose Transporter GLUT4. *J. Biol. Chem.* 274 (25), 17742–17747. doi:10.1074/jbc.274.25.17742
- Kaps, S., Kettner, K., Migotti, R., Kanashova, T., Krause, U., Rödel, G., et al. (2015). Protein Kinase Ymr291w/Tda1 Is Essential for Glucose Signaling in *Saccharomyces cerevisiae* on the Level of Hexokinase Isoenzyme SchXk2 Phosphorylation*. *J. Biol. Chem.* 290 (10), 6243–6255. doi:10.1074/jbc.M114.595074
- Katakura, Y., Sano, R., Hashimoto, T., Ninomiya, K., and Shioya, S. (2010). Lactic Acid Bacteria Display on the Cell Surface Cytosolic Proteins that Recognize Yeast Mannan. *Appl. Microbiol. Biotechnol.* 86 (1), 319–326. doi:10.1007/s00253-009-2295-y
- Kelley, N., Jeltama, D., Duan, Y., and He, Y. (2019). The NLRP3 Inflammasome: An Overview of Mechanisms of Activation and Regulation. *Ijms* 20 (13), 3328. doi:10.3390/ijms20133328
- Kennedy, M. C., Mende-Mueller, L., Blondin, G. A., and Beinert, H. (1992). Purification and Characterization of Cytosolic Aconitase from Beef Liver and its Relationship to the Iron-Responsive Element Binding Protein. *Proc. Natl. Acad. Sci.* 89 (24), 11730–11734. doi:10.1073/pnas.89.24.11730
- Kim, D. J., Vo, M.-T., Choi, S. H., Lee, J.-H., Jeong, S. Y., Hong, C. H., et al. (2019). Tristetraprolin-mediated Hexokinase 2 Expression Regulation Contributes to Glycolysis in Cancer Cells. *MBoC* 30 (5), 542–553. doi:10.1091/mbc.E18-09-0606
- Kim, H.-B., Cho, J.-I., Ryoo, N., Shin, D.-H., Park, Y.-I., Hwang, Y.-S., et al. (2016). Role of rice Cytosolic Hexokinase OsHXK7 in Sugar Signaling and Metabolism. *J. Integr. Plant Biol.* 58 (2), 127–135. doi:10.1111/jipb.12366
- Kim, H., Smith, J. E., Ridenour, J. B., Woloshuk, C. P., and Bluhm, B. H. (2011). HXK1 Regulates Carbon Catabolism, Sporulation, Fumonisin B1 Production and Pathogenesis in *Fusarium Verticillioides*. *Microbiol.* 157, 2658–2669. doi:10.1099/mic.0.052506-0
- Kim, J.-w., and Dang, C. V. (2005). Multifaceted Roles of Glycolytic Enzymes. *Trends Biochem. Sci.* 30 (3), 142–150. doi:10.1016/j.tibs.2005.01.005
- Kim, M., Lim, J.-H., Ahn, C. S., Park, K., Kim, G. T., Kim, W. T., et al. (2006). Mitochondria-Associated Hexokinases Play a Role in the Control of Programmed Cell Death in Nicotiana Benthamiana. *Plant Cell* 18 (9), 2341–2355. doi:10.1105/tpc.106.041509
- Kim, S.-C., Guo, L., and Wang, X. (2020). Nuclear Moonlighting of Cytosolic Glyceraldehyde-3-Phosphate Dehydrogenase Regulates Arabidopsis Response to Heat Stress. *Nat. Commun.* 11 (1), 3439. doi:10.1038/s41467-020-17311-4
- Kim, S. G., Hoffman, G. R., Poulgiannis, G., Buel, G. R., Jang, Y. J., Lee, K. W., et al. (2013a). Metabolic Stress Controls mTORC1 Lysosomal Localization and Dimerization by Regulating the TTT-Ruvb1/2 Complex. *Mol. Cell* 49, 172–185. doi:10.1016/j.molcel.2012.10.003
- Kim, Y.-M., Heinzel, N., Giese, J.-O., Koeber, J., Melzer, M., Rutten, T., et al. (2013b). A Dual Role of Tobacco Hexokinase 1 in Primary Metabolism and Sugar Sensing. *Plant Cell Environ* 36 (7), 1311–1327. doi:10.1111/pce.12060
- Kimura, A., Sakurai, T., Koumura, A., Yamada, M., Hayashi, Y., Tanaka, Y., et al. (2010). High Prevalence of Autoantibodies against Phosphoglycerate Mutase 1 in Patients with Autoimmune central Nervous System Diseases. *J. Neuroimmunology* 219 (1-2), 105–108. doi:10.1016/j.jneuroim.2009.11.014
- Kinby, B., Booth, N. A., and Svensäter, G. (2008). Plasminogen Binding by Oral Streptococci from Dental Plaque and Inflammatory Lesions. *Microbiology* 154 (3), 924–931. doi:10.1099/mic.0.2007/013235-0
- Kiri, A., and Goldspink, G. (2002). RNA-protein Interactions of the 3' Untranslated Regions of Myosin Heavy Chain Transcripts. *J. Muscle Res. Cell Motil.* 23, 119–129. doi:10.1023/a:1020211729728
- Kotiadis, V. N., Duchon, M. R., and Osellame, L. D. (2014). Mitochondrial Quality Control and Communications with the Nucleus Are Important in Maintaining Mitochondrial Function and Cell Health. *Biochim. Biophys. Acta (Bba) - Gen. Subjects* 1840 (4), 1254–1265. doi:10.1016/j.bbagen.2013.10.041
- Kriegel, T. M., Rush, J., Vojtek, A. B., Clifton, D., and Fraenkel, D. G. (1994). *In Vivo* phosphorylation Site of Hexokinase 2 in *Saccharomyces cerevisiae*. *Biochemistry* 33, 148–152. doi:10.1021/bi00167a019
- Kuettner, E. B., Kettner, K., Keim, A., Svergun, D. I., Volke, D., Singer, D., et al. (2010). Crystal Structure of Hexokinase Khxk1 of *Kluyveromyces Lactis*: A Molecular Basis for Understanding the Control of Yeast Hexokinase Functions via Covalent Modification and Oligomerization. *J. Biol. Chem.* 285, 41019–41033. doi:10.1074/jbc.M110.185850
- Kundu, M. (2014). Too Sweet for Autophagy: Hexokinase Inhibition of mTORC1 Activates Autophagy. *Mol. Cell* 53 (4), 517–518. doi:10.1016/j.molcel.2014.02.009
- Kuser, P. R., Krauchenco, S., Antunes, O. A. C., and Polikarpov, I. (2000). The High Resolution Crystal Structure of Yeast Hexokinase PII with the Correct Primary Sequence Provides New Insights into its Mechanism of Action. *J. Biol. Chem.* 275 (27), 20814–20821. doi:10.1074/jbc.M910412199
- Larion, M., Hansen, A. L., Zhang, F., Bruschweiler-Li, L., Tugarinov, V., Miller, B. G., et al. (2015). Kinetic Cooperativity in Human Pancreatic Glucokinase Originates from Millisecond Dynamics of the Small Domain. *Angew. Chem. Int. Ed.* 54, 8129–8132. doi:10.1002/anie.201501204
- Larion, M., and Miller, B. G. (2012). Homotropic Allosteric Regulation in Monomeric Mammalian Glucokinase. *Arch. Biochem. Biophys.* 519 (2), 103–111. doi:10.1016/j.abb.2011.11.007
- Laurian, R., Dementhon, K., Doumèche, B., Soulard, A., Noel, T., Lemaire, M., et al. (2019). Hexokinase and Glucokinases Are Essential for Fitness and Virulence in the Pathogenic Yeast *Candida Albicans*. *Front. Microbiol.* 10, 327. doi:10.3389/fmicb.2019.00327
- Lay, A. J., Jiang, X.-M., Kisker, O., Flynn, E., Underwood, A., Condron, R., et al. (2000). Phosphoglycerate Kinase Acts in Tumour Angiogenesis as a Disulphide Reductase. *Nature* 408 (6814), 869–873. doi:10.1038/35048596
- Lee, S.-K., Kim, H., Cho, J.-I., Nguyen, C. D., Moon, S., Park, J. E., et al. (2019). Deficiency of rice Hexokinase HXK5 Impairs Synthesis and Utilization of Starch in Pollen Grains and Causes Male Sterility. *J. Exp. Bot.* 71 (1), 116–125. doi:10.1093/jxb/erz436
- Li, S., Swanson, S. K., Gogol, M., Florens, L., Washburn, M. P., Workman, J. L., et al. (2015). Serine and SAM Responsive Complex SESAME Regulates Histone Modification Crosstalk by Sensing Cellular Metabolism. *Mol. Cell* 60 (3), 408–421. doi:10.1016/j.molcel.2015.09.024
- Ling, Q., and Jarvis, P. (2015). Regulation of Chloroplast Protein Import by the Ubiquitin E3 Ligase SP1 Is Important for Stress Tolerance in Plants. *Curr. Biol.* 25 (19), 2527–2534. doi:10.1016/j.cub.2015.08.015
- Lionaki, E., Gkikas, I., and Tavernarakis, N. (2016). Differential Protein Distribution between the Nucleus and Mitochondria: Implications in Aging. *Front. Genet.* 7, 162. doi:10.3389/fgene.2016.00162
- Liu, H.-E., Shi, H.-H., and Luo, X.-J. (2020). Upregulated Long Noncoding RNA UCA1 Enhances Warburg Effect via miR-203/HK2 Axis in Esophageal Cancer. *J. Oncol.* 2020, 1–11. doi:10.1155/2020/8847687

- Liu, H., Zeng, H., Yao, Q., Yuan, J., Zhang, Y., Qiu, D., et al. (2012). Steinernema Glaseri Surface Enolase: Molecular Cloning, Biological Characterization, and Role in Host Immune Suppression. *Mol. Biochem. Parasitol.* 185 (2), 89–98. doi:10.1016/j.molbiopara.2012.06.006
- Lu, G.-T., Xie, J.-R., Chen, L., Hu, J.-R., An, S.-Q., Su, H.-Z., et al. (2009). Glyceraldehyde-3-phosphate Dehydrogenase of *Xanthomonas Campestris* P.v. *Campestris* Is Required for Extracellular Polysaccharide Production and Full Virulence. *Microbiology* 155 (5), 1602–1612. doi:10.1099/mic.0.023762-0
- Lu, M., Holliday, L. S., Zhang, L., Dunn, W. A., Jr, and Gluck, S. L. (2001). Interaction between Aldolase and Vacuolar H⁺-ATPase. *J. Biol. Chem.* 276 (32), 30407–30413. doi:10.1074/jbc.M008768200
- Ludvik, A. E., Pusec, C. M., Priyadarshini, M., Angueira, A. R., Guo, C., Lo, A., et al. (2016). HKDC1 Is a Novel Hexokinase Involved in Whole-Body Glucose Use. *Endocrinology* 157 (9), 3452–3461. doi:10.1210/en.2016-1288
- Lykke-Andersen, J., and Wagner, E. (2005). Recruitment and Activation of mRNA Decay Enzymes by Two ARE-Mediated Decay Activation Domains in the Proteins TTP and BRF-1. *Genes Development* 19, 351–361. doi:10.1101/gad.1282305
- Ma, R., Wu, Y., Li, S., and Yu, X. (2021). Interplay between Glucose Metabolism and Chromatin Modifications in Cancer. *Front. Cell Dev. Biol.* 9, 654337. doi:10.3389/fcell.2021.654337
- Maccallini, P., Bonin, S., and Trevisan, G. (2018). Autoimmunity against a Glycolytic Enzyme as a Possible Cause for Persistent Symptoms in Lyme Disease. *Med. Hypotheses* 110, 1–8. doi:10.1016/j.mehy.2017.10.024
- Machida, K., Ohta, Y., and Osada, H. (2006). Suppression of Apoptosis by Cyclophilin D via Stabilization of Hexokinase II Mitochondrial Binding in Cancer Cells. *J. Biol. Chem.* 281, 14314–14320. doi:10.1074/jbc.M513297200
- Madureira, P., Baptista, M., Vieira, M., Magalhães, V., Camelo, A., Oliveira, L., et al. (2007). *Streptococcus agalactiae* GAPDH Is a Virulence-Associated Immunomodulatory Protein. *J. Immunol.* 178 (3), 1379–1387. doi:10.4049/jimmunol.178.3.1379
- Magrys, A., Anekonda, T., Ren, G., and Adamus, G. (2007). The Role of Anti- α -enolase Autoantibodies in Pathogenicity of Autoimmune-Mediated Retinopathy. *J. Clin. Immunol.* 27 (2), 181–192. doi:10.1007/s10875-006-9065-8
- Majewski, N., Nogueira, V., Bhaskar, P., Coy, P. E., Skeen, J. E., Gottlob, K., et al. (2004). Hexokinase-mitochondria Interaction Mediated by Akt Is Required to Inhibit Apoptosis in the Presence or Absence of Bax and Bak. *Mol. Cell* 16 (5), 819–830. doi:10.1016/j.molcel.2004.11.014
- Mani, M., Chen, C., Amblee, V., Liu, H., Mathur, T., Zwicke, G., et al. (2015). MoonProt: a Database for Proteins that Are Known to Moonlight. *Nucleic Acids Res.* 43, D277–D282. doi:10.1093/nar/gku954
- Marcos, C. M., Oliveira, H. C. d., da Silva, J. d. F., Assato, P. C. A., Fusco-Almeida, A. M., and Mendes-Giannini, M. J. S. (2014). The Multifaceted Roles of Metabolic Enzymes in the Paracoccidioides Species Complex. *Front. Microbiol.* 5 (719), 1–10. doi:10.3389/fmicb.2014.00719
- Mathupala, S. P., Ko, Y. H., and Pedersen, P. L. (2006). Hexokinase II: Cancer's Double-Edged Sword Acting as Both Facilitator and Gatekeeper of Malignancy when Bound to Mitochondria. *Oncogene* 25 (34), 4777–4786. doi:10.1038/sj.onc.1209603
- Matschinsky, F. M., and Randle, P. J. (1993). Evolution of the Glucokinase Glucose Sensor Paradigm for Pancreatic Beta Cells. *Diabetologia* 36 (11), 1215–1217. doi:10.1007/BF00401072
- Matschinsky, F. M., and Wilson, D. F. (2019). The central Role of Glucokinase in Glucose Homeostasis: a Perspective 50 Years after Demonstrating the Presence of the Enzyme in Islets of Langerhans. *Front. Physiol.* 10, 148. doi:10.3389/fphys.2019.00148
- Matta, S. K., Agarwal, S., and Bhatnagar, R. (2010). Surface Localized and Extracellular Glyceraldehyde-3-Phosphate Dehydrogenase of *Bacillus Anthracis* Is a Plasminogen Binding Protein. *Biochim. Biophys. Acta (Bba) - Proteins Proteomics* 1804 (11), 2111–2120. doi:10.1016/j.bbapap.2010.08.004
- McConnell, E. W., Berg, P., Westlake, T. J., Wilson, K. M., Popescu, G. V., Hicks, L. M., et al. (2019). Proteome-wide Analysis of Cysteine Reactivity during Effector-Triggered Immunity. *Plant Physiol.* 179 (4), 1248–1264. doi:10.1104/pp.18.01194
- Meyer-Siegler, K., Mauro, D. J., Seal, G., Wurzer, J., deRiel, J. K., and Sirover, M. A. (1991). A Human Nuclear Uracil DNA Glycosylase Is the 37-kDa Subunit of Glyceraldehyde-3-Phosphate Dehydrogenase. *Proc. Natl. Acad. Sci.* 88 (19), 8460–8464. doi:10.1073/pnas.88.19.8460
- Meyerhof, O. (1927). Über die enzymatische Milchsäurebildung im Muskelextrakt. III. Mitteilung: Die Milchsäurebildung aus den gärfähigen Hexosen. *Biochem. Z.* 183, 176–215.
- Meyerhof, O. (1935). Über die Wirkungsweise der Hexokinase. *Naturwissenschaften* 23, 850–851. doi:10.1007/bf01491989
- Michels, P. A., Bringaud, F., Herman, M., and Hannaert, V. (2006). Metabolic Functions of Glycosomes in Trypanosomatids. *Biochim. Biophys. Acta* 1763, 1463–1477. doi:10.1016/j.bbamcr.2006.08.019
- Modun, B., Morrissey, J., and Williams, P. (2000). The Staphylococcal Transferrin Receptor: a Glycolytic Enzyme with Novel Functions. *Trends Microbiol.* 8 (5), 231–237. doi:10.1016/s0966-842x(00)01728-5
- Monaghan, R. M., and Whitmarsh, A. J. (2015). Mitochondrial Proteins Moonlighting in the Nucleus. *Trends Biochem. Sci.* 40 (12), 728–735. doi:10.1016/j.tibs.2015.10.003
- Moore, B. d. (2004). Bifunctional and Moonlighting Enzymes: Lighting the Way to Regulatory Control. *Trends Plant Sci.* 9 (5), 221–228. doi:10.1016/j.tplants.2004.03.005
- Moore, B., Zhou, L., Rolland, F., Hall, Q., Cheng, W. H., Liu, Y. X., et al. (2003). Role of the Arabidopsis Glucose Sensor HXK1 in Nutrient, Light, and Hormonal Signaling. *Science* 300 (5617), 332–336. doi:10.1126/science.1080585
- Moreno, F., and Herrero, P. (2002). The Hexokinase 2-dependent Glucose Signal Transduction Pathway of *Saccharomyces Cerevisiae*. *FEMS Microbiol. Rev.* 26, 83–90. doi:10.1111/j.1574-6976.2002.tb00600.x
- Mouveaux, T., Oria, G., Werkmeister, E., Slomianny, C., Fox, B. A., Bzik, D. J., et al. (2014). Nuclear Glycolytic Enzyme Enolase of *Toxoplasma Gondii* Functions as a Transcriptional Regulator. *PLoS ONE* 9 (8), e105820. doi:10.1371/journal.pone.0105820
- Nilsson, A., Olsson, T., Ulfstedt, M., Thelander, M., and Ronne, H. (2011). Two Novel Types of Hexokinases in the moss *Physcomitrella Patens*. *BMC Plant Biol.* 11 (1), 32. doi:10.1186/1471-2229-11-32
- Nogueira, S. V., Fonseca, F. L., Rodrigues, M. L., Mundodi, V., Abi-Chacra, E. A., Winters, M. S., et al. (2010). Paracoccidioides Brasiliensis Enolase Is a Surface Protein that Binds Plasminogen and Mediates Interaction of Yeast Forms with Host Cells. *Iai* 78 (9), 4040–4050. doi:10.1128/IAI.00221-10
- Nojima, H., Tokunaga, C., Eguchi, S., Oshiro, N., Hidayat, S., Yoshino, K.-i., et al. (2003). The Mammalian Target of Rapamycin (mTOR) Partner, Raptor, Binds the mTOR Substrates P70 S6 Kinase and 4E-BP1 through Their TOR Signaling (TOS) Motif. *J. Biol. Chem.* 278 (18), 15461–15464. doi:10.1074/jbc.C200665200
- Olsson, T., Thelander, M., and Ronne, H. (2003). A Novel Type of Chloroplast Stromal Hexokinase Is the Major Glucose-Phosphorylating Enzyme in the moss *Physcomitrella Patens*. *J. Biol. Chem.* 278 (45), 44439–44447. doi:10.1074/jbc.M306265200
- Osawa, H., Sutherland, C., Robey, R. B., Printz, R. L., and Granner, D. K. (1996). Analysis of the Signaling Pathway Involved in the Regulation of Hexokinase II Gene Transcription by Insulin. *J. Biol. Chem.* 271, 16690–16694. doi:10.1074/jbc.271.28.16690
- Ozcan, S., Leong, T., and Johnston, M. (1996). Rgt1p of *Saccharomyces cerevisiae*, a Key Regulator of Glucose-Induced Genes, Is Both an Activator and a Repressor of Transcription. *Mol. Cell Biol.* 16 (11), 6419–6426. doi:10.1128/mcb.16.11.6419
- Palomino, A., Herrero, P., and Moreno, F. (2005). Rgt1, a Glucose Sensing Transcription Factor, Is Required for Transcriptional Repression of the HXK2 Gene in *Saccharomyces cerevisiae*. *Biochem. J.* 388, 697–703. doi:10.1042/BJ20050160
- Pancholi, V., and Fischetti, V. A. (1992). A Major Surface Protein on Group A Streptococci Is a Glyceraldehyde-3-Phosphate-Dehydrogenase with Multiple Binding Activity. *J. Exp. Med.* 176 (2), 415–426. doi:10.1084/jem.176.2.415
- Parkison, C., Ashizawa, K., McPhie, P., Lin, K. H., and Cheng, S. Y. (1991). The Monomer of Pyruvate Kinase, Subtype M1, Is Both a Kinase and a Cytosolic Thyroid Hormone Binding Protein. *Biochem. Biophys. Res. Commun.* 179 (1), 668–674. doi:10.1016/0006-291x(91)91424-b
- Pastorino, J. G., and Hoek, J. B. (2008). Regulation of Hexokinase Binding to VDAC. *J. Bioenerg. Biomembr.* 40 (3), 171–182. doi:10.1007/s10863-008-9148-8
- Pastorino, J. G., Shulga, N., and Hoek, J. B. (2002). Mitochondrial Binding of Hexokinase II Inhibits Bax-Induced Cytochrome C Release and Apoptosis. *J. Biol. Chem.* 277 (9), 7610–7618. doi:10.1074/jbc.M109950200

- Peláez, R., Fernández-García, P., Herrero, P., and Moreno, F. (2012). Nuclear Import of the Yeast Hexokinase 2 Protein Requires α/β -Importin-dependent Pathway. *J. Biol. Chem.* 287, 3518–3529. doi:10.1074/jbc.M111.317230
- Peláez, R., Herrero, P., and Moreno, F. (2009). Nuclear export of the Yeast Hexokinase 2 Protein Requires the Xpo1 (Crm1)-dependent Pathway. *J. Biol. Chem.* 284, 20548–20555. doi:10.1074/jbc.M109.013730
- Pereira, L. A., Bãjo, S. n. N., Barbosa, M. n. S., da Silva, J. L. M., Felipe, M. S. S., de Santana, J. M., et al. (2007). Analysis of the Paracoccidiosis Brasiiliensistriosephosphate Isomerase Suggests the Potential for Adhesin Function. *FEMS Yeast Res.* 7 (8), 1381–1388. doi:10.1111/j.1567-1364.2007.00292.x
- Petit, F. M., Serres, C., Bourgeon, F., Pineau, C., and Auer, J. (2013). Identification of Sperm Head Proteins Involved in Zona Pellucida Binding. *Hum. Reprod.* 28 (4), 852–865. doi:10.1093/humrep/des452
- Petit, P., Antoine, M., Ferry, G., Boutin, J. A., Lagarde, A., Gluais, L., et al. (2011). The Active Conformation of Human Glucokinase Is Not Altered by Allosteric Activators. *Acta Crystallogr. D Biol. Cryst.* 67 (11), 929–935. doi:10.1107/S0907444911036729
- Piatigorsky, J., and Wistow, G. J. (1989). Enzyme/crystallins: Gene Sharing as an Evolutionary Strategy. *Cell* 57 (2), 197–199. doi:10.1016/0092-8674(89)90956-2
- Popanda, O., Fox, G., and Thielmann, H. W. (1998). Modulation of DNA Polymerases α , δ and ϵ by Lactate Dehydrogenase and 3-phosphoglycerate Kinase. *Biochim. Biophys. Acta (Bba) - Gene Struct. Expr.* 1397 (1), 102–117. doi:10.1016/s0167-4781(97)00229-7
- Prior, C., Mamessier, P., Fukuhara, H., Chen, X. J., and Wesolowski-Louvel, M. (1993). The Hexokinase Gene Is Required for Transcriptional Regulation of the Glucose Transporter Gene RAG1 in *Kluyveromyces Lactis*. *Mol. Cel. Biol.* 13 (7), 3882–3889. doi:10.1128/mcb.13.7.3882
- Ramajo-Hernández, A., Pérez-Sánchez, R., Ramajo-Martín, V., and Oleaga, A. (2007). Schistosoma Bovis: Plasminogen Binding in Adults and the Identification of Plasminogen-Binding Proteins from the Worm Tegument. *Exp. Parasitol.* 115 (1), 83–91. doi:10.1016/j.exppara.2006.07.003
- Ramiah, K., van Reenen, C. A., and Dicks, L. M. T. (2008). Surface-Bound Proteins of *Lactobacillus Plantarum* 423 that Contribute to Adhesion of Caco-2 Cells and Their Role in Competitive Exclusion and Displacement of *Clostridium sporogenes* and *Enterococcus faecalis*. *Res. Microbiol.* 159, 470–475. doi:10.1016/j.resmic.2008.06.002
- Randez-Gil, F., Herrero, P., Sanz, P., Prieto, J. A., and Moreno, F. (1998). Hexokinase PII Has a Double Cytosolic-Nuclear Localisation in *Saccharomyces Cerevisiae*. *FEBS Lett.* 425, 475–478. doi:10.1016/s0014-5793(98)00289-0
- Ritterson Lew, C., and Tolan, D. R. (2013). Aldolase Sequesters WASP and Affects WASP/Arp2/3-stimulated Actin Dynamics. *J. Cel. Biochem.* 114 (8), 1928–1939. doi:10.1002/jcb.24538
- Roberts, D. J., and Miyamoto, S. (2015). Hexokinase II Integrates Energy Metabolism and Cellular protection: Aktung on Mitochondria and TORCing to Autophagy. *Cell Death Differ* 22, 248–257. doi:10.1038/cdd.2014.173
- Roberts, D. J., Tan-Sah, V. P., Smith, J. M., and Miyamoto, S. (2013). Akt Phosphorylates HK-II at Thr-473 and Increases Mitochondrial HK-II Association to Protect Cardiomyocytes. *J. Biol. Chem.* 288, 23798–23806. doi:10.1074/jbc.M113.482026
- Roberts, D. J., Tan-Sah, V. P., Ding, E. Y., Smith, J. M., and Miyamoto, S. (2014). Hexokinase-II Positively Regulates Glucose Starvation-Induced Autophagy through TORC1 Inhibition. *Mol. Cel* 53, 521–533. doi:10.1016/j.molcel.2013.12.019
- Rodríguez-Enríquez, S., Marín-Hernández, Á., Gallardo-Pérez, J. C., Pacheco-Velázquez, S. C., Belmont-Díaz, J. A., Robledo-Cadena, D. X., et al. (2019). Transcriptional Regulation of Energy Metabolism in Cancer Cells. *Cells* 8 (10), 1225. doi:10.3390/cells8101225
- Rolland, F., Winderickx, J., and Thevelein, J. M. (2002). Glucose-sensing and -signalling Mechanisms in Yeast. *FEMS Yeast Res.* 2 (2), 183–201. doi:10.1111/j.1567-1364.2002.tb00084.x
- Ros, S., and Schulze, A. (2013). Glycolysis Back in the Limelight: Systemic Targeting of HK2 Blocks Tumor Growth. *Cancer Discov.* 3 (10), 1105–1107. doi:10.1158/2159-8290.CD-13-0565
- Roth, M. S., Westcott, D. J., Iwai, M., and Niyogi, K. K. (2019). Hexokinase Is Necessary for Glucose-Mediated Photosynthesis Repression and Lipid Accumulation in a green Alga. *Commun. Biol.* 2 (2), 1–10. doi:10.1038/s42003-019-0577-1
- Ryu, J.-Y., Jeong, S.-W., Kim, S. Y., Ko, Y., Yoon, S., Choi, S.-B., et al. (2008). Cyanobacterial Glucokinase Complements the Glucose Sensing Role of *Arabidopsis thaliana* Hexokinase I. *Biochem. Biophysical Res. Commun.* 374 (3), 454–459. doi:10.1016/j.bbrc.2008.07.041
- Sahoo, S., Murugavel, S., Devi, I. K., Vedamurthy, G. V., Gupta, S. C., Singh, B. P., et al. (2013). Glyceraldehyde-3-phosphate Dehydrogenase of the Parasitic nematode *Haemonchus contortus* binds to Complement C3 and Inhibits its Activity. *Parasite Immunol.* 35 (12), 457–467. doi:10.1111/pim.12058
- Saltman, P. (1953). Hexokinase in Higher Plants. *J. Biol. Chem.* 200, 145–154. doi:10.1016/s0021-9258(18)38447-3
- Sarwar, S., Lee, J.-Y., Ahn, E.-R., and Pai, H.-S. (2008). A Role of Hexokinases in Plant Resistance to Oxidative Stress and Pathogen Infection. *J. Plant Biol.* 51 (5), 341–346. doi:10.1007/bf03036136
- Sawyer, G. M., Monzingo, A. F., Poteet, E. C., O'Brien, D. A., and Robertus, J. D. (2008). X-ray Analysis of Phosphoglycerate Kinase 2, a Sperm-specific Isoform from *Mus musculus*. *Proteins* 71 (3), 1134–1144. doi:10.1002/prot.21801
- Schalm, S. S., and Blenis, J. (2002). Identification of a Conserved Motif Required for mTOR Signaling. *Curr. Biol.* 12 (8), 632–639. doi:10.1016/s0960-9822(02)00762-5
- Schulz, G. E., and Schirmer, R. H. (1979). “The Covalent Structure of Proteins,” in *Principles of Protein Structure*. Editor C. H. Cantor (Boston, MA: Springer), 47.
- Segovia-Gamboa, N. C., Chávez-Munguía, B., Medina-Flores, Y., Cázares-Raga, F. E., Hernández-Ramírez, V. I., Martínez-Palomo, A., et al. (2010). Entamoeba Invadens, Encystation Process and Enolase. *Exp. Parasitol.* 125 (2), 63–69. doi:10.1016/j.exppara.2009.12.019
- Sha, J., Erova, T. E., Alyea, R. A., Wang, S., Olano, J. P., Pancholi, V., et al. (2009). Surface-expressed Enolase Contributes to the Pathogenesis of Clinical Isolate SSU of *Aeromonas Hydrophila*. *Jb* 191 (9), 3095–3107. doi:10.1128/JB.00005-09
- Sharma, P., and Guptasarma, P. (2017). Endoglucanase Activity at a Second Site in *Pyrococcus Furiosus* triosephosphate Isomerase-Promiscuity or Compensation for a Metabolic Handicap? *FEBS Open Bio* 7 (8), 1126–1143. doi:10.1002/2211-5463.12249
- Shaw, G., and Kamen, R. (1986). A Conserved AU Sequence from the 3' Untranslated Region of GM-CSF mRNA Mediates Selective mRNA Degradation. *Cell* 46, 659–667. doi:10.1016/0092-8674(86)90341-7
- Sheikh, T., Gupta, P., Gowda, P., Patrick, S., and Sen, E. (2018). Hexokinase 2 and Nuclear Factor Erythroid 2-related Factor 2 Transcriptionally Coactivate Xanthine Oxidoreductase Expression in Stressed Glioma Cells. *J. Biol. Chem.* 293 (13), 4767–4777. doi:10.1074/jbc.M117.816785
- Shetty, S., Muniyappa, H., Halady, P. K. S., and Idell, S. (2004). Regulation of Urokinase Receptor Expression by Phosphoglycerate Kinase. *Am. J. Respir. Cel Mol. Biol.* 31 (1), 100–106. doi:10.1165/rcmb.2003-0104OC
- Shyu, A.-B., and Wilkinson, M. F. (2000). The Double Lives of Shuttling mRNA Binding Proteins. *Cell* 102, 135–138. doi:10.1016/s0092-8674(00)00018-0
- Šimčíková, D., Kockocová, L., Vackárová, K., Tešinsky, M., and Heneberg, P. (2017). Evidence-based Tailoring of Bioinformatics Approaches to Optimize Methods that Predict the Effects of Nonsynonymous Amino Acid Substitutions in Glucokinase. *Sci. Rep.* 7, 9499. doi:10.1038/s41598-017-09810-0
- Singh, N., and Bhalla, N. (2020). Moonlighting Proteins. *Annu. Rev. Genet.* 54, 265–285. doi:10.1146/annurev-genet-030620-102906
- Sriram, G., Martínez, J. A., McCabe, E. R. B., Liao, J. C., and Dipple, K. M. (2005). Single-gene Disorders: what Role Could Moonlighting Enzymes Play? *Am. J. Hum. Genet.* 76 (6), 911–924. doi:10.1086/430799
- Steitz, T. A., Shoham, M., and Bennett, W. S., Jr. (1981). Structural Dynamics of Yeast Hexokinase during Catalysis. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 293, 43–52. doi:10.1098/rstb.1981.0058
- Subramanian, A., and Miller, D. M. (2000). Structural Analysis of α -Enolase. *J. Biol. Chem.* 275 (8), 5958–5965. doi:10.1074/jbc.275.8.5958
- Sun, M.-H., Ma, Q.-J., Hu, D.-G., Zhu, X.-P., You, C.-X., Shu, H.-R., et al. (2018). The Glucose Sensor MdHXK1 Phosphorylates a Tonoplast Na⁺/H⁺ Exchanger to Improve Salt Tolerance. *Plant Physiol.* 176 (4), 2977–2990. doi:10.1104/pp.17.01472
- Sun, Y.-J., Chou, C.-C., Chen, W.-S., Wu, R.-T., Meng, M., and Hsiao, C.-D. (1999). The crystal Structure of a Multifunctional Protein: Phosphoglucose Isomerase/autocrine Motility Factor/neuroleukin. *Proc. Natl. Acad. Sci.* 96 (10), 5412–5417. doi:10.1073/pnas.96.10.5412
- Tan, V. P., and Miyamoto, S. (2016). Nutrient-sensing mTORC1: Integration of Metabolic and Autophagic Signals. *J. Mol. Cell Cardiol.* 95, 31–41. doi:10.1016/j.yjmcc.2016.01.005

- Tayser, K. A., and Colowick, S. P. (1961). Properties of Crystalline Hexokinase from Yeast IV. Multiple Forms of the Enzyme. *Arch. Biochem. Biophys.* 94, 177–181. doi:10.1016/0003-9861(61)90026-1
- Terrier, B., Degand, N., Guilpain, P., Servettaz, A., Guillevin, L., and Mouthon, L. (2007). Alpha-enolase: a Target of Antibodies in Infectious and Autoimmune Diseases. *Autoimmun. Rev.* 6 (3), 176–182. doi:10.1016/j.autrev.2006.10.004
- Tsai, H. J., and Wilson, J. E. (1996). Functional Organization of Mammalian Hexokinases: Both N- and C-Terminal Halves of the Rat Type II Isozyme Possess Catalytic Sites. *Arch. Biochem. Biophys.* 329, 17–23. doi:10.1006/abbi.1996.0186
- Tsang, C. K., Liu, Y., Thomas, J., Zhang, Y., and Zheng, X. F. S. (2014). Superoxide Dismutase 1 Acts as a Nuclear Transcription Factor to Regulate Oxidative Stress Resistance. *Nat. Commun.* 5 (3446), 1–4. doi:10.1038/ncomms4446
- Tsuge, S., Ochiai, H., Inoue, Y., Oku, T., Tsuno, K., Kaku, H., et al. (2004). Involvement of Phosphoglucose Isomerase in Pathogenicity of *Xanthomonas Oryzae* Pv. *Oryzae. Phytopathology* 94 (5), 478–483. doi:10.1094/PHYTO.2004.94.5.478
- Tunio, S. A., Oldfield, N. J., Ala'Aldeen, D. A., Wooldridge, K. G., and Turner, D. P. (2010a). The Role of Glyceraldehyde 3-phosphate Dehydrogenase (GapA-1) in *Neisseria Meningitidis* Adherence to Human Cells. *BMC Microbiol.* 10, 280. doi:10.1186/1471-2180-10-280
- Tunio, S. A., Oldfield, N. J., Berry, A., Ala'Aldeen, D. A., Wooldridge, K. G., and Turner, D. P. J. (2010b). The Moonlighting Protein Fructose-1, 6-bisphosphate Aldolase of *Neisseria Meningitidis*: Surface Localization and Role in Host Cell Adhesion. *Mol. Microbiol.* 76 (3), 605–615. doi:10.1111/j.1365-2958.2010.07098.x
- Ulfstedt, M., Hu, G.-Z., Eklund, D. M., and Ronne, H. (2018). The Ability of a Charophyte Alga Hexokinase to Restore Glucose Signaling and Glucose Repression of Gene Expression in a Glucose-Insensitive *Arabidopsis* Hexokinase Mutant Depends on its Catalytic Activity. *Front. Plant Sci.* 9 (1887), 1–15. doi:10.3389/fpls.2018.01887
- Van Schaftingen, E. (2013). "Hexokinase/Glucokinase," in *Encyclopedia of Biological Chemistry*. 2nd edition (Academic Press), 543–547. doi:10.1016/b978-0-12-378630-2.00073-6
- Vanegas, G., Quiñones, W., Carrasco-López, C., Concepción, J. L., Albericio, F., and Avilán, L. (2007). Enolase as a Plasminogen Binding Protein in *Leishmania Mexicana*. *Parasitol. Res.* 101 (6), 1511–1516. doi:10.1007/s00436-007-0668-7
- Veiga-Malta, I., Duarte, M., Dinis, M., Tavares, D., Videira, A., and Ferreira, P. (2004). Enolase from *Streptococcus sobrinus* is an Immunosuppressive Protein. *Cell Microbiol.* 6, 79–88. doi:10.1046/j.1462-5822.2003.00344.x
- Vega, M., Riera, A., Fernández-Cid, A., Herrero, P., and Moreno, F. (2016). Hexokinase 2 Is an Intracellular Glucose Sensor of Yeast Cells that Maintains the Structure and Activity of Mig1 Protein Repressor Complex. *J. Biol. Chem.* 291 (14), 7267–7285. doi:10.1074/jbc.M115.711408
- Velho, G., Froguel, P., Clement, K., Pueyo, M. E., Rakotoambinina, B., Zouali, H., et al. (1992). Primary Pancreatic Beta-Cell Secretory Defect Caused by Mutations in Glucokinase Gene in Kindreds of Maturity Onset Diabetes of the Young. *The Lancet* 340, 444–448. doi:10.1016/0140-6736(92)91768-4
- Veramendi, J., Fernie, A. R., Leisse, A., Willmitzer, L., and Trethewey, R. N. (2002). Potato Hexokinase 2 Complements Transgenic *Arabidopsis* Plants Deficient in Hexokinase 1 but Does Not Play a Key Role in Tuber Carbohydrate Metabolism. *Plant Mol. Bio.* 49 (5), 491–501. doi:10.1023/a:1015528014562
- Vermeulen, N., Vermeire, S., Arijis, I., Michiels, G., Ballet, V., Derua, R., et al. (2011). Seroreactivity against Glycolytic Enzymes in Inflammatory Bowel Disease. *Inflamm. Bowel Dis.* 17 (2), 557–564. doi:10.1002/ibd.21388
- Von Euler, H., and Adler, E. (1935). Über die komponenten der dehydrasesysteme. 6. dehydrierung von hexosen unter mitwirkung von adenosintriphosphorsäure. *Z. Physiol. Chem.* 23, 122–153.
- Wang, F., Ogasawara, M. A., and Huang, P. (2010). Small Mitochondria-Targeting Molecules as Anti-cancer Agents. *Mol. Aspects Med.* 31 (1), 75–92. doi:10.1016/j.mam.2009.12.003
- Wang, H., Xin, H., Guo, J., Gao, Y., Liu, C., Dai, D., et al. (2019). Genome-wide Screening of Hexokinase Gene Family and Functional Elucidation of HXK2 Response to Cold Stress in *Jatropha Curcas*. *Mol. Biol. Rep.* 46 (2), 1649–1660. doi:10.1007/s11033-019-04613-0
- Wang, L., Wang, F.-S., and Gershwin, M. E. (2015). Human Autoimmune Diseases: a Comprehensive Update. *J. Intern. Med.* 278 (4), 369–395. doi:10.1111/joim.12395
- Wang, X.-Q., Zheng, L.-L., Lin, H., Yu, F., Sun, L.-H., and Li, L.-M. (2017). Grape Hexokinases Are Involved in the Expression Regulation of Sucrose Synthase- and Cell wall Invertase-Encoding Genes by Glucose and ABA. *Plant Mol. Biol.* 94 (2), 61–78. doi:10.1007/s11103-017-0593-9
- Watanabe, H., Takehana, K., Date, M., Shinozaki, T., and Raz, A. (1996). Tumor Cell Autocrine Motility Factor Is the Neuroleukin/phosphohexose Isomerase Polypeptide. *Cancer Res.* 56 (13), 2960–2963.
- Wen, H., Ting, J. P.-Y., and O'Neill, L. A. J. (2012). A Role for the NLRP3 Inflammasome in Metabolic Diseases-Did Warburg Miss Inflammation? *Nat. Immunol.* 13, 352–357. doi:10.1038/ni.2228
- Wilson, J. E. (2003). Isozymes of Mammalian Hexokinase: Structure, Subcellular Localization and Metabolic Function. *J. Exp. Biol.* 206 (12), 2049–2057. doi:10.1242/jeb.00241
- Wistow, G. J., Lietman, T., Williams, L. A., Stapel, S. O., de Jong, W. W., Horwitz, J., et al. (1988). Tau-crystallin/alpha-enolase: One Gene Encodes Both an Enzyme and a Lens Structural Protein. *J. Cel Biol.* 107, 2729–2736. doi:10.1083/jcb.107.6.2729
- Wistow, G. J., and Piatigorsky, J. (1988). Lens Crystallins: the Evolution and Expression of Proteins for a Highly Specialized Tissue. *Annu. Rev. Biochem.* 57, 479–504. doi:10.1146/annurev.bi.57.070188.002403
- Wolf, A. J., Reyes, C. N., Liang, W., Becker, C., Shimada, K., Wheeler, M. L., et al. (2016). Hexokinase Is an Innate Immune Receptor for the Detection of Bacterial Peptidoglycan. *Cell* 166, 624–636. doi:10.1016/j.cell.2016.05.076
- Xiang, L., Le Roy, K., Bolouri-Moghaddam, M.-R., Vanhaecke, M., Lammens, W., Rolland, F., et al. (2011). Exploring the Neutral Invertase-Oxidative Stress Defence Connection in *Arabidopsis thaliana*. *J. Exp. Bot.* 62 (11), 3849–3862. doi:10.1093/jxb/err069
- Xu, L., Carrie, C., Law, S. R., Murcha, M. W., and Whelan, J. (2013). Acquisition, Conservation, and Loss of Dual-Targeted Proteins in Land Plants. *Plant Physiol.* 161 (2), 644–662. doi:10.1104/pp.112.210997
- Yego, E. C. K., and Mohr, S. (2010). siah-1 Protein Is Necessary for High Glucose-Induced Glyceraldehyde-3-Phosphate Dehydrogenase Nuclear Accumulation and Cell Death in Müller Cells. *J. Biol. Chem.* 285 (5), 3181–3190. doi:10.1074/jbc.M109.083907
- Yuan, W., Tuttle, D. L., Shi, Y. J., Ralph, G. S., and Dunn, W. A., Jr (1997). Glucose-induced Microautophagy in *Pichia pastoris* Requires the Alpha-Subunit of Phosphofructokinase. *J. Cel Sci.* 110 (16), 1935–1945. doi:10.1242/jcs.110.16.1935
- Zhang, L., Li, B., Zhang, Y., Jia, X., and Zhou, M. (2016). Hexokinase Plays a Critical Role in Deoxynivalenol (DON) Production and Fungal Development in *Fusarium Graminearum*. *Mol. Plant Pathol.* 17 (1), 16–28. doi:10.1111/mpp.12258
- Zheng, L.-F., Jiang, S., and Liu, M.-F. (2017). MicroRNA Regulation and Analytical Methods in Cancer Cell Metabolism. *Cell. Mol. Life Sci.* 74, 2929–2941. doi:10.1007/s00018-017-2508-y
- Zheng, L., Roeder, R. G., and Luo, Y. (2003). S Phase Activation of the Histone H2B Promoter by OCA-S, a Coactivator Complex that Contains GAPDH as a Key Component. *Cell* 114 (2), 255–266. doi:10.1016/s0092-8674(03)00552-x
- Zhu, W., Zhang, Q., Li, J., Wei, Y., Cai, C., Liu, L., et al. (2017). Glyceraldehyde-3-phosphate Dehydrogenase Acts as an Adhesin in *Erysipelothrix Rhusiopathiae* Adhesion to Porcine Endothelial Cells and as a Receptor in Recruitment of Host Fibronectin and Plasminogen. *Vet. Res.* 48 (1), 16. doi:10.1186/s13567-017-0421-x
- Ziveri, J., Tros, F., Guerrero, I. C., Chhuon, C., Audry, M., Dupuis, M., et al. (2017). The Metabolic Enzyme Fructose-1,6-Bisphosphate Aldolase Acts as a Transcriptional Regulator in Pathogenic *Francisella*. *Nat. Commun.* 8 (1), 1–15. doi:10.1038/s41467-017-00889-7

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Rodríguez-Saavedra, Morgado-Martínez, Burgos-Palacios, King-Díaz, López-Coria and Sánchez-Nieto. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.