



Corrigendum: Feasibility of Functional Near-Infrared Spectroscopy (fNIRS) to Investigate the Mirror Neuron System: An Experimental Study in a Real-Life Situation

Pei-Pei Sun¹, Fu-Lun Tan¹, Zong Zhang¹, Yi-Han Jiang¹, Yang Zhao¹ and Chao-Zhe Zhu^{1,2,3*}

¹ State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing, China, ² IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing, China, ³ Center for Collaboration and Innovation in Brain and Learning Sciences, Beijing Normal University, Beijing, China

Keywords: social interaction, mirror neuron system, fNIRS, real-life situation, double density, channel-based group analysis, ROI-based group analysis

A Corrigendum on

Feasibility of Functional Near-Infrared Spectroscopy (fNIRS) to Investigate the Mirror Neuron System: An Experimental Study in a Real-Life Situation

by Sun, P.-P., Tan, F.-L., Zhang, Z., Jiang, Y.-H., Zhao, Y., and Zhu, C.-Z. (2018). *Front. Hum. Neurosci.* 12:86. doi: 10.3389/fnhum.2018.00086

OPEN ACCESS

Edited and reviewed by:

Martin J. Herrmann,
University of Wuerzburg, Germany

*Correspondence:

Chao-Zhe Zhu
czzhu@bnu.edu.cn

Received: 27 March 2019

Accepted: 18 April 2019

Published: 08 May 2019

Citation:

Sun P-P, Tan F-L, Zhang Z, Jiang Y-H,
Zhao Y and Zhu C-Z (2019)
Corrigendum: Feasibility of Functional
Near-Infrared Spectroscopy (fNIRS) to
Investigate the Mirror Neuron System:
An Experimental Study in a Real-Life
Situation.
Front. Hum. Neurosci. 13:148.
doi: 10.3389/fnhum.2019.00148

In the original article, there was a mistake in **Table 2** as published. The last two lines in the first two columns of the table were presented in reverse order. The corrected table appears below.

In the original article, there was also an error in the text. In one passage, there were two commas missing and one redundant word “and”. A correction has been made to the section Materials and Methods, sub-section Traditional Channel-Based Group Analysis. The corrected sentence is presented below.

“The reason why we chose one-sided *t*-test was because we had a hypothesis about the direction of the effect, that was, the task period yielded an increased hemodynamic response compared with the rest period in both conditions.”

The authors apologize for these errors and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2019 Sun, Tan, Zhang, Jiang, Zhao and Zhu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 2 | Activation when using the ROI-based group analysis.

ROI	Definition in the Brodmann atlas	t-value	p-value	Significance (FDR-corrected)
PMC	BA6	$t_{(exe)} = 6.98$	$p_{(exe)} = 0.0000$	***
		$t_{(obs)} = 3.62$	$p_{(obs)} = 0.0006$	**
IFG	BA44, BA45	$t_{(exe)} = 6.53$	$p_{(exe)} = 0.0000$	***
		$t_{(obs)} = 2.00$	$p_{(obs)} = 0.0274$	*
SPL	BA7	$t_{(exe)} = 5.44$	$p_{(exe)} = 0.0000$	***
		$t_{(obs)} = 7.29$	$p_{(obs)} = 0.0000$	***
Rostral IPL	BA40	$t_{(exe)} = 2.24$	$p_{(exe)} = 0.0165$	*
		$t_{(obs)} = 1.68$	$p_{(obs)} = 0.0519$	

The first column shows the four ROIs. The second column defines the ROI in the Brodmann atlas. PMC, premotor cortex; IFG, inferior frontal gyrus; SPL, superior parietal lobule; IPL, inferior parietal lobule; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.