

# Corrigendum: Combination Treatment of Icariin and L-DOPA Against 6-OHDA-Lesioned Dopamine Neurotoxicity

OPEN ACCESS

Di-Sheng Lu $^{\dagger}$ , Ce Chen $^{\dagger}$ , Ya-Xin Zheng $^{\dagger}$ , Dai-Di Li, Guo-Qing Wang, Jie Liu, Jingshan Shi and Feng Zhang $^{*}$ 

## Edited and reviewed by:

Jean-Marc Taymans, Institut National de la Santé et de la Recherche Médicale (INSERM), France

### \*Correspondence:

Feng Zhang zhangfengzmc@163.com

<sup>†</sup>These authors have contributed equally to this work

Received: 13 September 2020 Accepted: 30 September 2020 Published: 29 October 2020

### Citation:

Lu D-S, Chen C, Zheng Y-X, Li D-D, Wang G-Q, Liu J, Shi J and Zhang F (2020) Corrigendum: Combination Treatment of Icariin and L-DOPA Against 6-OHDA-Lesioned Dopamine Neurotoxicity. Front. Mol. Neurosci. 13:605722. doi: 10.3389/fnmol.2020.605722 Key Laboratory of Basic Pharmacology of Ministry of Education and Joint International Research Laboratory of Ethnomedicine of Ministry of Education, Zunyi Medical University, Zunyi, China

Keywords: Parkinson's disease, L-DOPA, dyskinesia, Icariin, combination treatment

1

### A Corrigendum on

# Combination Treatment of Icariin and L-DOPA Against 6-OHDA-Lesioned Dopamine Neurotoxicity

by Lu, D.-S., Chen, C., Zheng, Y.-X., Li, D.-D., Wang, G.-Q., Liu, J., et al. (2018). Front. Mol. Neurosci. 11:155. doi: 10.3389/fnmol.2018.00155

In the original article, there was a mistake in **Figure 6** as published. One western blot band of  $\beta$ -actin corresponding to TNF- $\alpha$  protein level 21 days after daily L-DOPA combined with ICA treatment in **Figure 6B** was inserted incorrectly. The corrected **Figure 6** appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Lu, Chen, Zheng, Li, Wang, Liu, Shi and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 6 | ICA inhibited the neuroinflammation in LID after 6-OHDA-induced DA neuronal damage. The protein levels of neuroinflammatory factors, such as COX-2, IL-1β and TNF-α, in the SN were detected 7 (A) and 21 (B) days after daily L-DOPA (25 mg/kg, i.p.) combined with ICA (20 mg/kg, p.o.) treatment by western blotting. Data were shown as mean  $\pm$  SEM from six rats (n = 6). \*p < 0.05 compared with the control group; #p < 0.05 compared with 6-OHDA+L-DOPA group.