



RETRACTED: An Antifungal Mechanism of Protolichesterinic Acid from the Lichen *Usnea albopunctata* Lies in the Accumulation of Intracellular ROS and Mitochondria-Mediated Cell Death Due to Apoptosis in *Candida tropicalis*

S. N. Kumar* and C. Mohandas

Division of Crop Protection, Central Tuber Crops Research Institute, Sreehariyam, India

OPEN ACCESS

Edited by:

Nicola Clementi,
Vita-Salute San Raffaele University,
Italy

Reviewed by:

Shamala Devisekaran,
University of Malaya, Malaysia
Carlos Henrique Gomes Martins,
Universidade de Franca, Brazil

*Correspondence:

S. N. Kumar
nishanthctcri@gmail.com;
micronishanth@rediffmail.com

Specialty section:

This article was submitted to
Experimental Pharmacology and Drug
Discovery,
a section of the journal
Frontiers in Pharmacology

Received: 02 March 2017

Accepted: 10 May 2017

Published: 29 May 2017

Citation:

Kumar SN and Mohandas C (2017)
An Antifungal Mechanism
of Protolichesterinic Acid from
the Lichen *Usnea albopunctata* Lies
in the Accumulation of Intracellular
ROS and Mitochondria-Mediated Cell
Death Due to Apoptosis in *Candida*
tropicalis. *Front. Pharmacol.* 8:301.
doi: 10.3389/fphar.2017.00301

Candida species causes superficial and life-threatening systemic infections and are difficult to treat due to the resistance of these organism to various clinically used drugs. Protolichesterinic acid is a well-known lichen compound. Although the antibacterial activity of protolichesterinic acid has been reported earlier, the antifungal property and its mechanism of action are still largely unidentified. The goal of the present investigation is to explore the anticandidal activity and mechanism of action of protolichesterinic acid, especially against *Candida tropicalis*. The Minimum Inhibitory Concentration (MIC) value was established through microdilution techniques against four *Candida* species and out of four species tested, *C. tropicalis* showed a significant effect (MIC: 2 µg/ml). In the morphological interference assay, we observed the enhanced inhibition of hyphae when the cells were treated with protolichesterinic acid. Time-kill assay demonstrated that the maximum rate of killing was recorded between 2 and 6 h. *C. tropicalis* exposed to protolichesterinic acid exhibited an increased ROS production, which is one of the key factors of fungal death. The rise in ROS was due to the dysfunction of mitochondria caused by protolichesterinic acid. We confirmed that protolichesterinic acid-induced dysfunction of mitochondria in *C. tropicalis*. The damage of cell membrane due to protolichesterinic acid treatment was confirmed by the influx of propidium iodide and was further confirmed by the release of potassium ions. The treatment of protolichesterinic acid also triggered calcium ion signaling. Moreover, it commenced apoptosis which is clearly evidenced by Annexin V and propidium iodide staining. Interestingly protolichesterinic acid recorded excellent immunomodulatory property when tested against lymphocytes. Finally protolichesterinic acid showed low toxicity toward a normal human cell line Foreskin (FS) normal fibroblast. In *in vivo* test, protolichesterinic acid significantly enhanced the survival of *C. tropicalis* infected *Caenorhabditis elegans*. This investigation proposes that the protolichesterinic acid

induces apoptosis in *C. tropicalis* via the enhanced accumulation of intracellular ROS and mitochondrial damage, which leads fungal cell death via apoptosis. Our work revealed a new key aspect of mechanisms of action of protolichesterinic acid in *Candida* species. This article is the first study on the antifungal and mechanism of action of protolichesterinic acid in *Candida* species.

Keywords: protolichesterinic acid, *Candida*, mechanism of action, apoptosis, ROS

INTRODUCTION

The occurrence of infections due to pathogenic fungi has been gradually growing as assessed to occur in more than a billion humans every year worldwide (Lee and Lee, 2014). Diseases caused by pathogenic *Candida* spp. and other fungi continue to embody a noteworthy health burden, especially to human beings. The *Candida* species represent second most numerous agents causing fungal infection worldwide (Brown et al., 2012). Here some *Candida* spp. is highly resistant to various clinically used azoles and polyenes drugs. Moreover, *Candida* species is the fourth most common source of hospital-acquired systemic fungal infections with more than 50% mortality rates (Hwang et al., 2014). It is extensively believed that more than 70% of all the fungal infections in humans are caused by the *Candida* species viz. *Candida albicans*, *Candida tropicalis*, *Candida glabrata*, and *Candida parapsilosis* (Hwang et al., 2014). Investigation on the development of novel antifungal agents over the past few decades has resulted in only a few drugs, which are being clinically used (Bhattacharya et al., 2015). Five major groups of compounds are used to treat various the fungal infections and this includes azoles, polyenes, fluoropyrimidines, echinocandins, and allylamines (Bhattacharya et al., 2015). But, most of the compounds used to treat pathogenic fungi are usually synthetic in nature. But a few drugs to treat fungal infections like polyenes and echinocandins are purely natural compounds, which isolated from microbial sources. Fluconazole which comes under azole family, is the backbone in treating various infections caused by *Candida* species due to the great efficacy, reduced toxicity and inexpensive (Liu et al., 2016). However, with the widespread usage of fluconazole, the occurrence of drug resistance to many other azole derivatives emerged (Liu et al., 2016). Moreover, prolonged use of antifungals other than fluconazole in treating infections caused by *Candida* species has led to the emergence of resistance to almost all drugs used clinically. This leads to drug resistance in clinical isolates of *Candida* species mostly resulted through cross-resistance to many other drugs, a phenomenon termed as multidrug resistance (MDR). MDR is a serious problem during the treatment of opportunistic fungal infections which poses severe concern given the limited number of clinically useful antifungal drugs available in the market. Despite the accessibility of various other classes of antifungals, inherent toxicity, and development of multi drug resistance continue major problems in managing various deadly pathogenic fungi and the availability of novel antifungal drugs to overcome these situations are still limited (Sharma et al., 2015). These problems point to an urgent and unmet need for the development of novel antifungal agents or searching for new

antifungals from natural sources to fight against deadly fungal diseases.

Candida spp. are normal commensal microbes of humans, commonly found in the oral, gastrointestinal, urinary, and vaginal mucosa (Alves et al., 2014), and also produces superficial and severe systemic infections. Due to this, *Candida* spp. are also known as opportunistic fungal pathogens. *Candida* genus is most commonly recovered from hospital fungal infections and is known as candidaemia (Alves et al., 2014). *Candida* spp. are present in approximately 50% of human population and around 80% of women have suffered from *Candida* infections, with around 5% of these infections are recurring (de Castro et al., 2013). Through a failure of human defense systems, *Candida* can cause severe infections in almost any part of the body, including skin, nails, urogenital, or can be systemic (Wang et al., 2014).

Wide spread use of various immunosuppressive drugs, the use of broad-spectrum antibiotics to treat various infections are responsible for an increasing number of immunocompromised peoples and this in turn leads to various opportunistic infections worldwide. A debilitated or reduced immune system offers favorable situations for infectious diseases caused by various pathogenic microorganisms. AIDS due to HIV (human immunodeficiency virus) is one of the main causative features for the rising figure of peoples with various deadly fungal infections, especially by *Candida* spp. (Nissapatorn et al., 2003; Kothavade et al., 2010). The widespread use of antifungal drugs for prophylaxis in AIDS patients became one of the foremost reason of establishment of pathogenic *Candida non-albicans* species (CNA) and increased resistance to various antifungals (Kothavade et al., 2010). Surprisingly, in India, *C. tropicalis*, an important CNA, is one of the major cause of severe nosocomial candidaemia in many patients (Kothavade et al., 2010).

Medicinal plants are well-known natural sources for the treatment of various diseases since ancient times (Srivastava et al., 2013). Lichens are among the most mesmerizing organisms on our planet. Lichens are a well-known, self-supporting, mutualistic association between fungi (mycobiont) and algae (photobionts) which form a unique symbiotic structure, known as lichen thallus (Suzuki et al., 2016). Lichen can adapt to grow in many adverse environmental conditions. Interestingly, there are no reports on the destructive microbial infections in lichens and this clearly indicated the presence of many antimicrobial compounds. The ability of lichen to produce a wide range of biologically active metabolites help to explain the fact that lichens have also been traditionally used as sources of human medicines, pigments, perfumes etc. (Suzuki et al., 2016). Various chemical compounds have consistently been associated with lichens, and these compounds may have pharmaceutically important




FIGURE 1 | Chemical structure of protolichesterinic acid isolated from *U. albopunctata*. The chemical structure of protolichesterinic acid was confirmed based on the NMR and MS spectral analyses.

properties. Interestingly, most of the lichen compounds are phenolic in nature (Suzuki et al., 2016).

Protolichesterinic acid (Figure 1), a higher aliphatic acid is a typical lichen compound exhibiting a wide range of pharmaceutically interesting properties. In spite of the pharmacological effects that have already been recognized, various reports have shown that protolichesterinic acid exhibits several interesting biological properties such as antimicrobial, antitumor, and antioxidant activities (Ingolfssdottir et al., 1997; Turk et al., 2003; Mitrović et al., 2011; Russo et al., 2012). All these properties were suggesting that this compound is important classes of perfect natural bioactive compound. Even though the antibacterial and anticancer properties of this compound have been explored extensively and data about the antifungal activity especially against *Candida* species lacks in the literature. The reported biological properties of protolichesterinic acid clearly highlighted the importance of this compound as leading bioactive compound with pharmacological and medical potential in the development of novel drug formulations in nearby future (Nguyen et al., 2014; Bessadóttir et al., 2015).

Recently we have reported the purification and antimicrobial property of protolichesterinic acid from *Usnea albopunctata* (Kumar et al., 2014). We also reported the preliminary anticandidal activity of protolichesterinic acid against *C. albicans* and *C. tropicalis* in the above cited paper. In the present investigation, we studied the detailed antifungal properties of protolichesterinic acid by metabolic and morphological analyses to confirm its mechanism of action in *Candida* cells, especially *C. tropicalis*. Here, we examined both the *Candida* cell proliferation and the production of intracellular reactive oxygen species (ROS) by protolichesterinic acid as ROS play a vital role in cell death due to apoptosis. Besides, we also observed apoptotic topographies caused by protolichesterinic acid, which includes mitochondrial membrane depolarization and externalization of phosphatidylserine and calcium signals. Further, we also tested the toxicity profile of this compound against Foreskin (FS) normal fibroblast cell line. Finally, we showed protolichesterinic acid greatly prolonged the survival of *C. tropicalis* infected *C. elegans*.

MATERIALS AND METHODS

Extraction and Isolation of Protolichesterinic Acid

Usnea albopunctata were collected from Trivandrum, Kerala, India and were identified through morphological analysis and a voucher specimen has been deposited in the Crop Protection

Division of CTCRI for future reference. Finely dry ground thalli of *U. albopunctata* (50 g) were extracted using acetone: methanol (1: 1) in a Soxhlet apparatus. The final extracts were filtered and then concentrated under reduced pressure in a rotary evaporator at 40°C. The dried extract was stored at −20°C for future experiments. The dried extract of the lichen *U. albopunctata* (1 g) was dissolved in ethyl acetate. After filtration, the filtrate was fractioned using silica gel column chromatography (silica gel 60, 230–400 mesh). The silica gel column was eluted with methanol-ethyl acetate solvent gradient (10: 1 and 5: 1) which yield five major fractions. The fourth fraction yielded pure protolichesterinic acid (60 mg). The chemical structure of protolichesterinic acid was confirmed based on the detailed spectral analyses (NMR and HRMS) (Kumar et al., 2014).

Preparation of Protolichesterinic Acid and the Fungal Strains

A stock solution of protolichesterinic acid (50 mg/ml) dissolved in 0.5% dimethyl sulfoxide (DMSO) was used for all assays. The *Candida* strains *Candida albicans* MTCC 277, *Candida tropicalis* MTCC 230, *Candida glabrata* MTCC 3019, and *Candida parapsilosis* MTCC 6510 were used for antifungal studies. All the test fungi were obtained from MTCC (Microbial Type Culture Collection Centre), Institute of Microbial Technology (IMTECH, Chandigarh, India). The test fungi were preserved on potato dextrose agar (PDA) slants. The fungal cells were cultured in potato dextrose broth (PDB) with aeration at 37°C under agitation at 150 rpm. Initially, the fungal growth was monitored by measuring the OD at 600 nm with a microtiter ELISA plate reader. After incubation, the *Candida* cells were harvested by centrifugation at 5000 × g for 5 min at 4°C and washed twice in 10 ml of 0.1 M phosphate-buffered saline (PBS; pH 7). Cell pellets thus formed were resuspended in 10 ml PDB and the cell density was adjusted to 1 × 10⁶ cells/ml using a hemocytometer.

Determination of Minimum Inhibitory Concentration (MIC) of Protolichesterinic Acid

Minimum inhibitory concentrations (MICs) were done by the microdilution method conferring to the reference of CLSI (Clinical and Laboratory Standards Institute [CLSI], 2005) using multi-well ELISA plates altered according to the commendations that have been mentioned for the more efficient determination of antimicrobial potential of natural compounds (Cos et al., 2006). In brief, 180 µl of PDB was dispensed aseptically to the wells of a 96-well microdilution plates. After that, 10 µl each of the serial dilutions of protolichesterinic acid and amphotericin B (AmpB) (reference antimycotic drug) were prepared in the PDB (1 to 2000 µg/ml). About 10 µl of *Candida* suspension at a density of 10⁶ CFU/ml was inoculated in each well and the plates were incubated for 48 h at 37°C. The Candidal growth was measured by taking the absorbance at 600 nm using a microtiter plate reader. Dimethyl sulfoxide and sterile water (deionized) were used as a blank control which did not inhibit the growth of *Candida* species tested.

Minimum inhibitory concentration was defined as the lowest concentrations of any agent that inhibited the growth of the test microbes by $\geq 90\%$ compared with that of the control. Minimum inhibitory concentrations were attained from three independent tests that performed in triplicate (Wonyoung and Dong Gun, 2014).

Minimum Fungicidal Concentration (MFC) of Protolichesterinic Acid

To determine the minimum fungicidal concentrations (MFCs) of protolichesterinic acid and amphotericin B, an aliquot (10 μ l) taken from each well that recorded total inhibition of fungal growth was plated onto PDA plates. The presence of fungal colonies was monitored after 24 h of incubation at 37°C. MFC was defined as the lowest concentration of protolichesterinic acid and amphotericin B that produced a total reduction in CFU (99.99% inhibition compared to non-treated microbial growth). The experiment was repeated for three times.

Agar Disk Diffusion Assay

Hundred microliter of overnight *Candida* cultures were spread over PDA plates. Six millimeter sterile filter paper disks (Hi-Media) containing MIC concentration of protolichesterinic acid and amphotericin B were placed on the agar plates. Filter paper disks incorporated with 10 μ l of DMSO served as negative control. Plates were incubated at 37°C for 24 h. After incubation, the diameter of zone of inhibition was envisaged around the disks containing test substances. The antimicrobial activity of the test compounds was determined by measuring the zone of inhibition (diameter) and expressed in mm. The experiment was repeated for three times.

Cell Viability Assay

Antifungal activity of protolichesterinic acid was tested with the colony-forming unit (CFU) assay. *Candida* species were treated with different concentrations of protolichesterinic acid (0.5 \times MIC, MIC, and 2 \times MIC) and incubated for 8 h at 37°C. About 100 μ l samples from each treatment were collected at 0, 2, 4, 6, and 8 h and cells were washed with PBS and diluted serially. Dilutions were spread on PDA plates individually and incubated for 24 h at 37°C. To measure the cell viability, CFUs were counted by eye. The experiment was repeated time times.

Hyphal Induction Assay

Candida hyphae were induced as reported previously (Yu et al., 2014), except *C. glabrata* (As this species does not form any hyphae). Briefly, 24 h *Candida* cells were washed thoroughly with PBS and suspended in RPMI-1640 medium comprising the protolichesterinic acid with the specified concentrations (0.5 \times MIC, MIC, and 2 \times MIC). After that, the tubes were incubated in a shaking incubator for 4 h at 37°C. The percentage of hyphal induction was calculated as the previously reported method (Yu et al., 2015).

Time-Kill Kinetic Analysis

Exponential-phase *Candida* cells (1×10^6 cells/ml) were incubated with different concentration of protolichesterinic acid

(0.5 \times MIC, MIC, and 2 \times MIC) and amphotericin B (MIC). Control cells were incubated with DMSO and sterile deionized water. About 10 μ l samples from each treatment were collected at 0, 1, 2, 4, 6, 12, 24, and 48 h and serially diluted in PBS, and 50 μ l aliquots were gently plated on the surface of PDA plates. The viable *Candida* colonies were counted after incubation at 37°C for 48 h. All of the experiments were performed in triplicate.

After the incubation period, the number of viable *Candida* cells was counted and expressed in log CFU/ml. The results were scrutinized and represented graphically, that is a fungal death curve vs. function of time. Data analysis for the protolichesterinic acid was measured as showing fungicidal activity when there was a decrease greater than or equal to 3 log₁₀ CFU/ml of the initial inoculum, resulting in the reduction of 99.99% or more CFU/ml in 48 h compared with the initial inoculum. Fungistatic activity was considered as the decrease in growth lower than 99.99% or <3 log₁₀ in CFU/ml from the original inoculum (Leite et al., 2015).

Mechanism Action on Protolichesterinic Acid on *Candida*

Out of four *Candida* species tested, *C. tropicalis* recorded the best activity and thus this strain was selected for further detailed studies including mechanism of action.

Detection of Reactive Oxygen Species (ROS) in *C. tropicalis* after Treatment with Protolichesterinic Acid

5-(and-6)-chloromethyl-2',7'-dichlorodihydrofluorescein diacetate acetyl ester (CMH₂DCFDA) effortlessly diffuses through the microbial cell membrane, and then it is freely hydrolyzed by intracellular esterases to produce non-fluorescent dichlorofluorescein (DCFH), which is then very quickly oxidized to form extremely fluorescent 2',7'-dichlorofluorescein (DCF) by various intracellular oxidative stressors (da Silva et al., 2014). The DCF fluorescence intensity thus formed is directly proportional to the sum of intracellular ROS formed by the action of any test compounds (da Silva et al., 2014). For revealing of ROS produced over a 24 h culture period, when treated with protolichesterinic acid, *C. tropicalis* was incubated with 20 μ M CMH₂DCFDA at 37°C for 2 h in the dark. Then, the *C. tropicalis* cells were collected, washed three times thoroughly with PBS buffer, resuspended in PBS, and immediately examined using a confocal microscopy (BD Pathways). The fluorescence was further measured using a spectrofluorophotometer (Shimadzu RF-5301PC, Shimadzu, Kyoto, Japan) at an excitation wavelength of 495 nm and an emission wavelength of 525 nm. 2.5 mM H₂O₂ was used as positive control.

N-Acetyl Cysteine (NAC) Assay

For ROS quenching, NAC (Sigma–Aldrich, St. Louis, MO, United States) was added to the *C. tropicalis* culture (1×10^6 cells/ml) to attain a final concentration of 5 mmol/L. *C. tropicalis* culture

was then treated with protolichsterinic acid and amphotericin B followed by the addition of NAC. The concentrations of NAC were determined to reduce the growth inhibition of *C. tropicalis*. Colony count (CFU/ml) was monitored after 4 h exposure to the protolichsterinic acid and amphotericin B. For measuring the colony count, 100 μ l of the *C. tropicalis* were collected, thoroughly washed with PBS buffer. This was then serially diluted in PBS buffer. All dilutions were plated onto PDA plates, and incubated at 37°C for 18 h. Serial dilutions that produced between 10 and 100 fungal colonies were counted, and the CFU/ml was stated as *Candida* survival (expressed in percentage) using the formula: [(CFU/ml of *C. tropicalis* treated with test compound)/(CFU/ml of non-treated control) \times 100] (Hwang et al., 2014). The results were express as mean \pm standard deviation for three independent tests.

Determination of Membrane Permeabilization in *C. tropicalis*

To investigate the alteration of membrane permeabilization under protolichsterinic acid treatment, cell membrane impermeable fluorescent dye propidium iodide (PI) was used. *C. tropicalis* was treated with the concentrations of 0 (control), 0.5 \times MIC, MIC, and 2 \times MIC concentration of protolichsterinic acid in PDB with 1×10^6 *Candida* cells/ml for 4 h. The amphotericin B treatment served as the control. 2.5 mM H₂O₂ was used as the positive control. After incubation at 37°C for 3 h, the *Candida* cells were stained with 5 μ g/ml of PI at 37°C for 30 min in the dark. The *Candida* cell membrane permeabilization was directly analyzed by confocal microscopy (BD Pathways). The fluorescence was further measured using a spectrofluorophotometer (Shimadzu RF-5301PC, Shimadzu, Kyoto, Japan) at an excitation wavelength of 535 nm and an emission wavelength of 617 nm. The test was performed three times (Li et al., 2015).

Determination of the Release of Potassium Ions after Protolichsterinic Acid Treatment

The variation in the ion concentration due to the treatment of protolichsterinic acid was determined by the release of potassium ions. *C. tropicalis* cells (OD approximately 1) were centrifuged at 5000 rpm for 5 min and the pellet thus obtained was resuspended in 5 ml PBS and treated with the protolichsterinic acid (0.5 \times MIC, MIC, and 2 \times MIC). Amphotericin B serves as the drug control. After incubation for 5-min intervals at 37°C, the cells were centrifuged at 13,000 rpm for 10 min. The supernatant was measured using an ion-selective electrode (ISE) meter (Orion Star A214, Thermo Scientific, Singapore). The cells were sonicated to determine 100% potassium release. The percentage of potassium release caused by the compounds was calculated as follows: potassium release (%) = $100 \times ([K^+]_t - [K^+]_0) / ([K^+]_t - [K^+]_0)$, where $[K^+]_t$ represents the potassium release achieved after addition of the compounds and $[K^+]_0$ and $[K^+]_t$ represent the potassium release

without the compounds and with sonication, respectively (Yun et al., 2015).

Determination of Changes in Mitochondrial Membrane Potential

The fluorescent dye rhodamine 123 or RHO123 (2-[6-amino-3-imino-3Hxanthen-9-yl] benzoic acid methyl ester) was used to study the mitochondrial membrane potential ($\Delta\Psi_m$) of *C. tropicalis* after the treatment of test compound. RHO123 is a positively charged dye that usually enters any cells by diffusion. The RHO123 accumulation and retention in cells mainly depend on the $\Delta\Psi_m$ value in response to the result of protolichsterinic acid treatment on mitochondrial transmembrane. Briefly, *C. tropicalis* (1×10^6 cells/ml) in the mid-log phase were centrifuged and resuspended in PBS (pH 7.4) and incubated for 15 min at 37°C with rhodamine 123 (10 mM). The cells were again washed with PBS and treated with 0.5 \times MIC, MIC, and 2 \times MIC concentration of protolichsterinic acid for 2 h at 37°C. 2.5 mM H₂O₂ was used as positive control. The microscopic examination of the delivery of rhodamine 123 in the *C. tropicalis* cells was achieved through confocal laser scanning microscopy (BD Pathways). The fluorescence was further measured using a spectrofluorophotometer (Shimadzu RF-5301PC, Shimadzu, Kyoto, Japan) at an excitation wavelength of 511 nm and an emission wavelength of 534 nm. Untreated *C. tropicalis* cells were used as controls.

Detection of Phosphatidylserine Externalization through Annexin-V Staining

Protoplast Preparation

The *C. tropicalis* cell wall was digested with lyticase (1 mg/g cells) in different washing steps in protoplast buffers (pH 7.4) containing 1 M sorbitol, DTT (buffer 1–30 mM; buffer 2–1 mM; buffer 3–0 mM), 50 mM tris base and 10 mM MgCl₂. Briefly, cells were harvested and washed thrice for 5 min each with buffer 1 (3 ml/g cells). Cells were then incubated in buffer 2 (5 ml/g cells; supplemented with lyticase) for 2 h at 37°C. After removing buffer 2 by centrifugation, cells were incubated with buffer 3 (5 ml/g cells) for 15 min and then again centrifuged to remove buffer 3. Protoplasts were finally washed once with PBS and re-suspended in the same.

Annexin-V Staining

Phosphatidylserine externalization due to apoptosis after protolichsterinic acid treatment was studied by FITC-Annexin V apoptosis detection kit (Sigma–Aldrich). For this experiment, the protoplasts were incubated with 0.5 \times MIC, MIC, and 2 \times MIC concentration of protolichsterinic acid for 2 h at 37°C and incubated for 15–20 min in an annexin binding buffer containing 5 μ l of FITC-Annexin-V/ml and 5 μ l of PI. After this the cells were then analyzed by confocal laser scanning microscopy. 2.5 mM H₂O₂ was used as positive control. The experiment was performed three times, and the results were shown as mean values \pm SDs.

Studying the Cytosolic and Mitochondrial Ca^{2+} Level Up on the Treat of Protolicheterinic Acid

To examine cytosolic and mitochondrial Ca^{2+} levels, Fura-2AM (Molecular Probes) and Rhod-2AM (Molecular Probes) were used, respectively. *C. tropicalis* cells (1×10^6 cells/ml) were incubated with protolicheterinic acid ($0.5 \times \text{MIC}$, MIC , and $2 \times \text{MIC}$) and amphotericin B at 37°C for 4 h. Ca^{2+} level was studied according to the previously reported method (Choi and Lee, 2015). $2.5 \text{ mM H}_2\text{O}_2$ was used as positive control.

Protolicheterinic Acid Toxicity to Leukocytes

The toxicity of protolicheterinic acid to leukocytes was evaluated using MTT assay. Briefly, the blood was collected in a sterile heparinized tube under sterile environment. Blood was diluted with a double volume of PBS and mixed well. Using a 10 ml syringe with needle attached, 3 ml Ficoll was transferred to a clean sterile 50 ml vial. The blood was diluted on to the gradient without mixing using a sterile pipette. After this it was centrifuged for 10 min at 3000 rpm. Following centrifugation, the test tube was carefully removed and kept inside the laminar. Using a sterile Pasteur pipette, it was removed gradually and the upper layer of the solution was discarded down till within $\sim 1 \text{ cm}$ of the interphase. After that, middle layer was carefully removed down till $\sim 0.5 \text{ cm}$ now leaving a little material there. An equal volume of PBS was added and centrifuged for 5 min at 3000 rpm. After that the supernatant solution was removed carefully and the pellet was diluted using RPMI medium. The cells were counted using a hemocytometer. About 1×10^5 cells/ml for culture was diluted and MTT experiment was performed as reported previously (Aravind et al., 2014). Proliferation Rate was calculated using formula $\text{T/C} \times 100$.

Foreskin (FS) Normal Fibroblast Cell Proliferation Inhibition Using the MTT Test

Cultivation of FS Normal Fibroblast Cell

FS cells were cultivated under standard culture conditions in minimal essential medium with Earle's salts. FS normal fibroblast growth was analyzed on the basis of the ability of living FS cells to reduce the yellow color of MTT dye (Sigma-Aldrich) to the purple formazan product. FS cells were plated in 96-well plates (3000 cells/well), and protolicheterinic acid ($1\text{--}200 \mu\text{g/ml}$) dissolved in 0.1% DMSO were added to each well, followed by incubation for 24 h under standard cultivation conditions. Afterward, the plates were centrifuged and the medium was replaced with fresh medium ($150 \mu\text{l}$) containing 0.5 mg/ml MTT. Three hours later, the MTT formazan product was dissolved in $150 \mu\text{l}$ DMSO and the absorbance was measured using an ELISA plate reader. The effects of the test compounds were quantified as the percentage of control absorbance of the reduced dye at 595 nm. Cisplatin was used as positive control. Experiments were carried out in triplicate and expressed mean \pm standard deviation.

In Vitro Antifungal Property of Protolicheterinic Acid in *C. elegans* Infected with *C. tropicalis*

Caenorhabditis elegans – *C. tropicalis* assay was performed according to previously reported method (Chang et al., 2012). Briefly, nematodes were transferred on 48 h old *C. tropicalis* lawns [on Brain Heart Infusion agar (BHI) plates] for 2 h. The nematodes were washed off the plates with screen medium [30% BHI broth in M9 buffer containing kanamycin ($100 \mu\text{g/ml}$), ampicillin ($200 \mu\text{g/ml}$), and streptomycin ($200 \mu\text{g/ml}$)]. The screen buffer containing nematodes were re-suspended at a density of two nematodes/ μl screen medium. The suspension of preinfected nematodes ($20 \mu\text{l}$) was added to wells of ELISA plates. $80 \mu\text{l}$ of screen medium containing the various concentration of protolicheterinic acid ($0.5 \times \text{MIC}$, MIC , and $2 \times \text{MIC}$) was dispensed into the indicated well. Amphotericin B serves as the positive control. The nematodes' survival in all treatment was checked each day until 5 days (Chang et al., 2015). The nematode survival rates were calculated by counting the live and dead nematodes based on shape (live nematodes usually appear as sinusoidal while the dead nematodes appear rod shape).

Statistical Analysis

In vitro susceptibility tests were repeated at least thrice on different days. All the results were presented as means \pm standard deviations. Geometric means were used to compare the MIC results. Significant differences between the treatments were established using one-way ANOVA test ($P < 0.05$).

RESULTS

Protolicheterinic Acid Recorded Strong Antifungal Effect

Although previous studies conducted by various researchers on protolicheterinic acid have mainly focused on studying the preliminary antimicrobial property, there is a shortage of information on the antifungal effects especially against *Candida* species of this compound in literature. The foremost emphasis of the current investigation was to elucidate the antifungal activity and preliminary mechanism of action of protolicheterinic acid in *Candida*. Our early results clearly indicated that protolicheterinic acid has significant antifungal activity against *Candida* species, especially to *C. tropicalis* and therefore, we further conducted detailed experiments to determine its mechanism of action on *C. tropicalis*.

The results of the sensitivity test of protolicheterinic acid and the standard drug (amphotericin B) against the *Candida* species are presented in **Table 1**. The MIC values of protolicheterinic acid ranged from 2 to $16 \mu\text{g/ml}$, whereas the MFC values range from 4 to $32 \mu\text{g/ml}$. But the antifungal activity of protolicheterinic acid was inferior when compared to the activity of amphotericin B (**Table 1**). Out of four *Candida* species tested, *C. tropicalis* recorded best MIC value ($2 \mu\text{g/ml}$). Zone of inhibition obtained from agar disk diffusion experiment was

presented in **Table 2** and significant inhibition of zone was also observed for *C. tropicalis* (26 mm).

The antifungal activity of protolichesterinic acid was also analyzed by CFU assay. Protolichesterinic acid treated *Candida* recorded significant cell survival percentages (**Figure 2**). Maximum inhibition was recorded at 4 h (**Figure 2**). Here also *C. tropicalis* recorded significant inhibition as evidenced by low cell survival percentages.

Protolichesterinic Acid Strongly Inhibits the Formation of Hyphae in *Candida* Species

Hyphal development is an important determinant for *Candida* species to effectively invade host tissues. So, we further tested the effect of the protolichesterinic acid on the inhibition of hyphal development. Excitingly, protolichesterinic acid exhibited stronger activity against the hyphal development of three *Candida* species tested (**Figure 3**). This result clearly indicated the high efficiency of the protolichesterinic acid against *Candida* hyphal development.

Time-Kill Assay

In the time-kill experiment, the outcome presented regarding the changes in the CFU/ml of viable *Candida* colonies indicated that the protolichesterinic acid exhibited potent anticandidal property (**Figure 4**). From the figure, it is very clear that the net growth rates sharply decreased from 2 h onward and the significant reduction in the number of *Candida* cells was recorded between 2 and 6 h when compared with untreated controls at 37°C. Interestingly, the protolichesterinic acid showed significantly better activity than amphotericin B regarding the growth of the yeast cells until 48 h incubation at 37°C. In the time-kill assay, protolichesterinic acid recorded best activity against *C. tropicalis* (**Figure 4**).

TABLE 1 | Minimum Inhibitory Concentration (MIC) of protolichesterinic acid.

Test fungi	MIC/MFC (μg/ml)	
	Protolichesterinic acid	Amphotericin B
<i>C. albicans</i>	8/16	0.5/1
<i>C. tropicalis</i>	2/4	1/1
<i>C. glabrata</i>	16/32	0.5/0.5
<i>C. parapsilosis</i>	16/32	1/1

TABLE 2 | Disk diffusion data of protolichesterinic acid.

Test fungi	Zone of inhibition (mm)	
	Protolichesterinic acid	Amphotericin B
<i>C. albicans</i>	21 ± 0.57	26 ± 2.21
<i>C. tropicalis</i>	26 ± 1.52	27 ± 1.77
<i>C. glabrata</i>	24 ± 1.52	28 ± 1.12
<i>C. parapsilosis</i>	22 ± 1	28 ± 0.53

In the time kill curve experiment, significant killing kinetics was recorded by 2× MIC concentrations, followed by MIC and 0.5× MIC concentration of protolichesterinic acid. Interestingly regrowth was not recorded for protolichesterinic acid.

Mechanism of Action of Protolichesterinic Acid on *C. tropicalis*

Out of four *Candida* species tested, *C. tropicalis* recorded the best activity in the preliminary experiments and thus this strain was selected for further detailed studies including mechanism of action.

Protolichesterinic Acid Increases Total Intracellular ROS Levels in *C. tropicalis*

Candida tropicalis treated with protolichesterinic acid exhibited elevated ROS levels compared with untreated *C. tropicalis*. In the H₂O₂ (positive control) treated *C. tropicalis*, there was a significant increase in fluorescence. As shown in **Figures 5A,B**, cells exposed to protolichesterinic acid showed a significant raise in the intracellular ROS level when compared to untreated control *C. tropicalis* cells. Our results clearly showed that accumulation of ROS induced by protolichesterinic acid was increased in a dose-dependent manner (**Figures 5A,B**).

In addition, changes in *C. tropicalis* viability after adding the ROS scavenger NAC were also studied. NAC (5 mM) did not record any adverse result on *C. tropicalis*. NAC enhances the survival of the *C. tropicalis* from 46.3 to 84.25% for protolichesterinic acid (**Figure 5C**).

Loss of Membrane Integrity by Protolichesterinic Acid as Evidenced by Propidium Iodide Staining

Propidium iodide has been reported to be a useful fluorescent dye for staining membrane impermeable nucleic acid. Propidium iodide enters the cells with compromised permeability only, and then it binds tightly to the double stranded DNA. After binding it produces a significant red fluorescence when excited at 480 nm. We examined the propidium iodide uptake by *Candida* cells in the presence of the protolichesterinic acid at 0.5×, 1×, and 2× MIC by confocal microscopy (**Figure 6**). Here protolichesterinic acid disrupted the membrane of *C. tropicalis* as evidenced by the staining of propidium iodide. As shown in **Figure 6**, cells treated with protolichesterinic acid displayed an increase of red fluorescence level when compared to that of untreated cells. This result indicated that the fungal cell membrane was damaged by the treatment of protolichesterinic acid.

Efflux of Potassium Ions

Figure 7 clearly showed that the concentration of potassium ion outside of the *C. tropicalis* cell was significantly enhanced after protolichesterinic acid treatment. Usually the concentration of potassium ions within the cell is high and for maintaining the cell homeostasis it must remain as such when there are no changes in the surroundings. If any changes like cell membrane damage happen to the cells, the ions levels in the surroundings get increased. *C. tropicalis* treated with protolichesterinic acid




FIGURE 2 | Antifungal effect of protolichesterinic acid. *Candida* species cells were treated with 0.5 × MIC, MIC, and 2 × MIC concentration of protolichesterinic acid and incubated for 8 h. After 0, 2, 4, 6, and 8 h the samples were collected, serially diluted, spreading in PDA agar plates and cell colonies were counted after 24 h incubation at 37°C.

and amphotericin B for 25 min recorded excellent potassium ions release when compared to non-treated cells. Based on the results, we confirmed that protolichesterinic acid affects *C. tropicalis* cells by destructing the cell membrane, which intern cause the permeabilization of cell membrane, and leads to the increased concentration of potassium ion outside the cells.

Protolichesterinic Acid Induces Collapse of Mitochondrial Membrane Potential ($\Delta\Psi_m$) in *C. tropicalis*

To further conclude whether the mitochondria-mediated pathway is associated with protolichesterinic acid induced apoptosis in *C. tropicalis*, we studied the changes of $\Delta\Psi_m$ by using the membrane sensitive fluorescence dye rhodamine 123 (Rho 123) that usually aggregates in mitochondria and produces an intense green fluorescence. The depolarization of mitochondrial membrane was significantly visible after treatment with H_2O_2 (which was used as the positive control). In agreement with this, protolichesterinic acid disrupted the $\Delta\Psi_m$ as evidenced by an increase in the intensity of green fluorescence by Rho 123 (Figure 8). Moreover our results clearly showed

that protolichesterinic acid enhances the intensity of fluorescent *C. tropicalis* cells in a dose-dependent manner. Therefore, protolichesterinic acid treated *C. tropicalis* were injured in permeability of mitochondria by the collapse of $\Delta\Psi_m$, which is a significant step in *C. tropicalis* cells undergoing programmed cell death (apoptosis).

Phosphatidylserine Externalization by Protolichesterinic Acid in *Candida*

To differentiate between apoptotic and necrotic *Candida* cell death induced by protolichesterinic acid, Annexin V double staining assay using the FITC Annexin V and PI was selected. As revealed in Figure 9, *C. tropicalis* exposed to protolichesterinic acid were considerably stained green fluorescence (positive FITC-Annexin V and negative PI) at the edge of the cell after digestion of the cell wall, representing a very clear externalization of phosphatidylserine. FITC-Annexin V stained *C. tropicalis* cells were not recorded in experiments conducted without protolichesterinic acid (control set). The exposure phosphatidylserine from the inner to the outer leaflet of plasma membrane in cells is an early morphological marker of apoptosis.




FIGURE 3 | Inhibition of *Candida* hyphae when treated with protolichesterinic acid. Protolichesterinic acid, AmpB. Different letters in the superscript were statistically different according to Duncan's multiple range test ($p < 0.05$).

Therefore, these results suggest that protolichesterinic acid induced early apoptosis in *C. tropicalis* cells, as shown by a significant staining in Annexin V-positive apoptotic cells. From our results, we established that protolichesterinic acid induces the production and accumulation of intracellular ROS, and which finally induces apoptotic features in *C. tropicalis*.

Influence on the Cytoplasmic Free Ca^{2+} Concentration

Intracellular calcium levels were evaluated using two membrane-permeable derivatives of the ratio metric calcium indicator, Fura-2AM and Rhod-2AM to detect whether calcium signaling was involved in cell death. The concentration of cytoplasmic free Ca^{2+} in *C. tropicalis* cells treated protolichesterinic acid

was higher than those in the control cells. The fluorescence intensity of Rhod-2AM was increased in cells treated with protolichesterinic acid (Figure 10A). Free calcium levels were also increased in the mitochondria of cells treated with protolichesterinic acid (Figure 10B), indicating that calcium signaling was involved in fungal cell death.

Immunomodulatory Properties of Protolichesterinic Acid

The proliferation of lymphocyte was evaluated to assess the immunomodulatory properties of protolichesterinic acid. The proliferation of lymphocyte was slightly improved in cells treated with protolichesterinic acid when compared to that of the cells treated without test compound (control). Lymphocyte




FIGURE 4 | Time-kill assay of protolichesterinic acid against *Candida* species. An overnight culture of *Candida* species were treated with 0.5× MIC, MIC, and 2× MIC of protolichesterinic acid and incubated at 37°C. Samples were collected at indicated time points and plated onto PDA plates. Plates were incubated for 48 h prior to counting the colony forming units (CFU). —■— Control, —●— AmpB, —▲— 0.5× MIC, —▼— MIC, —◆— 2× MIC.

proliferation index was recorded to be 1.19 for 10 µg/ml of protolichesterinic acid in the presence of PHA (Figure 11).

Protolichesterinic Acid Recorded no Toxicity toward Normal Cell Line

Interestingly protolichesterinic acid recorded no toxicity toward FS normal fibroblast when tested up to 200 µg/ml (Figure 12).

The *In Vivo* Efficacy of Protolichesterinic Acid

In vivo studies with *C. elegans* establish that protolichesterinic acid provided substantial protection for the worms infected with

C. tropicalis (Figure 13). Protolichesterinic acid (0.5× MIC, MIC, and 0.5× MIC) enhanced the survival of *C. tropicalis* infected nematodes with comparable efficacy as amphotericin B treatment (Figure 13).

DISCUSSION

As an opportunistic, deep-infective fungus, *Candida* species is an important infective microorganism for the patients mainly with immune dysfunction [for e.g., chronic diabetes, malignant tumors, leukemia, organ transplantations, and acquired immunodeficiency syndrome (AIDS)]. *Candida* species are a




FIGURE 5 | Intracellular ROS accumulation in *C. tropicalis* cell when treated with protolichesterinic acid. Protolichesterinic acid increased ROS accumulation in a dose-dependent manner. Protolichesterinic acid treated *C. tropicalis* cells were incubated with H₂DCFDA dye for 2 h at 37°C. The stained cells were measured using confocal microscope and spectrofluorophotometer. **(A)** Confocal image of *C. tropicalis*. **(B)** Fluorescence intensity of various treatments measured by spectrofluorophotometer. **(C)** The viability of cell was checked by counting (expresses as CFU/ml) after incubation with the test compound for 4 h and presented as the “percentage of cell survivals.” The result represents the mean \pm standard deviation from three independent assays. Different letters in the superscript were significantly different according to Duncan’s multiple range test ($p < 0.05$). Where PA is protolichesterinic acid.

widespread opportunistic fungal pathogen of human beings. The aggressive candidiasis has a probable mortality rate of over 50%, even with the employ of various antifungal agents mainly due to increased drug resistance often termed as MDR *Candida* species. Therefore, the discovery of new successful antifungal drugs and therapies especially from natural sources is urgently needed (Ding et al., 2016). Antifungal compounds like azoles

(fluconazole, ketoconazole, and clotrimazole) and polyenes (amphotericin B, nystatin, and natamycin) are frequently used to treat and control infections caused by various *Candida* species. The majority of the antifungal drugs are phenolic structure. The compounds with phenolic group show the highest anti-candidal property when compared to that of other class of compounds (Khodavandi et al., 2014). In the current years,




FIGURE 6 | Protolichesterinic acid leads to the membrane permeabilization as evidenced propidium iodide (PI) uptake. (A) Confocal image of *C. tropicalis*. **(B)** Fluorescence intensity of various treatments measured by spectrofluorophotometer. The data represent the average, standard deviation (SD), and different letters in the superscript were statistically different according to DMRT ($p < 0.05$) test analyzed by SPSS 17.




FIGURE 7 | The percentage of potassium released from *C. tropicalis* induced by the compounds was measured every 5 min up to 25 min. The results are expressed as mean \pm standard deviation from three independent experiments, performed in triplicate.

in the market (Khodavandi et al., 2014). Hence, there is a very urgent need to discover novel methods to enhance the efficiency of antifungal drugs with minimum resistance to drug for treating the infections caused by pathogenic *Candida* species. Toward this aim, finding novel compounds from natural sources and understanding its mechanism of actions are very much necessary to discover and develop new antifungal drugs to control various infectious diseases caused by pathogenic fungi.

The coming out of opportunistic fungal infections, especially in immunocompromised patients, highlights the necessity to expound novel therapeutic options. Most of the established antifungal drugs have been proved less effective against *Candida* species (Ma et al., 2015). Therefore, the research and development of new antifungal agents against *Candida* species are urgent needed (Ma et al., 2015). In the present investigation, we mainly focused on the antifungal property of protolichesterinic acid against four human pathogenic *Candida* species (*C. albicans*, *C. tropicalis*, *C. glabrata*, and *C. parapsilosis*). Protolichesterinic acid is an excellent therapeutic molecule that has been established to have many interesting biological activities (Thorsteinsdottir et al., 2016). However, the effect of protolichesterinic acid against *Candida* species has not been elucidated. Therefore, in the present investigation, we focused on studying the antifungal property of protolichesterinic acid against *Candida* species and exploring the underlying mechanism of action. In this study, we found that protolichesterinic acid had high antifungal property against the four *Candida* species tested and out of the four *Candida* species tested, *C. tropicalis* recorded significant effect.

Lichen metabolites (depsides and depsidone) exert manifold biological activity (Müller, 2001; Mitrović et al., 2011). The

failure of drug treatment against various fungal infections such as candidiasis is usually rising due to drug-resistance and the poor efficacy of anticandidal drugs that are available




FIGURE 8 | protolichesterinic acid induces mitochondrial membrane depolarization ($\Delta\psi_m$) of *C. tropicalis*. (A) Confocal image of *C. tropicalis*. (B) Fluorescence intensity of various treatments measured by spectrofluorophotometer. The data represent the average, SD, and different letters in the superscript were statistically different according to DMRT ($p < 0.05$) test analyzed by SPSS 17.





FIGURE 9 | Phosphatidylserine externalization showed by FITC-Annexin-V/PI staining in protolichesterinic acid-treated *C. tropicalis*. Phosphatidylserine externalization, which is observed at an early stage of apoptosis, shown by Annexin V staining. The probe enabled the detection of alteration of phosphatidylserine localization from the inner membrane to the outer membrane. The fluorescence intensity indicates the amount of exposed phosphatidylserine in cells treated with protolichesterinic acid at the 0.5 × MIC, MIC, and 2 × MIC. (A) Confocal image of *C. tropicalis*. (B) Percentage of Annexin-V positive cells. The data represent the average, SD, and different letters in the superscript were statistically different according to DMRT ($p < 0.05$) test analyzed by SPSS 17.

structural diversity of depsides and depsidone and their occurrence in lichens have evoked substantial interest in their pharmaceutically interesting biological properties (Shrestha and Clair, 2013). Lichens have been used as the main ingredients in many folk medicines for several decades to treat various

diseases as part of their traditional medicines (Dayan and Romagni, 2001). The Ayurvedic and Unani systems mentioned the medicinal properties of some lichens which are used to treat a wide array of common ailments (Shrestha and Clair, 2013). Recent studies in the field of medical have resulted





in the investigation of the biological property of a limited number of lichen compounds with some investigation suggesting that certain lichen compounds possibly offer a promising new source of future drug development programs (Kingston, 2011). More than 1000 compounds have been isolated and identified from lichens, relatively small number (~50 species) has been screened for antibiotic activity (Shrestha and Clair, 2013).

Recent investigations have reported that pathogenic fungal infections have considerably increased (Richardson, 2005) and natural compounds especially phytochemicals exerted significant antimicrobial effects (Joshi et al., 2011). Phytochemicals have comparatively less toxicity to human beings and have a range of interesting biological properties (Ahmadiani et al., 1998). These phytochemicals are essential to a plant's reproductive system and usually prevent the entry of pathogenic microorganism which is spreading throughout the plants. Among the phytochemicals from plant, the antimicrobial property and mode of actions of certain compounds such as (+)-Medioresinol (Hwang et al., 2012) and amentoflavone (Hwang et al., 2013) have been well reported earlier. In this study, we reported the antifungal property of protolichesterinic acid from *U. albopunctata* and explained its mode of action against *C. tropicalis* for the first time.

Reactive oxygen species is essential for cell signaling and cell functions. Several reports have reported that the increase of ROS induces and regulates the induction of cellular apoptosis (Hildeman et al., 1999; Hwang et al., 2011; Chen et al., 2016). As a result, to establish the intracellular ROS production and accumulation by protolichesterinic acid, we chose to employ the ROS-sensitive dye CMH₂DCFDA, which


has been used previously as a common marker for detecting the level of cellular ROS. Multiple ROS in cells directly oxidizes CMH₂DCFDA to the very highly stable, fluorescent substance DCFDA in such a way that an enhancement in the fluorescent signal is directly proportional to the ROS production (Hwang et al., 2011). Here in our study, protolichesterinic acid induces significant ROS production as evidenced by green fluorescent through confocal microscopy and spectrofluorophotometer.

Phytochemicals usually have a membrane active effect by the disturbance of the cell membrane integrity (Abad et al., 2007; Khan et al., 2014). We studied the outcome of protolichesterinic acid on the cell membrane integrity by using the fluorescence dye propidium iodide (PI). To know whether protolichesterinic acid exerts an antifungal activity by targeting the cell membrane, we investigated its effect on the integrity of cell membrane using the propidium iodide probe. Propidium iodide enters only through membrane compromised cells. After entering, PI intercalates between the guanine and the cytosine pair of DNA or with a stoichiometry of one unit dye per 3–5 base pairs. After DNA binding, the red fluorescent color of PI enhances more than 20-fold (Suzuki et al., 1997; Choi et al., 2012). From our study, we showed the distribution of the PI inside the *C. tropicalis* through the injured plasma membrane, and finally establish the cell membrane active mechanisms of protolichesterinic acid in *Candida* cells. Maintaining the cytoplasmic membrane integrity is essential to various important functions of microorganisms, including the formation of gradient and discriminating permeability, cellular energetic, synthesis of various biopolymers, and other key virulence determinants (Choi et al., 2012). In this admiration,



any changes in the physical state of the cell membrane induced by protolichesterinic acid may hinder with one or more functions directly or indirectly. From the propidium iodide assay, we showed the diffusion of the PI into the *C. tropicalis* mainly through the injured plasma membrane. These results clearly indicated that protolichesterinic acid permeabilized the *C. tropicalis* cell membrane. To further confirm the membrane permeabilization in *Candida* cells treated with protolichesterinic acid, a potassium ion release assay was conducted. The potassium ion concentration within the cell is generally high and it must remain as such when there are no changes in the environment for maintaining the homeostasis (Yun et al., 2015). After treatment with protolichesterinic acid, the potassium ion concentration outside of the *Candida* cell was increased significantly (Figure 7). Cells treated with varying concentration of protolichesterinic acid for 25 min recorded significant release of potassium ions, as compared to untreated control cells. When any changes such as membrane damage and permeabilization occur, potassium ions are released from the cell and level of potassium ions outside the cell is directly proportional to the damaged cells. From the results, we established that protolichesterinic acid affects *C. tropicalis* by mainly destructing the cell membrane, triggering the permeabilization of the membrane, and this intern alters the potassium ion concentration inside and outside the *Candida* cells (Yun et al., 2015).

The mitochondria play an important role in the cell, for example apoptosis, pH homeostasis, redox and ATP production (Chen et al., 2013). In the mitochondria, ATP is usually formed from the phosphorylation of mitochondrial oxidative mechanism, in which $\Delta\Psi_m$ plays a vital function (Chen et al., 2013). $\Delta\Psi_m$ is a key factor indicating the energetic state of the cell and other cellular organelles such as mitochondria (Chen et al., 2013). Mitochondrial electron transport inhibitors,



such as antimycin A and potassium cyanide, reduce $\Delta\Psi_m$ by inhibiting the function of respiratory chain proton pumping, which leads to reduced production of ATP and this finally leads to cell death. In an earlier study presented that the

antifungal activity of farnesol involves $\Delta\Psi_m$ hyperpolarization in *C. albicans* (Machida and Tanaka, 1999). In the current study, RHO123 was used to study the effect of protolichsterinic acid on mitochondrial potentials. RHO123 is a cationic and lipophilic dye that pervades the negatively charged mitochondria and reflects $\Delta\Psi_m$. Exposure to different concentrations of protolichsterinic acid showed significant hyperpolarization of *C. tropicalis* $\Delta\Psi_m$ in a dose-dependent manner, suggesting mitochondrial dysfunction. Moreover recent reports showed that mitochondria play a vital role in the induction of apoptosis; the indulgence of $\Delta\Psi_m$ is a primary and crucial cellular event during apoptosis induced by numerous stimuli, and this disturbance leads to the opening of transition pore in mitochondrial membrane and, internally, it leads to the release of many apoptogenic factors from mitochondria into the cytosol (Barroso et al., 2006).

Annexin V is a 35–36 kDa, a phospholipid binding protein with very high affinity for phosphatidylserine in the attendance of calcium ions (Ca^{2+}) (Chen et al., 2008). In the present assay, *Candida* cells with apoptosis can be stained by Annexin V-FITC, which usually binds to phosphatidylserine which are externalized on the outer side of the cells, while necrotic *Candida* cells accrue only PI into the nucleus through the permeabilized plasma membrane (Smrz et al., 2007). As shown in **Figure 9**, *Candida* exposed to protolichsterinic acid is considerably stained with green fluorescent (Annexin V-FITC: +ve, PI: –ve) at the periphery of cell after digesting the cell wall, representing clear externalization of phosphatidylserine. Annexin V-FITC stained *Candida* cells were not noticed in control experiments without protolichsterinic acid treatment. A primary morphological indicator of apoptosis is the externalization of phosphatidylserine from the inner to the outer plasma membrane in cells (Weghuber et al., 2010; Hwang et al., 2011). This very clearly indicated that protolichsterinic acid induces cell apoptosis through mitochondrial disruption *C. tropicalis*.

Calcium signaling plays a role in the regulation of cellular processes, including the initiation and execution of apoptosis (Yoon et al., 2012). Similar to mammalian cells, organelles of yeast such as the vacuoles, endoplasmic reticulum, and Golgi bodies store Ca^{2+} ions and maintenance of calcium signaling is important for normal cell survival (Bonill et al., 2002). We confirmed that protolichsterinic acid induced the release of intracellular Ca^{2+} storage into the cytosol and an influx of Ca^{2+} into the mitochondria. During the apoptotic

response, imbalance in the plasma membrane Ca^{2+} influx and Ca^{2+} export from the ER is known to elevate cytoplasmic calcium ion concentration, inducing a progressive increase in the mitochondrial Ca^{2+} uptake (Hajnóczky et al., 2006). This mitochondrial Ca^{2+} overload is one of the main factors that induce mitochondrial damage, mediated by perturbation or rupture due to a decrease in mitochondrial membrane potential (Lee and Lee, 2014). Interestingly protolichsterinic acid recorded excellent immunomodulatory property when tested against lymphocytes. Finally protolichsterinic acid showed low toxicity toward a normal human cell line FS normal fibroblast up to 200 $\mu\text{g/ml}$.

CONCLUSION

To our knowledge, this is the first study to evaluate the physiologic mechanisms of *C. tropicalis* cell death induced by protolichsterinic acid, a depside compound. We showed that cellular apoptosis accounted for protolichsterinic acid-induced cell death through the production ROS and dysfunction of mitochondria. Moreover protolichsterinic acid also triggered calcium ion signaling in *C. tropicalis*. Our findings suggest that protolichsterinic acid exerts their powerful antifungal activity against *Candida* species by generating ROS, inhibiting cell wall integrity in cells, which then initiates dysfunction of mitochondria, calcium ion signaling and this internally initiate apoptosis cell death programmed (i.e., programmed cell death). This compound is also non-toxic to lymphocytes and FS normal fibroblast (normal human cell). Finally *in vivo* test showed protolichsterinic acid greatly prolonged the survival of *C. tropicalis* infected *C. elegans*.

AUTHOR CONTRIBUTIONS

All authors listed, have made substantial, direct and intellectual contribution to the work, and approved it for publication.

ACKNOWLEDGMENT

SK thanks, ICMR, New Delhi for providing Senior Research fellowship.

REFERENCES

- Abad, M. J., Ansuategui, M., and Bermejo, P. (2007). Active antifungal substances from natural sources. *ARKIVOC* 7, 116–145. doi: 10.3998/ark.5550190.0008.711
- Ahmadiani, A., Fereidoni, M., Semnani, S., Kamalinejad, M., and Saremi, S. (1998). Antinociceptive and antiinflammatory effects of *Sambucus ebulus* rhizome extract in rats. *J. Ethnopharmacol.* 61, 229–235. doi: 10.1016/S0378-8741(98)00043-9
- Alves, C. T., Ferreira, I. C., Barros, L., Silva, S., Azeredo, J., and Henriques, M. (2014). Antifungal activity of phenolic compounds identified in flowers from North Eastern Portugal against *Candida* species. *Future Microbiol.* 9, 139–146. doi: 10.2217/fmb.13.147
- Aravind, S. R., Sreelekha, T. T., Dileep Kumar, B. S., Nishanth Kumar, S., and Mohandas, C. (2014). Characterization of three depside compounds from a Western Ghat lichen *Parmelia erumpens* Kurok with special reference to antimicrobial and anticancer activity. *RSC Adv.* 4, 34632–34643. doi: 10.1039/C4RA04137B
- Barroso, G., Taylor, S., Morshedi, M., Manzur, F., Gavino, F., and Oehninger, S. (2006). Mitochondrial membrane potential integrity and plasma membrane translocation of phosphatidylserine as early apoptotic markers: a comparison of two different sperm subpopulations. *Fertil. Steril.* 85, 149–154. doi: 10.1039/C4RA04137B
- Bessadóttir, M., Eiríksson, F. F., Becker, S., Ögmundsdóttir, M. H., Ómarsdóttir, S., Thorsteinsdóttir, M., et al. (2015). Anti-proliferative and pro-apoptotic effects

- of lichen-derived compound protolichsterinic acid are not mediated by its lipoxygenase-inhibitory activity. *Prostaglandins Leukot. Essent. Fatty Acids* 98, 39–47. doi: 10.1016/j.plefa.2015.04.009
- Bhattacharya, A. K., Chand, H. R., John, J., and Deshpande, M. V. (2015). Clerodane type diterpene as a novel antifungal agent from *Polyalthia longifolia* var. pendula. *Eur. J. Med. Chem.* 94, 1–7. doi: 10.1016/j.ejmech.2015.02.054
- Bonill, M., Nastase, K. K., and Cunningham, K. W. (2002). Essential role of Calcineurin in response to endoplasmic reticulum stress. *EMBO J.* 21, 2343–2353. doi: 10.1093/emboj/21.10.2343
- Brown, G. D., Denning, D. W., Gow, N. A. R., Levitz, S. M., Netea, M. G., and White, T. C. (2012). Hidden killers: human fungal infections. *Med. Mycol. Sci. Transl. Med.* 4:165rv13. doi: 10.1126/scitranslmed.3004404
- Chang, W., Li, Y., Zhang, L., Cheng, A., and Lou, H. (2012). Retigeric acid B attenuates the virulence of *Candida albicans* via inhibiting adenylyl cyclase activity targeted by enhanced farnesol production. *PLoS ONE* 7:e41624. doi: 10.1371/journal.pone.0041624
- Chang, W., Zhang, M., Ying, L., Li, X., Gao, Y., Xie, Z., et al. (2015). Lichen endophyte derived pyridoxatin inactivates *Candida* growth by interfering with ergosterol biosynthesis. *Biochim. Biophys. Acta* 1850, 1762–1771. doi: 10.1016/j.bbagen.2015.05.005
- Chen, X., Song, M., Zhang, B., and Zhang, Y. (2016). Reactive oxygen species regulate T cell immune response in the tumor microenvironment. *Oxid. Med. Cell. Longev.* 2016:1580967. doi: 10.1155/2016/1580967
- Chen, Y., Hong, Z., Jun, T., Xiaoquan, B., Bingxin, M., and Youwei, W. (2013). Antifungal mechanism of essential oil from *Anethum graveolens* seeds against *Candida albicans*. *J. Med. Microbiol.* 62, 1175–1183. doi: 10.1099/jmm.0.055467-0
- Chen, Z., Tong, L., Li, Z., Yoon, K. C., Qi, H., Farley, W., et al. (2008). Hyperosmolarity-induced cornification of human corneal epithelial cells is regulated by JNK MAPK. *Invest. Ophthalmol. Vis. Sci.* 49, 539–549. doi: 10.1167/iovs.07-0569
- Choi, H., Jaeyong, C., Qinglong, J., Eun-Rhan, W., and Dong Gun, L. (2012). Antifungal property of dihydrodehydrodiconiferyl alcohol 9'-O- β -D-glucoside and its pore-forming action in plasma membrane of *Candida albicans*. *Biochim. Biophys. Acta* 1818, 1648–1655. doi: 10.1016/j.bbame.2012.02.026
- Choi, H., and Lee, D. G. (2015). Lycopene induces apoptosis in *Candida albicans* through reactive oxygen species production and mitochondrial dysfunction. *Biochimie* 115, 108–115. doi: 10.1016/j.biochi.2015.05.009
- Clinical and Laboratory Standards Institute [CLSI] (2005). *Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts, as the Document is M27-S4*. Wayne, PA: Clinical and Laboratory Standards Institute.
- Cos, P., Vlietinck, A. J., Berghe, D. V., and Maes, L. (2006). Anti-infective potential of natural products: how to develop a stronger in vitro 'proof-of-concept'. *J. Ethnopharmacol.* 106, 290–302. doi: 10.1016/j.jep.2006.04.003
- da Silva, C. R., Joao Batista de, A. N., Rosana de, S. C., Narjara Silvestre, F., Leticia Serpa, S., Hemerson Iury, F. M., et al. (2014). Synergistic effect of the flavonoid catechin, quercetin, or epigallocatechin gallate with fluconazole induces apoptosis in *Candida tropicalis* resistant to fluconazole. *Antimicrob. Agents Chemother.* 58, 1468–1478. doi: 10.1128/AAC.00651-13
- Dayan, F. E., and Romagni, J. G. (2001). Lichens as a potential source of pesticides. *Pestic. Outlook* 12, 229–232. doi: 10.1039/b110543b
- de Castro, P. A., Bom, L. P., Brown, N. A., de Almeida, R. S. C., Ramalho, L. N., Savoldi, M., et al. (2013). Identification of the cell targets important for propolis-induced cell death in *Candida albicans*. *Fungal Genet. Biol.* 60, 74–86. doi: 10.1016/j.fgb.2013.07.001
- Ding, Y., Li, Z., Li, Y., Lu, C., Wang, H., Shen, Y., et al. (2016). HSAF-induced antifungal effects in *Candida albicans* through ROS-mediated apoptosis. *RSC Adv.* 6, 30895–30904. doi: 10.1039/C5RA26092B
- Hajnóczky, G., Csordás, G., Das, S., Garcia-Perez, C., Saotome, M., SinhaRoy, S., et al. (2006). Mitochondrial calcium signaling and cell death: approaches for assessing the role of mitochondrial Ca^{2+} uptake in apoptosis. *Cell Calcium* 40, 553–560. doi: 10.1016/j.ceca.2006.08.016
- Hildeman, D. A., Mitchell, T., Teague, T. K., Henson, P., Day, B. J., Kappler, J., et al. (1999). Reactive oxygen species regulate activation-induced T cell apoptosis. *Immunity* 10, 735–744. doi: 10.1016/S1074-7613(00)80072-2
- Hwang, B., Hwang, J. S., Lee, J., and Lee, D. G. (2011). The antimicrobial peptide, psacothasin induces reactive oxygen species and triggers apoptosis in *Candida albicans*. *Biochem. Biophys. Res. Commun.* 405, 267–271. doi: 10.1016/j.bbrc.2011.01.026
- Hwang, J. H., Choi, H., Kim, A. R., Yun, J. W., Yu, R., Woo, E. R., et al. (2014). Hibicislide C-induced cell death in *Candida albicans* involves apoptosis mechanism. *J. Appl. Microbiol.* 117, 1400–1411. doi: 10.1111/jam.12633
- Hwang, J. H., Hwang, I. S., Liu, Q. H., Woo, E. R., and Lee, D. G. (2012). (+)-Medioresinol leads to intracellular ROS accumulation and mitochondria-mediated apoptotic cell death in *Candida albicans*. *Biochimie* 94, 1784–1793. doi: 10.1016/j.biochi.2012.04.010
- Hwang, J. H., Jin, Q., Woo, E. R., and Lee, D. G. (2013). Antifungal property of hibicislide C and its membrane-active mechanism in *Candida albicans*. *Biochimie* 95, 1917–1922. doi: 10.1016/j.biochi.2013.06.019
- Ingolfssdottir, K., Hjalmarsdottir, M. A., Sigurdsson, A., Gudjonsdottir, G. A., Brynjolfssdottir, A., and Steingrimsdottir, O. (1997). In vitro susceptibility of *Helicobacter pylori* to protolichsterinic acid from the lichen *Cetraria islandica*. *Antimicrob. Agents Chemother.* 41, 215–217.
- Joshi, B., Sah, G. P., Basnet, B. B., Bhatt, M. R., Sharma, D., Subedi, K., et al. (2011). Phytochemical extraction and antimicrobial properties of different medicinal plants: *Ocimum sanctum* (Tulsi), *Eugenia caryophyllata* (Clove), *Achyranthes bidentata* (Datiwan) and *Azadirachta indica* (Neem). *J. Microbiol. Antimicrob.* 3, 1–7.
- Khan, A., Ahmad, A., Khan, L. A., and Manzoor, N. (2014). *Ocimum sanctum* (L.) essential oil and its lead molecules induce apoptosis in *Candida albicans*. *Res. Microbiol.* 165, 411–419.
- Khodavandi, A., Alizadeh, F. A., Vanda, N. A., Karimi, G., and Chong, P. P. (2014). Possible mechanisms of the antifungal activity of fluconazole in combination with terbinafine against *Candida albicans*. *Pharm. Biol.* 52, 1505–1509. doi: 10.3109/13880209.2014.900808
- Kingston, D. G. I. (2011). Modern natural products drug discovery and its relevance to biodiversity conservation. *J. Nat. Prod.* 74, 496–511. doi: 10.1021/np100550t
- Kothavade, R. J., Kura, M. M., Yaland, A. G., and Panthaki, M. H. (2010). *Candida tropicalis*: its prevalence, pathogenicity and increasing resistance to fluconazole. *J. Med. Microbiol.* 59, 873–880. doi: 10.1099/jmm.0.013227-0
- Kumar, S. N., Sreerag, R. S., Ragesh, L., Jubi, J., Dileep Kumar, B. S., and Nambisan, B. (2014). Protolichsterinic acid: a prominent broad spectrum antimicrobial compound from the lichen *Usnea albobunctata*. *Int. J. Antibiot.* 2014:302182.
- Lee, J., and Lee, D. G. (2014). Melittin triggers apoptosis in *Candida albicans* through the reactive oxygen species-mediated mitochondria/caspase-dependent pathway. *FEMS Microbiol. Lett.* 355, 36–42. doi: 10.1111/1574-6968.12450
- Leite, M. C., de Brito Bezerra, A. P., de Sousa, J. P., and de Oliveira, L. E. (2015). Investigating the antifungal activity and mechanism(s) of geraniol against *Candida albicans* strains. *Med. Mycol.* 53, 275–284. doi: 10.1093/mmy/nyu078
- Li, Y., Chang, W., Zhang, M., Li, X., Jiao, Y., and Lou, H. (2015). Diorninol D exerts fungicidal action against *Candida albicans* through cytoplasm membrane destruction and ROS accumulation. *PLoS ONE* 10:e0128693. doi: 10.1371/journal.pone.0128693
- Liu, S., Yue, L., Gu, W., Li, X., Zhang, L., and Sun, S. (2016). Synergistic effect of fluconazole and calcium channel blockers against resistant *Candida albicans*. *PLoS ONE* 11:e0150859. doi: 10.1371/journal.pone.0150859
- Ma, C., Faya, D., Lan, Y., Gonghao, H., Jianchang, H., Chengying, W., et al. (2015). Potent activities of Roemerine against *Candida albicans* and the underlying mechanisms. *Molecules* 20, 17913–17928. doi: 10.3390/molecules20107913
- Machida, K., and Tanaka, T. (1999). Farnesol-induced generation of reactive oxygen species dependent on mitochondrial transmembrane potential hyperpolarization mediated by F_0F_1 -ATPase in yeast. *FEBS Lett.* 462, 108–112. doi: 10.1016/S0014-5793(99)01506-9
- Mitrović, T., Stamenković, S., Cvetković, V., Nikolić, M., Tošić, S., and Stojičić, D. (2011). Lichens as source of versatile bioactive compounds. *Biol. Nyssana* 2, 1–6.
- Müller, K. I. (2001). Pharmaceutically relevant metabolites from lichens. *Appl. Microbiol. Biotechnol.* 56, 9–16. doi: 10.1007/s002530100684
- Nguyen, T. T., Yoon, S., Yang, Y., Lee, H. B., Oh, S., Jeong, M. H., et al. (2014). Lichen secondary metabolites in *Flavocetraria cucullata* exhibit anticancer effects on human cancer cells through the induction of apoptosis

- and suppression of tumorigenic potentials. *PLoS ONE* 9:e11157. doi: 10.1371/journal.pone.0111575
- Nissapatorn, V., Lee, C., Fatt, Q. K., and Abdullah, K. A. (2003). AIDS related opportunistic infections in Hospital Kuala Lumpur. *Jpn. J. Infect. Dis.* 56, 187–192.
- Richardson, M. D. (2005). Changing patterns and trends in systemic fungal infections. *J. Antimicrob. Chemother.* 56, i5–i11. doi: 10.1093/jac/dki218
- Russo, A., Caggia, S., Piovano, M., Garbarin, J., and Cardile, V. (2012). Effect of vicanicin and protolichsterinic acid on human prostate cancer cells: role of Hsp70 protein. *Chem. Biol. Interact.* 195, 1–10. doi: 10.1016/j.cbi.2011.10.005
- Sharma, M., Biswas, D., Kotwal, A., Thakuria, B., Kakati, B., and Chauhan, B. S. (2015). Ibuprofen-mediated reversal of fluconazole resistance in clinical isolates of *Candida*. *J. Clin. Diagn. Res.* 9, DC20–DC22. doi: 10.7860/JCDR/2015/10094.5494
- Shrestha, G., and Clair, L. L. S. (2013). Lichens: a promising source of antibiotic and anticancer drugs. *Phytochem. Rev.* 12, 229–244. doi: 10.1007/s11101-013-9283-7
- Smrz, D., Draberova, L., and Draber, P. (2007). Non-apoptotic phosphatidylserine externalization induced by engagement of glycosylphosphatidylinositol-anchored proteins. *J. Biol. Chem.* 282, 10487–10497. doi: 10.1074/jbc.M611090200
- Srivastava, P., Upreti, D. K., Dhole, T. N., Srivastava, A. K., and Nayak, M. T. (2013). Antimicrobial property of extracts of Indian lichen against human pathogenic bacteria. *Int. Perspect. Infect. Dis.* 2013, 709348. doi: 10.1155/2013/709348
- Suzuki, M. T., Parrot, D., Berg, G., Grube, M., and Tomasi, S. (2016). Lichens as natural sources of biotechnologically relevant bacteria. *Appl. Microbiol. Biotechnol.* 100, 583–595.
- Suzuki, T., Fujikura, K., Higashiyama, T., and Takata, K. (1997). DNA staining for fluorescence and laser confocal microscopy. *J. Histochem. Cytochem.* 45, 49–53. doi: 10.1177/002215549704500107
- Thorsteinsdottir, U. A., Thorsteinsdottir, M., and Lambert, I. H. (2016). Protolichsterinic acid, isolated from the lichen *Cetraria islandica*, reduces LRRC8A expression and volume-sensitive release of organic osmolytes in human lung epithelial cancer cells. *Phytother. Res.* 30, 97–104. doi: 10.1002/ptr.5507
- Turk, A. O., Yilmaz, M., Kivanc, M., and Turk, H. (2003). The antimicrobial activity of extracts of the lichen *Cetraria aculeata* and its protolichsterinic acid constituent. *Z. Naturforsch. C* 58, 850–854. doi: 10.1515/znc-2003-11-1219
- Wang, X., Xiumin, W., Da, T., Yong, Z., Ruoyu, M., Di, X., et al. (2014). Candidacidal mechanism of the arenicin-3-derived peptide NZ17074 from *Arenicola marina*. *Appl. Microbiol. Biotechnol.* 98, 7387–7398. doi: 10.1007/s00253-014-5784-6
- Weghuber, J., Lipp, A. M., Stadlbauer, J., Aichinger, M. C., Ruprecht, V., Sonnleitner, A., et al. (2010). Antimicrobial and immunostimulatory peptide, KLK, induces an increase in cytosolic Ca^{2+} concentration by mobilizing Ca^{2+} from intracellular stores. *Cell Biol. Int.* 34, 1109–1112. doi: 10.1042/CBI20100408
- Wonyoung, L., and Dong Gun, L. (2014). An antifungal mechanism of curcumin lies in membrane-targeted action within *Candida albicans*. *IUBMB Life* 66, 780–785. doi: 10.1002/iub.1326
- Yoon, M. J., Kim, E. H., Kwon, T. K., Park, S. A., and Choi, K. S. (2012). Simultaneous mitochondrial Ca^{2+} overload and proteasomal inhibition are responsible for the induction of paraptosis in malignant breast cancer cells. *Cancer Lett.* 324, 197–209. doi: 10.1016/j.canlet.2012.05.018
- Yu, Q., Ding, X., Zhang, B., Xu, N., Jia, C., Mao, J., et al. (2014). Inhibitory effect of verapamil on *Candida albicans* hyphal development, adhesion and gastrointestinal colonization. *FEMS Yeast Res.* 34, 633–641. doi: 10.1111/1567-1364.12150
- Yu, Q., Zhang, B., Ma, F., Jia, C., Xiao, C., Zhang, B., et al. (2015). Novel mechanisms of surfactants against *Candida albicans* growth and morphogenesis. *Chem. Biol. Interact.* 227, 1–6. doi: 10.1016/j.cbi.2014.12.014
- Yun, J., Lee, H., Kob, H. J., Woo, E. R., and Lee, D. G. (2015). Fungicidal effect of isoquercitrin via inducing membrane disturbance. *Biochim. Biophys. Acta* 1848, 695–701. doi: 10.1016/j.bbame.2014.11.019

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Kumar and Mohandas. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.