



Efficacy and Safety of Tenueligliptin in Patients With Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis of Randomized Controlled Trials

Xiaoxuan Li^{1†}, Xuefei Huang^{2†}, Chongfei Bai^{1,3†}, Dalian Qin¹, Shousong Cao⁴, Qibing Mei¹, Yun Ye^{2,5*} and Jianming Wu^{1*}

¹ Laboratory of Chinese Materia Medica, Department of Pharmacology, School of Pharmacy, Southwest Medical University, Luzhou, China, ² Department of Clinical Pharmacy, School of Pharmacy, Southwest Medical University, Luzhou, China, ³ Department of Chinese Materia Medica, School of Pharmacy, Chengdu University of Traditional Chinese Medicine, Chengdu, China, ⁴ Laboratory of Cancer Pharmacology, Department of Pharmacology, School of Pharmacy, Southwest Medical University, Luzhou, China, ⁵ Department of Pharmacy, Affiliated Hospital of Southwest Medical University, Luzhou, China

OPEN ACCESS

Edited by:

Brian Godman,
Karolinska Institute (KI), Sweden

Reviewed by:

Godfrey Mutashambara Rwegerera,
University of Botswana, Botswana
Gustavo Laine Araújo De Oliveira,
Ministry of Health (Brazil), Brazil

*Correspondence:

Jianming Wu
jianmingwu@swmu.edu.cn
Yun Ye
yeyun8622@163.com

[†]These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Pharmaceutical Medicine and
Outcomes Research,
a section of the journal
Frontiers in Pharmacology

Received: 12 February 2018

Accepted: 17 April 2018

Published: 04 May 2018

Citation:

Li X, Huang X, Bai C, Qin D, Cao S,
Mei Q, Ye Y and Wu J (2018) Efficacy
and Safety of Tenueligliptin in Patients
With Type 2 Diabetes Mellitus: A
Systematic Review and Meta-Analysis
of Randomized Controlled Trials.
Front. Pharmacol. 9:449.
doi: 10.3389/fphar.2018.00449

Background: Tenueligliptin is a 3rd-generation dipeptidyl peptidase-4 (DPP-4) inhibitor. There is a limited evidence regarding the effect of tenueligliptin. Therefore, this study is to assess the efficacy and safety of tenueligliptin in type 2 diabetes mellitus (T2DM) patients with inadequately glycemic controlled.

Methods: A search of PubMed, Medline, Embase, and The Cochrane Library during 2000.01–2018.03 was performed for randomized controlled trials of tenueligliptin compared to placebo in patients with T2DM with monotherapy or add-on treatment.

Results: Ten trials with 2119 patients were analyzed. Tenueligliptin produced absolute reductions in glycated hemoglobin A1c (HbA1c) levels (weighted mean difference (WMD) 0.82%, 95% confidence interval (CI) [−0.91 to −0.72], $p < 0.00001$) compared with placebo. However, after 36–42 weeks of follow-up (open-label), HbA1c level rise higher than duration (double-blind) in tenueligliptin group. Tenueligliptin led to greater decrease of fasting plasma glucose (FPG) level (vs. placebo, WMD −18.32%, 95% CI [−21.05 to −15.60], $p < 0.00001$). Tenueligliptin also significantly decreased the 2 h post-prandial plasma glucose (2 h PPG) (WMD −46.94%, 95% CI [−51.58 to −42.30], $p < 0.00001$) and area under the glucose plasma concentration-time curve from 0 to 2 h (AUC_{0–2h}) for PPG (WMD −71.50%, 95% CI [−78.09 to −64.91], $p < 0.00001$) compared with placebo. Patients treated with tenueligliptin achieved increased homeostasis model assessment of β cell function (HOMA- β) with 9.31 (WMD, 95% CI [7.78–10.85], $p < 0.00001$). However, there was no significant difference between tenueligliptin and placebo in overall adverse effects (0.96 risk ratio (RR), 95% CI [0.87, 1.06], $p = 0.06$). The risks of hypoglycemia were not significantly different between tenueligliptin and placebo (1.16 RR, 95% CI [0.59, 2.26], $p = 0.66$).

Conclusions: Teleniglipitin improved blood glucose levels and β -cells function with low risk of hypoglycemia in patients with T2DM. Common adverse effects of teleniglipitin including hypoglycemia were identified and reviewed. Risks of cardiovascular events are less certain, and more data for long-term effects are needed.

Keywords: teleniglipitin, dipeptidyl peptidase-4 (DPP-4) inhibitor, type 2 diabetes mellitus (T2DM), systematic review, glycemic control

INTRODUCTION

Approximately 425 million adults (one in eleven) are living with diabetes and one in two adults remains undiagnosed worldwide (2017)¹. Type 2 diabetes mellitus (T2DM) accounts for more than 90% cases of diabetes. T2DM induces microvascular and macrovascular complications, which place a huge burden on patients, caregivers, and health care systems (Chatterjee et al., 2017). A recent statement of Standards of Medical Care in Diabetes by the American Diabetes Association (ADA) has recommended that initial treatment with metformin as monotherapy after inadequate life style modification, followed by sulfonylurea, thiazolidinedione, dipeptidyl peptidase-4 (DPP-4) inhibitor, sodium-glucose cotransporter 2 inhibitor (SGLT2-i), glucagon-like peptide 1 (GLP-1) receptor agonist and insulin alone or in combination (American Diabetes Association, 2017). However, it is still difficult to find an antihyperglycemic agent with long-term glucose control, minimal hypoglycemia, no weight gain and a relatively affordable price (Liao, 2011). DPP-4 inhibitors have been considered as a cornerstone in the management of T2DM because of their robust efficacy and favorable tolerability profiles (Deacon, 2011).

A large body of randomized controlled trials (RCTs) showed that the DPP4 inhibitors such as sitagliptin and vildagliptin were effective for blood glycemic control by declining the HbA1c, FPG, and PPG and improving the function of pancreatic α and β cells. In addition to targeting glycemic control, DPP4 inhibitors have low risk of hypoglycaemia with neutral effect of body weight with a favorable safety profile (Inzucchi et al., 2012; Garber et al., 2015). Clinical trials showed that the relative common adverse events of DPP4 inhibitors are gastrointestinal symptoms, nasopharyngitis, upper respiratory infections and headache in clinical trials. Other less common adverse event such as skin- and immune-related effects are reported in post-marketing surveillance (Filippatos et al., 2014).

Teleniglipitin, a 3rd-generation DPP-4 inhibitor, acts as a competitive reversible inhibitor of DPP-4 and decreases the degradation of incretins, especially GLP-1, consequently stimulating insulin secretion and suppressing glucagon secretion

in a glucose-dependent manner (Gallwitz, 2010). More importantly, teleniglipitin is effective and safe for patients with T2DM with renal impairment, or even end-stage renal disease, without dose adjustments (Abubaker et al., 2017). Teleniglipitin was synthesized in Japan and is available in Japan, Argentina, Korea and India. It is currently in phase I clinical trials in the USA and phase II clinical trials in Europe for management of T2DM (Kishimoto, 2013; Scott, 2015). A few clinical studies showed that teleniglipitin significantly improves glycemic control, is well tolerated, and causes a low incidence of hypoglycemia when used as monotherapy or combination therapy (Eto et al., 2012; Kadowaki and Kondo, 2013a,b,c; Kim et al., 2015; Bryson et al., 2016; Hong et al., 2016; Kadowaki et al., 2017a). However, there were only few reports of comprehensive profiles of the benefits and risks of teleniglipitin in patients with T2DM to date. Thus, we conducted a systematic review and meta-analysis to assess the efficacy and safety of teleniglipitin in the management of T2DM either as monotherapy or add-on treatment. In adults, teleniglipitin is primarily metabolized by cytochrome P450 (CYP) 3A4 and flavin monooxygenases (FMO) (Patel et al., 2016). Approximately 34% of each administered dose of teleniglipitin is excreted unchanged via the renal route, while 66% is metabolized and eliminated via the hepatic and renal routes (Sharma et al., 2016). Teleniglipitin is usually orally administered at 20 mg once daily and increased to 40 mg once daily if the dosage is insufficient (Kishimoto, 2013). Thus, we evaluated 20 mg of teleniglipitin once daily in the treatment of T2DM patients.

MATERIALS AND METHODS

Search Strategy and Study Selection

According to the preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement (Liberati et al., 2009), we systematically searched PubMed, Medline, Embase and the Cochrane Library from inception to March 2018 without a language restriction. Clinical Trials (<http://www.clinicaltrials.gov>) were also searched. We used the terms “Teleniglipitin,” “MP-513,” “T2DM,” and “Type 2 diabetes mellitus” for the searches, and the terms were adjusted to conform to the relevant rules in each database. We also used (“Diabetes Mellitus, Type 2”[Mesh] AND teleniglipitin) to search the data. The PROSPERO registration number is CRD42018091232.

Titles and abstracts of all retrieved citations were screened by two independent reviewers (L.X.X and H.X.F) to identify all potentially eligible studies. Full texts were retrieved for relevant

Abbreviations: T2DM, type 2 diabetes mellitus; DPP-4, dipeptidyl peptidase-4; HbA1c, glycated hemoglobin A1c; WMD, weighted mean difference; CI, confidence interval; RR, risk ratio; FPG, fasting plasma glucose; PPG, post-prandial plasma glucose; HOMA- β , Homeostasis model assessment of β cell function; HOMA-IR, homeostasis model assessment of insulin resistance; AEs, adverse events; RCTs, randomized controlled trials; AUC_{0–2h}, area under the glucose plasma concentration-time curve from 0 to 2 h.

¹IDF diabetes atlas - 8th edition [EB/OL]. <http://www.diabetesatlas.org/>

records. Any resulting discrepancies were resolved by discussion, with involvement of a third reviewer if necessary.

Inclusion and Exclusion Criteria

Trials of teneligliptin for the treatment of T2DM according to the WHO diagnostic criteria were included if they met the following criteria: (1) Patients: any ethnic origin and aged over 18. (2) Interventions: any use of teneligliptin as monotherapy or

combination therapy, with a duration of intervention of at least 4 weeks. (3) Comparison: placebo or active comparators with or without background therapy. (4) Outcomes: at least one of the following indicators was reported: (a) glycated hemoglobin A1c (HbA1c), (b) fasting plasma glucose (FPG), (c) 2 h post-prandial plasma glucose (2 h PPG), (d) area under the glucose plasma concentration-time curve from 0 to 2 h (AUC_{0-2h}) PPG, (e) homeostasis model assessment of β cell function (HOMA- β),




TABLE 1 | Characteristics of randomized controlled trials.

Study, Year	Participants	Interventions	Age (years)	BMI (kg/m ²)	HbA1c (%)	FPG (mg/dL)	Body weight (kg)	T2DM duration (years)	Background therapy	Phase	Duration (extension)
Kadowaki et al., 2017a	N: 154	Teneeligliptin(20 mg)	55.9 ± 8.3	25.53 ± 3.95	7.98 ± 0.80	148.5 ± 21.2	72.32 ± 12.08	8.15 ± 5.86	Add on to canagliflozin	III	24 w (2 w)
		Placebo	54.1 ± 10.2	26.50 ± 4.82	8.09 ± 0.85	151.5 ± 25.3	73.68 ± 15.75	7.34 ± 5.34			
Kadowaki et al., 2017b	N: 148	Teneeligliptin(20 mg)	60.1 ± 11.3	24.87 ± 3.14	8.70 ± 0.81	16.9 ± 40.4	67.62 ± 12.98	11.85 ± 8.14	Add on to insulin	IV	16 w (36 w)
		Placebo	57.4 ± 12	25.11 ± 3.66	8.73 ± 0.81	162.8 ± 45.0	68.61 ± 13.68	12.97 ± 9.22			
Bryson et al., 2016	N: 447	Teneeligliptin(20 mg)	58.3 ± 9.5	NR	NR	NR	NR	NR	Add on to metformin	II	24 w (28 w)
		Placebo	58.9 ± 8.2								
Hong et al., 2016	N: 142	Teneeligliptin(20 mg)	56.64 ± 10.07	24.96 ± 2.51	7.74 ± 0.61	155.96 ± 26.14	65.81 ± 11.39	4.59 ± 3.87	Monotherapy	III	24 w (2 w)
		Placebo	57.93 ± 11.90	25.07 ± 3.23	7.74 ± 0.53	162.09 ± 31.3	66.61 ± 12.15	4.59 ± 3.94			
Kim et al., 2015	N: 204	Teneeligliptin(20 mg)	55.7 ± 8.7	NR	7.79 ± 0.80	151.32 ± 35.55	NR	6.7 ± 4.8	Add on to metformin	III	16 w
		Placebo	56.4 ± 9.2		7.72 ± 0.65	151.31 ± 25.86	NR	8.0 ± 5.9			
NCT00988881 2014	N: 203	Teneeligliptin(20 mg)	58.5 ± 10.0	NR	NR	NR	NR	NR	Monotherapy	III	12 w
		Placebo	59.2 ± 10.1								
Kadowaki and Kondo, 2013a	N:324	Teneeligliptin(20 mg)	59.2 ± 9.5	24.9 ± 3.9	7.8 ± 0.7	143.0 ± 26.8	NR	6.3 ± 6.4	Monotherapy	II	12 w (2 w)
		Placebo	58.5 ± 9.6	25.2 ± 3.9	8.0 ± 0.7	150.0 ± 30.3	NR	5.8 ± 5.0			
Kadowaki and Kondo, 2013b	N: 204	Teneeligliptin(20 mg)	59.7 ± 9.7	26.2 ± 5.2	8.1 ± 0.9	150.7 ± 28.1	70.0 ± 16.6	7.2 ± 4.8	Add on to pioglitazone	III	12 w (42 w)
		Placebo	61.1 ± 8.9	25.6 ± 3.7	7.9 ± 0.8	145.7 ± 26.5	67.7 ± 13.0	7.7 ± 6.1			
Kadowaki and Kondo, 2013c	N: 194	Teneeligliptin(20 mg)	58.4 ± 8.6	24.9 ± 3.6	8.4 ± 0.8	165.1 ± 24.5	66.2 ± 12.1	9.3 ± 6.7	Add on to glimepiride	III	12 w (42 w)
		Placebo	60.3 ± 7.8	24.6 ± 3.6	8.4 ± 0.8	163.4 ± 31.3	65.9 ± 11.8	8.3 ± 6.2			
Eto et al., 2012	N: 99	Teneeligliptin(20 mg)	57.1 ± 8.7	24.8 ± 3.8	8.3 ± 0.8	163.1 ± 30.8	NR	6.4 ± 5.9	Monotherapy	NR	24 w
		Placebo	58.6 ± 8.9	25.7 ± 4.5	8.2 ± 1.1	153.6 ± 31.9	NR	7.8 ± 6.2			

BMI, body mass index; HbA1c, glycated hemoglobin; FPG, fasting plasma glucose; T2DM, type 2 diabetes mellitus; NR not report NCT00988881. This study have not been published. Inclusion criteria: patients in the study must have an HbA1c level between 6.5 and 10.0%.

homeostasis model assessment of insulin resistance (HOMA-IR), and (f) adverse events (AEs) such as hypoglycemia. (5) Trial design: randomized controlled trials (RCTs) published or searched in Clinical Trials without language restrictions. We excluded the studies of non-randomized trials, case reports, editorials, letters to the editors, and conference abstracts.

Data Extraction and Risk of Bias Assessment

Two reviewers (L.X.X and H.X.F) independently screened the title and abstract and extracted study characteristics, baseline characteristics, and prespecified outcomes of efficacy and safety. They also independently assessed the risk of bias of randomized controlled trials (Higgins and Green, 2011),

including random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, and other bias (i.e., design-specific risks of bias, baseline imbalance, blocked randomization in unblinded trials) with the Cochrane Risk of Bias tool. Any discrepancies were resolved by discussion. We mainly focused on the data for patients randomly assigned to teneigliptin 20 mg/day, which is the most common dose used in clinical practice.

Statistical Analysis

All outcomes were pooled using RevMan 5.3. The efficacy was evaluated by weighted mean difference (WMD, indicators changed from baseline), along with 95% confidence intervals




FIGURE 2 | Summary (A) and graph (B) of the risk of bias in the included trials by Cochrane risk of bias toll based upon reviewers' judgment of each domain.

(CIs). The safety was assessed by risk ratios (RRs, incidence of AEs or hypoglycemia), along with 95% CIs. Heterogeneity was assessed by the chi-square test and the I^2 statistic. If $I^2 < 50\%$, the fixed-effect model with the Mantel-Haenszel method was used; otherwise, the random-effect model was adopted. If the primary outcome data standard deviation (SD) were missing or incomplete, we e-mailed the corresponding authors or the sponsors to obtain them. When necessary, the values of SD were calculated from SE as described in the Cochrane Handbook.

RESULTS

Study Selection and Characteristics

We identified 251 publications in four databases, and 132 were excluded for duplication. Nineteen were remained after further removing due to non-human studies, irrelevant data, review or meta-analysis. Finally, a total of 10 RCTs (Eto et al., 2012; Kadowaki and Kondo, 2013a,b,c; Mitsubishi Tanabe Pharma Corporation, 2014; Kim et al., 2015; Bryson et al., 2016; Hong et al., 2016; Kadowaki et al., 2017a,b) ($n = 2119$) met the final inclusion criteria for meta-analysis after excluding 10 and adding one study (Figure 1).

The characteristics of the included RCTs are shown in Table 1. All included trials were double-blind RCTs; two were phase II (Kadowaki and Kondo, 2013c; Bryson et al., 2016), six were phase III (Kadowaki and Kondo, 2013a,b; Mitsubishi Tanabe Pharma Corporation, 2014; Kim et al., 2015; Hong et al., 2016; Kadowaki et al., 2017b), and one was phase IV (Kadowaki et al., 2017a). Trial durations ranged from 4 to 24 weeks. Seven trials had extension periods (ranging from 2 to 42 weeks) (Kadowaki and Kondo, 2013a,b,c; Bryson et al., 2016; Hong et al., 2016; Kadowaki et al., 2017a,b). Mean baseline HbA1c across the study populations

ranged from 7.72 to 8.73%; mean baseline FPG ranged from 143.0 to 165.1 mg/dL. Participants in most trials were mainly middle-aged and overweight adults who had T2DM for more than 4 years. Mean age ranged from 55.9 to 60.4 years. Body mass index (BMI) in most trials ranged from 24.8 to 26.5 kg/m².


Tenueligliptin was administered before a standard meal at a dose of 20 mg once daily and the efficacy and safety were compared to placebo in all trails. Four trials were monotherapy and six add-on treatments, and the background therapies were canagliflozin, insulin, metformin, pioglitazone, and glimepiride.

Risk of Bias

All the included studies were randomized, double-blind, placebo-controlled studies and had a low risk for bias, as evaluated by The Cochrane Collaboration's tool for assessing risk of bias (Figure 2). However, only four of ten studies elaborated the generation of random sequences and only one record of the allocation concealment.

HbA1c

The effects of tenueligliptin vs. placebo on the HbA1c change from baseline are shown in Figure 3. Tenueligliptin significantly reduced HbA1c (WMD -0.82% , 95% CI $[-0.91$ to $-0.72]$, $p < 0.00001$) as monotherapy (WMD -0.86% , 95% CI $[-0.95$ to $-0.76]$, $p < 0.00001$), or add-on treatment (WMD -0.79% , 95% CI $[-0.93$ to $-0.66]$, $p < 0.00001$) compared to placebo. Analyses of Subgroup didn't reduce the high level of heterogeneity with different drugs and different treatment duration (Figures 4, 5). Removing two studies (Kadowaki and Kondo, 2013c; Bryson et al., 2016) because of larger effect size than other trials, the heterogeneity and effect size of HbA1c reduced significantly






FIGURE 4 | Effect of tenueligliptin on HbA1c with different background therapy compared to placebo.

(-0.82%, -0.89 to -0.76; I² = 0%). 36–42 weeks of follow-up didn't show better decline of HbA1c in tenueligliptin group (Figure 6). A greater proportion of subjects received tenueligliptin achieved the target of HbA1c < 7% (RR 3.99, 95% CI [2.98–5.34], *p* < 0.00001) compared to placebo (Figure 7).

FPG

A significant decrease from the baseline in FPG level was also observed in the tenueligliptin group compared to placebo (WMD -18.32%, 95% CI [-21.05 to -15.60], *p* < 0.00001) (Figure 8) as monotherapy (WMD -17.47%, 95% CI [-20.70 to -14.24], *p* < 0.00001), or add-on treatment (WMD -18.85%, 95% CI [-23.38 to -14.31], *p* < 0.00001).

2h PPG and AUC_{0–2h} for PPG

Tenueligliptin significantly decreased the 2h PPG (WMD -46.94%, 95% CI [-51.58 to -42.30], *p* < 0.00001) and

AUC_{0–2h} of PPG (WMD -71.50%, 95% CI [-78.09 to -64.91], *p* < 0.00001) compared to placebo (Figures 9, 10). Tenueligliptin reduced 2h PPG by 47.28 mg/dl as monotherapy (WMD, 95% CI [-54.26 to -40.29], *p* < 0.00001) and by -46.67 mg/dl as combination therapy (WMD, 95% CI [-52.88 to -40.46], *p* < 0.00001). Similarly, the AUC_{0–2h} of PPG was diminished by 73.75 mg-h/dl in the tenueligliptin group compared to placebo as monotherapy (WMD, 95% CI [-83.54 to -63.96], *p* < 0.00001) and by 69.64 mg-h/dl as add-on (WMD, 95% CI [-78.54 to -60.74], *p* < 0.00001).

HOMA-β and HOMA-IR

Patients treated with tenueligliptin exhibited increased HOMA-β by 9.31 when all regimens were included (WMD, 95% CI [7.78–10.85], *p* < 0.00001), 9.18 when tenueligliptin was used as monotherapy (WMD, 95% CI [5.95–12.41], *p* < 0.00001) and




FIGURE 5 | Effect of teneligliptin on HbA1c with different duration of treatment compared to placebo.




FIGURE 6 | Comparative effect of teneligliptin in double-blind period vs. different follow-up time.




FIGURE 7 | The proportion of patients who achieved HbA1c < 7% treated with teneligliptin vs. placebo by meta-analysis.

9.35 when used as add-on treatment (WMD, 95% CI [7.61–11.09], $p < 0.00001$) (Figure 11). Only combined treatment showed statistical significance, with a decrease in HOMA-IR by -0.25 (WMD, 95% CI $[-0.47$ to $-0.03]$, $p = 0.03$) (Figure 12).

Overall AEs and Hypoglycemia

There was no significant difference between teneligliptin and placebo in overall AEs ($p > 0.05$; Figure 13). The incidence of hypoglycemia was low in all included patients, and there was no severe hypoglycemia in most studies. The risk of hypoglycemia




FIGURE 8 | Effects of teneligliptin and placebo on FPG change from the baseline by meta-analysis.




FIGURE 9 | Effects of teneligliptin and placebo on 2 h PPG change from the baseline by meta-analysis.

was similar between teneligliptin and placebo: 0.34 (RR, 95% CI [0.04,3.18], $p = 0.34$) in monotherapy and 1.36 (RR, 95% CI [0.66, 2.78], $p = 0.40$) in add-on treatment (Figure 14).

DISCUSSION

According to this meta-analysis, we found that teneligliptin was effective and safe in the treatment of T2DM. HbA1c level is a primary goal for glycemic control (Koren and Rapoport, 2014). Patients treated with teneligliptin had a greater decrease

in HbA1c levels from baseline than the placebo group when teneligliptin was used as monotherapy or add-on therapy to insulin or other antidiabetic drugs. Although we performed subgroup analyses, we found a high level of heterogeneity. We speculate that there are two reasons for the high level of heterogeneity. On one hand, accumulating evidence from clinical trials showed that incretin-based drugs were more effective in Asians due to the diverse pathophysiology of type 2 diabetes in different ethnic groups. Asian patients with type 2 diabetes are generally characterized by defective early phase




FIGURE 10 | Meta-analysis for Effects of teneligliptin and placebo on AUC_{0–2h} for PPG change from the baseline by meta-analysis.




FIGURE 11 | Effects of teneligliptin and placebo on HOMA-β change from the baseline by meta-analysis.

insulin secretion, DPP-4 inhibitors can improve impaired insulin secretion and then exert greater effects in HbA1c in patients with type 2 diabetes (Xueying and Jingbo, 2016). On the other hand, gene-lifestyle interplay in patients with type 2 diabetes from the complex interplay of lifestyle factors acts as a backdrop of inherited DNA risk variants (Paul and Jordi, 2018). It is reasonable to consider that an HbA1c range of 5.7–6.4% (39–47 mmol/mol) is a factor of increased risk for diabetes (American Diabetes Association, 2018). Ideally, the best glycemic control for diabetic patient is to decrease the HbA1c level as close to normal level as possible. Nevertheless, the HbA1c value that is commonly advocated as threshold value for satisfactory

glycemic control for most patients is 7%. This value, and less commonly a level of 6.5%, appears in guidelines from the United States, Canada, Europe, Australia and many other countries (Koren and Rapoport, 2014). We found that the effects of DPP4 inhibitors on HbA1c were different after 12 weeks of administration. Treatment of Sitagliptin showed an average decrease in HbA1c levels of 0.65% after 12 weeks, 0.84% after 18 weeks, 0.85% after 24 weeks, 1.0% after 30 weeks, and 0.67% after 52 weeks (Dror, 2011). In addition, omarigliptin reduced HbA1c levels after 66 weeks of follow-up, which was lower than 12 weeks (Philip and Steve, 2016). Our meta-analysis with 3 RCTs showed that the level of HbA1c increased




FIGURE 12 | Effects of teneligliptin and placebo on HOMA-IR change from the baseline by meta-analysis.




FIGURE 13 | The incidence of AEs of teneligliptin vs. placebo by meta-analysis.

after 36–42 weeks of follow-up time in teneligliptin group, and thus more high-quality and high volume samples were needed for reliable conclusion. A significantly larger proportion of patients treated with teneligliptin achieved an HbA1c level <7% compared to placebo, suggesting teneligliptin is effective in the treatment of patients with T2DM. Teneligliptin lowered

not only HbA1c but also other parameters of interest, such as FPG, 2h PPG and AUC_{0–2h} for PPG. These results indicate that teneligliptin improves glycemic control index and suggest it may be a useful treatment option for patients with T2DM who are inadequately glycemic controlled. For T2DM, body weight is an essential index. However, the included studies in




FIGURE 14 | The incidence of hypoglycemia in patients treated with teligliptin vs. placebo by meta-analysis.

this meta-analysis did not have enough data to assess body weight.

β -Cell destruction plays a key role in the pathophysiology of T2DM, and β -cell conservation delays disease progression (Hong et al., 2016). DPP-4 inhibitors are associated with enhanced β -cell function, making them a good therapeutic option in early disease when the patients still maintain sufficient levels of β -cell function (Kutuh et al., 2014). The HOMA model that is a method for assessing β -cell function and insulin resistance (IR) has proved as a robust clinical and epidemiological tool in evaluation of the pathophysiology of diabetes (Wallace et al., 2004). According to this study of meta-analysis, HOMA- β was significantly improved when teligliptin was used either as monotherapy or add-on treatment. These facts suggest that teligliptin has a beneficial effect on pancreatic β -cells and could modestly improve pancreatic function. However, no significant difference in HOMA-IR was observed in the present study.

Teligliptin has a unique structure with five consecutive rings. Thus, teligliptin acts on the S2 extensive subsite of DPP-4; this interaction enhances its potency and selectivity (Yoshida et al., 2012; Nabeno et al., 2013; Abubaker et al., 2017). Because of its high selectivity, most AEs of teligliptin were mild and led to few discontinuations in the meta-analysis. The incidences of AEs were not significantly different between the patients treated with teligliptin and placebo. Although hypoglycemia was the main adverse event for T2DM patients (Sharma et al., 2016), teligliptin led to a low risk of hypoglycemia. In addition, the cardiovascular effects of DPP-4 inhibitors remain controversial, while one DPP-4 inhibitor (saxagliptin) increased

the risk of hospitalization for heart failure in the overall population; another (alogliptin) showed inconsistent effects on heart failure hospitalization across subgroups of patients, and a third (sitagliptin) had no effect on heart failure (Secrest et al., 2017). The effect of teligliptin on cardiovascular complication remains unclear. Therefore, it is needed for rational design for longer period of dosing and more RCTs from multicenter for more reliable conclusions.

Teligliptin is still a relatively new drug, and published clinical studies concerning this drug are sparse (Kishimoto, 2013). The present study is the first meta-analysis about teligliptin, including 10 high qualities of RCTs. However, it still has several potential limitations. First, there were relatively few patients in the 10 RCTs, which limited our ability to reach clear conclusions about the effects and safety of teligliptin in the treatment of patients with T2DM. The use of GRADE system (Table 2) suggests that the classification is low or very low, and thus the estimates of the effects are insufficiently. Second, the durations and extensions of the 10 RCTs were relative short, and longer follow-up is needed to evaluate the long-term benefits and risks of teligliptin. Third, only placebo-controlled RCTs were conducted. Trials are need to be conducted to assess the therapeutic effect of teligliptin by comparing the effects of other active agents. Fourth, we did not assess the 24 h glucose fluctuations. Large fluctuations in glucose levels may increase the risk of complications, such as cardiovascular disease, so it is better to evaluate post-prandial glucose fluctuations over the entire 24 h dosing interval (Tanaka et al., 2014; Morishita and Nakagami, 2015). In addition,

TABLE 2 | Quality of included studies, assessed using the GRADE system.

No of studies	Design	Quality assessment					No of patients		Effect		Quality
		Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Tenepliptin vs. Placebo	Control	Relative (95% CI)	Absolute	
HbA1c											
9	Randomized trials	Not serious	Serious ^a	Serious ^b	Not serious	Reporting bias ^c	857	730	–	MD 0.82 lower (0.91–0.72 lower)	⊕○○○ VERY LOW
HbA1c<24 weeks											
6	Randomized trials	Not serious	Not serious	Serious ^b	Not serious	Reporting bias ^c	590	522	–	MD 0.84 lower (0.92–0.76 lower)	⊕⊕○○ LOW
HbA1c ≥24 weeks											
3	Randomized trials	Not serious	Not serious	Serious ^b	Serious ^d	Reporting bias ^c	267	208	–	MD 0.77 lower (1.08–0.45 lower)	⊕○○○ VERY LOW
HbA1c<7%											
5	Randomized trials	Not serious	Serious ^e	Not serious	Serious ^d	Reporting bias ^c	257/480 (53.5%)	43/347 (12.4%)	RR 3.99 (2.98–5.34)	371 more per 1000 (from 245 more to 538 more)	⊕○○○ VERY LOW
FPG											
10	Randomized trials	Not serious	Not serious	Serious ^b	Not serious	Reporting bias ^c	890	760	–	MD 18.32 lower (21.05–15.6 lower)	⊕⊕○○ LOW
2 h-PPG											
7	Randomized trials	Not serious	Not serious	Serious ^b	Not serious	Reporting bias ^c	547	526	–	MD 46.94 lower (51.58–42.3 lower)	⊕⊕○○ LOW
PPG AUC 0–2h											
7	Randomized trials	Not serious	Not serious	Serious ^b	Not serious	Reporting bias ^c	537	526	–	MD 71.5 lower (78.09–64.91 lower)	⊕⊕○○ LOW
HOMA-IR											
6	Randomized trials	Not serious	Serious ^e	Serious ^b	Not serious	Reporting bias ^c	604	475	–	MD 0.17 lower (0.34 lower to 0 higher)	⊕○○○ VERY LOW
HOMA-β											
7	Randomized trials	Not serious	Not serious	Serious ^b	Not serious	Reporting bias ^c	681	551	–	MD 9.31 higher (7.78–10.85 higher)	⊕⊕○○ LOW
AEs											
9	Randomized trials	Not serious	Not serious	Not serious	Not serious	Reporting bias ^c	376/790 (47.6%)	338/658 (51.4%)	RR 0.96 (0.87–1.06)	21 fewer per 1,000 (from 67 fewer to 31 more)	⊕⊕⊕○ MODERATE
Hypoglycemia											
8	Randomized trials	Not serious	Serious ^e	Not serious	Serious ^d	Reporting bias ^c	19/692 (2.7%)	14/615 (2.3%)	RR 1.16 (0.59–2.26)	4 more per 1,000 (from 9 fewer to 29 more)	⊕○○○ VERY LOW

^aHigh level of heterogeneity was observed despite performing subgroup analysis.
^bSurrogate marker.
^cPublication bias strongly suspected.
^dVery small sample size.
^eLarge variation in effect.

there is a lack of cost-effectiveness studies of teneeligliptin. More data will be necessary for a better, more comprehensive analysis.

CONCLUSION

This meta-analysis suggests that treatment of teneeligliptin provided clinically and statistically significant reductions in HbA1c and FPG levels in patients with T2DM. These effects were associated with significant improvements in β -cell function. Furthermore, the incidences of AEs were not significantly in patients treated with teneeligliptin compared to placebo. Therefore, the present study demonstrated that teneeligliptin exhibits beneficial effects in T2DM patients. However, it is warranted to further investigate with more rational, and longer duration of drug administration from multicenter study and ongoing monitoring due to the potential risks of unclear AEs.

REFERENCES

- Abubaker, M., Mishra, P., and Swami, O. (2017). Teneeligliptin in management of diabetic kidney disease: a review of place in therapy. *J. Clin. Diagn. Res.* 11, E5–E9. doi: 10.7860/JCDR/2017/25060.9228
- American Diabetes Association (2017). Pharmacologic approaches to glycemic treatment. *Diabetes Care* 40, S64–S74. doi: 10.2337/dc17-S011
- American Diabetes Association. (2018). Classification and diagnosis of diabetes: standards of medical care in diabetes. *Diabetes Care* 41, S13–S27. doi: 10.2337/dc18-s002
- Bryson, A., Jennings, P., Deak, L., Paveliu, F., and Lawson, M. (2016). The efficacy and safety of teneeligliptin added to ongoing metformin monotherapy in patients with type 2 diabetes: a randomized study with open label extension. *Expert Opin. Pharmacother.* 17, 1309–1316. doi: 10.1080/14656566.2016.1190334
- Chatterjee, S., Davies, M., and Khunti, K. (2017). Type 2 diabetes. *Lancet* 389, 2239–2251. doi: 10.1016/S0140-6736(17)30058-2
- Deacon, C. (2011). Dipeptidyl peptidase-4 inhibitors in the treatment of type 2 diabetes: a comparative review. *Diabetes Obes. Metab.* 13, 7–18. doi: 10.1111/j.1463-1326.2010.01306.x
- Dror, D. (2011). DPP-4 inhibitors: impact on glycemic control and cardiovascular risk factors. *Diabetes Care* 34, S276–S278. doi: 10.2337/dc11-s229
- Eto, T., Inoue, S., and Kadowaki, T. (2012). Effects of once-daily teneeligliptin on 24-h blood glucose control and safety in Japanese patients with type 2 diabetes mellitus: a 4-week, randomized, double-blind, placebo-controlled trial. *Diabetes Obes. Metab.* 14, 1040–1046. doi: 10.1111/j.1463-1326.2012.01662.x
- Filippatos, T., Athyros, V., and Elisaf, M. (2014). The pharmacokinetic considerations and adverse effects of DPP-4 inhibitors. *Expert Opin. Drug Met.* 10, 787–812. doi: 10.1517/17425255.2014.907274
- Gallwitz, B. (2010). The evolving place of incretin-based therapies in type 2 diabetes. *Pediatr. Nephrol.* 25, 1207–1217. doi: 10.1007/s00467-009-1435-z
- Garber, A., Abrahamson, M., Barzilay, J., Blonde, L., Bloomgarden, Z., and Bush, M. (2015). AACE/ACE comprehensive diabetes management algorithm 2015. *Endocr. Pract.* 21, 438–447. doi: 10.4158/EP15693.CS
- Higgins, J., and Green, S., (2011). *Cochrane Handbook for Systematic Reviews of Interventions*. The Cochrane Collaboration. Available online at: <http://handbook.cochrane.org>.
- Hong, S., Park, C., Han, K., Chung, C., Ku, B., Jang, H., et al. (2016). Efficacy and safety of teneeligliptin, a novel dipeptidyl peptidase-4 inhibitor, in Korean patients with type 2 diabetes mellitus: a 24-week multicentre, randomized, double-blind, placebo-controlled phase III trial. *Diabetes Obes. Metab.* 18, 528–532. doi: 10.1111/dom.12631

AUTHOR CONTRIBUTIONS

XL, XH, CB, YY, and JW conceived and designed the review; XL, XH, CB, DQ, SC, QM, YY, and JW reviewed the literature; XL, XH, CB, YY, SC, and JW wrote the manuscript.

FUNDING

This work was supported by Grants from the National Natural Science Foundation of China (No. 81774013), Science and Technology Program of Luzhou, China (Grant No. 2015LZCYD-S09(5/8), 2015-S-43, 2016LZXNYD-T03).

ACKNOWLEDGMENTS

The authors thank Yueshan Sun for assistance with literature search, Jing Yang for helping to modify format of the manuscript and Ling Fan and Xingxing Xie for providing meaningful advices.

- Inzucchi, S. E., Bergenstal, R. M., Buse, J. B., Diamant, M., Ferrannini, E., Nauck, M., et al. (2012). Management of hyperglycemia in type 2 diabetes: a patient-centered approach: position statement of the american diabetes association (ADA) and the european association for the study of diabetes (EASD). *Diabetes Care* 35, 1364–1379. doi: 10.2337/dc12-0413
- Kadowaki, T., Inagaki, N., Kondo, K., Nishimura, K., Kaneko, G., Maruyama, N., et al. (2017a). Efficacy and safety of teneeligliptin added onto canagliflozin monotherapy in Japanese patients with type 2 diabetes mellitus: a multicentre, randomised, double-blind, placebo-controlled, parallel-group comparative study. *Diabetes Obes. Metab.* 20, 453–457. doi: 10.1111/dom.13079
- Kadowaki, T., and Kondo, K. (2013a). Efficacy and safety of teneeligliptin added to glimepiride in Japanese patients with type 2 diabetes mellitus: a randomized, double-blind, placebo-controlled study with an open-label, long-term extension. *Diabetes Obes. Metab.* 16, 418–425. doi: 10.1111/dom.12235
- Kadowaki, T., and Kondo, K. (2013b). Efficacy and safety of teneeligliptin in combination with pioglitazone in Japanese patients with type 2 diabetes mellitus. *J. Diabetes Investig.* 4, 576–584. doi: 10.1111/jdi.12092
- Kadowaki, T., and Kondo, K. (2013c). Efficacy, safety and dose-response relationship of teneeligliptin, a dipeptidyl peptidase-4 inhibitor, in Japanese patients with type 2 diabetes mellitus. *Diabetes Obes. Metab.* 15, 810–818. doi: 10.1111/dom.12092
- Kadowaki, T., Kondo, K., Sasaki, N., Miyayama, K., Yokota, S., Terata, R., et al. (2017b). Efficacy and safety of teneeligliptin add-on to insulin monotherapy in Japanese patients with type 2 diabetes mellitus: a 16-week, randomized, double-blind, placebo-controlled trial with an open-label period. *Expert Opin. Pharmacother.* 18, 1291–1300. doi: 10.1080/14656566.2017.1359259
- Kim, M., Rhee, E., Han, K., Woo, A., Lee, M., Ku, B., et al. (2015). Efficacy and safety of teneeligliptin, a dipeptidyl peptidase-4 inhibitor, combined with metformin in Korean patients with type 2 diabetes mellitus: a 16-week, randomized, double-blind, placebo-controlled phase III trial. *Diabetes Obes. Metab.* 17, 309–312. doi: 10.1111/dom.12424
- Kishimoto, M. (2013). Teneeligliptin: a DPP-4 inhibitor for the treatment of type 2 diabetes. *Diabetes Metab. Syndr. Obes.* 6, 187–195. doi: 10.2147/DMSO.S35682
- Koren, S., and Rapoport, M. (2014). Aiming for better glucose control: is HbA1c of 7% the “holy grail” the clinical approach or seven clinical questions about 7. *Diabetes Metab. Res. Rev.* 30, 346–349. doi: 10.1002/dmrr.2532
- Kutogh, E., Hirate, M., and Ikeno, Y. (2014). Teneeligliptin as an initial therapy for newly diagnosed, drug naive subjects with type 2 diabetes. *J. Clin. Med. Res.* 6, 287–294. doi: 10.14740/jocmr1841e
- Liao, E. (2011). Management of type 2 diabetes: new and future developments in treatment. *Lancet* 125, S2–S3. doi: 10.1016/j.amjmed.2012.05.008

- Liberati, A., Altman, D., Tetzlaff, J., Mulrow, C., Gøtzsche, P., Ioannidis, J., et al. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J. Clin. Epidemiol.* 62, e1–e34. doi: 10.1016/j.jclinepi.2009.06.006
- Mitsubishi Tanabe Pharma Corporation. (2014). *Monotherapy Study of MP-513 in Patients With Type 2 Diabetes*. (NCT00998881). Available online at: <https://www.clinicaltrials.gov/>
- Morishita, R., and Nakagami, H. (2015). Teleniglipitin: expectations for its pleiotropic action. *Expert Opin. Pharmacol.* 16, 417–426. doi: 10.1517/14656566.2015.1000301
- Nabeno, M., Akahoshi, F., Kishida, H., Miyaguchi, I., Tanaka, Y., Ishii, S., et al. (2013). A comparative study of the binding modes of recently launched dipeptidyl peptidase IV inhibitors in the active site. *Biochem. Biophys. Res. Commun.* 434, 191–196. doi: 10.1016/j.bbrc.2013.03.010
- Patel, D., Sharma, R., Patel, H., and Barkate, H. (2016). Teleniglipitin: a review on cardio-renal safety. *IJBCP* 5, 229–234. doi: 10.18203/2319-2003.ijbcp20160738
- Paul, W., and Jordi, M. (2018). Gene-lifestyle interplay in type 2 diabetes. *ScienceDirect* 50, 35–40. doi: 10.1016/j.gde.2018.02.001
- Philip, M., and Steve, C. (2016). Omarigliptin for the treatment of Type 2 diabetes mellitus. *Expert Opin. Pharmacother.* 14, 1947–1952. doi: 10.1080/14656566.2016.1218472
- Scott, L. (2015). Teleniglipitin: a review in type 2 diabetes. *Clin. Drug Investig.* 35, 765–772. doi: 10.1007/s40261-015-0348-9
- Secrest, M., Udel, J., and Fillion, K. (2017). The cardiovascular safety trials of DPP-4 inhibitors, GLP-1 agonists, and SGLT2 inhibitors. *Trends Cardiovasc. Med.* 27, 194–202. doi: 10.1016/j.tcm.2017.01.009
- Sharma, S., Panneseelam, A., Singh, K., Parmar, G., Gadge, P., Swami, O., et al. (2016). Teleniglipitin in management of type 2 diabetes mellitus. *Diabetes Metab. Syndr. Obes.* 9, 251–260. doi: 10.2147/DMSO.S106133
- Tanaka, S., Suzuki, K., Aoki, C., Niitani, M., Kato, K., Tomotsune, T., et al. (2014). Add-on treatment with teleniglipitin ameliorates glucose fluctuations and improves glycemic control index in Japanese patients with type 2 diabetes on insulin therapy. *Diabetes Technol Ther.* 16, 840–845. doi: 10.1089/dia.2014.0095
- Wallace, T., Levy, J., and Matthews, D. (2004). Use and abuse of HOMA modeling. *Diabetes Care* 27, 1487–1495. doi: 10.2337/diacare.27.6.1487
- Xueying, T., and Jingbo, H. (2016). Evogliptin: a new dipeptidyl peptidase inhibitor for the treatment of type 2 diabetes. *Expert Opin. Pharmacother.* 17, 1285–1293. doi: 10.1080/14656566.2016.1183645
- Yoshida, T., Akahoshi, F., Sakashita, H., Kitajima, H., and Nakamura, M. (2012). Discovery and preclinical profile of teleniglipitin(3-[(2S,4S)-4-[4-(3-methyl-1-phenyl-1H-pyrazol-5-yl)piperazin-1-yl]pyrrolidin-2-ylcarbonyl]thiazolidine): a highly potent, selective, long-lasting and orally active dipeptidyl peptidase IV inhibitor for the treatment of type 2 diabetes. *Bioorg. Med. Chem.* 20, 5705–5719. doi: 10.1016/j.bmc.2012.08.012

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Li, Huang, Bai, Qin, Cao, Mei, Ye and Wu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.