



Ibogaine Administration Modifies GDNF and BDNF Expression in Brain Regions Involved in Mesocorticolimbic and Nigral Dopaminergic Circuits

OPEN ACCESS

Edited by:

Andrew Robert Gallimore,
Okinawa Institute of Science
and Technology Graduate University,
Japan

Reviewed by:

Wladyslaw Lason,
Institute of Pharmacology (PAN),
Poland
Sebastien Caricella,
INSERM U1216 Grenoble Institut des
Neurosciences (GIN), France
Walter Sandtner,
Medical University of Vienna, Austria
Deborah Carmen Mash,
University of Miami, United States

*Correspondence:

Cecilia Scorza
cscorza@ibce.edu.uy
Patricia Cassina
pcassina@fmed.edu.uy
Ignacio Carrera
icarrera@fq.edu.uy

† These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Neuropharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 28 November 2017

Accepted: 14 February 2019

Published: 05 March 2019

Citation:

Marton S, González B,
Rodríguez-Bottero S, Miquel E,
Martínez-Palma L, Pazos M, Prieto JP,
Rodríguez P, Sames D, Seoane G,
Scorza C, Cassina P and Carrera I
(2019) Ibogaine Administration
Modifies GDNF and BDNF Expression
in Brain Regions Involved
in Mesocorticolimbic and Nigral
Dopaminergic Circuits.
Front. Pharmacol. 10:193.
doi: 10.3389/fphar.2019.00193

Soledad Marton^{1†}, Bruno González^{2†}, Sebastián Rodríguez-Bottero¹, Ernesto Miquel¹, Laura Martínez-Palma¹, Mariana Pazos², José Pedro Prieto³, Paola Rodríguez², Dalibor Sames⁴, Gustavo Seoane², Cecilia Scorza^{3*}, Patricia Cassina^{1*} and Ignacio Carrera^{2*}

¹ Departamento de Histología y Embriología, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay,

² Laboratorio de Síntesis Orgánica, Departamento de Química Orgánica, Facultad de Química, Universidad de la República, Montevideo, Uruguay, ³ Departamento de Neurofarmacología Experimental, Instituto de Investigaciones Biológicas Clemente Estable, Montevideo, Uruguay, ⁴ Department of Chemistry, Columbia University, New York, NY, United States

Ibogaine is an atypical psychedelic alkaloid, which has been subject of research due to its reported ability to attenuate drug-seeking behavior. Recent work has suggested that ibogaine effects on alcohol self-administration in rats are related to the release of Glial cell Derived Neurotrophic Factor (GDNF) in the Ventral Tegmental Area (VTA), a mesencephalic region which hosts the soma of dopaminergic neurons. Although previous reports have shown ibogaine's ability to induce GDNF expression in rat midbrain, there are no studies addressing its effect on the expression of GDNF and other neurotrophic factors (NFs) such as Brain Derived Neurotrophic Factor (BDNF) or Nerve Growth Factor (NGF) in distinct brain regions containing dopaminergic neurons. In this work, we examined the effect of ibogaine acute administration on the expression of these NFs in the VTA, Prefrontal Cortex (PFC), Nucleus Accumbens (NAcc) and the Substantia Nigra (SN). Rats were i.p. treated with ibogaine 20 mg/kg (I₂₀), 40 mg/kg (I₄₀) or vehicle, and NFs expression was analyzed after 3 and 24 h. At 24 h an increase of the expression of the NFs transcripts was observed in a site and dose dependent manner. Only for I₄₀, GDNF was selectively upregulated in the VTA and SN. Both doses elicited a large increase in the expression of BDNF transcripts in the NAcc, SN and PFC, while in the VTA a significant effect was found only for I₄₀. Finally, NGF mRNA was upregulated in all regions after I₄₀, while I₂₀ showed a selective upregulation in PFC and VTA. Regarding protein levels, an increase of GDNF was observed in the VTA only for I₄₀ but no significant increase for BDNF was found in all the studied areas. Interestingly, an increase of proBDNF was detected in the NAcc for both doses. These results show for the first time a selective increase of GDNF specifically in the VTA for I₄₀ but not for I₂₀ after 24 h of administration, which agrees with the effective dose found in previous self-administration studies in rodents. Further research is needed to understand the contribution of these changes to ibogaine's ability to attenuate drug-seeking behavior.

Keywords: ibogaine, neurotrophic factors, GDNF, BDNF, NGF

INTRODUCTION

Ibogaine is the main indole alkaloid isolated from the root bark of the African shrub *Tabernanthe iboga* (Lavaud and Massiot, 2017). Traditionally used in African religious ceremonies as a psychedelic, ibogaine became a subject of interest to the scientific community due to its reported ability to reduce craving and self-administration of several drugs of abuse in humans (Brown, 2013). These effects found mainly in uncontrolled clinical trials and observational studies, have been reported to be long-lasting enduring weeks to months after a single administration of large doses of ibogaine (Schenberg et al., 2014; Brown and Alper, 2017; Noller et al., 2017; Corkery, 2018; Malcolm et al., 2018; Mash et al., 2018). In animal models for drug dependence, ibogaine also reduces the self-administration of morphine and heroin (Glick et al., 1991, 1994; Dworkin et al., 1995), cocaine (Cappendijk and Dzoljic, 1993; Glick et al., 1994), and alcohol (He et al., 2005), with long-lasting effects that persists beyond pharmacokinetic elimination of the drug (Alper, 2001). In addition, ibogaine administration to animals also reduces naloxone or naltrexone precipitated-withdrawal signs (Dzoljic et al., 1988; Glick et al., 1992; Leal et al., 2003).


Although a vast amount of research has been done regarding the pharmacology of ibogaine, the mechanism of action of its ability to attenuate drug-seeking behavior remains unresolved (Alper, 2001; Maciulaitis et al., 2008; Brown, 2013). Ibogaine binds to numerous central nervous system (CNS) targets at the micromolar range such as: nicotinic acetylcholine receptors (nAChR $\alpha 3\beta 4$ and $\alpha 2\beta 4$) (Fryer and Lukas, 1999; Arias et al., 2010, 2015), N-methyl-D-aspartate (NMDA) (Mash et al., 1995b), kappa and mu opioid (Antonio et al., 2013; Maillet et al., 2015), 5HT_{2A} and 5HT₃ receptors (Glick et al., 2000) and the dopamine and serotonin transporters (Mash et al., 1995a; Glick et al., 2001; Asjad et al., 2017). However, these ibogaine-receptor interactions do not seem to account for the long-lasting effects of ibogaine found in rodents which are described to last for 48 to 72 h after ibogaine administration (Glick et al., 1991, 1994; Cappendijk and Dzoljic, 1993). In rodents, ibogaine has a short half-life of 1–2 h raising the hypothesis that its longer-lived active metabolite, noribogaine, could be responsible for the enduring effects elicited by ibogaine. Both, the parent drug and its metabolite have differences in their binding profiles and affinities to the abovementioned CNS receptors (Staley et al., 1996). However, no appreciable amounts of noribogaine have been found in rodents' brain tissue 19 h after ibogaine intraperitoneal (i.p.) administration (Pearl et al., 1997), and only approximately 5% of the noribogaine C_{max} was detected in serum 24 h after the same treatment (Baumann et al., 2001b).

A few years ago, a novel hypothesis linking ibogaine's attenuation of alcohol self-administration in rodents to its ability to modulate the expression of Glial Cell Derived Neurotrophic Factor (GDNF) in the brain was proposed. It was shown that a single ibogaine i.p. administration (40 mg/kg) increased the expression of GDNF in the midbrain of rats and mice for up to 24 h (He et al., 2005). In addition, microinjection of ibogaine into the Ventral Tegmental Area (VTA), produced a long-lasting reduction of ethanol self-administration, a response that was

attenuated by the intra-VTA delivery of anti-GDNF neutralizing antibodies. These results suggested that ibogaine mediates its effects against ethanol consumption by increasing GDNF content in the VTA (He et al., 2005). Accordingly, another study from the same research group showed that the intra-VTA infusion of noribogaine induced a long-lasting decrease in ethanol self-administration (Carnicella et al., 2010). Further, ibogaine-derived synthetic derivatives were recently shown to induce the release of GDNF *in vitro*, in established cell line systems (Gassaway et al., 2016). These observations formed the basis for a new rationale to explain the long-lasting effects of ibogaine; i.e., the induction of GDNF by ibogaine/noribogaine may activate an autocrine loop, leading a long-term synthesis and release of GDNF (that persists beyond elimination of both substances). This mechanism may reverse the biochemical adaptations to chronic exposure to drugs of abuse in the reward system (He and Ron, 2006).

Neurotrophic Factors (NFs), such as GDNF and BDNF (Brain Derived Neurotrophic Factor) are small proteins that promote the growth, differentiation, synaptogenesis, and survival of neurons. Their expression in the nervous tissue is relatively high during the development of the CNS, where substantial growth, differentiation and remodeling of the nervous system occur (Barde, 1990; Lu and Figurov, 1997). More recently, it has been discovered that NFs play important roles in the adult brain where they modulate maintenance, protection, repair and plasticity of the nervous tissue (Reichardt, 2006; Schmidt and Duman, 2007). Furthermore, accumulating evidence has suggested that GDNF and BDNF mediate neuronal remodeling processes that occur during the development of substance use disorders (SUDs) (Bolaños and Nestler, 2004; McGough et al., 2004; Angelucci et al., 2007; Jeanblanc et al., 2009; Bie et al., 2012). Particularly, the role of GDNF and BDNF in the neuroadaptations in the mesocorticolimbic dopamine system (Prefrontal Cortex, PFC- VTA-Nucleus Accumbens, NAcc pathway) induced by repeated exposure to drugs of abuse has been extensively studied, including the impact of manipulating NFs levels on drug-seeking behavior in animal models (Russo et al., 2009; Ghitza et al., 2010; Koskela et al., 2017). It has been shown that the administration of BDNF or GDNF can either promote or inhibit drug-taking behaviors depending mainly on the brain site of administration, along with other several factors such as the drug type, the addiction phase (initiation, maintenance, abstinence or relapse), the time interval between site-specific NFs injections and the related behavioral assessments (Ghitza et al., 2010). For example, BDNF infusion into the NAcc increases cocaine-seeking behavior (Graham et al., 2007), while BDNF infusion into the medial pre-frontal cortex (mPFC) suppresses it (Berglind et al., 2007). Additionally, infusion of BDNF into the dorsolateral striatum decreases ethanol self-administration in rats (Jeanblanc et al., 2009).

Given the importance and the site-specificity of the elicited responses, we decided to analyze the effect of a single administration of ibogaine on the expression of GDNF and BDNF (mRNA transcripts and protein content) at two time points in those brain areas which define the mesocorticolimbic dopamine system such as VTA, PFC and NAcc (**Figure 1**). As the Substantia Nigra (SN) is a major nucleus of dopaminergic



neurons important in the basal ganglia functioning, the expression of these NFs in this region was also studied. In order to examine the impact of ibogaine administration on the expression of other relevant NFs (which impact on drug-seeking behaviors has been much less studied) the Nerve Growth Factor (NGF) transcript content was also analyzed in the abovementioned brain areas. Selected time points were chosen considering previous pharmacokinetics reports in rats using i.p. administration (Pearl et al., 1997; Zubaran et al., 1999; Baumann et al., 2001a,b). In this manner, we chose to study NFs expression/content in the selected brain areas at 3 h, where ibogaine and noribogaine are present in relevant concentrations (Baumann et al., 2001b), and at 24 h where ibogaine is no longer detected and no significant amounts of noribogaine would be present in the brain (Pearl et al., 1997). In this manner, is expected that the observed effects found at 24 h, would be due to long lasting mechanisms elicited by the drug which remain after it has been cleared from the brain, but not from the acute effects of ibogaine/noribogaine. Finally, a behavioral study recording the locomotor activity of the control and drug-treated animals was performed using an open field test for each time point.

MATERIALS AND METHODS

Ibogaine HCl

The ibogaine used in this study was chemically synthesized using voacangine as starting material, which was extracted from the root bark of *Voacanga africana* (purchased from CAPE LABS) using a modification of a previously described procedure (Jenks, 2002). Briefly, 100g of grounded root bark of *V. africana* was extracted with a 1% aqueous solution of HCl (6 × 500 mL). The combined aqueous extracts were basified by adding concentrated NH₄OH until pH 10–11. A brown precipitate was separated by centrifugation and dried at 60°C for 24 h. This solid was taken in acetone and filtered to discard root impurities. The solvent was evaporated *in vacuo* to afford a total alkaloid extract of 3.5–4.0 g. Column chromatography (SiO₂, Hex:EtOAc:NH₄OH, 90:10:0.01) allowed to obtain 1g of pure voacangine which was

analyzed by ¹H and ¹³C NMR (See **Supplementary Material**). Voacangine was decarboxylated as follows. To a solution of voacangine in EtOH (0.45 M) in a double necked round bottomed flask, KOH in pellets (5 equivalents) was added. The solution was heated to reflux until consumption of the starting material was evident by thin layer chromatography (TLC) analysis. EtOH was removed under reduced pressure, and the residue was dissolved at 0°C in a round bottomed flask using a 6% (v/v) aqueous solution of HCl (enough quantity to adjust pH to 1). The system was then heated to reflux for 5 min. Once the starting material consumption was evident by TLC analysis, the solution was carefully basified using 50% NaOH (pH 10–11). Precipitation of ibogaine as a white solid was observed. Ethyl acetate was added, and the resultant biphasic system was transferred into a separation funnel. The aqueous phase was extracted three times with EtOAc. The combined organic layers were dried under Na₂SO₄, and the solvent was removed *in vacuo*. Purification was carried out using column chromatography purification (SiO₂, hexanes: ethyl acetate 8:2 + 0.5% ammonium hydroxide). Ibogaine free base was obtained with an 86% and was analyzed by ¹H and ¹³C NMR (see **Supplementary Material**). Crystallization from EtOH afforded a crystalline solid which was converted to the corresponding hydrochloride by treatment with diethyl ether saturated with HCl(g). Purity of ibogaine-HCl was determined by GC-MS analysis as 98.3% (see **Supplementary Material**). Dissolution of ibogaine-HCl to prepare the samples for i.p. injection was carried out using warm saline that was previously degassed by nitrogen bubbling.

Experimental Animals

Thirty-six male Wistar adult rats (270–300 g) were used in this study and assigned to one of the following groups: Vehicle group at 3 and 24 h (*n* = 6 per each group); Ibogaine 20- (I₂₀) treated group at 3 and 24 h (*n* = 6 per each group) and Ibogaine 40- (I₄₀) treated group at 3 and 24 h (*n* = 6 per each group). Animals were housed four to five per cage and maintained on a 12-h light/dark cycle (lights on at 07.00 h) with food and water freely available before and after i.p. injection of vehicle or

ibogaine until behavioral testing and sacrifice. All experimental procedures were conducted in agreement with the National Animal Care Law (#18611) and with the “Guide to the care and use of laboratory animals” (8th edition, National Academy Press, Washington, DC, 2010). Furthermore, the local Institutional Animal Care Committee (IIBCE) approved the experimental procedures (Protocol Number 007/05/2014). Adequate measures were taken to minimize pain, discomfort or stress of the animals, and all efforts were made to use the minimal number of animals necessary to obtain reliable scientific data.

Behavioral Analysis

Animals were brought to the experimental room in their home cages, identified and weighed prior to the behavioral test. An open field (OF) apparatus consisting of a square area (45 cm wide \times 45 cm long \times 40 cm high) with transparent plastic walls indirectly illuminated (35 luxes) to avoid reflection and shadows were employed. The OF was placed in a quiet experimental room with controlled temperature ($22 \pm 2^\circ\text{C}$). As rats were not habituated to the OF before drug or vehicle administration, novelty-induced motor activity was automatically recorded by a camera connected to a computer equipped with the Ethovision XT 12.0 software (Noldus, Netherlands) located above the OF. Using this video tracking software, we specifically measured the total distance traveled in meters (m) during 30 min, starting 3 and 24 h after ibogaine or vehicle administration. Animals were randomly assigned to different experimental groups and were used only once. Taking into account that immediately after i.p. administration ibogaine can produce a dose-dependent unusual motor profile and some prototypical serotonergic syndrome-related behaviors (e.g., tremor, flat body posture, forepaw treading) (Gonzalez et al., 2018), these specific behaviors were assessed by a trained investigator every 5 min (for a total of 30 min) starting 3 and 24 h after ibogaine administration. During all experiments, the OF was cleaned with 30% alcohol before placing the following rat. All experiments were done between 9 AM and 3 PM.

Ex vivo Studies

Brain Dissection

Three or twenty-four hours after I_{20} , I_{40} or vehicle (i.p.) injection, animals were sacrificed by decapitation and the brains were carefully removed and chilled in ice cold saline. According to Paxinos and Watson (2005), the whole NAcc (shell and core), PFC (including mPFC), Substantia Nigra (SN, pars compacta-SNpc and pars reticulata-SNpr) and VTA were dissected out on ice and the tissue obtained was immediately frozen and rapidly stored at -80°C until the processing day (Scorza et al., 1997; Meikle et al., 2013). Representative examples of coronal section at the level of each dissected brain area are shown in the **Supplementary Material**.

Semiquantitative qPCR

For RT-PCR analysis total RNA was extracted from the different brain regions using Trizol reagent (Thermo Fisher Scientific) followed by chloroform extraction and isopropanol precipitation. Possible DNA contaminations were eliminated with DNase

treatment using DNase free Kit (Thermo Fisher Scientific). RNA quality was evaluated by agarose gel electrophoresis followed by ethidium bromide staining and quantified using a NanoDrop 1000 Spectrophotometer (Thermo Fisher Scientific). 500 ng of this total RNA was reverse-transcribed using 200 U M-MLV-reverse transcriptase (Thermo Fisher Scientific) following manufacturer instructions. 25 ng of the resulting cDNA was diluted in Biotools Quantimix Easy master mix (Biotools) in 10 μl volume. All reactions were performed in triplicates in strip tubes (Axygen[®] Brand Products), using specific forward and reverse primers. The sequences of the quantitative PCR primers (IDT, Integrated DNA Technologies) used are as follows: for GAPDH F: 5'-CAC TGA GCA TCT CCC TCA CAA-3' and R: 5'-TGG TAT TCG AGA GAA GGG AGG-3', for BDNF F: 5'-GAG GGG TAG ATT TCT GTT TGT T-3' and R: 5'-TTG CCT TAA TTT TTA TTC GTT T-3', for GDNF F: 5'-AAA TCG GGG GTG CGT CTT AAC T-3' and R: 5'-AAC ATG GGC CTA CCT TGT C-3', for NGF F: 5'-AAG TTA TCC CAG CCA AAC TA-3' and R: 5'-ATG TCA GTG TTG GGA GTA GG-3'. According to the sample, we used cycles 15–23 (the threshold cycle, Ct), in order to calculate the relative amounts of our gene of interest. PCR amplification was done over 40 cycles using a Rotor-Gene 6000 System (Corbett Life Science) and data were analyzed using Rotor Gene 6000 software (Corbett Life Science). Quantification was performed with $\Delta\Delta\text{Ct}$ method using rats treated with vehicle as a negative control, and GAPDH mRNA as reference.

Western Blot Analysis

The selected brain regions were sonicated in a lysis buffer containing 50 mM NaCl, 50 mM HEPES, 2 mM sodium orthovanadate, 1% Triton X-100, and SigmaFAST Protease inhibitor cocktail (Sigma-Aldrich). After quantification and denaturation, the samples were loaded and separated by 12% SDS-PAGE gels and then transferred into a nitrocellulose membrane. The membranes were incubated for 1 h in blocking solution (BS: 5% Bovine serum albumin, 1% Tween 20 in PBS), and incubated overnight at 4°C with primary antibodies to GDNF (1:500 in BS; Abcam ab119473), BDNF (1:400 in BS; Promega G1641), or proBDNF (1:500 in BS; Invitrogen PA1-18360), together with anti-alpha-tubulin (1:3000 in BS; Abcam ab184613) as loading control. Afterward, the membranes were washed and incubated for 1 h at room temperature with IRDye 680RD/IRDye 800CW-Conjugated Goat Anti-Mouse IgG/Goat Anti-Rabbit IgG/Donkey Anti-Chicken IgG secondary antibodies (1:15000 in PBS each, LI-COR Biosciences #926-68070, #926-32210, #926-68071, #926-32211, and #925-32218). The Odyssey system (LI-COR Biosciences) was used to detect the bands. Quantification of band intensity was performed using Image Studio software version 5.2.5.

Data Analysis

GraphPad Prism software 5 was used to design figure graphs and data analysis. Data are presented as mean \pm SEM values. Six animals per group were assessed for behavioral and PCR studies. In some cases, some data was excluded from the analysis due to insufficient sample or high deviation from the mean of the group, rendering a lower n , but never smaller than 4. The total

sample size (N) is given in figure legends and the sample for each treatment (n) can be observed in the scatter plot graphs in each figure. For western blot analysis, samples from 4 animals per group were assessed. Data from qPCR and western blot were analyzed and compared by one-way ANOVA followed by *post hoc* Tukey's Multiple Comparison Test. In all cases, statistical significance was set at $P < 0.05$. General P and F values from ANOVA, and p values from Tukey's multiple comparison test are provided in figure legends for each data set when significance is reached. Also, eta squared values (η^2) accounting for effect size are provided. Data from motor activity were analyzed by two-way (treatment, time, and interaction between factors) ANOVA for repeated measures followed by Newman-Keuls multiple comparison *post hoc* test and Unpaired- t -test.


RESULTS

In a previous study, we reported a very high impact of the I_{40} treatment on novelty-induced locomotion after 2 h of ibogaine administration and the concomitant transient induction of some of the behavioral signs related to the serotonergic syndrome (Gonzalez et al., 2018). Thus, we decided to analyze the behavioral effect of ibogaine treatment in the time points used in the present study (3 and 24 h). The behavioral response induced by ibogaine administration is shown in **Figure 2**. Compared to the control group, novelty-induced locomotion was not altered by I_{20} at any evaluated time (data not shown). Whereas I_{40} was not effective to induce any behavioral alterations 3 h after i.p. administration, it elicited a significantly reduction of the animal locomotion 24 h after injection (**Figures 2A,B** respectively). No abnormal behaviors were present for both time points and animals were qualitatively indistinguishable from the vehicle group animals (data not shown). Immediately after each behavioral test, animals were sacrificed to pursue brain dissection for the qPCR and Western Blot studies.

qPCR Quantification of NFs mRNA


qPCR results for the GDNF (**Figure 3**) showed that ibogaine acute administration differentially regulated GDNF mRNA expression levels in the selected brain regions in a dose and time-dependent manner. At 3 h, no changes in the GDNF mRNA expression was found for both doses of ibogaine in all the studied areas. In contrast, after 24 h of treatment, changes in the expression of GDNF were found in a dose and site-specific manner. While the I_{20} dose did not affect the GDNF expression in any of the studied areas, the I_{40} dose selectively increased GDNF mRNA content in the midbrain regions: VTA (12-fold increase compared to the control group) and SN (6-fold increase vs. the control group) with no appreciable effects in the PFC and NAcc.

For BDNF, ibogaine treatment produced an appreciable downregulation of its expression in the PFC at 3 h after injection (1.7 and 2-fold decrease for I_{20} and I_{40} , respectively, compared to control, control = 1.000 ± 0.099 , $I_{20} = 0.596 \pm 0.045$, $I_{40} = 0.492 \pm 0.094$), while no response was seen for the other brain areas at this time point (**Figure 4**). At 24 h, ibogaine administration upregulated the mRNA expression of BDNF in all



the brain regions studied in a dose-dependent manner (**Figure 4**). A large effect was found in the NAcc for both doses of ibogaine (220-fold increase compared to the control for I_{20} , and 340-fold increase for I_{40}). The I_{20} dose increased BDNF expression in PFC (55-fold increase compared to the control) but not in the VTA or SN. On the other hand, in addition to the NAcc, the I_{40} dose also upregulated BDNF expression in PFC (107-fold increase compared to the control), VTA (43-fold increase compared to the control) and SN (21-fold increase compared to the control).

For NGF (**Figure 5**), no difference in the content of mRNA was found 3 h after ibogaine treatments. At 24 h, an upregulation



of NGF mRNA content was found in: PFC (14-fold increase compared to the control), NAcc (15-fold increase compared to the control), VTA (11-fold increase compared to the control), and SN (4-fold increase compared to the control). For the I_{20} dose a significant effect was only found in the PFC (7-fold increase compared to the control) and VTA (5-fold increase compared to the control). However, the levels of increase in the NGF mRNA were not as high as those for BDNF.


GDNF, BDNF and proBDNF Protein Content by Western Blot

Considering the changes found for the expression of NFs after 24 h of ibogaine administration, we decided to analyze the content of mature proteins BDNF and GDNF for all the studied brain regions, because of their previously mentioned well studied involvement in the addictive behavior. Precursor of BDNF, proBDNF was also considered since it is well described that it shows opposite effects to the mature protein because of a higher affinity to the p75 receptor (Woo et al., 2005; Xu et al., 2011; Sun et al., 2012). For GDNF, a single dose of ibogaine affected mature protein content in a region- and dose-dependent manner (Figure 6). While no changes in GDNF content were observed for I_{20} in any of the studied regions, GDNF content was increased in VTA for the I_{40} dose (2-fold increase compared to the control group). No effect was observed in the GDNF content at the NAcc, SN, and PFC in comparison to the control group. For BDNF no significant change in the mature protein content was detected for

all the studied regions for both doses of ibogaine. Nevertheless, in the case of proBDNF we found a selective increase in the protein content for I_{20} and I_{40} in the NAcc (2.7 and 2.8-fold increase for I_{20} and I_{40} doses, respectively, compared to control), while no significant change was detected in the other brain areas.

DISCUSSION

In the present study, we have demonstrated that ibogaine administration simultaneously alters the transcripts levels of GDNF and BDNF (which have been extensively related to drug-seeking behaviors) in a dose- and time-dependent manner. Additionally, NGF expression was also modified, showing potential effects of ibogaine administration on the expression of other relevant NFs. Regarding the protein content, we showed that after 24 h of treatment, I_{40} selectively increased mature GDNF in the VTA, while proBDNF content was increased selectively in NAcc by both doses. Since as mentioned before, ibogaine is rapidly metabolized to produce noribogaine, further experiments are needed to elucidate if the metabolite and/or the parent drug produced these effects. Considering that dopamine neurotransmission, specifically in the mesocorticolimbic pathway, is related to rewarding/reinforcing and motivational actions of most drugs of abuse (Di Chiara and Imperato, 1988; Koob and Bloom, 1988; Kalivas and Volkow, 2005) our findings contribute to shed light on a mechanism




underlying the ability of ibogaine administration to attenuate drug-seeking behavior.

Regarding the motor function, a decrease in the novelty-related motor activity was observed 24 h after I_{40} (while 3 h after the same treatment, animals displayed a similar activity than the control). There is no evidence at this point to establish a potential connection between this intriguing behavior and the observed changes in NFs expression. In this regard, considering the changes in the expression of NFs at 24 h in the SN, it is plausible that a neurochemical imbalance in the basal ganglia output may underlie the changes in the motor activity (Day et al., 2008; Calabresi et al., 2014). Moreover, we cannot rule out that this acute motor impairment is related to this neurochemical effect eliciting a decrease in the animal overall motivation. Behavioral studies using valid and reliable experimental paradigms for studying the effect of ibogaine on reward-related behaviors should be done to understand these observations. On the other hand, we cannot discard the participation of other factors which may be altered in ibogaine-treated animals at this time point.

At 3 h after I_{20} and I_{40} treatments, no alteration of the GDNF transcript content was found in all the studied brain areas. While in a previous report by He et al. (2005), a significant GDNF upregulation was found 3 h after I_{40} treatment in the midbrain of rats, our results show that this increase doesn't occur in the NAcc and in the specific midbrain areas studied (VTA, NAcc, and SN). On the other hand, after 24 h, we found that the I_{40} dose increased GDNF expression and mature protein content specifically in the

rat VTA, which was also found in the whole midbrain at this time point in the mentioned previous report. In this manner our study identifies the VTA as the key brain region of the mesocorticolimbic system where GDNF is upregulated after 24 h of ibogaine administration. This finding is important since the ability of ibogaine to attenuate ethanol self-administration had previously been proposed to be mediated, at least in part, by the increase in GDNF content in the VTA. (He et al., 2005; He and Ron, 2006) Furthermore, we show that I_{20} administration does not increase GDNF expression in any of the studied brain areas, which is in accordance with the observation that this dose was not effective in reducing drug self-administration in the majority of previous studies in rodents (Glick et al., 1991, 1994; Cappendijk and Dzoljic, 1993; Dworkin et al., 1995). In addition, our results are in line with the reports indicating that GDNF infusions into the VTA has been effective in reducing drug self-administration or conditioned place preference for cocaine and alcohol (Messer et al., 2000; He and Ron, 2006; Carnicella et al., 2008, 2009), and with evidence that shows that GDNF mediates negative regulatory effects on chronic morphine-induced neuroadaptations in VTA of rodents (Li et al., 2014; Koskela et al., 2017). Additionally, the selective increase found in this study for GDNF in the VTA by I_{40} , could account, at least in part, to the anti-addictive properties of ibogaine considering that upregulation of the GDNF pathway has been proposed a potential strategy to treating SUDs (Carnicella and Ron, 2009). Lastly, I_{40} administration increases GDNF expression in the



SN, which was not accompanied with a significant increase of the GDNF protein content at this time point. Given the relevant role of the nigro-striatal pathway in the neuropathology of neurodegenerative disorders like Parkinson's Disease (PD) (Dauer and Przedborski, 2003), it would be interesting to study if ibogaine is able to attenuate the cell loss in the SN and the biochemical changes at the striatum throughout the NFs expression using an experimental model of PD.

With regard to BDNF, a selective downregulation of its expression in the PFC for both doses of ibogaine was found after 3 h of administration, while no changes in other areas were observed. Ibogaine and noribogaine administration in rats stimulate the secretion of corticosterone, being ibogaine a more potent releaser (Baumann et al., 2001b). Since corticosterone decreases BDNF expression in the frontal cortex (Dwivedi et al., 2006; Huang et al., 2011), ibogaine induced corticosterone secretion during the first hours after treatment (where ibogaine concentrations in blood are high), could be the reason behind this result. In contrast, at 24 h, an impressive upregulation of BDNF expression was found, which was much more pronounced compared to the effect on GDNF and NGF expression in all the studied brain areas at this time point. Nevertheless, this high effect on BDNF expression was not reflected on an increase in the content of BDNF mature protein (no significant differences were found between both doses and the control group at this time point, although trending toward increased BDNF protein levels in NAcc and VTA for both doses) (Figure 6). Since BDNF is synthesized in a precursor form, we included

proBDNF in our experimental design. A selective increase in the proBDNF content was selectively found for NAcc for both ibogaine doses. It is known that the mature BDNF protein and its precursor proBDNF have opposite effects on neuronal protection, axonal growth, maturation of dendrites and synaptic plasticity, owing to different affinities of each form to the TrkB and p75 receptors (Lu et al., 2005; Teng et al., 2005; Benarroch, 2015; Borodinova and Salozhin, 2016; Li et al., 2017). In this regard, since it is well-documented that an increase in BDNF content in the NAcc increases cocaine-seeking behavior (Graham et al., 2007; Bahi et al., 2008) and vulnerability to substance abuse (Krishnan et al., 2007; Burke and Miczek, 2015), an increase in proBDNF in this brain area could have an opposite impact. In this line of reasoning, the increase in proBDNF content in NAcc generated by ibogaine after 24 h of administration in rats could also be implicated in ibogaine's effect in drug self-administration paradigms. Further experiments are required to address this hypothesis.

Despite implicit assumption that differentially expressed mRNAs are reflected in protein content, numerous previous studies comparing mRNA and protein levels concluded that the correlation is poor (de Sousa Abreu et al., 2009; Maier et al., 2009). While the increase in GDNF mRNA expression was linked to augmented mature protein content, our data showing an impressive increase in BDNF mRNA expression and no changes in mature protein are intriguing. The possibility exists that the time frame of protein synthesis is different for both NFs, however, many other factors should be considered to explain this




FIGURE 6 | Effects of ibogaine administration on GDNF, BDNF, and proBDNF protein levels in specific brain areas. Western blot analysis of GDNF (A,B), BDNF (C,D) and proBDNF (E,F) protein levels in the indicated brain areas after 24 h of vehicle (0), 20, or 40 mg/kg ibogaine administration. A representative image from immunostained membrane of each condition is shown (A,C,E) with the corresponding quantification below (B,D,F). Data represent mean ± SEM of $n = 4$ biological replicates assayed in triplicate. For GDNF/VTA, $N = 12$, $P < 0.05$, $F_{2,9} = 6.86$, $\eta^2 = 0.60$; For proBDNF/NAcc, $N = 12$, $P < 0.05$, $F_{2,9} = 5.87$, $\eta^2 = 0.57$; * $P < 0.05$ between indicated groups.

incongruousness. These include post-transcriptional regulation, for example miRNA-based translation repression or alternative splicing, or translational and post-translational modifications. Indeed, it has been previously described that sortilin, an intracellular chaperon, acts as a regulatory switch for delivery of BDNF to the regulatory secretory pathway or to degradation in the lysosome, modulating in this way the neurotrophic factor availability (Evans et al., 2011). Interestingly, BDNF levels have been shown to be modified in PFC after chronic ethanol exposure (Yang et al., 2017).

How does ibogaine administration produce this long-term upregulation of GDNF and BDNF? It is well established that an increase in serotonin transmission leads to an increase in BDNF expression/signaling both *in vitro* and *in vivo* (Rantamaki et al., 2007; Popova et al., 2017). In addition, serotonin and SSRIs (Selective Serotonin Re-uptake Inhibitors) induce GDNF expression *in vitro* (Hisaoka et al., 2001; Mercier et al., 2004; Tsuchioka et al., 2008; Golan et al., 2011), and recently it has been shown that chronic treatment in mice using SSRIs induce GDNF content in SN and Striatum (Shadfar et al., 2018). It is well-established that ibogaine and noribogaine increase serotonin transmission (Wei et al., 1998; Wells et al., 1999; Baumann et al., 2001b). Both substances are serotonin-reuptake inhibitors (Jacobs et al., 2007; Bulling et al., 2012), and noribogaine is more potent at increasing serotonin levels in the NAcc than ibogaine, which correlates with the ability of both compounds to inhibit SERT *in vitro* (IC₅₀ of 3.85 and 0.18 μM for ibogaine and noribogaine, respectively) (Baumann et al., 2001b). In this manner, a sustained enhancement on serotonin transmission due to ibogaine and its long-lasting metabolite noribogaine could account, at least in part, for the observed effect on BDNF and GDNF expression after 24 h of ibogaine administration.

Finally, in addition to GDNF and BDNF, ibogaine also modulated the expression of other NF as NGF, 24 h after treatment, while no changes were found at 3 h. The effect of NGF administration in specific brain areas on drug-seeking behavior has been much less studied in comparison to GDNF and BDNF, and scarce data is available on the effects of NGF in brain regions related to the dopaminergic mesocorticolimbic circuitry. Nevertheless, NGF (as other neurotrophins) is likely involved in mediating important responses related to chronic intake of drugs of abuse, as illustrated by previous studies that show that NGF content decreases in the hippocampus and hypothalamus of alcohol-treated mice (Aloe et al., 1993) and in the serum of chronic heroin and cocaine users (Angelucci et al., 2007). Also, NGF administration into the central nucleus of the amygdala mimicked the morphine reward sensitization (Bie et al., 2012).

The modifications found in NFs levels induced by ibogaine/noribogaine, may underlie neuroplasticity processes in the discrete brain regions analyzed as has been described by several drugs used in clinical practice including drugs of abuse (Castren and Antila, 2017). Most of these drugs regulate the expression of NFs, reactivating a process defined as induced plasticity (iPlasticity), which allows networks reorganization in the adult brain (Castren and Antila, 2017). This is in accordance with the fact that recently noribogaine has been recently classified as a “psychoplastogen,” since it is capable to promote

neurogenesis in cultured rat cortical neurons (Ly et al., 2018). In this manner, neuroplastic changes generated by the selective increase in NFs expression after ibogaine administration could explain, at least in part, the ability of ibogaine to attenuate drug-seeking behavior in rodents (which could be related to its effects on drug craving and reinstatement in humans).

CONCLUSION AND FUTURE PERSPECTIVES

This study demonstrates for the first time that ibogaine administration simultaneously alters the expression of GDNF, BDNF, and NGF transcripts in rat brain regions related to the dopamine neurotransmission in a dose- and time-dependent manner. Our results add relevant information concerning specific brain areas involved in the increment of GDNF levels (VTA) as a putative mechanism of action underlying the anti-addictive effect of ibogaine. In addition, we showed that only I₄₀ promoted this increase in GDNF content, which is in accordance with previous reports where the I₂₀ treatment was not effective in reducing drug self-administration in rodents (Glick et al., 1991, 1994; Cappendijk and Dzoljic, 1993; Dworkin et al., 1995). Also, we found that both doses of ibogaine produced an increase in the proBDNF content in NAcc after 24 h of treatment, which could be another factor mediating long-lasting effects of ibogaine related to attenuate drug dependence, in addition to the already highlighted increase in GDNF. Future experiments are needed to clarify these important implications in order to elucidate ibogaine's biological mechanism to attenuate drug seeking behavior. Considering safety concerns raised by adverse effects found in humans after ibogaine intake, such as prolongation of the QT_C interval in the EKG (which has been associated with sudden death cases after ibogaine intake) (Koenig and Hilber, 2015), contributions to the understanding of ibogaine's mechanism of action will provide basis for the preparation of safer and more effective analogs in the future.

AUTHOR CONTRIBUTIONS

SM, BG, and SR-B performed the qPCR experiments. EM and SR-B performed the Western Blot experiments. SM, BG, LM-P, and MP contributed in the experiments and analyzed the data. BG prepared the ibogaine HCl used in this study. JP, PR, and CS performed the experiments with animals and the brain dissection. IC, PC, GS, and CS provided the funding for the experiments. MP, PC, GS, CS, DS, and IC planned the experiments and wrote the manuscript. All the authors participated in critical revision the manuscript, added important intellectual content, and approved the definitive version.

FUNDING

This work was supported by the Agencia Nacional de Investigación e Innovación (ANII, Montevideo – Uruguay)

Project Fondo María Viñas 103488, Comisión Sectorial de Investigación Científica (UdelaR) – Projects Grupos I+D 981 and 1104, and Programa de Desarrollo de las Ciencias Básicas (PEDECIBA).

ACKNOWLEDGMENTS

We would like to thank Agencia Nacional de Investigación e Innovación (ANII) and Comisión Sectorial de Investigación Científica (CSIC-UdelaR) for financial support. We also thank

Inés Carrera, Analia Richeri, and Juan Pablo Rodríguez for their help in designing the manuscript figures and Dr. Kenneth Alper for important discussions regarding ibogaine and noribogaine pharmacology.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphar.2019.00193/full#supplementary-material>

REFERENCES

- Aloe, L., Bracci-Laudiero, L., and Tirassa, P. (1993). The effect of chronic ethanol intake on brain NGF level and on NGF-target tissues of adult mice. *Drug Alcohol Depend.* 31, 159–167. doi: 10.1016/0376-8716(93)90068-2
- Alper, K. R. (2001). Ibogaine: a review. *Alkaloids Chem. Biol.* 56, 1–38. doi: 10.1016/S0099-9598(01)56005-8
- Angelucci, F., Ricci, V., Pomponi, M., Conte, G., Mathe, A. A., Attilio Tonalì, P., et al. (2007). Chronic heroin and cocaine abuse is associated with decreased serum concentrations of the nerve growth factor and brain-derived neurotrophic factor. *J. Psychopharmacol.* 21, 820–825. doi: 10.1177/0269881107078491
- Antonio, T., Childers, S. R., Rothman, R. B., Dersch, C. M., King, C., Kuehne, M., et al. (2013). Effect of Iboga alkaloids on micro-opioid receptor-coupled G protein activation. *PLoS One* 8:e77262. doi: 10.1371/journal.pone.0077262
- Arias, H. R., Rosenberg, A., Targowska-Duda, K. M., Feuerbach, D., Yuan, X. J., Jozwiak, K., et al. (2010). Interaction of ibogaine with human alpha3beta4-nicotinic acetylcholine receptors in different conformational states. *Int. J. Biochem. Cell Biol.* 42, 1525–1535. doi: 10.1016/j.biocel.2010.05.011
- Arias, H. R., Targowska-Duda, K. M., Feuerbach, D., and Jozwiak, K. (2015). Coronaridine congeners inhibit human alpha3beta4 nicotinic acetylcholine receptors by interacting with luminal and non-luminal sites. *Int. J. Biochem. Cell Biol.* 65, 81–90. doi: 10.1016/j.biocel.2015.05.015
- Asjad, H. M. M., Kasture, A., El-Kasaby, A., Sackel, M., Hummel, T., Freissmuth, M., et al. (2017). Pharmacochaperoning in a Drosophila model system rescues human dopamine transporter variants associated with infantile/juvenile parkinsonism. *J. Biol. Chem.* 292, 19250–19265. doi: 10.1074/jbc.M117.797092
- Bahi, A., Boyer, F., Chandrasekar, V., and Dreyer, J. L. (2008). Role of accumbens BDNF and TrkB in cocaine-induced psychomotor sensitization, conditioned-place preference, and reinstatement in rats. *Psychopharmacology* 199, 169–182. doi: 10.1007/s00213-008-1164-1
- Barde, Y. A. (1990). The nerve growth factor family. *Prog. Growth Factor Res.* 2, 237–248. doi: 10.1016/0955-2235(90)90021-B
- Baumann, M. H., Pablo, J., Ali, S. F., Rothman, R. B., and Mash, D. C. (2001a). Comparative neuropharmacology of ibogaine and its O-desmethyl metabolite, noribogaine. *Alkaloids Chem. Biol.* 56, 79–113. doi: 10.1016/S0099-9598(01)56009-5
- Baumann, M. H., Rothman, R. B., Pablo, J. P., and Mash, D. C. (2001b). In vivo neurobiological effects of ibogaine and its O-desmethyl metabolite, 12-hydroxyibogamine (noribogaine), in rats. *J. Pharmacol. Exp. Therap.* 297, 531–539.
- Benarroch, E. E. (2015). Brain-derived neurotrophic factor: regulation, effects, and potential clinical relevance. *Neurology* 84, 1693–1704. doi: 10.1212/WNL.0000000000001507
- Berglind, W. J., See, R. E., Fuchs, R. A., Ghee, S. M., Whitfield, TW Jr, Miller, S. W., et al. (2007). A BDNF infusion into the medial prefrontal cortex suppresses cocaine seeking in rats. *Eur. J. Neurosci.* 26, 757–766. doi: 10.1111/j.1460-9568.2007.05692.x
- Bie, B., Wang, Y., Cai, Y. Q., Zhang, Z., Hou, Y. Y., and Pan, Z. Z. (2012). Upregulation of nerve growth factor in central amygdala increases sensitivity to opioid reward. *Neuropsychopharmacology* 37, 2780–2788. doi: 10.1038/npp.2012.144
- Bolaños, C. A., and Nestler, E. J. (2004). Neurotrophic mechanisms in drug addiction. *Neuromol. Med.* 5, 69–83. doi: 10.1385/NMM:5:1:069
- Borodina, A. A., and Salozhin, S. V. (2016). [Diversity of proBDNF and mBDNF functions in the central nervous system]. *Zh Vyssh Nerv Deiat Im I P Pavlova* 66, 3–23.
- Brown, T. K. (2013). Ibogaine in the treatment of substance dependence. *Curr. Drug Abuse Rev.* 6, 3–16. doi: 10.2174/15672050113109990001
- Brown, T. K., and Alper, K. (2017). Treatment of opioid use disorder with ibogaine: detoxification and drug use outcomes. *Am. J. Drug Alcohol Abuse* 44, 24–36. doi: 10.1080/00952990.2017.1320802
- Bulling, S., Schicker, K., Zhang, Y. W., Steinkellner, T., Stockner, T., Gruber, C. W., et al. (2012). The mechanistic basis for noncompetitive ibogaine inhibition of serotonin and dopamine transporters. *J. Biol. Chem.* 287, 18524–18534. doi: 10.1074/jbc.M112.343681
- Burke, A. R., and Miczek, K. A. (2015). Escalation of cocaine self-administration in adulthood after social defeat of adolescent rats: role of social experience and adaptive coping behavior. *Psychopharmacology* 232, 3067–3079. doi: 10.1007/s00213-015-3947-5
- Calabresi, P., Picconi, B., Tozzi, A., Ghiglieri, V., and Di Filippo, M. (2014). Direct and indirect pathways of basal ganglia: a critical reappraisal. *Nat. Neurosci.* 17, 1022–1030. doi: 10.1038/nn.3743
- Cappendijk, S. L., and Dzoljic, M. R. (1993). Inhibitory effects of ibogaine on cocaine self-administration in rats. *Eur. J. Pharmacol.* 241, 261–265. doi: 10.1016/0014-2999(93)90212-Z
- Carnicella, S., Ahmadiantehrani, S., Janak, P. H., and Ron, D. (2009). GDNF is an endogenous negative regulator of ethanol-mediated reward and of ethanol consumption after a period of abstinence. *Alcohol. Clin. Exp. Res.* 33, 1012–1024. doi: 10.1111/j.1530-0277.2009.00922.x
- Carnicella, S., He, D. Y., Yowell, Q. V., Glick, S. D., and Ron, D. (2010). Noribogaine, but not 18-MC, exhibits similar actions as ibogaine on GDNF expression and ethanol self-administration. *Addict. Biol.* 15, 424–433. doi: 10.1111/j.1369-1600.2010.00251.x
- Carnicella, S., Kharazia, V., Jeanblanc, J., Janak, P. H., and Ron, D. (2008). GDNF is a fast-acting potent inhibitor of alcohol consumption and relapse. *Proc. Natl. Acad. Sci. U.S.A.* 105, 8114–8119. doi: 10.1073/pnas.0711755105
- Carnicella, S., and Ron, D. (2009). GDNF—a potential target to treat addiction. *Pharmacol. Ther.* 122, 9–18. doi: 10.1016/j.pharmthera.2008.12.001
- Castren, E., and Antila, H. (2017). Neuronal plasticity and neurotrophic factors in drug responses. *Mol. Psychiatry* 22, 1085–1095. doi: 10.1038/mp.2017.61
- Corkery, J. M. (2018). Ibogaine as a treatment for substance misuse: potential benefits and practical dangers. *Prog. Brain Res.* 242, 217–257. doi: 10.1016/bs.pbr.2018.08.005
- Dauer, W., and Przedborski, S. (2003). Parkinson's disease: mechanisms and models. *Neuron* 39, 889–909. doi: 10.1016/S0896-6273(03)00568-3
- Day, M., Wokosin, D., Plotkin, J. L., Tian, X., and Surmeier, D. J. (2008). Differential excitability and modulation of striatal medium spiny neuron dendrites. *J. Neurosci.* 28, 11603–11614. doi: 10.1523/JNEUROSCI.1840-08.2008
- de Sousa Abreu, R., Penalva, L. O., Marcotte, E. M., and Vogel, C. (2009). Global signatures of protein and mRNA expression levels. *Mol. Biosyst.* 5, 1512–1526. doi: 10.1039/b908315d
- Di Chiara, G., and Imperato, A. (1988). Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely

- moving rats. *Proc. Natl. Acad. Sci. U.S.A.* 85, 5274–5278. doi: 10.1073/pnas.85.14.5274
- Dwivedi, Y., Rizavi, H. S., and Pandey, G. N. (2006). Antidepressants reverse corticosterone-mediated decrease in brain-derived neurotrophic factor expression: differential regulation of specific exons by antidepressants and corticosterone. *Neuroscience* 139, 1017–1029. doi: 10.1016/j.neuroscience.2005.12.058
- Dworkin, S. I., Gleeson, S., Meloni, D., Koves, T. R., and Martin, T. J. (1995). Effects of ibogaine on responding maintained by food, cocaine and heroin reinforcement in rats. *Psychopharmacology* 117, 257–261. doi: 10.1007/BF02246099
- Dzolic, E. D., Kaplan, C. D., and Dzolic, M. R. (1988). Effect of ibogaine on naloxone-precipitated withdrawal syndrome in chronic morphine-dependent rats. *Arch. Int. Pharmacodyn. Ther.* 294, 64–70.
- Evans, S. F., Irmady, K., Ostrow, K., Kim, T., Nykjaer, A., Saftig, P., et al. (2011). Neuronal brain-derived neurotrophic factor is synthesized in excess, with levels regulated by sortilin-mediated trafficking and lysosomal degradation. *J. Biol. Chem.* 286, 29556–29567. doi: 10.1074/jbc.M111.219675
- Fryer, J. D., and Lukas, R. J. (1999). Noncompetitive functional inhibition at diverse, human nicotinic acetylcholine receptor subtypes by bupropion, phencyclidine, and ibogaine. *J. Pharmacol. Exp. Ther.* 288, 88–92.
- Gassaway, M. M., Jacques, T. L., Kruegel, A. C., Karpowicz, R. J., Li, X., Li, S., et al. (2016). Deconstructing the iboga alkaloid skeleton: potentiation of FGF2-induced glial cell line-derived neurotrophic factor release by a novel compound. *ACS Chem. Biol.* 11, 77–87. doi: 10.1021/acscmbio.5b00678
- Ghitza, U. E., Zhai, H., Wu, P., Airavaara, M., Shaham, Y., and Lu, L. (2010). Role of BDNF and GDNF in drug reward and relapse: a review. *Neurosci. Biobehav. Rev.* 35, 157–171. doi: 10.1016/j.neubiorev.2009.11.009
- Glick, S. D., Kuehne, M. E., Raucchi, J., Wilson, T. E., Larson, D., Keller, R. W., et al. (1994). Effects of iboga alkaloids on morphine and cocaine self-administration in rats: relationship to tremorigenic effects and to effects on dopamine release in nucleus accumbens and striatum. *Brain Res.* 657, 14–22. doi: 10.1016/0006-8993(94)90948-2
- Glick, S. D., Maisonneuve, I. M., and Szumlinski, K. K. (2000). 18-Methoxycoronaridine (18-MC) and ibogaine: comparison of antiaddictive efficacy, toxicity, and mechanisms of action. *Ann. N. Y. Acad. Sci.* 914, 369–386. doi: 10.1111/j.1749-6632.2000.tb05211.x
- Glick, S. D., Maisonneuve, I. M., and Szumlinski, K. K. (2001). Mechanisms of action of ibogaine: relevance to putative therapeutic effects and development of a safer iboga alkaloid congener. *Alkaloids Chem. Biol.* 56, 39–53. doi: 10.1016/S0099-9598(01)56006-X
- Glick, S. D., Rossman, K., Rao, N. C., Maisonneuve, I. M., and Carlson, J. N. (1992). Effects of ibogaine on acute signs of morphine withdrawal in rats: independence from tremor. *Neuropharmacology* 31, 497–500. doi: 10.1016/0028-3908(92)90089-8
- Glick, S. D., Rossman, K., Steindorf, S., Maisonneuve, I. M., and Carlson, J. N. (1991). Effects and aftereffects of ibogaine on morphine self-administration in rats. *Eur. J. Pharmacol.* 195, 341–345. doi: 10.1016/0014-2999(91)90474-5
- Golan, M., Schreiber, G., and Avissar, S. (2011). Antidepressants elevate GDNF expression and release from C(6) glioma cells in a beta-arrestin1-dependent, CREB interactive pathway. *Int. J. Neuropsychopharmacol.* 14, 1289–1300. doi: 10.1017/S1461145710001550
- Gonzalez, J., Prieto, J. P., Rodríguez, P., Cavelli, M., Benedetto, L., Mondino, A., et al. (2018). Ibogaine acute administration in rats promotes wakefulness, long-lasting REM sleep suppression, and a distinctive motor profile. *Front. Pharmacol.* 9:374. doi: 10.3389/fphar.2018.00374
- Graham, D. L., Edwards, S., Bachtell, R. K., Dileone, R. J., Rios, M., and Self, D. W. (2007). Dynamic BDNF activity in nucleus accumbens with cocaine use increases self-administration and relapse. *Nat. Neurosci.* 10, 1029–1037. doi: 10.1038/nn1929
- He, D.-Y., Mcgough, N. N. H., Ravindranathan, A., Jeanblanc, J., Logrip, M. L., Phamluong, K., et al. (2005). Glial cell line-derived neurotrophic factor mediates the desirable actions of the anti-addiction drug ibogaine against alcohol consumption. *J. Neurosci.* 25, 619–628. doi: 10.1523/JNEUROSCI.3959-04.2005
- He, D.-Y., and Ron, D. (2006). Autoregulation of glial cell line-derived neurotrophic factor expression: implications for the long-lasting actions of the anti-addiction drug, Ibogaine. *FASEB J.* 20, 2420–2422. doi: 10.1096/fj.06-6394fje
- Hisaoka, K., Nishida, A., Koda, T., Miyata, M., Zensho, H., Morinobu, S., et al. (2001). Antidepressant drug treatments induce glial cell line-derived neurotrophic factor (GDNF) synthesis and release in rat C6 glioblastoma cells. *J. Neurochem.* 79, 25–34. doi: 10.1046/j.1471-4159.2001.00531.x
- Huang, Z., Zhong, X. M., Li, Z. Y., Feng, C. R., Pan, A. J., and Mao, Q. Q. (2011). Curcumin reverses corticosterone-induced depressive-like behavior and decrease in brain BDNF levels in rats. *Neurosci. Lett.* 493, 145–148. doi: 10.1016/j.neulet.2011.02.030
- Jacobs, M. T., Zhang, Y. W., Campbell, S. D., and Rudnick, G. (2007). Ibogaine, a noncompetitive inhibitor of serotonin transport, acts by stabilizing the cytoplasm-facing state of the transporter. *J. Biol. Chem.* 282, 29441–29447. doi: 10.1074/jbc.M704456200
- Jeanblanc, J., He, D. Y., Carnicella, S., Kharazia, V., Janak, P. H., and Ron, D. (2009). Endogenous BDNF in the dorsolateral striatum gates alcohol drinking. *J. Neurosci.* 29, 13494–13502. doi: 10.1523/JNEUROSCI.2243-09.2009
- Jenks, C. W. (2002). Extraction studies of *Tabernanthe iboga* and *Voacanga africana*. *Nat. Prod. Lett.* 16, 71–76. doi: 10.1080/10575630290014881
- Kalivas, P. W., and Volkow, N. D. (2005). The neural basis of addiction: a pathology of motivation and choice. *Am. J. Psychiatry* 162, 1403–1413. doi: 10.1176/appi.ajp.162.8.1403
- Koenig, X., and Hilber, K. (2015). The anti-addiction drug ibogaine and the heart: a delicate relation. *Molecules* 20, 2208–2228. doi: 10.3390/molecules20022208
- Koob, G. F., and Bloom, F. E. (1988). Cellular and molecular mechanisms of drug dependence. *Science* 242, 715–723. doi: 10.1126/science.2903550
- Koskela, M., Bäck, S., Vöikar, V., Richie, C. T., Domanskyi, A., Harvey, B. K., et al. (2017). Update of neurotrophic factors in neurobiology of addiction and future directions. *Neurobiol. Dis.* 97(Pt B), 189–200. doi: 10.1016/j.nbd.2016.05.010
- Krishnan, V., Han, M. H., Graham, D. L., Berton, O., Renthal, W., Russo, S. J., et al. (2007). Molecular adaptations underlying susceptibility and resistance to social defeat in brain reward regions. *Cell* 131, 391–404. doi: 10.1016/j.cell.2007.09.018
- Lavaud, C., and Massiot, G. (2017). The iboga alkaloids. *Prog. Chem. Org. Nat. Prod.* 105, 89–136. doi: 10.1007/978-3-319-49712-9_2
- Leal, M. B., Michelin, K., Souza, D. O., and Elisabetsky, E. (2003). Ibogaine attenuation of morphine withdrawal in mice: role of glutamate N-methyl-D-aspartate receptors. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 27, 781–785. doi: 10.1016/S0278-5846(03)00109-X
- Li, J. Y., Liu, J., Manaph, N. P. A., Bobrovskaya, L., and Zhou, X. F. (2017). ProBDNF inhibits proliferation, migration and differentiation of mouse neural stem cells. *Brain Res.* 1668, 46–55. doi: 10.1016/j.brainres.2017.05.013
- Li, L., Cao, J., Zhang, S., Wang, C., Wang, J., Song, G., et al. (2014). NCAM signaling mediates the effects of GDNF on chronic morphine-induced neuroadaptations. *J. Mol. Neurosci.* 53, 580–589. doi: 10.1007/s12031-013-0224-0
- Lu, B., and Figurov, A. (1997). Role of neurotrophins in synapse development and plasticity. *Rev. Neurosci.* 8, 1–12. doi: 10.1515/REVNEURO.1997.8.1.1
- Lu, B., Pang, P. T., and Woo, N. H. (2005). The yin and yang of neurotrophin action. *Nat. Rev. Neurosci.* 6, 603–614. doi: 10.1038/nrn1726
- Ly, C., Greb, A. C., Cameron, L. P., Wong, J. M., Barragan, E. V., Wilson, P. C., et al. (2018). Psychedelics promote structural and functional neural plasticity. *Cell Rep.* 23, 3170–3182. doi: 10.1016/j.celrep.2018.05.022
- Maciulaitis, R., Kontrimaviciute, V., Bressolle, F. M., and Briedis, V. (2008). Ibogaine, an anti-addictive drug: pharmacology and time to go further in development. A narrative review. *Hum. Exp. Toxicol.* 27, 181–194. doi: 10.1177/0960327107087802
- Maier, T., Guell, M., and Serrano, L. (2009). Correlation of mRNA and protein in complex biological samples. *FEBS Lett.* 583, 3966–3973. doi: 10.1016/j.febslet.2009.10.036
- Maillet, E. L., Milon, N., Heghinian, M. D., Fishback, J., Schurer, S. C., Garamszegi, N., et al. (2015). Noribogaine is a G-protein biased kappa-opioid receptor agonist. *Neuropharmacology* 99, 675–688. doi: 10.1016/j.neuropharm.2015.08.032
- Malcolm, B. J., Polanco, M., and Barsuglia, J. P. (2018). Changes in withdrawal and craving scores in participants undergoing opioid detoxification utilizing ibogaine. *J. Psychoactive Drugs* 50, 256–265. doi: 10.1080/02791072.2018.1447175
- Mash, D. C., Duque, L., Page, B., and Allen-Ferdinand, K. (2018). Ibogaine detoxification transitions opioid and cocaine abusers between dependence and abstinence: clinical observations and treatment outcomes. *Front. Pharmacol.* 9:529. doi: 10.3389/fphar.2018.00529

- Mash, D. C., Staley, J. K., Baumann, M. H., Rothman, R. B., and Hearn, W. L. (1995a). Identification of a primary metabolite of ibogaine that targets serotonin transporters and elevates serotonin. *Life Sci.* 57, L45–L50.
- Mash, D. C., Staley, J. K., Pablo, J. P., Holohean, A. M., Hackman, J. C., and Davidoff, R. A. (1995b). Properties of ibogaine and its principal metabolite (12-hydroxyibogamine) at the MK-801 binding site of the NMDA receptor complex. *Neurosci. Lett.* 192, 53–56.
- McGough, N. N., He, D. Y., Logrip, M. L., Jeanblanc, J., Phamluong, K., Luong, K., et al. (2004). RACK1 and brain-derived neurotrophic factor: a homeostatic pathway that regulates alcohol addiction. *J. Neurosci.* 24, 10542–10552. doi: 10.1523/JNEUROSCI.3714-04.2004
- Meikle, M. N., Prieto, J. P., Urbanavicius, J., Lopez, X., Abin-Carriquiry, J. A., Prunell, G., et al. (2013). Anti-aggressive effect elicited by coca-paste in isolation-induced aggression of male rats: influence of accumbal dopamine and cortical serotonin. *Pharmacol. Biochem. Behav.* 110, 216–223. doi: 10.1016/j.pbb.2013.07.010
- Mercier, G., Lennon, A. M., Renouf, B., Dessouroux, A., Ramaugue, M., Courtin, F., et al. (2004). MAP kinase activation by fluoxetine and its relation to gene expression in cultured rat astrocytes. *J. Mol. Neurosci.* 24, 207–216. doi: 10.1385/JMN:24:2:207
- Messer, C. J., Eisch, A. J., Carlezon, W. A., Whisler, K., Shen, L., Wolf, D. H., et al. (2000). Role for GDNF in biochemical and behavioral adaptations to drugs of abuse. *Neuron* 26, 247–257. doi: 10.1016/S0896-6273(00)81154-X
- Noller, G. E., Frampton, C. M., and Yazar-Klosinski, B. (2017). Ibogaine treatment outcomes for opioid dependence from a twelve-month follow-up observational study. *Am. J. Drug Alcohol Abuse* 44, 37–46. doi: 10.1080/00952990.2017.1310218
- Paxinos, G., and Watson, C. (2005). *The Rat Brain in Stereotaxic Coordinates*. Sydney, NSW: Academic Press.
- Pearl, S. M., Hough, L. B., Boyd, D. L., and Glick, S. D. (1997). Sex differences in ibogaine antagonism of morphine-induced locomotor activity and in ibogaine brain levels and metabolism. *Pharmacol. Biochem. Behav.* 57, 809–815. doi: 10.1016/S0091-3057(96)00383-8
- Popova, N. K., Ilchibaeva, T. V., and Naumenko, V. S. (2017). Neurotrophic factors (BDNF and GDNF) and the serotonergic system of the brain. *Biochemistry* 82, 308–317. doi: 10.1134/S0006297917030099
- Rantamaki, T., Hendolin, P., Kankaanpaa, A., Mijatovic, J., Piepponen, P., Domenici, E., et al. (2007). Pharmacologically diverse antidepressants rapidly activate brain-derived neurotrophic factor receptor TrkB and induce phospholipase-Cgamma signaling pathways in mouse brain. *Neuropsychopharmacology* 32, 2152–2162. doi: 10.1038/sj.npp.1301345
- Reichardt, L. F. (2006). Neurotrophin-regulated signalling pathways. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 361, 1545–1564. doi: 10.1098/rstb.2006.1894
- Russo, S. J., Mazei-Robison, M. S., Ables, J. L., and Nestler, E. J. (2009). Neurotrophic factors and structural plasticity in addiction. *Neuropharmacology* 56(Suppl. 1), 73–82. doi: 10.1016/j.neuropharm.2008.06.059
- Schenberg, E. E., De Castro Comis, M. A., Chaves, B. R., and Da Silveira, D. X. (2014). Treating drug dependence with the aid of ibogaine: a retrospective study. *J. Psychopharmacol.* 28, 993–1000. doi: 10.1177/0269881114552713
- Schmidt, H. D., and Duman, R. S. (2007). The role of neurotrophic factors in adult hippocampal neurogenesis, antidepressant treatments and animal models of depressive-like behavior. *Behav. Pharmacol.* 18, 391–418. doi: 10.1097/FBP.0b013e3282ee2aa8
- Scorza, M. C., Carrau, C., Silveira, R., Zapata-Torres, G., Cassels, B. K., and Reyes-Parada, M. (1997). Monoamine oxidase inhibitory properties of some methoxylated and alkylthio amphetamine derivatives: structure-activity relationships. *Biochem. Pharmacol.* 54, 1361–1369. doi: 10.1016/S0006-2952(97)00405-X
- Shadfar, S., Kim, Y. G., Katila, N., Neupane, S., Ojha, U., Bhurtel, S., et al. (2018). Neuroprotective effects of antidepressants via upregulation of neurotrophic factors in the MPTP model of Parkinson's disease. *Mol. Neurobiol.* 55, 554–566. doi: 10.1007/s12035-016-0342-0
- Staley, J. K., Ouyang, Q., Pablo, J., Hearn, W. L., Flynn, D. D., Rothman, R. B., et al. (1996). Pharmacological screen for activities of 12-hydroxyibogamine: a primary metabolite of the indole alkaloid ibogaine. *Psychopharmacology* 127, 10–18. doi: 10.1007/BF02805969
- Sun, Y., Lim, Y., Li, F., Liu, S., Lu, J. J., Haberberger, R., et al. (2012). ProBDNF collapses neurite outgrowth of primary neurons by activating RhoA. *PLoS One* 7:e35883. doi: 10.1371/journal.pone.0035883
- Teng, H. K., Teng, K. K., Lee, R., Wright, S., Tevar, S., Almeida, R. D., et al. (2005). ProBDNF induces neuronal apoptosis via activation of a receptor complex of p75NTR and sortilin. *J. Neurosci.* 25, 5455–5463. doi: 10.1523/JNEUROSCI.5123-04.2005
- Tsuchioka, M., Takebayashi, M., Hisaoka, K., Maeda, N., and Nakata, Y. (2008). Serotonin (5-HT) induces glial cell line-derived neurotrophic factor (GDNF) mRNA expression via the transactivation of fibroblast growth factor receptor 2 (FGFR2) in rat C6 glioma cells. *J. Neurochem.* 106, 244–257. doi: 10.1111/j.1471-4159.2008.05357.x
- Wei, D., Maisonneuve, I. M., Kuehne, M. E., and Glick, S. D. (1998). Acute ibogaine effects on extracellular serotonin (5-HT) levels in nucleus accumbens and striatum in rats. *Brain Res.* 800, 260–268. doi: 10.1016/S0006-8993(98)00527-7
- Wells, G. B., Lopez, M. C., and Tanaka, J. C. (1999). The effects of ibogaine on dopamine and serotonin transport in rat brain synaptosomes. *Brain Res. Bull.* 48, 641–647. doi: 10.1016/S0361-9230(99)00053-2
- Woo, N. H., Teng, H. K., Siao, C. J., Chiaruttini, C., Pang, P. T., Milner, T. A., et al. (2005). Activation of p75NTR by proBDNF facilitates hippocampal long-term depression. *Nat. Neurosci.* 8, 1069–1077. doi: 10.1038/nn1510
- Xu, Z. Q., Sun, Y., Li, H. Y., Lim, Y., Zhong, J. H., and Zhou, X. F. (2011). Endogenous proBDNF is a negative regulator of migration of cerebellar granule cells in neonatal mice. *Eur. J. Neurosci.* 33, 1376–1384. doi: 10.1111/j.1460-9568.2011.07635.x
- Yang, J. W., Ma, W., Yang, Y. L., Wang, X. B., Li, X. T., Wang, T. T., et al. (2017). Region-specific expression of precursor and mature brain-derived neurotrophic factors after chronic alcohol exposure. *Am. J. Drug Alcohol Abuse* 43, 602–608. doi: 10.1080/00952990.2016.1263642
- Zubaran, C., Shoab, M., Stoleran, I. P., Pablo, J., and Mash, D. C. (1999). Noribogaine generalization to the ibogaine stimulus: correlation with noribogaine concentration in rat brain. *Neuropsychopharmacology* 21, 119–126. doi: 10.1016/S0893-133X(99)00003-2

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Marton, González, Rodríguez-Bottero, Miquel, Martínez-Palma, Pazos, Prieto, Rodríguez, Sames, Seoane, Scorza, Cassina and Carrera. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.