



# **Corrigendum: Vitamin B12 Enhances Nerve Repair and Improves Functional Recovery After Traumatic Brain Injury by Inhibiting ER Stress-Induced Neuron Injury**

# OPEN ACCESS

## Edited and reviewed by:

Nicholas M. Barnes, University of Birmingham, United Kingdom

#### \*Correspondence:

Daqing Chen cdq1965@126.com Aiping Hu hap@wmu.edu.cn <sup>†</sup>These authors contributed equally to this work

#### Specialty section:

This article was submitted to Neuropharmacology, a section of the journal Frontiers in Pharmacology

Received: 24 August 2020 Accepted: 22 February 2021 Published: 12 April 2021

#### Citation:

Wu F, Xu K, Liu L, Zhang K, Xia L, Zhang M, Teng C, Tong H, He Y, Xue Y, Zhang H, Chen D and Hu A (2021) Corrigendum: Vitamin B12 Enhances Nerve Repair and Improves Functional Recovery After Traumatic Brain Injury by Inhibiting ER Stress-Induced Neuron Injury. Front. Pharmacol. 12:598335. doi: 10.3389/fphar.2021.598335 Fangfang Wu<sup>1,2†</sup>, Ke Xu<sup>3†</sup>, Lei Liu<sup>1†</sup>, Kairui Zhang<sup>1</sup>, Leilei Xia<sup>4</sup>, Man Zhang<sup>1</sup>, Chenhuai Teng<sup>1</sup>, Heyan Tong<sup>2</sup>, Yifang He<sup>2</sup>, Yujie Xue<sup>2</sup>, Hongyu Zhang<sup>2</sup>, Daqing Chen<sup>1\*</sup> and Aiping Hu<sup>1,2\*</sup>

<sup>1</sup>Department of Emergency, The Second Affiliated Hospital and Yuying Children's Hospital, Wenzhou Medical University, Wenzhou, China, <sup>2</sup>School of Pharmaceutical Sciences, Wenzhou Medical University, Wenzhou, China, <sup>3</sup>Institute of Life Sciences, Wenzhou University, Wenzhou, China, <sup>4</sup>Department of Emergency, Wenzhou People's Hospital, The Third Clinical Institute Affiliated to Wenzhou Medical University, Wenzhou, China

Keywords: vitamin B12, traumatic brain injury, endoplasmic reticulum stress, microtubule, myelin

### A Corrigendum on

## Vitamin B12 Enhances Nerve Repair and Improves Functional Recovery after Traumatic Brain Injury by Inhibiting ER Stress-Induced neuron injury

by Wu, F., Xu K., Liu, L., Zhang, K.-R., Xia, L.-L., Zhang, M., Teng, C.-H., Tong, H.-Y., He, Y.-F., Xue, Y.-J., Zhang, H.-Y., Chen, D.-Q., Hu, A.-P. (2019). Front. Pharmacol. 10, 406. doi:10.3389/fphar.2019. 00406

In the original article, there was a mistake in Figure 4P as published. The corrected Figure 4 appear below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way.

Copyright © 2021 Wu, Xu, Liu, Zhang, Xia, Zhang, Teng, Tong, He, Xue, Zhang, Chen and Hu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

1



**FIGURE 4** | Vitamin B12 inhibits microtubule damage through ER stress after TBI. (A) Representative western blots of the expression of MAP-2, Tau. (B, C) Quantification from (A).##p < 0.01, #p < 0.05 vs. the sham group. \*p < 0.05, && p < 0.01 vs. the TBI group. \*\*p < 0.01 vs. the indicated group. Values represent the mean ± SEM, n = 5. (D) Representative fluorescent images depicting MAP-2 (green) with NeuN (red) in the cortex after TBI. Scale bar = 50 µm, scale bar = 10 µm. (E) Representative fluorescent images depicting Ace-tubulin(green) with Tyr-tubulin(red) in the PC 12 cells. Scale bar = 20 µm. (F) Representative western blots of the expression of IRE1 $\alpha$ , GRP78, XBP-1, Chop, Cleaved-cas12. (G–K) Quantification from (F)\*p < 0.05, \*\*p < 0.01 and \*\*\*p < 0.01 vs. the indicated group. Values represent the mean ± SEM, n = 4. (L) Representative western blots of the expression of MAP-2, Ace-tubulin. (M–N) Quantification from (L).\*p < 0.05 vs. the indicated group. Values represent the mean ± SEM, n = 4. (O) Representative fluorescent images depicting Ace-tubulin(green) with Tyr-tubulin(red) in the PC 12 cells. Scale bar = 20 µm. (P). Representative fluorescent images depicting MAP-2 (green) *in vivo*. Scale bar = 50 µm.