



## OPEN ACCESS

## EDITED BY

Sanjay Kalra,  
Independent researcher, Karnal, India

## REVIEWED BY

Djordje S. Popovic,  
University of Novi Sad, Serbia  
Maria Dolores Garcia De Lucas,  
Agencia Sanitaria Costa del Sol, Spain  
Alina Yu Babenko,  
Almazov National Medical Research  
Centre, Russia

## \*CORRESPONDENCE

Xuening Li,  
li.xuening@zs-hospital.sh.cn  
Guo Ma,  
mg0328@fudan.edu.cn

## SPECIALTY SECTION

This article was submitted to Drugs  
Outcomes Research and Policies,  
a section of the journal  
Frontiers in Pharmacology

RECEIVED 20 July 2022

ACCEPTED 28 September 2022

PUBLISHED 14 October 2022

## CITATION

Guan R, Yang Q, Yang X, Du W, Li X and  
Ma G (2022), Efficacy and safety of  
tirzepatide in patients with type  
2 diabetes mellitus: A bayesian  
network meta-analysis.  
*Front. Pharmacol.* 13:998816.  
doi: 10.3389/fphar.2022.998816

## COPYRIGHT

© 2022 Guan, Yang, Yang, Du, Li and Ma.  
This is an open-access article  
distributed under the terms of the  
[Creative Commons Attribution License  
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or  
reproduction in other forums is  
permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original  
publication in this journal is cited, in  
accordance with accepted academic  
practice. No use, distribution or  
reproduction is permitted which does  
not comply with these terms.

# Efficacy and safety of tirzepatide in patients with type 2 diabetes mellitus: A bayesian network meta-analysis

Ruifang Guan<sup>1,2</sup>, Qing Yang<sup>1</sup>, Xiaolei Yang<sup>1</sup>, Wandu Du<sup>1</sup>,  
Xuening Li<sup>2\*</sup> and Guo Ma<sup>1\*</sup>

<sup>1</sup>Department of Clinical Pharmacy, School of Pharmacy, Fudan University, Shanghai, China,

<sup>2</sup>Department of Clinical Pharmacology, Zhongshan Hospital, Fudan University, Shanghai, China

**Background:** In light of clinical trials comparing different doses of tirzepatide with selective glucagon-like peptide-1 receptor agonist (GLP1-RA) or insulin analogue, a bayesian network meta-analysis was conducted to investigate the efficacy and safety of tirzepatide in patients with type 2 diabetes mellitus (T2DM).

**Methods:** We systematically searched PubMed, Embase, Web of science, Cochrane Central Register of Controlled Trials, and ClinicalTrials.gov from their inception to 2 May 2022. Final included studies met the eligibility criteria and methodological quality recommendations. Data analysis was performed using Stata 15.1 software. Each outcome was presented as a mean difference or an odds ratio, and the surface under the cumulative ranking curve value (SCURA).

**Results:** Ultimately, eight eligible RCTs involving 7245 patients were included. Generally speaking, compared with basal insulin (glargine or degludec); selective GLP1-RA (dulaglutide or semaglutide once weekly), 10 and 15 mg of tirzepatide exhibited better antidiabetic and weight-loss effect, especially, 15 mg of tirzepatide was dominant on reducing glycated hemoglobin (SCURA probability: 93.5%), body weight (99.7%), and fasting serum glucose (86.6%). As for safety, insulin caused less gastrointestinal events (93.5%), and there was no statistical difference between GLP1-RA and tirzepatide.

**Conclusion:** Compare with insulin and GLP1-RA, tirzepatide display favorable efficacy and acceptable safety for T2DM patients. More well-designed RCTs are needed to evaluate its clinical performance with higher doses of GLP1-RA and determine its potential cardiovascular benefits.

**Abbreviations:** GLP1-RA, glucagon-like peptide-1 receptor agonist; T2DM, type 2 diabetes mellitus; SCURA, the surface under the cumulative ranking curve value; GIP, glucose-dependent insulinotropic polypeptide; HbA1c, glycated hemoglobin; FSG, fasting serum glucose; RCT, randomized controlled trials; OR, odds ratio; CI, confidence interval.

## KEYWORDS


tirzepatide, type 2 diabetes mellitus, network meta-analysis, efficacy, safety

## 1 Introduction

Type 2 diabetes mellitus (T2DM) is a complex metabolic disorder, which not only characterized by hyperglycemia, but also associated with insulin resistance, impaired insulin secretion, hypertension, dyslipidemia and so on (Ferrannini and Cushman, 2012; DeFronzo et al., 2015; Taskinen and Borén, 2015). The International Diabetes Federation (IDF) estimated there were 537 million patients with diabetes around the world in 2021, and it will reach 783 million by 2045 (Sun et al., 2021). Many risk factors could lead to increased risk for T2DM, e.g., genetic history, unhealthy lifestyle, and obesity. Obesity is believed to be a strong promoter (Malone and Hansen, 2019), and it was also proved that T2DM interlinked with obesity in a pathophysiological way (Pappachan et al., 2019). Poor management of T2DM and obesity will increase the risk of cardiovascular disease (CVD). Some evidence has verified the importance of obesity in the progression of T2DM and CVD, and the

association between weight-loss and long-term decrease in cardiovascular risk (Lavie et al., 2009). Thus, ideal medication of T2DM should be efficacious in lowering glucose as well as promoting weight loss, which have proven cardiovascular benefits and low risk of adverse events (Min and Bain, 2021).

Glucagon-like peptide one receptor agonists (GLP1-RAs) have demonstrated remarkable glycemic control, favorable weight-loss, and cardiorenal outcomes, thus are recommended as first-line injectable therapy for T2DM patients by current guidelines (Draznin et al., 2022). Glucose-dependent insulinotropic polypeptide (GIP) is another incretin hormone, is responsible for the amplification of insulin secretion, and regulates glucose homeostasis together with glucagon-like peptide 1 (GLP-1) (Nauck and Meier, 2016). Yet, unlike GLP1, GIP does not inhibit appetite or food intake (Holst and Rosenkilde, 2020), and provides potential protection against hypoglycemia (Christensen et al., 2011). Moreover, GLP1 and GIP may improve  $\beta$ -cells functionality through synergistical





**FIGURE 2**  
Assessment of the risk of bias in included studies.

pharmacological activation (Bastin and Andreelli, 2019). Therefore, GIP-based therapy has become an attractive candidate to be combined with GLP1-RAs (Papachristou et al., 2021). As a novel GLP1/GIP receptor co-agonist allowing once-weekly subcutaneous administration, tirzepatide was developed in this context.

According to a published systematic review, tirzepatide showed robust reductions of glycated hemoglobin (HbA1c, -1.94%), fasting serum glucose (FSG, -54.72 mg/dl) and body weight (-8.47 kg). It was a safety profile similar to long-acting GLP1-RAs (Bhagavathula et al., 2021). Up to now, there are two reported Phase 2 clinical trials and six Phase 3 clinical trials (SURPASS 1-5, J-mono) to compare the efficacy and safety of tirzepatide with placebo or active comparators (i.e., semaglutide (Frías et al., 2021), dulaglutide (Frias et al., 2018; Inagaki et al., 2022), basal insulin analogue degludec (Ludvik et al., 2021) and glargine (Del Prato et al., 2021; Dahl et al., 2022)). In these studies, the weight-loss and

glucose-lowering effect was greater than the comparators. The gastrointestinal tolerance of tirzepatide was comparable to the GLP1-RAs. Network meta-analysis is a well-established approach which allows the available comparison of a complete set of interventions so as to assess their comparative efficacy and safety (Laws et al., 2019). Accordingly, we conducted a network meta-analysis to systematically estimate the efficacy and safety of tirzepatide to provide basis for its future use.

## 2 Methods

This systematic review conformed to PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analysis) (Moher et al., 2009) and its extension for the network meta-analyses (Hutton et al., 2015).

## 2.1 Search strategy

The literature search was conducted in PubMed, Embase, Web of Science, Cochrane Central Register of Controlled Trials, and [ClinicalTrials.gov](https://www.clinicaltrials.gov) from their inception to 2 May 2022 without language restriction. The databases were searched with the following MeSH (Medical Subject Headings) terms or keywords: 1) “diabetes mellitus, type 2” OR “diabetes mellitus, type II” OR “noninsulin dependent diabetes” OR “non-insulin dependent diabetes” OR “NIDDM” OR “type II diabetes” OR “type 2 diabetes” OR “T2DM” OR “mature onset diabetes” OR “late onset diabetes” OR “adult onset diabetes”; AND 2) “tirzepatide” OR “LY3298176”. Furthermore, the reference lists from the retrieved articles were screened to search for additional relevant studies.

## 2.2 Study selection

Studies were included in the systematic review if they met all of the following criteria: 1) Randomized controlled trials (RCTs) comparing tirzepatide with placebo or active therapeutic interventions in the patient with T2DM; 2) Minimum intervention period of 12 weeks; 3) Investigating the efficacy of tirzepatide on blood glucose parameters or body weight and its safety profile. Secondary analyses and phase I studies were excluded. Two reviewers, working independently, screened citations and evaluated full text records for eligible studies. Discordances were discussed with the third reviewer and resolved by consensus.

## 2.3 Data extraction and quality assessment

For each eligible study, two reviewers (RG and QY) independently extracted the following information: study characteristics (year of publication, study sites, study design), population (sample size, patient demographics, duration of disease; baseline HbA1c), description of interventions (drug class, name, dose), duration of treatment, and measured outcomes. The reviewers resolved disagreements by discussion or consulting the third reviewer (XY).

Since 1 mg was far below the effective dose, and only 29 participants who received 12 mg tirzepatide, these two dosing regimens were excluded. Three doses (5 mg, 10 mg, 15 mg) were included in the meta-analysis. Efficacy outcomes consisted of the mean changes in glycated hemoglobin (HbA1c), fasting serum glucose (FSG), and body weight. The number of gastrointestinal adverse events was chosen to represent safety outcomes. For all the outcomes, we extracted data for the modified intention-to-treat (mITT)

population, which was defined as all the randomly assigned participants who received at least one dose. We assessed the risk of bias for all the included RCTs with the Cochrane risk of bias tool (Sterne et al., 2019), which includes random sequence generation, allocation concealment, blinding, missing outcome data, and selective reporting of outcomes.

## 2.4 Data synthesis and analysis


A network meta-analysis was performed with Bayesian approach using Stata 15.1 (Stata Corp, College Station, TX, United States) in the present study. Effect estimates included odds ratio (OR) for dichotomous outcomes and mean difference (MD) for continuous outcomes. Semaglutide and dulaglutide were combined in a single group (GLP1-RA), and insulin degludec and glargine in another group (Insulin). Point estimates and 95% confidence interval (CI) were assessed using Markov Chain Monte method in Stata with a random-effects model. The surface under the cumulative ranking curve (SUCRA) was also calculated to rank the effectiveness of each treatment. The SUCRA was used to estimate the ranking probabilities for different interventions. A higher SUCRA value indicates a better treatment. Cochran Q test and  $I^2$  statistic were used to assess heterogeneity levels, agreement between direct and indirect estimates in every closed loop of evidence using node splitting approaches.  $p$ -value was used to test the degree of inconsistency. When  $p > 0.05$ , the heterogeneity is not obvious, and the difference within a group is considered as small. Publication bias was evaluated using funnel plots.

## 3 Results

### 3.1 Study selection and characteristics

The electronic search yielded 294 unique records. After removal of 112 duplicate records and screening 182 titles and abstracts, 16 reports were left for full-text assessment. After further selection (Figure 1), eight unique RCTs fulfilled the inclusion criteria (Frias et al., 2018; Frias et al., 2020; Del Prato et al., 2021; Frías et al., 2021; Ludvik et al., 2021; Rosenstock et al., 2021; Dahl et al., 2022; Inagaki et al., 2022). RCTs were published between 2018 and 2022 including 7245 participants (range, 111–1995 participants) with T2DM in total; six (75%) of them were multinational RCTs. Weighted means of baseline HbA1c, weight, and age were 8.2%, 90.3 kg, and 57.9 years, respectively; the duration of interventions ranged from 12 to 52 weeks. The detailed characteristics of individual study are provided in [Supplementary Table S1](#).

The risk of bias in eligible RCTs was shown in Figure 2. Among the eight RCTs, all of them had low risk for bias of incomplete



outcome data and selective reporting, seven RCTs had low risk for bias in random sequence generation, five RCTs had unclear risk for bias in allocation concealment, six RCTs had low risk for bias in blinding participants and personnel, and four RCTs had low risk for bias in blinding the outcome assessment. One study was deemed to possess high performance risk because it did not provide details of blinding the participants (Rosenstock et al., 2021). Overall, these studies had a low or moderate level of risk.

### 3.2 Network meta-analyses

The data on HbA1c, FSG, body weight and safety profile were available from all the eight RCTs. There was no evidence of global inconsistency in any network.

All *p*-values > 0.05, indicating the heterogeneity was not obvious. The treatment network was illustrated in Figure 3.

#### 3.2.1 HbA1c

The results suggested that, compared with Insulin, all the doses of tirzepatide exhibited statistically considerable improvements in reducing HbA1c. 10 mg and 15 mg of tirzepatide were more effective than GLP1-RA. (Table 1). The SUCRA was used to estimate the ranking probabilities for different interventions. A higher SUCRA value indicates a better treatment. Figure 4A and Table 2 presented the SUCRA values of different interventions. 15 mg of tirzepatide was associated with the highest probability of being the best option for the reduction of HbA1c (93.5%), followed by 10 mg of tirzepatide (80.9%), 5 mg of tirzepatide (62.4%), GLP1-RA (36.4%), Insulin (26.5%), and placebo (0.3%).

#### 3.2.2 Body weight

As shown in Table 1, compared with Insulin, all the doses of tirzepatide presented higher effectiveness in controlling body weight. Compared with GLP1-RA, 10 mg and 15 mg of tirzepatide showed statistically significant reductions in body weight. As shown in Table 2, 15 mg of tirzepatide seemed to be the best intervention for controlling body weight with a SUCRA value of 99.7%, followed by 10 mg of tirzepatide (80.1%), 5 mg of tirzepatide (57.9%), GLP1-RA (40.2%), placebo (21.9%), and Insulin (0.1%) (Figure 4B).


#### 3.2.3 FSG

The results indicated that with respect to reducing FSG, there were no differences between all the doses of tirzepatide and Insulin; compared with GLP1-RA, 10 mg and 15 mg of tirzepatide demonstrated more favorable effect (Table 3). The results of the SUCRA analysis (Table 2) showed that 15 mg of tirzepatide (86.6%) seemed to be the most effective option for significantly reducing FSG, followed by 10 mg of tirzepatide (71.6%), Insulin (65.4%), 5 mg of tirzepatide (51.1%), GLP1-RA (25.2%), and placebo (0.1%) (Figure 4C).

TABLE 1 Results of the network meta-analysis of the HbA1c (upper right quarter) and the body weight (lower left quarter).

<b>Tirzepatide 5 mg</b>	<b>-0.80 (-1.57, -0.03)</b>	<b>-0.40 (-1.17,0.37)</b>	<b>-0.13 (-1.03,0.78)</b>	<b>0.70 (0.24,1.15)</b>	<b>0.50 (-0.26,1.26)</b>
6.50 (3.03,9.97)	<b>Tirzepatide 15 mg</b>	0.40 (-0.38,1.18)	1.70 (0.93,2.47)	1.50 (0.60,2.40)	1.30 (0.53,2.07)
3.90 (0.51,7.29)	-2.60 (-6.09,0.89)	<b>Tirzepatide 10 mg</b>	0.11 (-0.78,0.99)	1.10 (0.20,1.99)	0.90 (0.13,1.67)
-2.26 (-6.28,1.76)	-4.40 (-7.80, -1.00)	-1.45 (-5.36,2.46)	<b>Placebo</b>	-1.00 (-1.90, -0.11)	-1.20 (-1.97, -0.43)
-9.37 (-11.33, -7.41)	-15.87 (-19.85, -11.89)	-13.27 (-17.19, -9.36)	-4.97 (-8.89, -1.05)	<b>Insulin</b>	-0.20 (-1.08,0.69)
-2.10 (-5.47,1.27)	-8.60 (-12.08, -5.12)	-6.00 (-9.40, -2.60)	2.30 (-1.11,5.71)	7.27 (3.37,11.17)	<b>GLP1-RA</b>

The differences between the compared groups were deemed as significant when the 95% CI, did not contain 0.00, which is marked with grey background.



**FIGURE 4** Plot of the surface under the cumulative ranking curves for outcomes. (A) HbA1c; (B) Body weight; (C) FSG; (D) gastrointestinal adverse events. Note: The surface under the cumulative ranking curve (SUCRA) was used to estimate the ranking probabilities for different treatments, which ranged from 0% to 100%. A better treatment is indicated by a higher SUCRA value.

**TABLE 2** Ranking probability of various interventions.

Intervention	HbA1c		Body weight		FSG		Safety	
	SUCRA (%)	Rank	SUCRA (%)	Rank	SUCRA (%)	Rank	SUCRA (%)	Rank
Placebo	0.3	6	21.9	5	0.1	6	86.3	2
Insulin	26.5	5	0.1	6	65.4	3	93.5	1
GLP1-RA	36.4	4	40.2	4	25.2	5	33.8	4
Tirzepatide 5 mg	62.4	3	57.9	3	51.1	4	56.1	3
Tirzepatide 10 mg	80.9	2	80.1	2	71.6	2	23.3	5
Tirzepatide 15 mg	93.5	1	99.7	1	86.6	1	7.0	6

### 3.2.4 Safety outcome

The number of gastrointestinal adverse events, which includes nausea, vomiting and diarrhea, was chosen to

summarize the overall safety in this network meta-analysis. As shown in Table 3, compared with GLP1-RA, there were no significant differences on the safety profile of

TABLE 3 Results of the network meta-analysis of the FSG (upper right quarter) and gastrointestinal adverse events (lower left quarter).

<b>Tirzepatide 5 mg</b>	<b>-0.90 (-2.49,0.69)</b>	<b>-1.10 (-2.66,0.46)</b>	<b>-1.16 (-2.99,0.66)</b>	<b>-0.22 (-1.10,0.66)</b>	<b>1.10 (-0.46,2.66)</b>
0.25 (0.09,0.67)	<b>Tirzepatide 15 mg</b>	-0.20 (-1.79,1.39)	3.20 (1.63,4.77)	0.68 (-1.14,2.49)	2.00 (0.40,3.60)
0.47 (0.18,1.24)	1.87 (0.70,4.98)	<b>Tirzepatide 10 mg</b>	-0.17 (-1.97,1.63)	0.88 (-0.91,2.66)	2.20 (0.64,3.76)
0.63 (0.16,2.46)	4.48 (1.31,15.25)	2.31 (0.77,6.91)	<b>Placebo</b>	-3.42 (-5.23, -1.62)	-2.10 (-3.68, -0.52)
6.00 (3.67,9.81)	23.97 (7.96,72.20)	12.82 (4.30,38.24)	1.34 (0.36,5.02)	<b>Insulin</b>	1.32 (-0.47,3.12)
0.66 (0.25,1.73)	2.62 (0.99,6.94)	1.40 (0.54,3.67)	0.15 (0.04,0.49)	0.11 (0.04,0.32)	<b>GLP1-RA</b>

The differences between the compared groups were deemed as significant when the 95% CI, did not contain 0.00 (upper right quarter) or 1.00 (lower left quarter), which is marked with grey background.

all the doses of tirzepatide. All the dosing regimens of tirzepatide caused more gastrointestinal adverse events than Insulin. The superiority of Insulin (93.5%), placebo (86.3%) and 5 mg of tirzepatide (56.1%) in overall safety was further confirmed by the SUCRA analysis (Figure 4D and Table 2). The risk of gastrointestinal adverse events increased along with increase of dose.

### 3.3 Publication bias

Publication bias was measured by the comparison-adjusted funnel plots in this network meta-analysis. Visual inspections indicated that distribution of the included RCTs were relatively symmetrically distributed based on the vertical zero line, and there was some angle between the adjusted auxiliary line and the




FIGURE 5 Comparison-adjusted funnel plot for (A) HbA1c; (B) Body weight; (C) FSG; (D) gastrointestinal adverse events.

horizontal zero line. Therefore, some publication bias may exist (Figure 5).

## 4 Discussion

GIP was not considered as an appealing target for the treatment of T2DM, because early research showed there was no reduction of GIP secretion in patients with diabetes, and GIP agonism seemed to have a less insulinotropic effect (Fisman and Tenenbaum, 2021). Until recently years, GIP agonism were proved effective under better glycemic control (Højberg et al., 2009). It could not only enhance meal-stimulated insulin secretion, but also modulate glucagon secretion in a glucose-dependent manner to facilitate postprandial lipid clearance and regulate food intake (Coskun et al., 2022). In addition, GIP amplifies adipose-tissue sensitivity to insulin, as well as enhancing GLP1-mediated central satiety (Nauck et al., 2021). Since glucagon can reduce body weight and appetite, delay gastrointestinal transit, and stimulate insulin secretion (Dailey and Moran, 2013). Thus, as a regulator of glucagon, GIP/GLP1 dual agonists provide bigger weight reduction than GLP1-RAs. Tirzepatide was developed based on those evidence. It was proved that the effects of tirzepatide improved biomarkers of  $\beta$ -cell function and insulin resistance (Thomas et al., 2021).

Our network meta-analysis identified eight RCTs that randomized T2DM patients to six interventions, including selective GLP-1RA (semaglutide and dulaglutide), basal insulin analogue (degludec and glargine), three different doses of tirzepatide (5 mg, 10 mg, 15 mg), and placebo. In terms of efficacy outcomes, compared with Insulin or GLP1-RA, high doses (10, 15 mg) of tirzepatide showed robust effectiveness in reducing HbA1c, body weight and FSG. As for safety, tirzepatide had similar safety profile with GLP1-RA. Comparing with Insulin, it may increase gastrointestinal adverse events. The GLP1-RA group in the network meta-analysis contained 0.75 mg and 1.5 mg of dulaglutide, and 1 mg of semaglutide, yet higher dosing of regimen had already either received marketing approval (dulaglutide 3 mg and 4.5 mg) or received recommendations of authority (semaglutide 2 mg) (Karagiannis et al., 2022). The comparison of efficacy and safety between tirzepatide and higher doses of GLP1-RA is unclear, thus further RCTs are needed. According to the regulatory guidance of US Food and Drug Administration (FDA), hypoglycemic agents are required to demonstrate cardiovascular safety. Yet, to date, only SURPASS-4 reported that tirzepatide was not associated with excess cardiovascular risk (Del Prato et al., 2021). Another ongoing clinical trial SURPASS-CVOT (NCT04255433) will provide more data on comparing the cardiovascular outcomes of tirzepatide with dulaglutide which has demonstrated cardiovascular benefits in T2DM patients with high cardiovascular risk (Gerstein et al., 2019). Now, management

of obesity is believed as a primary treatment goal of T2DM, and weight-loss is in favor of reversing the underlying metabolic abnormalities of T2DM (Lingvay et al., 2022). Tirzepatide showed robust weight-loss effect even when it was used in conjunction with insulin glargine, which was associated with weight-gain. Tirzepatide seems to be a novel and promising therapeutic option for the management of T2DM, obesity and its related cardiometabolic diseases. In addition to better health outcomes and fewer adverse effects, the patients with T2DM may prefer to minimizing multiple insulin injections, avoiding hypoglycemia, lowering weight or reducing risk of weight gain. Tirzepatide exactly display these characteristics and advantages.

There were several published meta-analyses of tirzepatide (Bhagavathula et al., 2021; Dutta et al., 2021; Karagiannis et al., 2022; Sattar et al., 2022), which only included six or seven RCTs. Our article was the first bayesian network meta-analysis of tirzepatide including eight RCTs, and first combined the participants who received the same dose from different RCTs in one group, then compared and evaluated the relative efficacy and safety of different doses of tirzepatide with Insulin and selective GLP1-RA. In the present study, a sensitive search comprehensively covering the latest research findings, independent study identification, selection and data extraction was performed by two reviewers. It needs to be noted that, heterogeneity in clinical settings, enrolled population, and follow-up duration cannot be avoided. Moreover, publication bias may exist. Especially, as a brand-new drug, at present there are very few RCTs of tirzepatide. In the future, more multicenter RCTs of tirzepatide will provide more information of this promising hypoglycemic drug.

## 5 Conclusion

In summary, efficacy and safety of tirzepatide were systematically evaluated based on the available eight RCTs in this bayesian network meta-analysis. Compared with the marketed antidiabetic drugs (e.g., Insulin, GLP1-RA), tirzepatide exhibited robust efficacy in reducing HbA1c, body weight, and FSG with an acceptable safety profile. In the future, effectiveness, safety, economics, suitability and accessibility of tirzepatide as well as its potential cardiovascular benefits should be further evaluated with evidence-based medicine and real-world evidence.

## Author contributions

RG, XL and GM designed the network meta-analysis; RG, QY, XY and WD performed the network meta-analysis and assessed its quality; RG and GM drafted the manuscript; XL

and GM revised the manuscript. All authors read and approved the final version of the manuscript.

## Funding

This work was supported by the National Natural Science Funds of China (Grants 82074109, 81873078, and 81374051) and MHHFDU-SPFDU Joint Research Fund (Grants 2021MHJC11).

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## References

- Bastin, M., and Andreelli, F. (2019). Dual GIP-GLP1-receptor agonists in the treatment of type 2 diabetes: A short review on emerging data and therapeutic potential. *Diabetes Metab. Syndr. Obes.* 12, 1973–1985. doi:10.2147/dmso.S191438
- Bhagavathula, A. S., Vidyasagar, K., and Tesfaye, W. (2021). Efficacy and safety of tirzepatide in patients with type 2 diabetes mellitus: A systematic review and meta-analysis of randomized phase II/III trials. *Pharm. (Basel)* 14 (10), 991. doi:10.3390/ph14100991
- Christensen, M., Vedtofte, L., Holst, J. J., Vilsbøll, T., and Knop, F. K. (2011). Glucose-dependent insulinotropic polypeptide: A bifunctional glucose-dependent regulator of glucagon and insulin secretion in humans. *Diabetes* 60 (12), 3103–3109. doi:10.2337/db11-0979
- Coskun, T., Urva, S., Roell, W. C., Qu, H., Loghin, C., Moyers, J. S., et al. (2022). LY3437943, a novel triple glucagon, GIP, and GLP-1 receptor agonist for glycemic control and weight loss: From discovery to clinical proof of concept. *Cell. Metab.* 34, 1234–1247.e9. doi:10.1016/j.cmet.2022.07.013
- Dahl, D., Onishi, Y., Norwood, P., Huh, R., Bray, R., Patel, H., et al. (2022). Effect of subcutaneous tirzepatide vs placebo added to titrated insulin glargine on glycemic control in patients with type 2 diabetes: The SURPASS-5 randomized clinical trial. *Jama* 327 (6), 534–545. doi:10.1001/jama.2022.0078
- Dailey, M. J., and Moran, T. H. (2013). Glucagon-like peptide 1 and appetite. *Trends Endocrinol. Metab.* 24 (2), 85–91. doi:10.1016/j.tem.2012.11.008
- DeFronzo, R. A., Ferrannini, E., Groop, L., Henry, R. R., Herman, W. H., Holst, J. J., et al. (2015). Type 2 diabetes mellitus. *Nat. Rev. Dis. Prim.* 1, 15019. doi:10.1038/nrdp.2015.19
- Del Prato, S., Kahn, S. E., Pavo, I., Weerakkody, G. J., Yang, Z., Doupis, J., et al. (2021). Tirzepatide versus insulin glargine in type 2 diabetes and increased cardiovascular risk (SURPASS-4): A randomised, open-label, parallel-group, multicentre, phase 3 trial. *Lancet* 398 (10313), 1811–1824. doi:10.1016/s0140-6736(21)02188-7
- Draznin, B., Aroda, V. R., Bakris, G., Benson, G., Brown, F. M., Freeman, R., et al. (2022). 9. Pharmacologic approaches to glycemic treatment: Standards of medical care in diabetes-2022. *Diabetes Care* 45, S125–S143. doi:10.2337/dc22-S009
- Dutta, D., Surana, V., Singla, R., Aggarwal, S., and Sharma, M. (2021). Efficacy and safety of novel twincretin tirzepatide a dual GIP and GLP-1 receptor agonist in the management of type-2 diabetes: A Cochrane meta-analysis. *Indian J. Endocrinol. Metab.* 25 (6), 475–489. doi:10.4103/ijem.ijem\_423\_21
- Ferrannini, E., and Cushman, W. C. (2012). Diabetes and hypertension: The bad companions. *Lancet* 380 (9841), 601–610. doi:10.1016/s0140-6736(12)60987-8
- Fisman, E. Z., and Tenenbaum, A. (2021). The dual glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide-1 (GLP-1) receptor agonist tirzepatide: A novel cardiometabolic therapeutic prospect. *Cardiovasc. Diabetol.* 20 (1), 225. doi:10.1186/s12933-021-01412-5
- Frias, J. P., Davies, M. J., Rosenstock, J., Pérez Manghi, F. C., Fernández Landó, L., Bergman, B. K., et al. (2021). Tirzepatide versus semaglutide once weekly in patients

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphar.2022.998816/full#supplementary-material>

with type 2 diabetes. *N. Engl. J. Med.* 385 (6), 503–515. doi:10.1056/NEJMoa2107519

Frias, J. P., Nauck, M. A., Van, J., Benson, C., Bray, R., Cui, X., et al. (2020). Efficacy and tolerability of tirzepatide, a dual glucose-dependent insulinotropic peptide and glucagon-like peptide-1 receptor agonist in patients with type 2 diabetes: A 12-week, randomized, double-blind, placebo-controlled study to evaluate different dose-escalation regimens. *Diabetes Obes. Metab.* 22 (6), 938–946. doi:10.1111/dom.13979

Frias, J. P., Nauck, M. A., Van, J., Kutner, M. E., Cui, X., Benson, C., et al. (2018). Efficacy and safety of LY3298176, a novel dual GIP and GLP-1 receptor agonist, in patients with type 2 diabetes: A randomised, placebo-controlled and active comparator-controlled phase 2 trial. *Lancet* 392 (10160), 2180–2193. doi:10.1016/s0140-6736(18)32260-8

Gerstein, H. C., Colhoun, H. M., Dagenais, G. R., Diaz, R., Lakshmanan, M., Pais, P., et al. (2019). Dulaglutide and cardiovascular outcomes in type 2 diabetes (REWIND): A double-blind, randomised placebo-controlled trial. *Lancet* 394 (10193), 121–130. doi:10.1016/s0140-6736(19)31149-3

Højberg, P. V., Vilsbøll, T., Rabøl, R., Knop, F. K., Bache, M., Krarup, T., et al. (2009). Four weeks of near-normalisation of blood glucose improves the insulin response to glucagon-like peptide-1 and glucose-dependent insulinotropic polypeptide in patients with type 2 diabetes. *Diabetologia* 52 (2), 199–207. doi:10.1007/s00125-008-1195-5

Holst, J. J., and Rosenkilde, M. M. (2020). GIP as a therapeutic target in diabetes and obesity: Insight from incretin Co-agonists. *J. Clin. Endocrinol. Metab.* 105 (8), dgaa327–2716. doi:10.1210/clinem/dgaa327

Hutton, B., Salanti, G., Caldwell, D. M., Chaimani, A., Schmid, C. H., Cameron, C., et al. (2015). The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: Checklist and explanations. *Ann. Intern. Med.* 162 (11), 777–784. doi:10.7326/m14-2385

Inagaki, N., Takeuchi, M., Oura, T., Imaoka, T., and Seino, Y. (2022). Efficacy and safety of tirzepatide monotherapy compared with dulaglutide in Japanese patients with type 2 diabetes (SURPASS J-mono): A double-blind, multicentre, randomised, phase 3 trial. *Lancet. Diabetes Endocrinol.* 10 (9), 623–633. doi:10.1016/s2213-8587(22)00188-7

Karagiannis, T., Avgerinos, I., Liakos, A., Del Prato, S., Matthews, D. R., Tsapas, A., et al. (2022). Management of type 2 diabetes with the dual GIP/GLP-1 receptor agonist tirzepatide: A systematic review and meta-analysis. *Diabetologia* 65, 1251–1261. doi:10.1007/s00125-022-05715-4

Lavie, C. J., Milani, R. V., and Ventura, H. O. (2009). Obesity and cardiovascular disease: Risk factor, paradox, and impact of weight loss. *J. Am. Coll. Cardiol.* 53 (21), 1925–1932. doi:10.1016/j.jacc.2008.12.068

Laws, A., Tao, R., Wang, S., Padhiar, A., and Goring, S. (2019). A comparison of national guidelines for network meta-analysis. *Value Health* 22 (10), 1178–1186. doi:10.1016/j.jval.2019.05.013

- Lingvay, I., Sumithran, P., Cohen, R. V., and le Roux, C. W. (2022). Obesity management as a primary treatment goal for type 2 diabetes: Time to reframe the conversation. *Lancet* 399 (10322), 394–405. doi:10.1016/s0140-6736(21)01919-x
- Ludvik, B., Giorgino, F., Jódar, E., Frias, J. P., Fernández Landó, L., Brown, K., et al. (2021). Once-weekly tirzepatide versus once-daily insulin degludec as add-on to metformin with or without SGLT2 inhibitors in patients with type 2 diabetes (SURPASS-3): A randomised, open-label, parallel-group, phase 3 trial. *Lancet* 398 (10300), 583–598. doi:10.1016/s0140-6736(21)01443-4
- Malone, J. I., and Hansen, B. C. (2019). Does obesity cause type 2 diabetes mellitus (T2DM)? Or is it the opposite? *Pediatr. Diabetes* 20 (1), 5–9. doi:10.1111/pedi.12787
- Min, T., and Bain, S. C. (2021). The role of tirzepatide, dual GIP and GLP-1 receptor agonist, in the management of type 2 diabetes: The SURPASS clinical trials. *Diabetes Ther.* 12 (1), 143–157. doi:10.1007/s13300-020-00981-0
- Moher, D., Liberati, A., Tetzlaff, J., and Altman, D. G. Prisma Group (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Ann. Intern. Med.* 151 (4), 264–269. w264. doi:10.7326/0003-4819-151-4-200908180-00135
- Nauck, M. A., and Meier, J. J. (2016). The incretin effect in healthy individuals and those with type 2 diabetes: Physiology, pathophysiology, and response to therapeutic interventions. *Lancet. Diabetes Endocrinol.* 4 (6), 525–536. doi:10.1016/s2213-8587(15)00482-9
- Nauck, M. A., Quast, D. R., Wefers, J., and Pfeiffer, A. F. H. (2021). The evolving story of incretins (GIP and GLP-1) in metabolic and cardiovascular disease: A pathophysiological update. *Diabetes Obes. Metab.* 23, 5–29. doi:10.1111/dom.14496
- Papachristou, S., Popovic, D. S., and Papanas, N. (2021). The new dual gastric inhibitory peptide/glucagon-like peptide 1 agonist tirzepatide in type 2 diabetes: Is the future bright? *Diabetes. Metab. Res. Rev.* 37 (8), e3503. doi:10.1002/dmrr.3503
- Pappachan, J. M., Fernandez, C. J., and Chacko, E. C. (2019). Diabesity and antidiabetic drugs. *Mol. Asp. Med.* 66, 3–12. doi:10.1016/j.mam.2018.10.004
- Rosenstock, J., Wysham, C., Frias, J. P., Kaneko, S., Lee, C. J., Fernández Landó, L., et al. (2021). Efficacy and safety of a novel dual GIP and GLP-1 receptor agonist tirzepatide in patients with type 2 diabetes (SURPASS-1): A double-blind, randomised, phase 3 trial. *Lancet* 398 (10295), 143–155. doi:10.1016/s0140-6736(21)01324-6
- Sattar, N., McGuire, D. K., Pavo, I., Weerakkody, G. J., Nishiyama, H., Wiese, R. J., et al. (2022). Tirzepatide cardiovascular event risk assessment: A pre-specified meta-analysis. *Nat. Med.* 28 (3), 591–598. doi:10.1038/s41591-022-01707-4
- Sterne, J. A. C., Savović, J., Page, M. J., Elbers, R. G., Blencowe, N. S., Boutron, I., et al. (2019). RoB 2: A revised tool for assessing risk of bias in randomised trials. *Bmj* 366, 14898. doi:10.1136/bmj.14898
- Sun, H., Saeedi, P., Karuranga, S., Pinkepank, M., Ogurtsova, K., Duncan, B. B., et al. (2021). IDF diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res. Clin. Pract.* 183, 109119. doi:10.1016/j.diabres.2021.109119
- Taskinen, M. R., and Borén, J. (2015). New insights into the pathophysiology of dyslipidemia in type 2 diabetes. *Atherosclerosis* 239 (2), 483–495. doi:10.1016/j.atherosclerosis.2015.01.039
- Thomas, M. K., Nikooienejad, A., Bray, R., Cui, X., Wilson, J., Duffin, K., et al. (2021). Dual GIP and GLP-1 receptor agonist tirzepatide improves beta-cell function and insulin sensitivity in type 2 diabetes. *J. Clin. Endocrinol. Metab.* 106 (2), 388–396. doi:10.1210/clinem/dgaa863