



# Hydrodynamic Approach to Electronic Transport in Graphene: Energy Relaxation

B. N. Narozhny<sup>1,2\*</sup> and I. V. Gornyi<sup>3,4</sup>

<sup>1</sup>Institute for Theoretical Condensed Matter Physics, Karlsruhe Institute of Technology, Karlsruhe, Germany, <sup>2</sup>National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), Moscow, Russia, <sup>3</sup>Institute for Quantum Materials and Technologies, Karlsruhe Institute of Technology, Karlsruhe, Germany, <sup>4</sup>Ioffe Institute, St. Petersburg, Russia

In nearly compensated graphene, disorder-assisted electron-phonon scattering or “supercollisions” are responsible for both quasiparticle recombination and energy relaxation. Within the hydrodynamic approach, these processes contribute weak decay terms to the continuity equations at local equilibrium, i.e., at the level of “ideal” hydrodynamics. Here we report the derivation of the decay term due to weak violation of energy conservation. Such terms have to be considered on equal footing with the well-known recombination terms due to nonconservation of the number of particles in each band. At high enough temperatures in the “hydrodynamic regime” supercollisions dominate both types of the decay terms (as compared to the leading-order electron-phonon interaction). We also discuss the contribution of supercollisions to the heat transfer equation (generalizing the continuity equation for the energy density in viscous hydrodynamics).

**Keywords:** hydrodynamics, graphene, electronic transport, energy relaxation, supercollisions, quasiparticle recombination

## 1 INTRODUCTION

Electronic hydrodynamics is quickly growing into a mature field of condensed matter physics [1–3]. Similarly to the usual hydrodynamics [4, 5]; this approach offers a universal, long-wavelength description of collective flows in interacting many-electron systems. As a macroscopic theory of strongly interacting systems, hydrodynamics should appear to be extremely attractive for condensed matter theorists dealing with problems where strong correlations invalidate simple theoretical approaches. However, electrons in solids exist in the environment created by a crystal lattice and typically experience collisions with lattice imperfections (or “disorder”) and lattice vibrations (phonons). The former typically dominate electronic transport at low temperatures, while at high temperatures the electron-phonon interaction takes over. In both cases the electron motion is diffusive (unless the sample size is smaller than the mean free path in which case the motion is ballistic) since in both types of scattering the electronic momentum is not conserved. On the other hand, if a material would exist where the momentum-conserving electron-electron interaction would dominate at least in some non-negligible temperature range, then one could be justified in neglecting the momentum non-conserving processes and applying the hydrodynamic theory. In recent years, several extremely pure materials became available with graphene being the most studied [1, 3].

In nearly neutral (or compensated) graphene the electron system is non-degenerate (at least at relatively high temperatures where the hydrodynamic approach is justified) with both the conductance and valence bands contributing on equal footing. Although the electron system is

## OPEN ACCESS

### Edited by:

Alessandro Principi,  
The University of Manchester,  
United Kingdom

### Reviewed by:

Ke-Qiu Chen,  
Hunan University, China  
Giovanni Vignale,  
University of Missouri, United States

### \*Correspondence:

B. N. Narozhny  
boris.narozhny@kit.edu

### Specialty section:

This article was submitted to  
Condensed Matter Physics,  
a section of the journal  
Frontiers in Physics

**Received:** 11 December 2020

**Accepted:** 11 February 2021

**Published:** 23 April 2021

### Citation:

Narozhny BN and Gornyi IV (2021)  
Hydrodynamic Approach to Electronic  
Transport in Graphene:  
Energy Relaxation.  
Front. Phys. 9:640649.  
doi: 10.3389/fphy.2021.640649

not Lorenz-invariant, the linearity of the Dirac spectrum plays an important role. Firstly, the Auger processes are kinematically suppressed leading to the near-conservation of the number of particles in each band [2, 3, 6, 7]. Secondly, the so-called collinear scattering singularity [7–14] allows for a non-perturbative solution to the kinetic (Boltzmann) equation focusing on the three hydrodynamic modes [13, 15, 16]. As a result, one can determine the general form of the hydrodynamic equations and evaluate the kinetic coefficients [16–18]. To be of any practical value, the latter calculation has to be combined with the renormalization group approach [19] since the effective coupling constant in real graphene (either encapsulated or put on a dielectric substrate) is not too small,  $\alpha_g \approx 0.2 - 0.3$  [20, 21].

Next to the conservation laws, the main assumption of the hydrodynamic approach is local equilibrium [4, 22] established by means of interparticle collisions. Neglecting all dissipative processes, this allows (together with the conservation laws) for a phenomenological derivation of hydrodynamic equations [4, 5] that can be further supported by the kinetic theory, where the local equilibrium distribution function nullifies the collision integral in the Boltzmann equation [22]. The resulting ideal hydrodynamics is described by the Euler equation and the continuity equations. This is where the electronic fluid in graphene differs from conventional fluids (both Galilean- and Lorentz-invariant): as in any solid, conservation laws in graphene are only approximate, leaving the collision integrals describing scattering processes other than the electron-electron interaction to be nonzero even in local equilibrium. This leads to the appearance of weak decay terms in the continuity equations.

Two such terms have already been discussed in literature. Firstly, even if the electron-electron interaction is the dominant scattering process in the system, no solid is absolutely pure. Consequently, even ultra-pure graphene samples possess some degree of weak disorder. Disorder scattering violates momentum conservation and hence a weak decay term must appear in the generalized Euler equation [2, 3, 15, 16]. Secondly, conservation of the number of particles in each band is violated by a number of processes (e.g., the Auger and three-particle scattering). Although commonly assumed to be weak, they are manifested in the decay – or recombination – term in the corresponding continuity equation. This was first established in [6] in the context of thermoelectric phenomena (for the most recent discussion see [23]). Later, quasiparticle recombination was shown to lead to linear magnetoresistance in compensated semimetals [24–27] as well as to giant magnetodrag [28, 29].

In this paper, we report the derivation of the third weak decay term in the hydrodynamic theory in graphene due to weak violation of energy conservation. Indeed, the electron-phonon interaction may lead not only to the loss of electronic momentum (responsible for electrical resistivity in most metals at high temperatures), but also to the loss of energy. Although subdominant in the hydrodynamic regime, the electron-phonon interaction should be taken into account as one of the dissipative processes. In graphene, the linearity of the Dirac spectrum once again plays an important role: as the speed of sound is much smaller than the electron velocity  $v_g$ , leading-order scattering on acoustic phonons is kinematically suppressed.

Consequently, scattering on the optical branch is usually considered [30, 31]. In contrast, we argue that there is another process, the disorder-assisted electron-phonon scattering [32] or “supercollisions” [33–36], that is responsible for both quasiparticle recombination and energy relaxation. In the high-temperature hydrodynamic regime, supercollisions are expected to dominate both decay contributions [32]. Moreover, this process contributes weak decay terms to the continuity equations already at local equilibrium, i.e., at the level of “ideal” hydrodynamics.

## 2 RESULTS

Our arguments are based on the kinetic theory approach to electronic transport. In the spirit of [19], we assume the possibility of deriving the hydrodynamic equations from the kinetic equation in the weak coupling limit [16],  $\alpha_g \ll 1$ , with the subsequent renormalization of the kinetic coefficients to the realistic parameter regime [17]. Under these assumptions, we start with the kinetic equation

$$\mathcal{L}f_{\lambda k} = \text{St}_{ee}[f_{\lambda k}] + \text{St}_R[f_{\lambda k}] + \text{St}_{\text{dis}}[f_{\lambda k}], \tag{1a}$$

with the Liouville’s operator (in the left-hand side)

$$\mathcal{L} = \partial_t + \mathbf{v} \cdot \nabla_r + \left( e\mathbf{E} + \frac{e}{c} \mathbf{v} \times \mathbf{B} \right) \cdot \nabla_k, \tag{1b}$$

and the collision integrals describing the electron-electron interaction ( $\text{St}_{ee}$ ), disorder scattering ( $\text{St}_{\text{dis}}$ ), and quasiparticle recombination ( $\text{St}_R$ ), where in this paper we focus on “supercollisions.” We employ the following notations for the Dirac spectrum (the chirality  $\lambda = \pm 1$  distinguishes the conduction and valence bands)

$$\varepsilon_{\lambda k} = \lambda v_g k, \tag{2a}$$

and velocities

$$\mathbf{v}_{\lambda k} = \lambda v_g \frac{\mathbf{k}}{k}, \quad \mathbf{k} = \frac{\lambda k}{v_g} \mathbf{v}_{\lambda k} = \frac{\varepsilon_{\lambda k} \mathbf{v}_{\lambda k}}{v_g^2}. \tag{2b}$$

Hydrodynamics is the macroscopic manifestation of the conservation of energy, momentum, and the number of particles. In a two-band system, the latter comprises excitations in both bands. In the conduction band these are electron-like quasiparticles with the number density ( $N = 4$  reflects spin and valley degeneracy in graphene)

$$n_+ = N \int \frac{d^2 k}{(2\pi)^2} f_{+,k}, \tag{3a}$$

while in the valence band the quasiparticles are hole-like

$$n_- = N \int \frac{d^2 k}{(2\pi)^2} (1 - f_{-,k}), \tag{3b}$$

with the total “charge” (or “carrier”) density being

$$n = n_+ - n_-. \tag{3c}$$

Assuming the numbers of particles in the conduction and valence bands are conserved independently, we can also define the total quasiparticle (or “imbalance” [6]) density

$$n_I = n_+ + n_- \tag{3d}$$

Global charge conservation (or gauge symmetry) can be expressed in terms of the usual continuity equation. This can be obtained from **Eq. 1** by performing a summation over all quasiparticle states upon which all three collision integrals vanish [22].

$$\begin{aligned} N \int \frac{d^2k}{(2\pi)^2} St_{ee}[f_{\lambda k}] &= N \int \frac{d^2k}{(2\pi)^2} St_R[f_{\lambda k}] = N \int \frac{d^2k}{(2\pi)^2} St_{dis}[f_{\lambda k}] \\ &= 0. \end{aligned} \tag{4a}$$

As a result, the continuity equation has the usual form

$$\partial_t n + \nabla_r \cdot \mathbf{j} = 0, \tag{4b}$$

where the corresponding current is defined as

$$\mathbf{j} = \mathbf{j}_+ - \mathbf{j}_- = N \int \frac{d^2k}{(2\pi)^2} [\mathbf{v}_{+,k} f_{+,k} - \mathbf{v}_{-,k} (1 - f_{-,k})], \tag{5}$$

The rest of the conservation laws in graphene are approximate as manifested in the collision integrals not vanishing upon corresponding summations. The continuity equation expressing momentum conservation (i.e., the Euler equation) is obtained by multiplying the kinetic equation by the quasiparticle momentum  $\mathbf{k}$  and summing over all states. Since the electron-electron interaction conserves momentum, the corresponding collision integral vanishes

$$N \int \frac{d^2k}{(2\pi)^2} \mathbf{k} St_{ee}[f_{\lambda k}] = 0. \tag{6}$$

Weak disorder scattering is typically described within the simplest  $\tau$ -approximation [22].

$$N \int \frac{d^2k}{(2\pi)^2} \mathbf{k} St_{dis}[f_{\lambda k}] = \frac{\mathbf{n}_k}{\tau_{dis}}, \tag{7}$$

where the momentum density is defined as

$$\mathbf{n}_k = N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \mathbf{k} f_{\lambda k} = \mathbf{v}_g^{-2} \mathbf{j}_E. \tag{8}$$

The last equality represents the fact that in graphene the momentum density is proportional to the energy density (due to the properties of the Dirac spectrum **Eq. 2**).

Supercollisions contributing to the recombination collision integral also violate momentum conservation, however, in comparison to the above weak disorder scattering, this is a second-order process. Moreover, the disorder mean free time  $\tau_{dis}$  is typically determined from experimental data (see e.g., [21]) and hence can be assumed to include the contribution of supercollisions as well.

The remaining two continuity equations – energy and quasiparticle imbalance – are unaffected by the electron-electron interaction and weak disorder scattering. Indeed, the

electron-electron interaction conserves energy and – neglecting the Auger processes – particle number in each band:

$$N \int \frac{d^2k}{(2\pi)^2} \epsilon_{\lambda k} St_{ee}[f_{\lambda k}] = N \int \frac{d^2k}{(2\pi)^2} \lambda St_{ee}[f_{\lambda k}] = 0. \tag{9}$$

Same applies to the (Elastic) disorder scattering

$$N \int \frac{d^2k}{(2\pi)^2} \epsilon_{\lambda k} St_{dis}[f_{\lambda k}] = N \int \frac{d^2k}{(2\pi)^2} \lambda St_{dis}[f_{\lambda k}] = 0. \tag{10}$$

However, supercollisions do not conserve both quantities and hence lead to weak decay terms in the two continuity equations. Contribution of the recombination collision integral to the continuity equation for the quasiparticle imbalance (for derivation see **Section 4**) is given by

$$N \sum_k \lambda St_R[f_{\lambda k}] \approx -\mu_I n_{I,0} \lambda_Q \approx -\frac{n_I - n_{I,0}}{\tau_R}. \tag{11}$$

Here  $n_I$  is the imbalance density (3d) in local equilibrium, while  $n_{I,0}$  is the same quantity evaluated with the Fermi distribution function (17), i.e. for  $\mu_I = 0$  and  $\mathbf{u} = 0$ . The first equality in **Eq. 11** coincides with the expression used in [6] and serves as the definition of the dimensionless coefficient  $\lambda_Q$ , while the second (valid to the leading order) was suggested in [16, 24] and provides the definition of the “recombination time”  $\tau_R$  (see also [23]). The two expressions are equivalent since  $n_I - n_{I,0} \propto \mu_I$ . The resulting continuity equation for the quasiparticle imbalance has the form

$$\partial_t n_I + \nabla_r \cdot \mathbf{j}_I = -\frac{n_I - n_{I,0}}{\tau_R}, \quad \mathbf{j}_I = \mathbf{j}_+ + \mathbf{j}_-. \tag{12}$$

The same scattering process contributes a weak decay term to the continuity equation for the energy density. Defining the decay coefficients similarly to **Eq. 11**, we may present the result in the form

$$N \sum_k \epsilon_{\lambda k} St_R[f_{\lambda k}] = -\mu_I n_{E,0} \lambda_{QE} \approx -\frac{n_E - n_{E,0}}{\tau_{RE}}. \tag{13}$$

Here the equivalence of the two forms of the decay term stems from the fact that  $n_E - n_{E,0} \propto \mu_I$  assuming the electrons and holes are characterized by the same temperature.

Finally, once the dissipative processes due to electron-electron interaction are taken into account, one usually replaces the continuity equation for the energy density by the equivalent equation for the entropy density, the so-called “heat transfer equation” [4] (for derivation see **Section 4**; here  $\delta \mathbf{j}$  and  $\delta \mathbf{j}_I$  are the dissipative corrections to the electric and imbalance currents, respectively)

$$\begin{aligned} T \left[ \frac{\partial s}{\partial t} + \nabla_r \cdot \left( s \mathbf{u} - \delta \mathbf{j} \frac{\mu}{T} - \delta \mathbf{j}_I \frac{\mu_I}{T} \right) \right] &= \\ = \delta \mathbf{j} \cdot \left[ e \mathbf{E} + \frac{e}{c} \mathbf{u} \times \mathbf{B} - T \nabla \frac{\mu}{T} \right] - T \delta \mathbf{j}_I \cdot \nabla \frac{\mu_I}{T} + \frac{\eta}{2} (\nabla_{\alpha} u_{\beta} + \nabla_{\beta} u_{\alpha} - \delta_{\alpha\beta} \nabla \cdot \mathbf{u})^2 \\ - \frac{n_E - n_{E,0}}{\tau_{RE}} + \mu_I \frac{n_I - n_{I,0}}{\tau_R} + \frac{\mathbf{u} \cdot \mathbf{n}_k}{\tau_{dis}}. \end{aligned} \tag{14}$$

The obtained heat transfer equation is the final hydrodynamic equation in graphene. Together with the generalized Navier-Stokes equation derived in [16] and the continuity equation **Eqs. 4b** and **12**, the heat transfer equation forms a complete set of macroscopic equations describing electronic transport in graphene in hydrodynamic regime.

### 3 DISCUSSION

Supercollisions are not the only scattering process contributing to both quasiparticle recombination and energy relaxation. Clearly, direct (not impurity-assisted) electron-phonon interaction contributes to energy relaxation as well as to quasiparticle recombination (in the case of intervalley scattering) [6, 13, 14, 29, 32]. In addition, optical phonons may also contribute [30, 31], although within the hydrodynamic approach these contributions were discussed only at the level of dissipative (viscous) hydrodynamics [31]. The contribution of the direct [37, 38] and impurity assisted electron-phonon scattering to energy relaxation was compared in [32], where it was argued that at high enough temperatures,  $T \gtrsim T_{BG}$  (where  $T_{BG}$  is the Bloch-Grüneisen temperature) the supercollisions do dominate. In the degenerate regime, where  $T_{BG} \propto \sqrt{n}$ , Ref. [32] estimates  $T_{BG}$  as “few tens of Kelvin”. At charge neutrality, we estimate  $T_{BG} = (s/v_g)T \ll T$  (where  $s$  is the speed of sound), such that supercollisions always dominate over direct electron-phonon coupling. Moreover, taking into account the additional scattering processes will not change the form of the decay terms (11) and (13), but rather change the numerical values of the parameters  $\lambda_Q$  and  $\lambda_{QE}$ , which may have to be considered phenomenological while interpreting experimental data [29].

At charge neutrality and in the hydrodynamic regime, the coefficients  $\lambda_Q$  and  $\lambda_{QE}$  are of the same order of magnitude (both are dominated by the same supercollisions), but quantitatively different. Indeed, the continuity equation for the energy density should contain one more contribution of the similar form. “Quasiparticle recombination” typically refers to scattering between the quasiparticle states in different bands only. This is the only type of supercollisions contributing to  $\lambda_Q$ . Similar supercollisions may also take place within a single band [32]. While this process does not change the number of particles in the band, it does describe the energy loss as the electron may scatter from the higher energy state into the lower energy state (and losing its momentum to the impurity along the way). Consequently, this additional process does contribute to energy relaxation. Given that the form of the corresponding collision integral is very similar to **Eq. 15** – one only has to change to band indices to be the same – the algebra remains the same and thus we can treat **Eq. 13** as the final result that takes this additional intraband supercollisions into account making the numerical values of  $\lambda_Q$  and  $\lambda_{QE}$  substantially different – we do not expect any accidental cancellation or smallness should the difference  $\lambda_Q - \lambda_{QE}$  appear in a particular solution of hydrodynamic equations. At the same time, at low temperatures – i.e., below the hydrodynamic range – we expect the coefficients  $\lambda_Q$  and  $\lambda_{QE}$  to be parametrically

different: energy relaxation is now dominated by the direct electron-phonon interaction [32]; while the recombination is still governed by supercollisions (together with the phonon-induced intervalley scattering).

The order of magnitude of  $\tau_R$  could be estimated based on the calculations of [32]. Adapting the latter to charge neutrality, we find  $\tau_R^{-1} \sim D^2 T^2 / (\rho s^2 v_g^2)$  (where  $D \approx 20$  eV is the deformation potential [37, 38] and  $\rho$  is the mass density per unit area) yielding the corresponding length scale  $\ell_R^{-1} \approx 10 \mu\text{m}$  at the top of the hydrodynamic temperature range,  $T \approx 250$  K. This should be further compared to the contribution of three-particle collisions [3, 31],  $\tau_3^{-1} \sim \alpha_g^4 T$ . Assuming the common sample design where graphene is encapsulated in hexagonal boron nitride (with the dielectric constant  $\epsilon \approx 4$ ), the effective coupling constant (taking into account renormalizations) is  $\alpha_g \approx 0.3 - 0.4$  leading to the similar estimate at high temperatures. On the other hand, at the low end of the hydrodynamic range [39],  $T \approx 50$  K, the contribution of the three-body collisions should dominate (accounting for the empirical value  $\ell_R \approx 1.2 \mu\text{m}$  reported in [29], however preserving the functional form of the weak decay terms in the continuity equations).

The obtained **Eq. 2** should be compared to the corresponding equations in [3, 6, 31], where energy relaxation due to supercollisions was not taken into account. All other terms are present in all four equations with the following exceptions. The Eq. 54 of [3] is written in the relativistic notation omitting the imbalance mode, quasiparticle recombination, and disorder scattering, all of which are discussed separately elsewhere in [3]. Ref. [6] was the first to focus on the imbalance mode with the Eq. 2.6 containing all the terms of **Eq. 2** except for the viscous term (and energy relaxation). Finally, the Eq. 1c of [31] contains all of the terms in **Eq. 2** except for energy relaxation and in addition contains a term describing energy relaxation due to electrons scattering on the optical phonon branch that is neglected here (generalization of the resulting theory is straightforward).

To summarize, we have considered supercollisions as a mechanism of quasiparticle recombination and energy relaxation in graphene and derived the corresponding decay terms in the hydrodynamic continuity equations. Since the same scattering mechanism is responsible for both effects, one has to take into account energy relaxation while considering quasiparticle recombination. The latter is an indispensable feature of electronic hydrodynamics in graphene in constrained geometries, where homogenous solutions violate the boundary conditions [24].

## 4 MATERIALS AND METHODS

### 4.1 Collision Integral due to Supercollisions

An electron in the upper (conductance) band may scatter into an empty state in the lower (valence) band – effectively recombining with a hole – emitting a phonon (which carries away the energy) and losing its momentum to the impurity. Within the standard approach to the electron-phonon interaction, this process is described by the collision integral

$$\text{St}_R[f_{+k_2}] = 2\pi \sum_{k_1, q} W_q \delta(\varepsilon_{+k_2} - \varepsilon_{-k_1} - \omega_q) [f_{-k_1}(1 - f_{+k_2})n_q - f_{+k_2}(1 - f_{-k_1})(1 + n_q)], \quad (15a)$$

where  $n_q$  is the phonon (Plank's) distribution function (the phonons are assumed to be at equilibrium and play the role of a "bath"),  $W_q$  is the effective scattering rate that includes the Dirac factors and is averaged over the angles [32].

Similarly, an electron in the lower band may absorb a phonon and scatter into the upper band – effectively creating an electron-hole pair – while still losing its momentum to the impurity

$$\text{St}_R[f_{-k_2}] = 2\pi \sum_{k_1, q} W_q \delta(\varepsilon_{+k_1} - \varepsilon_{-k_2} - \omega_q) [f_{+k_1}(1 - f_{-k_2})(1 + n_q) - f_{-k_2}(1 - f_{+k_1})n_q]. \quad (15b)$$

The collision Integral (14) conserves the total charge

$$N \sum_k \text{St}_R[f_{\lambda k}] = N \sum_k (\text{St}_R[f_{+k_2}] + \text{St}_R[f_{-k_2}]) = 0, \quad (15c)$$

(see Eq. 4) and vanishes in global equilibrium

$$\text{St}_R[f^{(0)}] = 0, \quad (15d)$$

where the quasiparticle distribution is described by the Fermi function. This should be contrasted with *local equilibrium* described by

$$f_{\lambda k}^{(le)} = \left\{ 1 + \exp \left[ \frac{\varepsilon_{\lambda k} - \mu_{\lambda}(\mathbf{r}) - \mathbf{u}(\mathbf{r}) \cdot \mathbf{k}}{T(\mathbf{r})} \right] \right\}^{-1}, \quad (16)$$

where  $\mu_{\lambda}(\mathbf{r})$  is the local chemical potential and  $\mathbf{u}(\mathbf{r})$  is the hydrodynamic (or "drift") velocity. The local equilibrium distribution function (16) allows for independent chemical potentials in the two bands, which can be expressed in terms of the "thermodynamic" and "imbalance" chemical potentials

$$\mu_{\lambda} = \mu + \lambda \mu_I. \quad (17)$$

In global equilibrium

$$f^{(0)} = f_{\lambda k}^{(le)}(\mu_I = 0, \mathbf{u} = 0). \quad (18)$$

Now we show, that in local equilibrium, i.e. for nonzero  $\mu_I$ , the recombination collision integral remains finite (unlike Eq. 14d). As a scalar quantity, the collision integral (15) cannot depend on the hydrodynamic velocity  $\mathbf{u}$  in the first (linear) order. Consequently, to the leading order the integrated collision integral yielding the decay terms in the continuity equations is proportional to  $\mu_I$ .

## 4.2 Derivation of the Weak Decay Terms in Continuity Equations

To the leading order, we can describe the difference between the local equilibrium distribution function  $f_{\lambda k}^{(le)}$  and the Fermi function  $f^{(0)}$  similarly to the leading non-equilibrium correction in the standard derivation of hydrodynamics [22].

$$\delta f = f_{\lambda k} - f^{(0)} = -T \frac{\partial f^{(0)}}{\partial \varepsilon} h = f^{(0)}(1 - f^{(0)})h. \quad (19)$$

Now we re-write the collision integral (14) with the help of the relations

$$f_1(1 - f_2)(1 + n_q) - f_2(1 - f_1)n_q = (1 - f_1)(1 - f_2)(1 + n_q) \left[ \frac{f_1}{1 - f_1} \frac{n_q}{1 + n_q} - \frac{f_2}{1 - f_2} \right],$$

and

$$f_2^{(0)}(1 - f_1^{(0)})(1 + n_q) = -\frac{\partial n_q}{\partial \omega} (f_2^{(0)} - f_1^{(0)}),$$

and find (to the leading order in  $h$ )

$$\text{St}_R[f_{+k_2}] = -2\pi \sum_{k_1, q} W_q \delta(\varepsilon_{+k_2} - \varepsilon_{-k_1} - \omega_q) \frac{\partial n_q}{\partial \omega} (f_{+k_2}^{(0)} - f_{-k_1}^{(0)})(h_{-k_1} - h_{+k_2}), \quad (20a)$$

$$\text{St}_R[f_{-k_2}] = 2\pi \sum_{k_1, q} W_q \delta(\varepsilon_{-k_2} - \varepsilon_{+k_1} + \omega_q) \frac{\partial n_q}{\partial \omega} (f_{+k_1}^{(0)} - f_{-k_2}^{(0)})(h_{-k_2} - h_{+k_1}). \quad (20b)$$

Consider now the contribution of the recombination collision integral to the continuity equation for the quasiparticle imbalance

$$\begin{aligned} N \sum_k \lambda \text{St}_R[f_{\lambda k}] &= N \sum_{k_2} (\text{St}_R[f_{+k_2}] - \text{St}_R[f_{-k_2}]) = \\ &= -4\pi N \sum_{k_1, k_2, q} W_q \delta(\varepsilon_{+k_2} - \varepsilon_{-k_1} - \omega_q) \frac{\partial n_q}{\partial \omega} (f_{+k_2}^{(0)} - f_{-k_1}^{(0)})(h_{-k_1} - h_{+k_2}). \end{aligned} \quad (21)$$

To the leading order, the deviation  $h_{\lambda k}$  is proportional to  $\mu_I$

$$h_{\lambda k} \approx \frac{\lambda \mu_I}{T}. \quad (22)$$

The remaining integral has dimensions of particle density divided by time and therefore the result can be written in the form Eq. 11. The same scattering process contributes a weak decay term to the continuity equation for the energy density. Indeed, multiplying the collision integral Eq. 15 by the quasiparticle energy and summing over all states, we find after similar algebra

$$\begin{aligned} N \sum_k \varepsilon_{\lambda k} \text{St}_R[f_{\lambda k}] &= N \sum_{k_2} (\varepsilon_{+k_2} \text{St}_R[f_{+k_2}] + \varepsilon_{-k_2} \text{St}_R[f_{-k_2}]) \sum_k = \\ &= -2\pi N \sum_{k_2, q} W_q \delta(\varepsilon_{+k_2} - \varepsilon_{-k_1} - \omega_q) \frac{\partial n_q^{(0)}}{\partial \omega} \omega_q (f_{+k_2}^{(0)} - f_{-k_1}^{(0)})(h_{-k_1} - h_{+k_2}). \end{aligned}$$

Defining the decay coefficients similarly to Eq. 11 above, we may present the result in the form Eq. 13.

## 4.3 Derivation of the Heat Transfer Equation

The entropy density of a system of fermions is defined in terms of the distribution function as



$$s = -N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} [f_{\lambda k} \ln f_{\lambda k} + (1 - f_{\lambda k}) \ln(1 - f_{\lambda k})]. \quad (23)$$

Treating this integral as

$$s = N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \mathcal{S}[f_{\lambda k}],$$

any derivative of  $s$  can be represented in the form

$$\frac{\partial s}{\partial z} = N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} \frac{\partial f_{\lambda k}}{\partial z}.$$

Consider now each term of the kinetic equation multiplied by the derivative  $\partial \mathcal{S}[f_{\lambda k}]/\partial f_{\lambda k}$  and summed over all states. Using the above relation with  $z \rightarrow t$ , one finds for the time derivative term

$$N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} \frac{\partial f_{\lambda k}}{\partial t} = \frac{\partial s}{\partial t}.$$

The gradient term yields similarly

$$N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} \mathbf{v}_{\lambda k} \cdot \nabla_r f_{\lambda k} = \nabla_r \cdot N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \mathbf{v}_{\lambda k} \mathcal{S}[f_{\lambda k}] = \nabla_r \cdot \mathbf{j}_S,$$

where the quantity

$$\mathbf{j}_S = N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \mathbf{v}_{\lambda k} \mathcal{S}[f_{\lambda k}], \quad (24)$$

can be interpreted as the entropy current. The electric field term vanishes as the total derivative

$$e\mathbf{E} \cdot N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} \nabla_k f_{\lambda k} = e\mathbf{E} \cdot N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \nabla_k \mathcal{S}[f_{\lambda k}] = 0,$$

while the Lorentz term vanishes for rotationally invariant systems upon integrating by parts [justified by the fact that  $\mathcal{S}(k \rightarrow \infty) \rightarrow 0$ ]

$$\begin{aligned} & \frac{e}{c} N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} [\mathbf{v}_{\lambda k} \times \mathbf{B}] \cdot \nabla_k f_{\lambda k} = \\ & = \frac{e}{c} N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} [\mathbf{v}_{\lambda k} \times \mathbf{B}] \cdot \nabla_k \mathcal{S}[f_{\lambda k}] \\ & = -\frac{e}{c} N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \mathcal{S}[f_{\lambda k}] \nabla_k \cdot [\mathbf{v}_{\lambda k} \times \mathbf{B}] = 0. \end{aligned}$$

The last equality follows from

$$\frac{\partial v_{\lambda k}^{\alpha}}{\partial k^{\beta}} = \frac{v_g}{\lambda k} \left( \delta_{\alpha\beta} - \frac{k^{\alpha} k^{\beta}}{k^2} \right).$$

Similar approach was used in [16] to derive the continuity equations (as outlined above). Combining all four terms, we conclude that integration with the factor  $\partial \mathcal{S}[f_{\lambda k}]/\partial f_{\lambda k}$  turns the left-hand side of the kinetic equation to the familiar form

$$N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} \mathcal{L}f_{\lambda k} = \frac{\partial s}{\partial t} + \nabla_r \cdot \mathbf{j}_S. \quad (25)$$

**Equation 25** is valid for an arbitrary distribution function.

Denoting the integral of the right-hand side of the kinetic equation by

$$\mathcal{I} = N \sum_{\lambda} \int \frac{d^2k}{(2\pi)^2} \frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} (\text{St}_{ee}[f] + \text{St}_R[f] + \text{St}_{dis}[f]), \quad (26)$$

we arrive at the ‘‘continuity equation for the entropy’’

$$\frac{\partial s}{\partial t} + \nabla_r \cdot \mathbf{j}_S = \mathcal{I}. \quad (27)$$

In the usual hydrodynamics [4] the only contribution to the collision integral is given by particle-particle scattering, i.e. the processes assumed to be responsible for establishing local equilibrium such that at  $\mathcal{I} = 0$  the ideal (Euler) hydrodynamic is isentropic. In the present case, local equilibrium is assumed to be achieved by means of the electron-electron interaction. Evaluating the derivative of  $\mathcal{S}$  explicitly, we find

$$\frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} = -\ln \frac{f_{\lambda k}}{1 - f_{\lambda k}} = \ln \left[ \frac{1}{f_{\lambda k}} - 1 \right].$$

For the Local Equilibrium Distribution Function

$$\frac{\partial \mathcal{S}[f_{\lambda k}]}{\partial f_{\lambda k}} = \frac{\epsilon_{\lambda k} - \mu_{\lambda} - \mathbf{u} \cdot \mathbf{k}}{T}.$$

Substituting this expression into **Eq. 26**, we find that the remaining integration is very similar to the above derivation of the continuity equations. The integral with the quasiparticle energy yields exactly the above **Eq. 13**. The integral with  $\lambda \mu_I$  yields **Eq. 11** multiplied by  $\mu_I$ . Finally, the term  $\mathbf{u} \cdot \mathbf{k}$  yields **Eq. 7** multiplied by the hydrodynamic velocity. The integral of this term with the recombination collision integral is assumed to be included into the definition of the mean free time, see the corresponding discussion above. As a result, we arrive at the following form of the integrated collision integral

$$\mathcal{I} = -\frac{1}{T} \frac{n_E - n_{E,0}}{\tau_{RE}} + \frac{\mu_I}{T} \frac{n_I - n_{I,0}}{\tau_R} + \frac{\mathbf{u} \cdot \mathbf{n}_k}{T \tau_{dis}}.$$

The decay terms in **Eq. 14** appear already at local equilibrium. To complete the heat transfer equation one has to take into account disipation. In graphene, this is most conveniently done by considering the classical limit of relativistic hydrodynamics since the form of dissipative corrections is determined by the symmetries of the quasiparticle spectrum. The result has been already reported in literature, therefore we combine the dissipative corrections with **Eq. 14** and write the heat transfer equation in graphene in the form **Eq. 14**.

### 4.3.1 Permission to Reuse and Copyright

Figures, tables, and images will be published under a Creative Commons CC-BY licence and permission must be obtained for

use of copyrighted material from other sources (including re-published/adapted/modified/partial figures and images from the internet). It is the responsibility of the authors to acquire the licenses, to follow any citation instructions requested by third-party rights holders, and cover any supplementary charges.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding author.

## AUTHOR CONTRIBUTIONS

Both authors have equally contributed to the reported research.

## REFERENCES

- Polini M, Geim AK. Viscous electron fluids. *Phys Today* (2020) 73:28–34. doi:10.1063/PT.3.4497
- Narozhny BN, Gornyi IV, Mirlin AD, Schmalian J. Hydrodynamic approach to electronic transport in graphene. *Ann Phys* (2017) 529:1700043. doi:10.1002/andp.201700043
- Lucas A, Fong KC. Hydrodynamics of electrons in graphene. *J Phys Condens Matter* (2018) 30:053001. doi:10.1088/1361-648x/aaa274
- Landau LD, Lifshitz EM. *Fluid mechanics*. London, UK: Pergamon Press (1959).
- Chaikin PM, Lubensky TC. *Principles of condensed matter physics*. Cambridge, UK: Cambridge University Press (1995).
- Foster MS, Aleiner IL. Slow imbalance relaxation and thermoelectric transport in graphene. *Phys Rev B* (2009) 79:085415. doi:10.1103/PhysRevB.79.085415
- Tomadin A, Brida D, Cerullo G, Ferrari AC, Polini M. Nonequilibrium dynamics of photoexcited electrons in graphene: collinear scattering, Auger processes, and the impact of screening. *Phys Rev B* (2013) 88:035430. doi:10.1103/PhysRevB.88.035430
- Müller M, Schmalian J, Fritz L. Graphene: a nearly perfect fluid. *Phys Rev Lett* (2009) 103:025301. doi:10.1103/PhysRevLett.103.025301
- Fritz L, Schmalian J, Müller M, Sachdev S. Quantum critical transport in clean graphene. *Phys Rev B* (2008) 78:085416. doi:10.1103/PhysRevB.78.085416
- Kashuba AB. Conductivity of defectless graphene. *Phys Rev B* (2008) 78:085415. doi:10.1103/PhysRevB.78.085415
- Müller M, Sachdev S. Collective cyclotron motion of the relativistic plasma in graphene. *Phys Rev B* (2008) 78:115419. doi:10.1103/PhysRevB.78.115419
- Schütt M, Ostrovsky PM, Gornyi IV, Mirlin AD. Coulomb interaction in graphene: relaxation rates and transport. *Phys Rev B* (2011) 83:155441. doi:10.1103/PhysRevB.83.155441
- Narozhny BN, Gornyi IV, Titov M, Schütt M, Mirlin AD. Hydrodynamics in graphene: linear-response transport. *Phys Rev B* (2015) 91:035414. doi:10.1103/PhysRevB.91.035414
- Schütt M, Ostrovsky PM, Titov M, Gornyi IV, Narozhny BN, Mirlin AD. Coulomb drag in graphene near the Dirac point. *Phys Rev Lett* (2013) 110:026601. doi:10.1103/PhysRevLett.110.026601
- Briskot U, Schütt M, Gornyi IV, Titov M, Narozhny BN, Mirlin AD. Collision-dominated nonlinear hydrodynamics in graphene. *Phys Rev B* (2015) 92:115426. doi:10.1103/PhysRevB.92.115426
- Narozhny BN. Electronic hydrodynamics in graphene. *Ann Phys* (2019a) 411:167979. doi:10.1016/j.aop.2019.167979

## FUNDING

This work was supported by the German Research Foundation DFG within FLAG-ERA Joint Transnational Call (Project GRANSPORT), by the European Commission under the EU Horizon 2020 MSCA-RISE-2019 program (Project 873028 HYDROTRONICS), by the German-Israeli Foundation for Scientific Research and Development (GIF) Grant no. I-1505-303.10/2019 (IVG) and by the Russian Science Foundation, Grant No. 20-12-00147 (IVG). BNN acknowledges the support by the MEFHI Academic Excellence Project, Contract No. 02.a03.21.0005.

## ACKNOWLEDGMENTS

The authors are grateful to P. Alekseev, A. Andreev, U. Briskot, A. Dmitriev, L. Golub, V. Kachorovskii, E. Kiselev, A. Mirlin, J. Schmalian, M. Schütt, A. Shnirman, K. Tikhonov, and M. Titov for fruitful discussions.

- Narozhny BN, Schütt M. Magnetohydrodynamics in graphene: shear and hall viscosities. *Phys Rev B* (2019) 100:035125. doi:10.1103/PhysRevB.100.035125
- Narozhny BN. Optical conductivity in graphene: hydrodynamic regime. *Phys Rev B* (2019b) 100:115434. doi:10.1103/PhysRevB.100.115434
- Sheehy DE, Schmalian J. Quantum critical scaling in graphene. *Phys Rev Lett* (2007) 99:226803. doi:10.1103/PhysRevLett.99.226803
- Kozikov AA, Savchenko AK, Narozhny BN, Shytov AV. Electron-electron interactions in the conductivity of graphene. *Phys Rev B* (2010) 82:075424. doi:10.1103/PhysRevB.82.075424
- Gallagher P, Yang C-S, Lyu T, Tian F, Kou R, Zhang H, et al. Quantum-critical conductivity of the Dirac fluid in graphene. *Science* (2019) 364:158. doi:10.1126/science.aat8687
- Lifshitz EM, Pitaevskii LP. *Physical kinetics*. London, United Kingdom: Pergamon Press (1981).
- Zarenia M, Principi A, Vignale G. Thermal transport in compensated semimetals: effect of electron-electron scattering on Lorenz ratio. *Phys Rev B* (2020) 102:214304. doi:10.1103/PhysRevB.102.214304
- Alekseev PS, Dmitriev AP, Gornyi IV, Kachorovskii VY, Narozhny BN, Schütt M, et al. Magnetoresistance in two-component systems. *Phys Rev Lett* (2015) 114:156601. doi:10.1103/PhysRevLett.114.156601
- Vasileva GY, Smirnov D, Ivanov YL, Vasilyev YB, Alekseev PS, Dmitriev AP, et al. Linear magnetoresistance in compensated graphene bilayer. *Phys Rev B* (2016) 93:195430. doi:10.1103/PhysRevB.93.195430
- Alekseev PS, Dmitriev AP, Gornyi IV, Kachorovskii VY, Narozhny BN, Schütt M, et al. Magnetoresistance of compensated semimetals in confined geometries. *Phys Rev B* (2017) 95:165410. doi:10.1103/PhysRevB.95.165410
- Alekseev PS, Dmitriev AP, Gornyi IV, Kachorovskii VY, Narozhny BN, Titov M. Nonmonotonic magnetoresistance of a two-dimensional viscous electron-hole fluid in a confined geometry. *Phys Rev B* (2018) 97:085109. doi:10.1103/PhysRevB.97.085109
- Gorbachev RV, Geim AK, Katsnelson MI, Novoselov KS, Tudorovskiy T, Grigorieva IV, et al. Strong Coulomb drag and broken symmetry in double-layer graphene. *Nat Phys* (2012) 8:896. doi:10.1038/nphys2441
- Titov M, Gorbachev RV, Narozhny BN, Tudorovskiy T, Schütt M, Ostrovsky PM, et al. Giant magnetodrag in graphene at charge neutrality. *Phys Rev Lett* (2013) 111:166601. doi:10.1103/PhysRevLett.111.166601
- Xie H-Y, Foster MS. Transport coefficients of graphene: interplay of impurity scattering, Coulomb interaction, and optical phonons. *Phys Rev B* (2016) 93:195103. doi:10.1103/PhysRevB.93.195103
- Xie H-Y, Levchenko A. Negative viscosity and eddy flow of the imbalanced electron-hole liquid in graphene. *Phys Rev B* (2019) 99:045434. doi:10.1103/PhysRevB.99.045434

32. Song JCW, Reizer MY, Levitov LS. Disorder-assisted electron-phonon scattering and cooling pathways in graphene. *Phys Rev Lett* (2012) 109:106602. doi:10.1103/PhysRevLett.109.106602
33. Graham MW, Shi S-F, Ralph DC, Park J, McEuen PL. Photocurrent measurements of supercollision cooling in graphene. *Nat Phys* (2013) 9:103–108. doi:10.1038/nphys2493
34. Betz AC, Jhang SH, Pallecchi E, Ferreira R, Fève G, Berroir J-M, et al. Supercollision cooling in undoped graphene. *Nat Phys* (2013) 9:109–112. doi:10.1038/nphys2494
35. Tikhonov KS, Gornyi IV, Kachorovskii VY, Mirlin AD. Resonant supercollisions and electron-phonon heat transfer in graphene. *Phys Rev B* (2018) 97:085415. doi:10.1103/PhysRevB.97.085415
36. Kong JF, Levitov L, Halbertal D, Zeldov E. Resonant electron-lattice cooling in graphene. *Phys Rev B* (2018) 97:245416. doi:10.1103/PhysRevB.97.245416
37. Bistritzer R, MacDonald AH. Electronic cooling in graphene. *Phys Rev Lett* (2009) 102:206410. doi:10.1103/PhysRevLett.102.206410
38. Tse W-K, Das Sarma S. Energy relaxation of hot Dirac fermions in graphene. *Phys Rev B* (2009) 79:235406. doi:10.1103/PhysRevB.79.235406
39. Bandurin DA, Shytov AV, Levitov LS, Kumar RK, Berdyugin AI, Ben Shalom M, et al. Fluidity onset in graphene. *Nat Commun* (2018) 9:4533. doi:10.1038/s41467-018-07004-4

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Narozhny and Gornyi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.