



The role of the *P1BS* element containing promoter-driven genes in Pi transport and homeostasis in plants

Lukasz Sobkowiak¹, Dawid Bielewicz¹, Ewelina M. Malecka¹, Iver Jakobsen², Merete Albrechtsen³, Zofia Szweykowska-Kulinska¹ and Andrzej Pacak^{1*}

¹ Department of Gene Expression, Adam Mickiewicz University, Poznan, Poland

² Department of Chemical and Biochemical Engineering, Technical University of Denmark, Lyngby, Denmark

³ Department of Plant Biology and Biotechnology, University of Copenhagen, Frederiksberg C, Denmark

Edited by:

Bernie Carroll, The University of Queensland, Australia

Reviewed by:

Chris Helliwell, Commonwealth Scientific and Industrial Research Organisation, Australia
Iain Robert Searle, Australian National University, Australia
Tony Millar, Australian National University, Australia

*Correspondence:

Andrzej Pacak, Department of Gene Expression, Adam Mickiewicz University, Umultowska 89, Poznan 61-614, Poland.
e-mail: apacak@amu.edu.pl

Inorganic phosphate (Pi) is an easily accessible form of phosphorus for plants. Plant Pi uptake is usually limited however by slow Pi diffusion through the soil which strongly adsorbs phosphate species. Plants have developed mechanisms to increase Pi availability. There are also abiotic (phosphate level) and biotic (e.g., mycorrhizal) factors regulating the expression of Pi-responsive genes. Transcription factors binding to the promoters of Pi-responsive genes activate different pathways of Pi transport, distribution, and homeostasis maintenance. Pi metabolism involves not only functional proteins but also microRNAs and other non-coding RNAs.

Keywords: phosphate, Pi-responsiveness, *P1BS*, microRNA


INTRODUCTION

Phosphorus (P) is one of the most important macronutrients in the plant lifecycle. It is involved in the synthesis of nucleic acids and phospholipids, in reactions of phosphorylation, and also in energy delivery (ATP). Lack of phosphate (inorganic phosphate, Pi) inhibits plant growth, whereas excessive phosphate levels are toxic to plants and stimulate algal bloom in water reservoirs (reviewed by Scott, 2008). Here we describe genes and their products involved in Pi transport and maintenance of Pi homeostasis (Figure 1A). Phosphate and its role in plant life have both scientific and economic importance. Figure 1B shows the effect of Pi-deprivation on plant growth: the barley plants grown without Pi in the absence (–Pi) or presence (–Pi, Myc) of mycorrhizal fungi had shoot weights 35.2 and 33% lower, respectively, than the plant grown in Pi replete soil (+Pi) at 23 days post-sowing.

TRANSCRIPTION FACTORS INVOLVED IN Pi ACQUISITION

The mechanisms of Pi acquisition include secretion of carbon dioxide, and plant enzymes, e.g., acid phosphatases and organic acids (citric and malic acids), which target organic and inorganic soil P species, respectively, and release Pi from the soil particles (Scott, 2008; Richardson et al., 2011). Phosphate starvation induces expression of transcription factors (TFs) like *phosphorus starvation response 1* (*Psr1*), which shows at least 10-fold increase of expression during Pi deficiency in *Chlamydomonas reinhardtii* (AF174532; Wykoff et al., 1999). In contrast, the related *Arabidopsis thaliana* ortholog *phosphate starvation response 1* (*AtP1H1*, At4g28610) is only weakly responsive to Pi starvation (Rubio et al., 2001). The key regulatory functions of AtP1H1 and other TFs involved in Pi response have been reviewed by Nilsson et al.

(2010) and Rouached et al. (2010). Signaling networks including TFs, Pi-responsive microRNAs, hormones, and sugars implicated in Pi sensing have been recently extensively reviewed by Chiou and Lin (2011). Located in the nucleus AtP1H1 can bind as a dimer via its MYB domain to the P1H1 binding site (*P1BS*, sequence GNATATNC), a motif present in the promoters of crucial Pi-responsive genes (Rubio et al., 2001; Schünmann et al., 2004a; Bustos et al., 2010; Nilsson et al., 2010; Oropeza-Aburto et al., 2012). A set of Pi-responsive genes with *P1BS* elements in their promoters is outlined in Figure 1A. This motif is frequent and considerably enriched in the promoters of Pi-responsive genes of *Arabidopsis* compared with the entire genome (Müller et al., 2007; Bustos et al., 2010). *AtP1H1* also regulates genes not directly involved in phosphate metabolism but possessing the *P1BS* element: e.g., *P1BS* occurs in the promoter of the *Arabidopsis sulfate transporter 1;3* gene (*AtSULTR1;3*, At1g22150), which is up-regulated during Pi-deficient conditions in wild-type plants but much less in *phr1* mutant plants (Rouached et al., 2011). Bustos et al. (2010) showed that expression of Pi-responsive genes in *Arabidopsis* also requires the P1H1-LIKE1 TF (*AtP1H1*, At5g29000), which also contains a MYB domain. The double mutant *phr1 phl1* exhibited lower expression levels of Pi transporter *AtPht1;1* (U62330, At5g43350, three *P1BS* elements) compared with either of the single mutants *phr1* and *phl1* or wild-type plants (Franco-Zorrilla et al., 2004; Bustos et al., 2010). Since the expression of neither *AtP1H1* nor *AtP1H1* is strongly induced by Pi starvation, it is still unclear how Pi-limitation influences Pi-responsive genes. It has been shown in *Arabidopsis* that there are at least four MYB-CC (CC – coiled coil domain) proteins highly similar to both AtP1H1 and AtP1H1, i.e., At2g20400, At3g04450, At3g13040, and At5g06800 (Bustos et al.,



2010). It is possible that these proteins are also involved in regulating the Pi-deprivation response. Furthermore, the AtPHR1 protein is a target for sumoylation by the SUMO E3 ligase, AtSIZ1 (At5g60410), which is also needed for Pi starvation-dependent

responses (Miura et al., 2005). The expression of *AtSIZ1* is not strongly induced by Pi starvation, however (Miura et al., 2005).


PHOSPHATE TRANSPORT

There are four Pi transporter classes, Pht1, Pht2, Pht3, and Pht4, which are responsible for phosphate transport across membranes of cells, chloroplasts, mitochondria, and Golgi, respectively (Karandashov and Bucher, 2005; Rouached et al., 2010). Analysis of six barley (*Hordeum vulgare*) *Pht1* gene promoters revealed that all analyzed promoters possess *P1BS* elements (Schünmann et al., 2004a). These genes encode proteins belonging to the Pht1 family of phosphate transporters, which represents plasma membrane phosphate-H⁺ symporters, containing 12 trans-membrane helices connected by a hydrophilic loop (reviewed by Karandashov and Bucher, 2005). Expression of the *AtPht1;1*, *HvPht1;1*, *HvPht1;2* (almost identical CDS to *HvPht1;1*), *Triticum aestivum Pht1;2* (AJ344241) gene members of the *Pht1* family, is largely restricted to roots (Muchhal et al., 1996; Davies et al., 2002; Schünmann et al., 2004a). The *HvPht1;1* (AF543197) and *HvPht1;2* (AY187019) promoters possess three and two *P1BS*-like motifs, respectively, and are Pi-limitation induced genes (Schünmann et al., 2004a,b; Glassop et al., 2005; Pacak et al., unpublished data; **Figure 1C**). Mutations in two out of three *P1BS* elements present in the *HvPht1;1* rearranged promoter completely abolished promoter completely abolished low-Pi induction (Schünmann et al., 2004b). Expression of the *TaPht1;2* gene is induced by Pi starvation in the wheat cultivar Dalcahue, although the full *P1BS* motif is not present in the promoter. Instead, four other conserved motifs have been identified. Three of them are also present in the promoter of *HvPht1;1*, whereas the fourth one is similar to the *P1BS* element (motif4; ATATRCA sequence; Tittarelli et al., 2007). Some phosphate transporter genes are expressed only in the presence of mycorrhizal fungi (reviewed by Javot et al., 2007; Smith et al., 2011). Mycorrhizal fungi do not colonize the roots of *Arabidopsis*, and no mycorrhiza-specific Pi transporter genes have been discovered in this species. They are, however, present in barley (*HvPht1;8*, AY187023; Glassop et al., 2005; Pacak et al., unpublished data; **Figure 1D**). Chen et al. (2011) described a set of mycorrhiza-activated phosphate transporters from eudicots. Apart from a mycorrhiza transcription factor binding sequence (MYCS), *P1BS* motifs were also found in the promoters of these genes. Mutation or deletion of either of these motifs resulted in a remarkable decrease or even the complete absence of gene expression (Chen et al., 2011).

Pi LEVEL REGULATION IN PLANTS BY microRNAs 399 AND IPS1

Among the Pi-responsive microRNAs which have been described in the latest review published by Kuo and Chiou (2011), *miR399* is especially interesting owing to its interaction with an RNA molecule named *induced by phosphate starvation 1 (IPS1)*. The *miR399* promoters possess the *P1BS* motif (Kuo and Chiou, 2011) and expression of their primary transcripts (pri-microRNAs) is activated by Pi starvation (Pant et al., 2009; Sobkowiak, unpublished data; **Figure 1E**). Two *P1BS* elements were found in the promoter of the *Arabidopsis* gene *AtIPS1* (AF236376, At3g09922; Martin et al., 2000; Rubio et al., 2001; Bustos et al., 2010). Originally *AtIPS1*

was described as a highly Pi starvation inducible gene belonging to the *Mt4/TPS1* family, which possesses a conserved nucleotide motif and encodes only short open reading frames (Martin et al., 2000). Excellent work done by Franco-Zorrilla et al. (2007) showed that *AtIPS1* sequesters miR399, which is bound to the mentioned conserved region, thereby preventing complete degradation of miR399 targets such as the mRNA for the phosphate2 protein (*PHO2*, At2g33770). *AtIPS1* itself is not cleaved by miR399 owing to the sequence mismatches (Figure 2A). As we showed in our previous report about miR gene structures and the processing of *Arabidopsis* HYL1-dependent pri-microRNAs, alternative splicing events and alternative polyadenylation of microRNA precursors are often observed (Szarzynska et al., 2009). Interestingly, the *Arabidopsis* pri-microRNA399a transcript undergoes alternative splicing, with 3' alternative splice site selection resulting in completely different sequences of the 3' exons in the two pri-miR399a isoforms as well as different polyadenylation sites (Figure 2B; Sobkowiak, unpublished data). The role of these processes in miR399a expression regulation is still elusive.



The six microRNA399 species (a–f) in *Arabidopsis* are considered as the immediate mediators of *PHO2* mRNA silencing. They are derived from six pre-microRNAs that differ in structure and sequence (Figures 2C–H). Mature miR399s can bind to five predicted target sites (differing in sequence from each other) present in the 5' UTR of the *PHO2* mRNA (Allen et al., 2005; Bari et al., 2006). Only miR399a and miR399f have full complementarity (including GU base pairings) with target sequences – sites 1 and 3 in the 5' UTR of *PHO2* mRNA, respectively (Lin et al., 2008). In *Arabidopsis*, an overexpression of miR399f leads to overaccumulation of Pi in shoots resembling the *pho2* mutant phenotype (Chiou et al., 2006). Suppression of *PHO2* by overexpressed miR399b/c is less efficient (Lin et al., 2008). The differences in microRNA affinity for the *PHO2* mRNA are probably owed to a nucleotide substitution at position 13 in the miR399b/c sequence, which reduces base pairing between microRNA399b/c and the target sites (Lin et al., 2008). Interestingly, the same nucleotide substitution improves base pairing of miR399b/c with the conserved region of the *At4/IPS1* family. A similar variation in miR399 sequences has been found in rice, *Medicago*, and *Populus*, where certain miR399 variants show reduced base pairing with target sequences in *PHO2* but improved base pairing with the *IPS1* homolog of the respective species (Lin et al., 2008).

The *Arabidopsis* *PHO2* protein is responsible for the Pi level decrease in shoots and Pi remobilization. *pho2* mutants grown under Pi replete conditions accumulate Pi in shoots, but not in roots, and show induction of some phosphate starvation-induced genes, e.g., acid phosphatase 5 (*AtACP5*, At3g17790) and *AtPht1;4* – At2g38940 (Delhaize and Randall, 1995; Bari et al., 2006). Down-regulation of *PHO2* expression in barley produced a similar effect of increased Pi levels in shoots and decreased Pi levels in roots (Pacak et al., 2010). *PHO2* possesses the Ubiquitin-Conjugating E2 enzyme catalytic domain (UBC domain). Since other UBC domain-containing proteins are usually smaller, 20 vs. 100 kDa (Bari et al., 2006), *PHO2* may contain unidentified additional domains. Such domains may interact with phosphate-related proteins, e.g., targeting them for degradation. Degradation of these proteins may affect the phosphate homeostasis. Bioinformatic analysis showed that, apart from *PHO2* mRNA, other transcripts derived from the following genes, At3g11130 (encoding Clathrin, heavy chain), At3g25905 (clavata3/ESR-related27 protein), At3g54700 (*AtPht1;7*), At4g00170 (vesicle-associated membrane protein, VAMP), and At4g09730 (DEAD-box protein), are potential targets for miR399 (Pant et al., 2009; Kuo and Chiou, 2011). These complex connections between the miR399 family members and their targets can explain the fact that miR399b-overexpressing plants and a *pho2* mutant exhibited not only Pi-related changes, but also an early flowering phenotype observed only at normal temperature (23°C; Kim et al., 2011).

FUTURE PROSPECTS

The data presented above show that the presence of one or more *P1BS* elements in a gene promoter is associated with low-Pi induction. Other factors, however, can modulate the response. In the promoter of the *Arabidopsis* phospholipase *DZ2* gene (*PLDZ2*, At3g05630), apart from five *P1BS* copies, elements such as *SRE* (*sugar-repressive element*) have been found. A 65-bp promoter

fragment spanning two of the *P1BS* motifs (the *EZ2* region) has been identified as particularly important for the Pi-limitation response but strong induction also required sucrose and was negatively affected by cytokinins (Oropeza-Aburto et al., 2012). Exactly how the information carried by the *P1BS* elements and by the various sugar and hormone responsive promoter elements is integrated by the MYB-CC TFs *PHR1* and *PHL1* and possibly others is still unknown. Furthermore, additional studies are necessary to find out how Pi-limitation directly affects the function and/or expression of Pi-related TFs. Finally, the influence of other factors which can modulate the Pi response of genes containing *P1BS* elements should be further investigated.

ACKNOWLEDGMENTS

We would like to thank Anne Olsen (DTU) for collaboration in barley cultivation in different phosphate conditions. Andrzej

Pacak and Zofia Szwejkowska-Kulinska were supported by the European Regional Development Fund through the Innovative Economy for Poland 2007–2013, project WND-POIG.01.03.01-00-101/08 POLAPGEN-BD “Biotechnological tools for breeding cereals with increased resistance to drought.” Andrzej Pacak and Merete Albrechtsen were supported by Marie Curie Intra-European Fellowships – BARPPO, 025110. Dawid Bielewicz was supported by the Foundation for Polish Science (FNP) – MPD 2010/3. Lukasz Sobkowiak was supported by a Ph.D. grant awarded by the Polish Ministry of Higher Education and Sciences no. NN301035839, a grant for scientific research from the Dean of Biology Faculty, Adam Mickiewicz University, Poznan, and the European Fund “Scholarship support for Ph.D. students specializing in majors strategic for Wielkopolska’s development,” Sub-measure 8.2.2, Human Capital Operational Programme. The work was also supported by grant MNiSW No. 3011/B/P01/2009/37.

REFERENCES

- Allen, E., Xie, Z., Gustafson, A. M., and Carrington, J. C. (2005). MicroRNA-directed phasing during trans-acting siRNA biogenesis in plants. *Cell* 121, 207–221.
- Bari, R., Pant, B. D., Stitt, M., and Scheible, W. R. (2006). PHO2, microRNA, and PHR1 define a phosphate-signaling pathway in plants. *Plant Physiol.* 141, 988–999.
- Bustos, R., Castrillo, G., Linhares, F., Puga, M. I., Rubio, V., Pérez-Pérez, J., Solano, R., Leyva, A., and Paz-Ares, J. (2010). A central regulatory system largely controls transcriptional activation and repression responses to phosphate starvation in *Arabidopsis*. *PLoS Genet.* 6, e1001102. doi:10.1371/journal.pgen.1001102
- Chen, A., Gu, M., Sun, S., Zhu, L., Hong, S., and Xu, G. (2011). Identification of two conserved cis-acting elements, MYCS and P1BS, involved in the regulation of mycorrhiza-activated phosphate transporters in eudicot species. *New Phytol.* 189, 1157–1169.
- Chiou, T. J., Aung, K., Lin, S. I., Wu, C. C., Chiang, S. F., and Su, C. I. (2006). Regulation of phosphate homeostasis by microRNA in *Arabidopsis*. *Plant Cell* 18, 412–421.
- Chiou, T. J., and Lin, S. I. (2011). Signaling network in sensing phosphate availability in plants. *Annu. Rev. Plant Biol.* 62, 185–206.
- Davies, T. G. E., Ying, J., Xu, Q., Li, Z. S., Li, J., and Gordon-Weeks, R. (2002). Expression analysis of putative high-affinity phosphate transporters in Chinese winter wheats. *Plant Cell Environ.* 25, 1325–1339.
- Delhaize, E., and Randall, P. J. (1995). Characterization of a phosphate-accumulator mutant of *Arabidopsis thaliana*. *Plant Physiol.* 107, 207–213.
- Franco-Zorrilla, J. M., González, E., Bustos, R., Linhares, F., Leyva, A., and Paz-Ares, J. (2004). The transcriptional control of plant responses to phosphate limitation. *J. Exp. Bot.* 55, 285–293.
- Franco-Zorrilla, J. M., Valli, A., Todesco, M., Mateos, I., Puga, M. I., Rubio-Somoza, I., Leyva, A., Weigel, D., Garcia, J. A., and Paz-Ares, J. (2007). Target mimicry provides a new mechanism for regulation of microRNA activity. *Nat. Genet.* 39, 1033–1037.
- Glassop, D., Smith, S. E., and Smith, F. W. (2005). Cereal phosphate transporters associated with the mycorrhizal pathway of phosphate uptake into roots. *Planta* 222, 688–698.
- Gu, M., Xu, K., Chen, A., Zhu, Y., Tang, G., and Xu, G. (2010). Expression analysis suggests potential roles of microRNAs for phosphate and arbuscular mycorrhizal signaling in *Solanum lycopersicum*. *Physiol. Plant.* 138, 226–237.
- Hansen, T. (2007). *Folder Version 1.11 Beta*. Denmark: MBI, University of Aarhus. Available at: <http://www.ncrnalab.dk/rnafolder/>
- Javot, H., Pumplun, N., and Harrison, M. J. (2007). Phosphate in the arbuscular mycorrhizal symbiosis: transport properties and regulatory roles. *Plant Cell Environ.* 30, 310–322.
- Karandashov, V., and Bucher, M. (2005). Symbiotic phosphate transport in arbuscular mycorrhizas. *Trends Plant Sci.* 10, 22–29.
- Kim, W., Ahn, H. J., Chiou, T. J., and Ahn, J. H. (2011). The role of the miR399-PHO2 module in the regulation of flowering time in response to different ambient temperatures in *Arabidopsis thaliana*. *Mol. Cells* 32, 83–88.
- Kuo, H. F., and Chiou, T. J. (2011). The role of microRNAs in phosphorus deficiency signaling. *Plant Physiol.* 156, 1016–1024.
- Lin, S. I., Chiang, S. F., Lin, W. Y., Chen, J. W., Tseng, C. Y., Wu, P. C., and Chiou, T. J. (2008). Regulatory network of microRNA399 and PHO2 by systemic signaling. *Plant Physiol.* 147, 732–746.
- Martin, A. C., del Pozo, J. C., Iglesias, J., Rubio, V., Solano, R., de La Peña, A., Leyva, A., and Paz-Ares, J. (2000). Influence of cytokinins on the expression of phosphate starvation responsive genes in *Arabidopsis*. *Plant J.* 24, 559–567.
- Miura, K., Rus, A., Sharkhuu, A., Yokoi, S., Karthikeyan, A. S., Raghothama, K. G., Baek, D., Koo, Y. D., Jin, J. B., Bressan, R. A., Yun, D. J., and Hasegawa, P. M. (2005). The *Arabidopsis* SUMO E3 ligase SIZ1 controls phosphate deficiency responses. *Proc. Natl. Acad. Sci. U.S.A.* 102, 7760–7765.
- Muchhal, U. S., Pardo, J. M., and Raghothama, K. G. (1996). Phosphate transporters from the higher plant *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U.S.A.* 93, 10519–10523.
- Müller, R., Morant, M., Jarmer, H., Nilsson, L., and Nielsen, T. H. (2007). Genome-wide analysis of the *Arabidopsis* leaf transcriptome reveals interaction of phosphate and sugar metabolism. *Plant Physiol.* 143, 156–171.
- Nilsson, L., Müller, R., and Nielsen, T. H. (2010). Dissecting the plant transcriptome and the regulatory responses to phosphate deprivation. *Physiol. Plant.* 139, 129–143.
- Oropeza-Aburto, A., Cruz-Ramírez, A., Acevedo-Hernández, G. J., Pérez-Torres, C. A., Caballero-Pérez, J., and Herrera-Estrella, L. (2012). Functional analysis of the *Arabidopsis* PLDZ2 promoter reveals an evolutionarily conserved low-Pi-responsive transcriptional enhancer element. *J. Exp. Bot.* 63, 2189–2202.
- Pacak, A., Geisler, K., Jørgensen, B., Barciszewska-Pacak, M., Nilsson, L., Nielsen, T. H., Johansen, E., Grønlund, M., Jakobsen, I., and Albrechtsen, M. (2010). Investigations of barley stripe mosaic virus as a gene silencing vector in barley roots and in *Brachypodium distachyon* and oat. *Plant Methods* 6, 26.
- Pant, B. D., Musialak-Lange, M., Nuc, P., May, P., Buhtz, A., Kehr, J., Walther, D., and Scheible, W. R. (2009). Identification of nutrient-responsive *Arabidopsis* and rapeseed microRNAs by comprehensive real-time polymerase chain reaction profiling and small RNA sequencing. *Plant Physiol.* 150, 1541–1555.
- Richardson, A. E., Lynch, J. P., Ryan, P. R., Delhaize, E., Smith, F. A., Smith, S. E., Harvey, P. R., Ryan, M. H., Veneklaas, E. J., Lambers, H., Oberon, A., Culvenor, R. A., and Simpson, R. J. (2011). Plant and microbial strategies to improve the phosphorus efficiency of agriculture. *Plant Soil* 349, 121–156.
- Rouached, H., Arpat, A. B., and Poirier, Y. (2010). Regulation of phosphate starvation responses in plants: signaling players and cross-talks. *Mol. Plant* 3, 288–299.
- Rouached, H., Secco, D., Bulak, A., and Poirier, Y. (2011). The transcription factor PHR1 plays a key role in the regulation of sulfate shoot-to-root flux upon phosphate starvation

- in *Arabidopsis*. *BMC Plant Biol.* 11, 19.
- Rubio, V., Linhares, F., Solano, R., Martín, A. C., Iglesias, J., Leyva, A., and Paz-Ares, J. (2001). A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. *Genes Dev.* 15, 2122–2133.
- Schünmann, P. H. D., Richardson, A. E., Smith, F. W., and Delhaize, E. (2004a). Characterization of promoter expression patterns derived from the Pht1 phosphate transporter genes of barley (*Hordeum vulgare* L.). *J. Exp. Bot.* 55, 855–865.
- Schünmann, P. H. D., Richardson, A. E., Vickers, C. E., and Delhaize, E. (2004b). Promoter analysis of the barley Pht1;1 phosphate transporter gene identifies regions controlling root expression and responsiveness to phosphate deprivation. *Plant Physiol.* 136, 4205–4214.
- Scott, P. (2008). *Physiology and Behaviour of Plants*. Chichester: John Wiley & Sons Ltd.
- Smith, S. E., Jakobsen, I., Grønlund, M., and Smith, F. A. (2011). Roles of arbuscular mycorrhizas in plant phosphorus nutrition: interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiol.* 156, 1050–1057.
- Stefanovic, A., Ribot, C., Rouached, H., Wang, Y., Chong, J., Belbahri, L., Delessert, S., and Poirier, Y. (2007). Members of the PHO1 gene family show limited functional redundancy in phosphate transfer to the shoot, and are regulated by phosphate deficiency via distinct pathways. *Plant J.* 50, 982–994.
- Szarzynska, B., Sobkowiak, L., Pant, B. D., Balazadeh, S., Scheible, W. R., Mueller-Roeber, B., Jarmolowski, A., and Szweykowska-Kulinska, Z. (2009). Gene structures and processing of *Arabidopsis thaliana* HYL1-dependent pri-miRNAs. *Nucleic Acids Res.* 37, 3083–3093.
- Tittarelli, A., Milla, L., Vargas, F., Morales, A., Neupert, C., Meisel, L. A., Salvo, G. H., Penaloza, E., Munoz, G., Corcuera, L. J., and Silva, H. (2007). Isolation and comparative analysis of the wheat TaPT2 promoter: identification in silico of new putative regulatory motifs conserved between monocots and dicots. *J. Exp. Bot.* 58, 2573–2582.
- Wykoff, D. D., Grossman, A. R., Weeks, D. P., Usuda, H., and Shimogawara, K. (1999). Psr1, a nuclear localized protein that regulates phosphorus metabolism in *Chlamydomonas*. *Proc. Natl. Acad. Sci. U.S.A.* 96, 15336–15341.
- could be construed as a potential conflict of interest.

Received: 30 November 2011; accepted: 07 March 2012; published online: 30 March 2012.

Citation: Sobkowiak L, Bielewicz D, Malecka EM, Jakobsen I, Albrechtsen M, Szweykowska-Kulinska Z and Pacak A (2012) The role of the P1BS element containing promoter-driven genes in Pi transport and homeostasis in plants. *Front. Plant Sci.* 3:58. doi: 10.3389/fpls.2012.00058

This article was submitted to *Frontiers in Plant Genetics and Genomics*, a specialty of *Frontiers in Plant Science*.

Copyright © 2012 Sobkowiak, Bielewicz, Malecka, Jakobsen, Albrechtsen, Szweykowska-Kulinska and Pacak. This is an open-access article distributed under the terms of the Creative Commons Attribution Non Commercial License, which permits non-commercial use, distribution, and reproduction in other forums, provided the original authors and source are credited.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that