



# Ethylene is Involved in Brassinosteroids Induced Alternative Respiratory Pathway in Cucumber (*Cucumis sativus* L.) Seedlings Response to Abiotic Stress

Li-Jie Wei<sup>†</sup>, Xing-Guang Deng<sup>†</sup>, Tong Zhu, Ting Zheng, Peng-Xu Li, Jun-Qiang Wu, Da-Wei Zhang\* and Hong-Hui Lin\*

State Key Laboratory of Hydraulics and Mountain River Engineering, Key Laboratory for Bio-Resource and Eco-Environment, College of Life Science, Ministry of Education, Sichuan University, Chengdu, China

## OPEN ACCESS

### Edited by:

David John Burritt,  
University of Otago, New Zealand

### Reviewed by:

Oksoo Han,  
Chonnam National University,  
South Korea  
David W. M. Leung,  
University of Canterbury, New Zealand

### \*Correspondence:

Da-Wei Zhang  
yuanmiao1892@163.com;  
Hong-Hui Lin  
hhlin@scu.edu.cn

<sup>†</sup>These authors have contributed  
equally to this work.

### Specialty section:

This article was submitted to  
Crop Science and Horticulture,  
a section of the journal  
Frontiers in Plant Science

Received: 08 August 2015

Accepted: 26 October 2015

Published: 10 November 2015

### Citation:

Wei L-J, Deng X-G, Zhu T, Zheng T,  
Li P-X, Wu J-Q, Zhang D-W and  
Lin H-H (2015) Ethylene is Involved in  
Brassinosteroids Induced Alternative  
Respiratory Pathway in Cucumber  
(*Cucumis sativus* L.) Seedlings  
Response to Abiotic Stress.  
*Front. Plant Sci.* 6:982.  
doi: 10.3389/fpls.2015.00982

Effects of brassinosteroids (BRs) on cucumber (*Cucumis sativus* L.) abiotic stresses resistance to salt, polyethylene glycol (PEG), cold and the potential mechanisms were investigated in this work. Previous reports have indicated that BRs can induce ethylene production and enhance alternative oxidase (AOX) pathway. The mechanisms whether ethylene is involved as a signal molecule which connected BR with AOX in regulating stress tolerance are still unknown. Here, we found that pretreatment with 1  $\mu$ M brassinolide (BL, the most active BRs) relieved stress-caused oxidative damage in cucumber seedlings and clearly enhanced the capacity of AOX and the ethylene biosynthesis. Furthermore, transcription level of ethylene signaling biosynthesis genes including ripening-related ACC synthase1 ( $C_S$ ACS1), ripening-related ACC synthase2 ( $C_S$ ACS2), ripening-related ACC synthase3 ( $C_S$ ACS3), 1-aminocyclopropane-1-carboxylate oxidase1 ( $C_S$ ACO1), 1-aminocyclopropane-1-carboxylate oxidase2 ( $C_S$ ACO2), and  $C_S$ AOX were increased after BL treatment. Importantly, the application of the salicylhydroxamic acid (SHAM, AOX inhibitor) and ethylene biosynthesis inhibitor aminoxyacetic acid (AOA) decreased plant resistance to environmental stress by blocking BRs-induced alternative respiration. Taken together, our results demonstrated that ethylene was involved in BRs-induced AOX activity which played important roles in abiotic stresses tolerance in cucumber seedlings.

**Keywords:** alternative oxidase, environmental stress, cucumber, brassinosteroid, ethylene

## INTRODUCTION

In both natural and agricultural environments, plants are challenged by heat, cold, drought, and high salinity throughout their life cycles, influencing plant growth and productivity (Xia et al., 2009). To survive from environmental stresses, plants have evolved biochemical and physiological mechanisms to activate tolerance against environmental stresses (Boyer, 1982). It is well known that stress conditions along with reactive oxygen species (ROS) generation, such as superoxide anion ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ) and other free radicals are harmful for plant organisms and making severely damage (Scandalios, 1993; Zhang et al., 2001). In order to

eliminate the ROS damage, plants have evolved the antioxidant enzymes that are responsible for scavenging superfluous ROS accumulation under environmental stresses, including guaiacol peroxidase (GPX), catalase (CAT), superoxide dismutase (SOD), ascorbate peroxidase (APX), and peroxidase (POD) (Zhang et al., 2015). However, salicylic acid (SA) and abscisic acid (ABA) treatments can lead to a temporary elevation of H<sub>2</sub>O<sub>2</sub>, resulting in an enhanced tolerance to heat, oxidative stress, high light and salt, which suggested that ROS may function as a second messenger in the signaling pathways of plant stress responses (Dat et al., 1998; Apel and Hirt, 2004).

Ethylene is a gaseous phytohormone that is synthesized in almost all plant organs. It not only plays a role in plant growth and development, but also is involved in plant responses to biotic and abiotic stresses such as salt, chilling and ozone (Kendrick and Chang, 2008). Methionine is the starting point in the ethylene biosynthetic pathway, which is converted into S-adenosylmethionine (SAM) by methionine adenosyltransferases. S-adenosylmethionine was changed to 1-aminocyclopropane-1-carboxylic acid (ACC) by ACC synthase (ACS) and then to ethylene by ACC oxidase (ACO) (Lin et al., 2010; Wang et al., 2012). Some studies have showed that ethylene induced the cyanide-resistant respiration in ripening fruits (Solomos and Laties, 1976). Several studies have also proposed that the increased ethylene production during fruit ripening might lead to increased HCN level, which in turn inhibited cytochrome oxidase and induced the activity of alternative oxidase (Mizutani et al., 1987; Pirrung and Brauman, 1987; Yip and Yang, 1988). It also has been showed that ethylene is closely connected with AOX induction under stress treatments, including salinity, chilling, pathogen infection, and ozone (Ederli et al., 2006; Wang et al., 2012). Recent studies reported that ethylene accumulation could be induced by BRs treatment in tomato fruit (Zhu et al., 2015).

Respiration plays a pivotal role in driving the cellular metabolism and transport processes. The electron transport system of the plant inner mitochondrial membrane contains two terminal oxidases, including cytochrome oxidase (COX) and alternative oxidase (AOX) (Clifton et al., 2006). It has been confirmed that AOX is the terminal oxidase in the alternative pathway (AP). AP is a non-phosphorylating electron transport chain (ETC) that diverts at the ubiquinone pool and uses AOX (McIntosh, 1994; Wang et al., 2012). As we know that the AP plays an important role in fruit ripening, plant thermogenesis and responses to stress conditions, including salinity, drought, low temperature, ozone, and pathogen invasion (Yoshida et al., 2007). Previous researches have investigated that AOX was involved in optimizing photosynthesis and it might has a pivotal role in relieving the over-reduction of chloroplasts (Yoshida et al., 2008; Giraud et al., 2009; Zhang et al., 2010). In addition, it indicated that AOX could also scavenge ROS to enhance the plant stress tolerance (Xu et al., 2012b).

Plant steroid hormones, brassinosteroids (BRs) were involved in a broad spectrum of plant growth and development. In addition, it played an important role in a variety of plant physiological processes and adaptation to various abiotic and

biotic stresses (Zhang et al., 2015). While, most studies on the role of BRs in stress responses depend on exogenous addition of BRs or their biosynthesis inhibitor. Recently, some studies found that BRs induce a transient increase in H<sub>2</sub>O<sub>2</sub> content. It has been suggested that BRs-induced ROS generation is important for BRs-induced stress tolerance in cucumber, tobacco and tomato (Nakashita et al., 2003; Zhou et al., 2014; Deng et al., 2015). So we hypothesized that there might be some connections among ethylene, ROS, AOX, and BR signaling during the induction of plant stress tolerance. Here, this assumption was examined and our experimental results demonstrated that ethylene might be involved in BRs-induced alternative pathway under environmental stress. We also testified the possible regulation and physiological function of the alternative pathway in BRs-induced stress tolerance.

## MATERIALS AND METHODS


### Plant Material and Growth Conditions

Plants growth as described (Xu et al., 2012b). Seeds of cucumber (*Cucumis sativus* L. cv. Jihong no. 2) were surface-sterilized for 15 min in 1% (w/v) NaClO, and then germinated on water-moistened filter paper. The plants were grown in a growth chamber at a 16-h photoperiod (100 μM m<sup>-2</sup> s<sup>-1</sup>), temperature of 25°C/17°C (day/night).

Cucumber seedlings were cultivated with half-strength Hoagland's nutrient solution until the three-leaf stage. Then, the cucumber plants were treated with Brassinolide (BL, the most active BRs; Chuo-Ku, Osaka, Japan) with 0.1, 0.5, 1, 5, or 10 μM solutions on leaves, while the control seedlings were sprayed with distilled water. Twelve hours after spraying, all the seedlings were exposed to salt (200 mM NaCl), PEG (16% PEG 6000), and cold stresses (at 4°C in a controlled growth chamber with a relative humidity of 70%) for 3 d. The third leaf of cucumber seedlings was used for the following experiments.

### Treatment

Salicylhydroxamic acid (SHAM, an inhibitor of the AOX pathway), dimethylthiourea (DMTU, an H<sub>2</sub>O<sub>2</sub> scavenger), DPI (an NADPH oxidase inhibitor), aminoxyacetic acid (AOA, an ethylene biosynthesis inhibitor), and ethylene were purchased from Sigma (StLouis, USA). 1 mM SHAM inhibits the AOX activity, as this concentration is sufficiently low to avoid the possible side effects (Møller et al., 1988). For BL+SHAM treatment, seedlings were pretreated with 1 mM SHAM, 24 h later were treated with 1 μM BL for another 12 h. Then, these plants were exposed to stress conditions as described earlier. To investigate the role of ROS in the resistance, leaves were pretreated with 100 μM DPI or 5 mM DMTU, then treated with 1 μM BL; 12 h later plants were treated with 10 mM H<sub>2</sub>O<sub>2</sub>. Then 1 mM AOA was sprayed to the seedlings for 12 h at room temperature. For the ethylene treatment, one lot seedling was incubated in 3.5 μl l<sup>-1</sup> ethylene solution in a closed container at room temperature for 12 h. Then, these plants were exposed to stress conditions as described earlier.



**FIGURE 1 |** At the three-leaf stage, cucumber seedlings were sprayed with 0.1, 0.5, 1, 5, or 10 μM BL solutions on leaves, while the control plants were sprayed with distilled water. 12 h later, the seedlings were challenged with salt (200 mM NaCl), PEG (16% PEG6000), or cold (4°C) stresses for 3d, the third leaf was used for H<sub>2</sub>O<sub>2</sub> content (A,C,E,G), electrolyte leakage (EL) (B,D,F,H). (I) Representative phenotypes of cucumber seedlings with or without stresses treatments. Data are the mean ± SD of three biological repeats; the significant difference was analyzed by Student's *t*-test (\**P* < 0.05, \*\**P* < 0.01) and compared to control (CK).

## Cucumber Leaf Respiration Measurements

Respiratory oxygen consumption was measured using Clark-type electrodes (Hansatech, King's Lynn, UK) as previously described (Xu et al., 2012b). Approximately 30 mg of leaves were cut into small pieces, then pretreated with 5 mL deionized water for 15 min in order to eliminate wound-induced respiration. Measurements were done at 25°C in a final volume of 1.5 mL phosphate buffer (pH 6.8) and the cuvette was tightly closed to prevent diffusion of oxygen from the air. Inhibitors of the cytochrome pathway (1 mM KCN) and the alternative pathway (0.5 mM *n*-propyl gallate, nPG) were used. The total respiration ( $V_t$ ) is defined as  $O_2$  uptake rate by cucumber leaves without any inhibitor. Next, 1 mM KCN was added to obtain the  $O_2$  uptake rate defined as  $V_0$ . Residual respiration ( $V_{res}$ ) was defined as  $O_2$  uptake in the presence of both 1 mM KCN and 0.5 mM nPG. The capacity of the cytochrome pathway ( $V_{cyt}$ ) and the alternative pathway ( $V_{alt}$ ) were calculated by the formula:  $V_{cyt} = V_t - V_0$ ;  $V_{alt} = V_0 - V_{res}$ . The  $V_{res}$  in our experiment was always low, and


was negligible relative to other respirations. Therefore, the  $V_{res}$  was not shown.

## Determination of Ethylene Emission

Ethylene was measured as previously described (Xu et al., 2012b). For ethylene production, intact seedlings were placed in 100 mL closed container and incubated at room temperature (25°C) for 1 h. Then, a 1 mL sample of gas from each container headspace was injected into a FID gas chromatograph (Agilent 6890 Series GC System, Salem, MA) equipped with an activated alumina SS column. The carrier gas (helium) flow rate was 0.5 mL  $s^{-1}$ . The detector and injector were operated at 100°C, and the oven was at 50°C.

## Analysis of Chlorophyll Fluorescence

Chlorophyll fluorescence was determined with an imaging pulse amplitude modulated fluorometer (IMAG-MINI; Heinz Walz, Germany). For measurement of  $F_v/F_m$ , plants were dark adapted



for 30 min. Minimal fluorescence ( $F_0$ ) was measured during the weak measuring pulses, and maximal fluorescence ( $F_m$ ) was measured by a 0.8 s pulse of light at about  $4000 \mu\text{mol m}^{-2} \text{s}^{-1}$ . An actinic light source was then applied to obtain steady-state fluorescence yield ( $F_s$ ), after which a second saturation pulse was applied for 0.7 s to obtain light adapted maximum fluorescence ( $F'_m$ ).  $F_v/F_m$  and non-photochemical quenching (NPQ) were calculated as  $F_m - F_0/F_m$  and  $(F_m/F'_m) - 1$ , respectively (Fu et al., 2014).


### RNA Extraction and Quantitative Real-time PCR (qRT-PCR)

Total RNA was extracted from cucumber leaves according to a previously described method (Zhang et al., 2010). RNA contents were calculated by measuring the absorbance

value taken at 260 nm. All gene expression levels were measured by qRT-PCR. The Ct (threshold cycle), defined as the PCR cycle at which a statistically significant increase of reporter fluorescence was first detected, was used as a measure for the starting copy numbers of the target gene. Relative quantitation of gene expression was performed using the comparative Ct method. Three biological replicates were performed for each experiment.  $\beta$ -actin was used as internal controls. All the qRT-PCR primers are listed in Supplemental Table S1.

### Histochemical Staining of Superoxide and $\text{H}_2\text{O}_2$

Superoxide and  $\text{H}_2\text{O}_2$  levels were visually detected with nitro blue tetrazolium (NBT) and 3, 3-diaminobenzidine (DAB),



**FIGURE 3 |** Effects of SHAM, AOA, or AOA + ET treatment on  $\text{H}_2\text{O}_2$  content (A), respiration rate (B,C), and  $CsAOX$  mRNA level (D) in the BL-treated cucumber seedlings. (E) NBT and DAB staining were used to detect the presence of superoxide and  $\text{H}_2\text{O}_2$  contents in leaves. Data are the mean  $\pm$  SD of three biological repeats; the significant difference was analyzed by Student's *t*-test ( $*P < 0.05$ ) and compared to control (CK).

respectively, as previously described (Yang et al., 2004). cucumber leaves were excised at the base with a razor blade and vacuum infiltrated with NBT ( $0.5 \text{ mg mL}^{-1}$ ) solution for 2 h, or DAB ( $2 \text{ mg mL}^{-1}$ ) solution for 8 h. Leaves were then decolorized in boiling ethanol (95%) for 20 min.

### Oxidative Damage Estimation

The  $\text{H}_2\text{O}_2$  content of leaves was measured as described (Xu et al., 2012b). Approximately 0.5 g of fresh cucumber leaves was cut into small pieces and homogenized in an ice bath with 5 mL 0.1% (w/v) trichloroacetic acid (TCA). The homogenate was centrifuged at 12,000 g for 20 min at  $4^\circ\text{C}$ . In addition, 0.5 mL of the supernatant was added to 0.5 mL 10 mM potassium phosphate buffer (pH 7.0) and 1 mL 1 M KI. The absorbance of the supernatant was read at 390 nm.

Electrolyte leakage (EL) was tested as previously described (Cao et al., 2009). After measuring the conductivity of the leaves, the leaves were boiled for 15 min to achieve 100% electrolyte leakage.

Lipid peroxidation was estimated by measuring the malondialdehyde (MDA) level as described previously (Velikova et al., 2000).

Cell death was showed by trypan blue staining. Leaves submerged in trypan blue ( $1.25 \text{ mg mL}^{-1}$ ; Sigma) were heated in a boiling water bath for 2 min. The samples were then decolorized in boiling ethanol (95%) for 15 min.

### Antioxidant Enzyme Extraction and Activity Assay

For the enzyme assays, 0.3 g of leaf were ground with 3 mL ice-cold 25 mM.

HEPES buffer (pH 7.8) containing 0.2 mM EDTA, 2 mM AsA, and 2% PVP. The homogenates were centrifuged at  $4^\circ\text{C}$  for 20 min at 12,000 g, and the resulting supernatants were used for the determination of enzymatic activity (Xia et al., 2009). Superoxide dismutase (SOD), ascorbate peroxidase (APX), catalase (CAT), guaiacol peroxidase (GPX), and peroxidase (POD) activities were assayed as described previously (Wang et al., 2011).


### Statistical Analysis


Means of three replicates were measured. Student's *t*-test was used for comparison between different treatments. A difference was considered to be statistically significant when  $P < 0.05$  or very significant when  $P < 0.01$ .

## RESULTS

### Suitable Concentration of BRs Pretreatment Improves Plant Abiotic Stresses Tolerance

$\text{H}_2\text{O}_2$  content and electrolyte leakage were detected following pretreatment with water, different concentrations of BL for 12 h.






As shown in **Figure 1A**, under normal growth condition, H<sub>2</sub>O<sub>2</sub> content increased significantly with increasing concentrations of BL pretreatment as compared with control, while there was no significant difference in electrolyte leakage (**Figure 1B**). In order to study the roles of BRs in plant stress tolerance, cucumber seedlings were exposed to different stress conditions, including salt (200 mM NaCl), PEG (16% PEG6000), and cold stresses (4°C), after being pretreated with different concentrations of BL for 12 h. Then 3 days later, we detected H<sub>2</sub>O<sub>2</sub> content and electrolyte leakage. As shown in **Figures 1C–H**, under abiotic stresses the content of H<sub>2</sub>O<sub>2</sub> and electrolyte leakage were repressed when seedlings were with 0.1, 0.5, 1, 5, or 10 μM BL treatment, especially under 1 μM BL treatment. In addition, pretreatment with 5 or 10 μM BL did not alleviate stress-induced injury (**Supplemental Figure S1A**). Cucumber seedlings pretreatment with 1 μM BL increased H<sub>2</sub>O<sub>2</sub> content in the early stage, 72 h abiotic stresses treatment later, the level of H<sub>2</sub>O<sub>2</sub> was lower than the control (**Supplemental Figure S1B**). The findings showed that seedlings pretreated with 1 μM BL displayed stronger tolerance to the three types of abiotic stresses (**Figure 1I**), and then the concentration of 1 μM BL was used in the following research.

## Effects of BL on the AOX Capacity and Ethylene Production

According to the recent reports (Xia et al., 2009; Jiang et al., 2013; Deng et al., 2015), we know that H<sub>2</sub>O<sub>2</sub> is induced by

BRs and then rapid generation of ROS induces the expression of AOX, which can eliminate ROS to maintain it in safe level. So we investigated whether BRs could further induce the alternative pathway capacity and ethylene biosynthesis under stress conditions in cucumber. Our results showed that both the alternative respiration and total respiration were increased in the BL-pretreated plants as compared to the control after abiotic stresses for 3 d (**Figure 2A**). Furthermore, under stress conditions, the cyanide-resistant respiration of BL-pretreated seedlings were about 2-fold higher than that of the control. As shown in **Figure 2A**, consistent with the determined alternative respiration rate, we found that compared with control, BL treatment resulted in an enhanced AOX expression in stress conditions, especially under salt and cold stresses. The content of ethylene was also increased with or without BL pretreatment under stress conditions (**Figure 2B**). But it worth noting that the production of ethylene was higher in BL-pretreated seedlings than that of the control plants after 3 d of stresses.

We further investigated the transcripts of ethylene biosynthesis genes. Our results suggested that the mRNA levels of *CsACS2*, *CsACS3*, and *CsACO1* were significantly increased in the BL-pretreated cucumber seedlings in comparison to control plants after 3 d of stresses (**Figures 2B,C**). However, the transcripts of *CsACS1* and *CsACO2* were higher in the BL treated plants than that of the control seedlings under PEG and cold stresses, respectively. Taken together, these results indicated that



both AOX and ethylene might be involved in the BR-induced stress tolerance.

## Ethylene is Involved in BR-induced AOX Capacity to Enhance Plant Abiotic Stresses Tolerance

We investigated the change of AOX activity on the plants against different abiotic stress conditions with or without AOA or AOX inhibitor treatment. Our results showed that AOA pretreatment could also inhibit cyanide-resistant respiration and ROS production which were induced by BL treatment. However, the decline was rescued in seedlings pretreated with ET (Figures 3A–E). These results showed that ethylene was involved in BRs-induced AOX activity. Most stress conditions have harmful effects on various biological processes in seedlings, including the photosystem II (PSII). Chlorophyll fluorescence measurements showed that the PSII photochemical efficiency was influenced by stress conditions (Liu et al., 2009).  $F_v/F_m$  and the non-photochemical quenching (NPQ) are indicators of PSII photochemistry. So, we investigated the function of

photosystem II (PSII) under different abiotic stresses treatments. As shown in Figures 4A,C, fluorescence images of  $F_v/F_m$  showed that compared with control, plants with BL treatment showed an increase in  $F_v/F_m$  in all three abiotic stresses treatments. In contrast, the non-photochemical quenching (NPQ) declined in the BL-pretreated seedlings in comparison with control plants under the same stress treatments for 3 d (Figures 4B,D). However, BL-induced tolerance to photo-oxidative stress was largely inhibited if the plants were pretreated with AOA or SHAM. Application of ET almost rescued the 1 mM AOA-decreased stress tolerance (Figure 4). These results showed that ethylene is involved in BRs-induced AOX activity and then to enhance plant stress tolerance.


We detected the accumulation of H<sub>2</sub>O<sub>2</sub> and superoxide using DAB and NBT staining procedures, respectively. Other physiological indicators such as EL, MDA content and cell death can also indicate the degree of oxidative damage in seedlings caused by abiotic stresses. The results showed that BL pretreatment can alleviate oxidative damage. However, these protective effects induced by BL were blocked in AOA or SHAM pretreated seedlings (Figures 3E, 5). Moreover, application of ET resumes these protective effects induced by BL in AOA pretreated plants. In a word, our results indicated that BL induced ethylene play a protective role in plant photosystem against stress conditions.

## Ethylene Cooperates with H<sub>2</sub>O<sub>2</sub> to Enhance the BL-induced Plant Abiotic Stresses Tolerance

On the basis of recent reports, we know that H<sub>2</sub>O<sub>2</sub> and ethylene might function in AOX pathway induction under stresses (Wang et al., 2010b). In order to investigate the relationship of ethylene and H<sub>2</sub>O<sub>2</sub> in the BL-induced AOX capacity, NADPH oxidase inhibitor DPI, H<sub>2</sub>O<sub>2</sub> scavenger DMTU, ethylene biosynthesis inhibitor AOA, H<sub>2</sub>O<sub>2</sub>, and ethylene were used in this study. As shown in Figure 6, application of 5 mM DMTU, 100 μM DPI or 1 mM AOA to cucumber seedlings had significantly down-regulated on H<sub>2</sub>O<sub>2</sub> content and ethylene generation, while EL was increased compared with that of the BL treatment plants. However, the decline of H<sub>2</sub>O<sub>2</sub> content and ethylene generation were rescued in seedlings pretreated with ET or H<sub>2</sub>O<sub>2</sub>. These results demonstrated that ethylene cooperates with H<sub>2</sub>O<sub>2</sub> played important roles in the BR-induced abiotic stresses tolerance.

## Ethylene and H<sub>2</sub>O<sub>2</sub> are Located in the Upstream of BR-induced AOX Pathway to Enhance Plant Abiotic Stresses Tolerance

To examine whether there were other factors involved in the BR-induced AOX pathway, the respiration rate was investigated in the cucumber seedlings treated with BL, DMTU/DPI, AOA, and DMTU/DPI+AOA. As shown in Figure 7, compared with BL-pretreated plants, BL induced cyanide-resistant respiration was decreased in BL+DMTU/DPI and BL+AOA treated plants under environment stresses. Besides, the cyanide-resistant respiration rate of the plants treated with BL+DMTU/DPI+AOA was strongly inhibited but still increased



**FIGURE 7 | Effects of  $H_2O_2$  and ethylene inhibitors on BRs-induced AOX capacity in response to environmental stress. (A–D)** Plants were pretreated with  $100\ \mu\text{M}$  DPI,  $5\ \text{mM}$  DMTU for 8 h or  $1\ \text{mM}$  AOA for 12 h, then plants were treated with  $1\ \mu\text{M}$  BL 12 h later, plants were challenged with stress conditions for 3 d. Data are the mean  $\pm$  SD of three biological repeats; the significant difference was analyzed by Student's *t*-test ( $*P < 0.05$ ) and compared to control (CK).

slightly as compared with the control. These results indicated that ethylene and hydrogen peroxide were involved in the BR-induced AOX pathway, as well as other factors.

### BL Alleviates Oxidative Damage and Activates Antioxidant System in Cucumber


$H_2O_2$  is a vital type of ROS in seedling cells, other indicators such as EL and MDA content can also show the degree of oxidative damage in seedlings caused by stress conditions. Consistent with the observed damage of cucumber plants in **Figure 5D**, the BL-pretreated cucumber showed lighter stress damage than that of the control. To demonstrate these results,  $H_2O_2$  level and superoxide content were detected by DAB and NBT staining (**Figure 3E**), which were consistent with the phenotypes of seedlings. Meanwhile, under the same stress condition, the BL-pretreated plants maintained lower MDA and EL than the control (**Figures 5A,B**).

According to the recent research, the antioxidant defense machinery protects plants against oxidative stress damages (Foyer and Noctor, 2009; Xu et al., 2012a). As shown in **Figure 8**, all antioxidant enzyme activities increased in the presence or absence of BL under stress treatments. However, the activities of APX, GPX, and CAT were pronouncedly increased with BL pretreatment, and their levels were approximately 1.3-fold higher compared with the control seedlings after 3 d of stresses. Besides, the activity of POD in BL treated plants showed a similar trend

to the control. In addition, we determined the mRNA levels of antioxidant enzyme genes. The results indicated the transcripts of  $C_3POD24$ ,  $C_3SOD1$ , and  $C_3SOD2$  were much higher in the BL-treated cucumber seedlings than in the non-treated after 3 d of stresses (**Supplemental Figure S2**). It has been showed that the antioxidative system was involved in polyunsaturated fatty acid (PUFA) metabolism (Lee et al., 2005; Cho et al., 2012). Whether, BR could regulate PUFA metabolism will be investigated in the future.

### DISCUSSION

Under salt, PEG and cold-stresses, cucumber seedlings displayed enhanced CN-resistant respiration and ethylene content after BL treatment. In addition, transcriptional levels of the AOX and ethylene biosynthesis related genes were also induced in the BL-pretreated seedlings as compared to the control (**Figure 2**). These results suggested that BRs could enhance ethylene content and AOX capacity. Our results confirmed the previous reports that the enhanced AOX and ethylene contribute to plant abiotic stresses tolerance (Kendrick and Chang, 2008; Xu et al., 2012b). It was reported that ethylene biosynthesis pathway was connected with AOX capacity in tomato (Xu et al., 2012a). When ethylene level is low under normal condition, it is not sufficient in itself to induce AOX activity (Wang et al., 2010b). On the contrary, when ethylene content maintains at high level in seedlings pretreated




with BL under stress conditions, the activated ethylene signal by BRs can activate downstream targets which induce AOX activity (Figures 2, 7). In this study, CN-resistant respiration was largely impaired by AOA treatment (Figure 7). Thus, our results, together with previous research, suggest that ethylene is involved in BRs-induced AOX capacity.

In the present work, BL-treated plants showed a significant increase in  $F_v/F_m$  and a decline in NPQ under abiotic stress conditions for 3 d (Figure 4). However, these BRs-induced protective effects, expressed as  $F_v/F_m$  and NPQ were mostly inhibited if the seedlings were pre-treated with SHAM or AOA (Figure 4). Addition of ET rescued the decreased stress tolerance tendency by AOA treatment (Figures 3E, 4, 5). Our study demonstrates that ethylene is involved in BRs-induced AOX capacity. Under abiotic stress conditions, the inhibition of AOX and ethylene resulted in more dissipated excitation energy and lower level of photochemical efficiency (Figure 4), similar to the previous study (Giraud et al., 2008). Previous studies

showed that the mitochondrial AOX pathway might play a pivotal role in the protection of plants photosystem by alleviating the inhibition of the repair of the photodamaged PSII (Zhang et al., 2011; Deng et al., 2015), and inhibition of the alternative pathway led to decreases in photosynthetic rate in seedlings (Yoshida et al., 2006; Zhang et al., 2010). Thus, these findings, together with previous research, suggest that the enhancement of AOX and ethylene by BRs might contribute to balancing chloroplast-to-mitochondria electron transfer, and thus decrease ROS accumulation in cucumber seedlings.

Xia et al. (2011) have proved that  $\text{H}_2\text{O}_2$  play an important role in BRs-induced stress tolerance. Firstly, BR-induced stress tolerance was promoted by increased  $\text{H}_2\text{O}_2$  level. Secondly, either scavenging of  $\text{H}_2\text{O}_2$  or inhibiting  $\text{H}_2\text{O}_2$  generation abolished BR-induced stress tolerance.  $\text{H}_2\text{O}_2$ , as a signaling molecule, is involved in multiple plant response to environmental stresses (Neill et al., 2003; Mittler et al., 2004; Zhang et al., 2007; Wang et al., 2010b). Feng et al. (2008) and Wang et al. (2010a) also



confirmed  $H_2O_2$  could induce AP and AOX1 expression under chilling and salinity conditions. Recently, we studied whether  $H_2O_2$  is involved in the BRs-induced AOX capacity (Deng et al., 2015). Here, we confirmed that  $H_2O_2$  may play a vital role in BR-induced AOX activity in cucumber. To demonstrate this assumption, the respiration rate was investigated in the cucumber seedlings pretreated with 5 mM DMTU or 100  $\mu$ M DPI. Moreover, compared with BL-treated plants, cyanide-resistant respiration was weaker in BL+ DMTU/DPI pretreated plants under environment stresses. Thus, it indicated that  $H_2O_2$  played a role in BR-induced AOX activity in cucumber (Figures 6, 7). Further, investigations were conducted to make clear the relationship between  $H_2O_2$  and ethylene in AOX induction under abiotic stress conditions. The results showed that ethylene cooperated with  $H_2O_2$  to induce AOX capacity under stresses (Figure 6). According to the results presented here, also those reported previously (Ederli et al., 2006; Giraud et al., 2008; Wang et al., 2009, 2010b; Xia et al., 2009), we present a hypothetical pattern in cucumber seedlings describing the interrelationships among BL, ethylene, ROS, the AOX and abiotic stress tolerance. In this pattern, BL induces  $H_2O_2$  and ethylene generation, which subsequently enhances AOX capacity. The enhanced AOX activity can eliminate superfluous  $H_2O_2$  generation to avoid oxidative damage in plant cells, and then improve stress tolerance. Besides, we don't know if there are other factors such as transcription factors and kinase in BR signaling pathway involved in AOX regulation in addition to ethylene and  $H_2O_2$  (Figure 9). BRs may also regulate plant stress tolerance through affecting other mechanisms. Exogenous application of BRs improved the Cu and Cr stress tolerance by inducing endogenous ABA, auxin, and polyamine profiles in radish (*Raphanus sativus*), which suggested the potential

crosstalk between BRs and ABA or auxin in stress responses (Choudhary et al., 2012). Furthermore, BRs may affect other pathways to activate antioxidative enzyme (Choudhary et al., 2010). Epibrassinolide induces the change in indole-3-acetic acid, abscisic acid and polyamine concentrations and enhances antioxidant potential of radish seedlings under copper stress (Choudhary et al., 2011). Epibrassinolide ameliorates Cr stress tolerance via influencing the levels of indole-3-acetic acid, abscisic acid, polyamines and antioxidant system of radish seedlings.

In brief, suitable concentrations of BL positively affected the tolerance of plant to stress conditions. AOX may play an important role in the plant resistance to environment stresses. Our study showed that both ethylene and  $H_2O_2$  were involved in BRs-induced AOX activity. Thus, our results reveal a novel role of BRs in plant against environmental stresses and clarify the relationships between ethylene, AOX and ROS during environmental stress, although the detailed mechanism relations need further investigation. Future, investigations should focus on whether BRs could directly induced AOX activity by BR signaling.

## ACKNOWLEDGMENTS

This study was supported by the National Natural Science Foundation of China (31470342, 91417305, 31400211, 31570237), the National Basic Research Program of China (973 Program) (2015CB150100) and Sichuan Natural Science Foundation (2015JY0101, 2015JY0223).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2015.00982>

**Supplemental Figure S1 | (A)** Representative phenotypes of cucumber seedlings under stresses for 3 days. Cucumber were pretreated with 0.1, 0.5, 1, 5, or 10  $\mu$ M BL while the control plants were pretreated with distilled water for 12 h, respectively, and then all the seedlings were subjected to salt (200 mM NaCl), PEG (16% PEG 6000) and cold stresses (at 4°C) for 3 days. **(B)** Change in the content of  $H_2O_2$  in the leaves of cucumber plants, plants were pretreated with 1  $\mu$ M BL for 12 h, then plants subjected to stresses.

**Supplemental Figure S2 | Changes in the expression of antioxidant enzyme related genes in cucumber seedlings under stress conditions for 3 days.** Transcripts of *CsSOD1*, *CsSOD2*, *CsPOD2*, *CsPOD24*, *CsCAT*, and *CsAPX* were determined by qRT-PCR. Data are the mean  $\pm$  SD of three biological repeats; the significant difference was analyzed by Student's *t*-test ( $*P < 0.05$ ) and compared to control (CK).

**Supplemental Table S1 | Sequences of primers used in real time qRT-PCR experiments.**

## REFERENCES

Apel, K., and Hirt, H. (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* 55, 373–399. doi: 10.1146/annurev.arplant.55.031903.141701

Boyer, J. S. (1982). Plant productivity and environment. *Science* 218, 443–448.  
Cao, Y., Zhang, Z. W., Xue, L. W., Du, J. B., Shang, J., Xu, F., et al. (2009). Lack of salicylic acid in NahG Arabidopsis protects plants against moderate salt stress. *Z. Naturforsch. C* 64, 231–238. doi: 10.1515/znc-2009-3-414

- Cho, K., Kim, Y. C., Woo, J. C., Rakwal, R., Agrawal, G. K., Yoeun, S., et al. (2012). Transgenic expression of dual positional maize lipoxygenase-1 leads to the regulation of defense-related signaling molecules and activation of the antioxidative enzyme system in rice. *Plant Sci.* 185, 238–245. doi: 10.1016/j.plantsci.2011.10.016
- Choudhary, S. P., Bhardwaj, R., Gupta, B. D., Dutt, P., Gupta, R. K., Biondi, S., et al. (2010). Epibrassinolide induces changes in indole-3-acetic acid, abscisic acid and polyamine concentrations and enhances antioxidant potential of radish seedlings under copper stress. *Physiol. Plant.* 140, 280–296. doi: 10.1111/j.1399-3054.2010.01403.x
- Choudhary, S. P., Kanwar, M., Bhardwaj, R., Gupta, B. D., and Gupta, R. K. (2011). Epibrassinolide ameliorates Cr (VI) stress via influencing the levels of indole-3-acetic acid, abscisic acid, polyamines and antioxidant system of radish seedlings. *Chemosphere* 84, 592–600. doi: 10.1016/j.chemosphere.2011.03.056
- Choudhary, S. P., Oral, H. V., Bhardwaj, R., Yu, J. Q., and Tran, L. S. P. (2012). Interaction of brassinosteroids and polyamines enhances copper stress tolerance in *Raphanus sativus*. *J. Exp. Bot.* 63, 5659–5675. doi: 10.1093/jxb/ers219
- Clifton, R., Millar, A. H., and Whelan, J. (2006). Alternative oxidases in Arabidopsis: a comparative analysis of differential expression in the gene family provides new insights into function of non-phosphorylating bypasses. *Biochim. Biophys. Acta* 1757, 730–741. doi: 10.1016/j.bbabi.2006.03.009
- Dat, J. F., Lopez-Delgado, H., Foyer, C. H., and Scott, I. M. (1998). Parallel changes in H<sub>2</sub>O<sub>2</sub> and catalase during thermotolerance induced by salicylic acid or heat acclimation in mustard seedlings. *Plant Physiol.* 116, 1351–1357. doi: 10.1104/pp.116.4.1351
- Deng, X. G., Zhu, T., Zhang, D. W., and Lin, H. H. (2015). The alternative respiratory pathway is involved in brassinosteroid-induced environmental stress tolerance in *Nicotiana benthamiana*. *J. Exp. Bot.* 66, 6219–6232. doi: 10.1093/jxb/erv328
- Ederli, L., Moretini, R., Borgogni, A., Wasternack, C., Miersch, O., Reale, L., et al. (2006). Interaction between nitric oxide and ethylene in the induction of alternative oxidase in ozone-treated tobacco plants. *Plant Physiol.* 142, 595–608. doi: 10.1104/pp.106.085472
- Feng, H., Li, X., Duan, J., Li, H., and Liang, H. (2008). Chilling tolerance of wheat seedlings is related to an enhanced alternative respiratory pathway. *Crop Sci.* 48, 2381–2388. doi: 10.2135/cropsci2007.04.0232
- Foyer, C. H., and Noctor, G. (2009). Redox regulation in photosynthetic organisms: signaling, acclimation, and practical implications. *Antioxid. Redox Signal.* 11, 861–905. doi: 10.1089/ars.2008.2177
- Fu, F. Q., Zhang, D. W., Deng, X. G., Li, J. Y., Peng, X. J., Tang, H., et al. (2014). Role of plastid signals in modulating Arabidopsis responses to Cucumber mosaic virus. *Plant Growth Regul.* 75, 761–769. doi: 10.1007/s10725-014-9979-8
- Giraud, E., Ho, L. H., Clifton, R., Carroll, A., Estavillo, G., Tan, Y. F., et al. (2008). The absence of ALTERNATIVE OXIDASE1a in Arabidopsis results in acute sensitivity to combined light and drought stress. *Plant Physiol.* 147, 595–610. doi: 10.1104/pp.107.115121
- Giraud, E., Van Aken, O., Ho, L. H., and Whelan, J. (2009). The transcription factor ABI4 is a regulator of mitochondrial retrograde expression of ALTERNATIVE OXIDASE1a. *Plant Physiol.* 150, 1286–1296. doi: 10.1104/pp.109.139782
- Jiang, Y. P., Huang, L. F., Cheng, F., Zhou, Y. H., Xia, X. J., Mao, W. H., et al. (2013). Brassinosteroids accelerate recovery of photosynthetic apparatus from cold stress by balancing the electron partitioning, carboxylation and redox homeostasis in cucumber. *Physiol. Plant.* 148, 133–145. doi: 10.1111/j.1399-3054.2012.01696.x
- Kendrick, M. D., and Chang, C. (2008). Ethylene signaling: new levels of complexity and regulation. *Curr. Opin. Plant Biol.* 11, 479–485. doi: 10.1016/j.pbi.2008.06.011
- Lee, S. H., Ahn, S. J., Im, Y. J., Cho, K., Chung, G. C., Cho, B. H., et al. (2005). Differential impact of low temperature on fatty acid unsaturation and lipoxygenase activity in figleaf gourd and cucumber roots. *Biochem. Biophys. Res. Commun.* 330, 1194–1198. doi: 10.1016/j.bbrc.2005.03.098
- Lin, L. C., Hsu, J. H., and Wang, L. C. (2010). Identification of novel inhibitors of 1-aminocyclopropane-1-carboxylic acid synthase by chemical screening in *Arabidopsis thaliana*. *J. Biol. Chem.* 285, 33445–33456. doi: 10.1074/jbc.M110.132498
- Liu, W. J., Chen, Y. E., Tian, W. J., Du, J. B., Zhang, Z. W., Xu, F., et al. (2009). Dephosphorylation of photosystem II proteins and phosphorylation of CP29 in barley photosynthetic membranes as a response to water stress. *Biochim. Biophys. Acta* 1787, 1238–1245. doi: 10.1016/j.bbabi.2009.04.012
- McIntosh, L. (1994). Molecular biology of the alternative oxidase. *Plant Physiol.* 105, 781–786. doi: 10.1104/pp.105.3.781
- Mittler, R., Vanderauwera, S., Gollery, M., and Van Breusegem, F. (2004). Reactive oxygen gene network of plants. *Trends Plant Sci.* 9, 490–498. doi: 10.1016/j.tplants.2004.08.009
- Mizutani, F., Hirota, R., and Kadota, K. (1987). Cyanide metabolism linked with ethylene biosynthesis in ripening apple fruit. *J. Jpn. Soc. Hortic. Sci.* 39, 1–7. doi: 10.2503/jjshs.56.31
- Møller, I. M., Bérczi, A., Plas, L. H., and Lambers, H. (1988). Measurement of the activity and capacity of the alternative pathway in intact plant tissues: identification of problems and possible solutions. *Physiol. Plant.* 72, 642–649. doi: 10.1111/j.1399-3054.1988.tb09176.x
- Nakashita, H., Yasuda, M., Nitta, T., Asami, T., Fujioka, S., Arai, Y., et al. (2003). Brassinosteroid functions in a broad range of disease resistance in tobacco and rice. *Plant J.* 33, 887–898. doi: 10.1046/j.1365-313X.2003.01675.x
- Neill, S. J., Desikan, R., and Hancock, J. T. (2003). Nitric oxide signalling in plants. *New Phytol.* 159, 11–35. doi: 10.1046/j.1469-8137.2003.00804.x
- Pirrung, M., and Brauman, J. (1987). Involvement of cyanide in the regulation of ethylene biosynthesis. *Plant Physiol. Biochem.* 25, 55–61.
- Scandalios, J. G. (1993). Oxygen stress and superoxide dismutases. *Plant Physiol.* 101, 7–12.
- Solomos, T., and Laties, G. (1976). Induction by ethylene of cyanide-resistant respiration. *Biochem. Biophys. Res. Commun.* 70, 663–671. doi: 10.1016/0006-291X(76)91098-6
- Velikova, V., Yordanov, I., and Edreva, A. (2000). Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Sci.* 151, 59–66. doi: 10.1016/S0168-9452(99)00197-1
- Wang, H., Huang, J., and Bi, Y. (2010a). Induction of alternative respiratory pathway involves nitric oxide, hydrogen peroxide and ethylene under salt stress. *Plant Signal. Behav.* 5, 1636–1637. doi: 10.4161/psb.5.12.13775
- Wang, H., Huang, J., Liang, X., and Bi, Y. (2012). Involvement of hydrogen peroxide, calcium, and ethylene in the induction of the alternative pathway in chilling-stressed *Arabidopsis callus*. *Planta* 235, 53–67. doi: 10.1007/s00425-011-1488-7
- Wang, H., Liang, X., Huang, J., Zhang, D., Lu, H., Liu, Z., et al. (2010b). Involvement of ethylene and hydrogen peroxide in induction of alternative respiratory pathway in salt-treated *Arabidopsis calluses*. *Plant Cell Physiol.* 51, 1754–1765. doi: 10.1093/pcp/pcq134
- Wang, H., Liang, X., Wan, Q., Wang, X., and Bi, Y. (2009). Ethylene and nitric oxide are involved in maintaining ion homeostasis in *Arabidopsis callus* under salt stress. *Planta* 230, 293–307. doi: 10.1007/s00425-009-0946-y
- Wang, S. D., Zhu, F., Yuan, S., Yang, H., Xu, F., Shang, J., et al. (2011). The roles of ascorbic acid and glutathione in symptom alleviation to SA-deficient plants infected with RNA viruses. *Planta* 234, 171–181. doi: 10.1007/s00425-011-1391-2
- Xia, X. J., Wang, Y. J., Zhou, Y. H., Tao, Y., Mao, W. H., Shi, K., et al. (2009). Reactive oxygen species are involved in brassinosteroid-induced stress tolerance in cucumber. *Plant Physiol.* 150, 801–814. doi: 10.1104/pp.109.138230
- Xia, X. J., Zhou, Y. H., Ding, J., Shi, K., Asami, T., Chen, Z., et al. (2011). Induction of systemic stress tolerance by brassinosteroid in *Cucumis sativus*. *New Phytol.* 191, 706–720. doi: 10.1111/j.1469-8137.2011.03745.x
- Xu, F., Yuan, S., Zhang, D. W., Lv, X., and Lin, H. H. (2012a). The role of alternative oxidase in tomato fruit ripening and its regulatory interaction with ethylene. *J. Exp. Bot.* 63, 5705–5716. doi: 10.1093/jxb/ers226
- Xu, F., Zhang, D. W., Zhu, F., Tang, H., Lv, X., Cheng, J., et al. (2012b). A novel role for cyanide in the control of cucumber (*Cucumis sativus* L.) seedlings response to environmental stress. *Plant Cell Environ.* 35, 1983–1997. doi: 10.1111/j.1365-3040.2012.02531.x
- Yang, Y., Qi, M., and Mei, C. (2004). Endogenous salicylic acid protects rice plants from oxidative damage caused by aging as well as biotic and abiotic stress. *Plant J.* 40, 909–919. doi: 10.1111/j.1365-313X.2004.02267.x
- Yip, W. K., and Yang, S. F. (1988). Cyanide metabolism in relation to ethylene production in plant tissues. *Plant Physiol.* 88, 473–476. doi: 10.1104/pp.88.2.473

- Yoshida, K., Terashima, I., and Noguchi, K. (2006). Distinct roles of the cytochrome pathway and alternative oxidase in leaf photosynthesis. *Plant Cell Physiol.* 47, 22–31. doi: 10.1093/pcp/pci219
- Yoshida, K., Terashima, I., and Noguchi, K. (2007). Up-regulation of mitochondrial alternative oxidase concomitant with chloroplast over-reduction by excess light. *Plant Cell Physiol.* 48, 606–614. doi: 10.1093/pcp/pcm033
- Yoshida, K., Watanabe, C., Kato, Y., Sakamoto, W., and Noguchi, K. (2008). Influence of chloroplastic photo-oxidative stress on mitochondrial alternative oxidase capacity and respiratory properties: a case study with *Arabidopsis* yellow variegated 2. *Plant Cell Physiol.* 49, 592–603. doi: 10.1093/pcp/pcn031
- Zhang, D. W., Deng, X. G., Fu, F. Q., and Lin, H. H. (2015). Induction of plant virus defense response by brassinosteroids and brassinosteroid signaling in *Arabidopsis thaliana*. *Planta* 241, 875–885. doi: 10.1007/s00425-014-2218-8
- Zhang, D. W., Xu, F., Zhang, Z. W., Chen, Y. E., Du, J. B., Jia, S. D., et al. (2010). Effects of light on cyanide-resistant respiration and alternative oxidase function in *Arabidopsis* seedlings. *Plant Cell Environ.* 33, 2121–2131. doi: 10.1111/j.1365-3040.2010.02211.x
- Zhang, F., Wang, Y., Yang, Y., Wu, H., Wang, D., and Liu, J. (2007). Involvement of hydrogen peroxide and nitric oxide in salt resistance in the calluses from *Populus euphratica*. *Plant Cell Environ.* 30, 775–785. doi: 10.1111/j.1365-3040.2007.01667.x
- Zhang, L. T., Zhang, Z. S., Gao, H. Y., Xue, Z. C., Yang, C., Meng, X. L., et al. (2011). Mitochondrial alternative oxidase pathway protects plants against photoinhibition by alleviating inhibition of the repair of photodamaged PSII through preventing formation of reactive oxygen species in *Rumex K-1* leaves. *Physiol. Plant.* 143, 396–407. doi: 10.1111/j.1399-3054.2011.01514.x
- Zhang, X., Zhang, L., Dong, F., Gao, J., Galbraith, D. W., and Song, C. P. (2001). Hydrogen peroxide is involved in abscisic acid-induced stomatal closure in *Vicia faba*. *Plant Physiol.* 126, 1438–1448. doi: 10.1104/pp.126.4.1438
- Zhou, J., Wang, J., Li, X., Xia, X. J., Zhou, Y. H., Shi, K., et al. (2014). H<sub>2</sub>O<sub>2</sub> mediates the crosstalk of brassinosteroid and abscisic acid in tomato responses to heat and oxidative stresses. *J. Exp. Bot.* 65, 4371–4383. doi: 10.1093/jxb/eru217
- Zhu, T., Tan, W. R., Deng, X. G., Zheng, T., Zhang, D. W., and Lin, H. H. (2015). Effects of brassinosteroids on quality attributes and ethylene synthesis in postharvest tomato fruit. *Postharvest Biol. Technol.* 100, 196–204. doi: 10.1016/j.postharvbio.2014.09.016

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Wei, Deng, Zhu, Zheng, Li, Wu, Zhang and Lin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.