



Genomic Tools in Pea Breeding Programs: Status and Perspectives

Nadim Tayeh¹, Grégoire Aubert¹, Marie-Laure Pilet-Nayel², Isabelle Lejeune-Hénaut³, Thomas D. Warkentin⁴ and Judith Burstin^{1*}

¹ INRA, UMR1347 Agroécologie, Dijon, France, ² INRA, UMR1349 Institut de Génétique Environnement et Protection des Plantes, Le Rheu, France, ³ INRA, USC Institut Charles Viollette-Adaptation au Froid du Pois, Estrées-Mons, France, ⁴ Crop Development Centre, College of Agriculture and Bioresources, University of Saskatchewan, Saskatoon, SK, Canada

OPEN ACCESS

Edited by:

Janila Pasupuleti,
International Crops Research Institute
for the Semi Arid Tropics, India

Reviewed by:

Vikas Kumar Singh,
International Crops Research Institute
for the Semi-Arid Tropics, India

Antonio M. De Ron,
Misión Biológica de Galicia, CSIC,
Spain

*Correspondence:

Judith Burstin
judith.burstin@dijon.inra.fr

Specialty section:

This article was submitted to
Plant Genetics and Genomics,
a section of the journal
Frontiers in Plant Science

Received: 19 June 2015

Accepted: 09 November 2015

Published: 27 November 2015

Citation:

Tayeh N, Aubert G, Pilet-Nayel M-L, Lejeune-Hénaut I, Warkentin TD and Burstin J (2015) Genomic Tools in Pea Breeding Programs: Status and Perspectives. *Front. Plant Sci.* 6:1037.
doi: 10.3389/fpls.2015.01037

Pea (*Pisum sativum* L.) is an annual cool-season legume and one of the oldest domesticated crops. Dry pea seeds contain 22–25% protein, complex starch and fiber constituents, and a rich array of vitamins, minerals, and phytochemicals which make them a valuable source for human consumption and livestock feed. Dry pea ranks third to common bean and chickpea as the most widely grown pulse in the world with more than 11 million tons produced in 2013. Pea breeding has achieved great success since the time of Mendel's experiments in the mid-1800s. However, several traits still require significant improvement for better yield stability in a larger growing area. Key breeding objectives in pea include improving biotic and abiotic stress resistance and enhancing yield components and seed quality. Taking advantage of the diversity present in the pea genepool, many mapping populations have been constructed in the last decades and efforts have been deployed to identify loci involved in the control of target traits and further introgress them into elite breeding materials. Pea now benefits from next-generation sequencing and high-throughput genotyping technologies that are paving the way for genome-wide association studies and genomic selection approaches. This review covers the significant development and deployment of genomic tools for pea breeding in recent years. Future prospects are discussed especially in light of current progress toward deciphering the pea genome.

Keywords: pea (*Pisum sativum* L.), breeding targets, genetic diversity, genomic resources, genotyping platforms, genetic maps, QTL and association mapping

INTRODUCTION

Thanks to significant technological breakthroughs, pea genetics is rapidly evolving from conventional to large-scale molecular-assisted approaches to uncover the molecular bases of important traits and enhance breeding. Several reviews have been published recently that dealt with pea systematics, seed quality, and breeding (Burstin et al., 2011; Dahl et al., 2012; Bohra et al., 2014; Arnoldi et al., 2015; Smýkal et al., 2015; Varshney et al., 2015; Warkentin et al., 2015). The present review focuses on the genomic toolkit that was developed recently in pea thanks in part to next-generation sequencing technologies. This includes transcriptome, genotyping, and mapping resources that will pave the way to renewed pea breeding programs.

ECONOMIC IMPORTANCE, NUTRITIVE VALUE, GROWING REGIONS

Pea (*Pisum sativum* L.) is a major cool-season pulse crop and an essential component of sustainable cropping systems (Nemecek et al., 2008; Duc et al., 2010; Jensen et al., 2012). Significant agro-ecological services linked with its ability to develop symbiotic nitrogen fixation as well as its role as a break crop for pest and pathogen pressure reduction have been described (Nemecek and Kägi, 2007; Hayer et al., 2010; Macwilliam et al., 2014). In 2013, the vegetable pea production amounted to 17.43 Mt worldwide (FAOSTAT)¹ and dry pea represented the third most important pulse crop production after common bean and chickpea with 11.16 Mt produced worldwide (FAOSTAT).

Pea seeds are an important source of proteins and provide an exceptionally varied nutrient profile (for a review, Burstin et al., 2011): major constituents are starch (from 18.6 to 54.1%) and proteins (15.8–32.1%), followed by fibers (5.9–12.7%), sucrose (1.3–2.1%), and oil (0.6–5.5%). Seeds also contain minerals, vitamins, and micro-nutrients such as polyphenolics, saponins, α -galactosides, and phytic acids whose health-promoting effects are being tested (Bastianelli et al., 1998; Mitchell et al., 2009; Dahl et al., 2012; Marles et al., 2013; Arnoldi et al., 2015). Peas enter in human nutrition in a wide diversity of forms: fresh seedlings, immature pods, and seeds provide a green vegetable, and whole or ground dry seeds are cooked in various dishes. High quality starch, protein, or oligoside isolates are being extracted from dry pea seeds and whole seed structural and functional characteristics have been assessed for food improvement (Brummer et al., 2015). Because dry seeds contain little anti-nutritional factors, they are also introduced as a protein source mainly in monogastric diets without affecting growth and production traits (Stein et al., 2006; Laudadio et al., 2012; Dotas et al., 2014). Pea hay is used as fodder in ruminant diets (Bastida Garcia et al., 2011).

Pea is mainly cultivated in temperate regions of the world on well-drained and fertile soils. However, being distributed over all continents, the pea production area is characterized by a large range of pedo-climatic conditions. Indeed, China is the largest producer of vegetable peas (10.60 Mt, FAOSTAT) followed by India (4 Mt). Canada is the main producer of dry peas (3.85 Mt) followed by China (1.6 Mt), the Russian Federation (1.35 Mt), USA (0.71 Mt), India (0.60 Mt), France (0.50 Mt), and Ethiopia (0.40 Mt).

VARIABILITY OF GERMPLASM USED FOR BREEDING

Pea was domesticated by Neolithic farmers in the Fertile Crescent some 10,000 years ago (Willcox et al., 2009; Weiss and Zohary, 2011; Smykal, et al., 2014). Pea then spread rapidly toward south-west Asia, the Mediterranean basin, and Europe (Zohary, 1999). Probably linked with their large range of cultivation and the diversity of their use as food, feed, or fodder, pea landraces and varieties now exhibit an incredible diversity of forms and growing types, adapted to diverse environments,

cropping systems, and end-uses (Burstin et al., 2015). This vast diversity of cultivated forms is the major reservoir for present crop improvement. Different types of pea varieties have been developed for vegetable pea production, varying at major genes controlling seed and plant traits. For example, wrinkled seeds are associated with significant changes of seed composition, linked with starch synthesis modification (Wang et al., 2003). Various types of dry peas are also available that differ by their cotyledon color, plant architecture, or flowering time. In addition to this cultivated reservoir of diversity, wild peas can be crossed with cultivated peas. Ben-Ze'Ev and Zohary (1973) showed that chromosomal rearrangements among accessions from the different *Pisum* species and subspecies could cause partial sterility in hybrids. Recently, Bogdanova et al. (2014) have identified a nucleo-cytoplasmic incompatibility between a *P. sativum elatius* accession and cultivated peas. However, within the *Pisum* genus, wild *P. fulvum*, wild subspecies *P. sativum elatius*, and *P. sativum humile* as well as *P. abyssinicum*, a taxon cultivated in Ethiopia, are in most cases inter-crossable with *P. sativum sativum* as long as the cultivated pea is used as female donor (Ben-Ze'Ev and Zohary, 1973; Ochatt et al., 2004). Different authors have thus used *P. fulvum* as well as wild *P. sativum* subspecies as a source of alleles for important breeding traits, such as resistance to various fungal diseases (Barilli et al., 2010; Fondevilla et al., 2011; Jha et al., 2012) or to *Bruchus pisorum* L. (Clement et al., 2009).

TARGET TRAITS AND ACHIEVEMENTS THROUGH CONVENTIONAL BREEDING

Continued grain yield improvement is necessary for pea to remain an attractive option compared to cereals and oilseeds in crop rotations. Improving yield involves addressing many biotic and abiotic stresses, using a large set of strategies including diverse germplasm as parents, making many crosses, selecting for major gene traits under conditions conducive to selection, and yield testing of a large number of breeding lines. These stresses are specific to each growing region and/or growing type. However, fungal diseases are the major biotic stress in most cases, followed by various insects, viruses, and parasitic plants such as broomrape. Drought and heat stress at flowering are the main abiotic stresses, while frost, salinity, and early season flooding are diversely important according to growing types.

Important achievements were obtained in pea cultivars through conventional breeding over the past 20 years. Yield gains of approximately 2% per year have been achieved (Warkentin et al., 2015). Lodging resistance has been improved through deployment of the *afila* gene for semi-leafless type (Kujala, 1953; Goldenberg, 1965) and secondarily through selection for increased stem strength (Banniza et al., 2005). Powdery mildew resistance based on the single recessive gene *er-1* (Harland, 1948) has been widely deployed. Partial resistance to the Ascochyta blight complex has been achieved through pyramiding of genes with minor effects (Kraft et al., 1998). Resistance to pea weevil (*Bruchus pisorum* L.) identified in the secondary gene pool (*P. fulvum*) (Clement et al., 2002) was transferred into cultivated pea through backcrossing (Clement et al., 2009; Aryamanesh et al.,

¹FAOSTAT. <http://faostat3.fao.org/>. 10/06/2015.

2012). Cultivars adapted to winter sowing have been developed and deployed in Europe and north-west USA giving the potential for better yields because of a longer growing season, higher biomass production, and earlier maturity to avoid late season drought and heat stress (Hanocq et al., 2009). The introgression of the *Hr* allele which delays flower initiation until after the main winter freezing periods have passed (Lejeune-Hénaut et al., 2008) permitted to obtain some cultivars with notably improved winter hardiness. Field pea production for whole seed food markets requires appropriate seed visual quality. Quantitative inheritance, transgressive segregation, and moderately high heritability were observed for seed color, shape, and surface dimpling (Ubayasena et al., 2011) allowing for good progress in breeding. Seed protein concentration has been maintained in pea cultivars, even though overall seed yield has increased (Jha et al., 2013).

In some cases, achievements are at an earlier stage of deployment. Useful germplasms such as sources of resistance to various biotic and abiotic stresses have been identified and are currently being evaluated and introgressed. Research in France and USA has led to the identification and introgression of useful variation for resistance to *Aphanomyces* root rot; partial resistance controlled by several quantitative trait loci (QTLs) is being deployed (Pilet-Nayel et al., 2002, 2005; McGee et al., 2012). Improved stress tolerance has been identified in landrace accessions for toxicity to boron (Bagheri et al., 1994), salinity (Leonforte et al., 2013a), iron deficiency (Kabir et al., 2012), and for heat tolerance during flowering (Petkova et al., 2009). Selection for major gene resistance to pea seed-borne mosaic virus and potyviruses is now incorporated into breeding strategies (van Leur et al., 2007). Diversity in pea seed micronutrient concentration (Ray et al., 2014) and an approach to improving iron bioavailability for humans (Liu et al., 2015) have been described.

Future targets in pea breeding include (i) the optimization of pea interactions with Rhizobia, Mycorrhiza and other beneficial microorganisms in view of crop resilience against stresses, (ii) the adaptation of plant morphology and phenology to novel cropping systems, and (iii) the adaptation of seed composition to novel end-use application possibilities.

AVAILABLE GENOMIC RESOURCES IN PEA

The pea genome is organized in 7 pairs of chromosomes ($2n = 2x = 14$). Its haploid size is estimated at 4.45 Gb (Dolezel et al., 1998; Dolezel and Greilhuber, 2010; Praca-Fontes et al., 2014). It is largely dominated by mobile elements, mainly of the Ty3/gypsy family (Macas et al., 2007). This large genome size and high transposable element content have undoubtedly contributed to delay the development and availability of genomic tools in pea. Recently, several national and international programs have developed diverse valuable genomic resources by taking advantage of cutting-edge sequencing and genotyping technologies. These programs indicate the determination of the pea community to make rapid progress toward targeted and efficient molecular breeding exploiting the rich diversity of pea

germplasm and its wild relatives. An international consortium has been initiated in order to generate the full-sequence of the pea genome (Madou et al., 2015).

Developing wide collections of mapped and easy-to-use molecular markers is among the first steps of gene tagging and gene introgression strategies. Pea genetic maps started to be developed early: Wellensiek (1925) constructed a six-linkage group (LG) map and Lamprecht (1948) published a full map with 7 LGs (see Rozov et al., 1999, for review). Later on, various marker types were developed and numerous linkage maps originating from intra- or inter-subspecific crosses have been generated (Table 1). Decisive progresses were achieved with the availability of mapped SSR and SNP markers. Thanks to their multi-allelic nature, genomic (Ford et al., 2002; Lordin et al., 2005; Sun et al., 2014) and EST-based SSR markers (Burstin et al., 2001; Gong et al., 2010; De Caire et al., 2012; Kaur et al., 2012; Mishra et al., 2012) have been widely used for studying germplasm diversity (Baranger et al., 2004; Smýkal et al., 2008; Zong et al., 2009; Sarikamis et al., 2010) and bridging between different genetic maps. Today, SNPs are the markers of choice because of their abundance, easy-scoring, and amenability to high-throughput genotyping. SNPs were identified based on sequencing data from 4 (Leonforte et al., 2013b), 6 (Sindhu et al., 2014), 8 (Duarte et al., 2014), and up to 16 (Tayeh et al., 2015a) pea genotypes. Illumina GoldenGate (Deulvot et al., 2010; Leonforte et al., 2013b; Duarte et al., 2014; Sindhu et al., 2014), Infinium (Tayeh et al., 2015a), and Sequenom MassARRAY (Cheng et al., 2015) platforms have been deployed for SNP genotyping (Table 2). In total, at least 52 genetic maps have been constructed for different F_2 or recombinant inbred line (RIL) populations, comprising up to 8503 markers (Table 1). Consensus maps have been built in order to offer higher mapping resolution and better genome coverage (Table 1). These maps combine molecular data from 2 (Aubert et al., 2006; Hamon et al., 2011), 3 (Lordin et al., 2005; Sudheesh et al., 2014), 4 (Hamon et al., 2013; Duarte et al., 2014), 5 (Sindhu et al., 2014), 6 (Bordat et al., 2011), or 12 (Tayeh et al., 2015a) populations.

Uncovering the molecular bases underlying agriculturally important traits requires knowledge of the gene content of genomic regions controlling these traits of interest. Besides recently available genomic resources such as the pea gene atlas (Alves-Carvalho et al., 2015), whole-genome polymorphism data for multiple genotypes (see Table 2 for full description), BAC libraries developed for the genotypes Cameor (<http://cnrgv.toulouse.inra.fr/fr>) and PI 269818 (Coyne et al., 2007), researchers can count on the conserved synteny between pea and close species with available genome sequences. Gene-based rich individual and consensus maps have revealed connections between pea and *Medicago truncatula* (Choi et al., 2004; Aubert et al., 2006; Bordat et al., 2011; Leonforte et al., 2013b; Duarte et al., 2014; Sindhu et al., 2014; Tayeh et al., 2015a), *Lotus japonicus*, soybean (Bordat et al., 2011; Leonforte et al., 2013b; Tayeh et al., 2015a), pigeon pea (Leonforte et al., 2013b), chickpea (Leonforte et al., 2013b; Tayeh et al., 2015a), and lentil (Sindhu et al., 2014). Comprehensive understanding of shared syntenic blocks was, for instance, reported to be of great

TABLE 1 | Available individual and consensus genetic maps constructed for bi-parental populations and quantitative trait loci positioned on these maps.

Original cross	# ind.	# LGs	# markers	Map length (cM)§	Mapped traits	References
(A) F2 POPULATIONS						
Erygel × 661	174	12	69	550	Plant height, flowering time, number of nodes, resistance to Ascochyta blight	Dinewanger et al., 1994
JI1089 × JI296	—	—	—	—	Leaf resistance to <i>Mycosphaerella pinodes</i>	Clulow et al., 1991
Primo × OSU442-15	227	11	108	1369§	Dry seed weight, seed color, seed yield, yield components, flowering nodes, total node number	Timmerman-Vaughan et al., 1996, 2005; McCallum et al., 1997
Vinco × Hurst's Greenshaft	—	—	—	—	Resistance to <i>Pseudomonas syringae</i> pv. <i>pisi</i>	Hunter et al., 2001
Partridge × Early Onward	—	—	—	—	Resistance to <i>Pseudomonas syringae</i> pv. <i>pisi</i>	Hunter et al., 2001
G0003973 × G0005527	190	11	157	1518	—	Sun et al., 2014
A26 × Rovar	148	13	99	930§	Ascochyta blight resistance	Timmerman-Vaughan et al., 2004
NGB5539 × JI1794 ^a	92	—	—	—	Node of flower initiation	Weller et al., 2012
IFPI3260 × IFPI3251	94	9	146	1283.3	Resistance to <i>Uromyces pisi</i> (Pers.) Wint.	Barili et al., 2010
(B) RIL POPULATIONS						
Puget × 90-2079	127	13	377	1523	Partial Resistance to <i>Aphanomyces euteiches</i>	Pilet-Nayel et al., 2002, 2005; Lordin et al., 2005; Hamon et al., 2013
A88 × Rovar	133	11	96	1050§	Resistance to Ascochyta blight field epidemics, plant reproductive maturity	Timmerman-Vaughan et al., 2002
Carneval × MP1401	88	10	207	1274§	Lodging reaction, Plant height, Resistance to Mycosphaerella blight, grain yield, seed protein concentration, days to maturity	Tar'an et al., 2003; Tar'An et al., 2004
Wt11238 × Wt0557	110	11	91	853	Yield components, seed protein content	Krajewski et al., 2012
Wt10245 × Wt11238	101	12	191	1086	Stem length, internode number, yield components, seed protein content	Irzykowska et al., 2002; Irzykowska and Wolko, 2004; Krajewski et al., 2012
JI296 × DP	135	7	5277	552.2§	Resistance to <i>Mycosphaerella pinodes</i> , plant height, flowering date	Prioul et al., 2004; Lordin et al., 2005; Tayeh et al., 2015a
Champagne × Terese	164	7	7072	888.2§	Photoperiod response, frost tolerance	Lordin et al., 2005; Aubert et al., 2006; Lejeune-Hénaut et al., 2008; Dumont et al., 2009; Tayeh et al., 2015a
VavD265 × Cameor	211	7	6952	752.6§	Seed protein quality/quantity (PQL)	Bourgeois et al., 2011; Tayeh et al., 2015a
Ballet × Cameor	207	7	6796	854.5§	Flowering time, leaf appearance rate, radiation use efficiency, leaf area, leaf chlorophyll content, shoot length, seed weight, plant N nutrition indices (biomass, nitrogen content, symbiotic N fixation and nodule efficiency), seed protein quality/quantity	Bourgeois et al., 2010; Bourgeois et al., 2011; Tayeh et al., 2015a
VavD265 × Ballet	211	7	6188	850.1§	Seed protein quality/quantity	Bourgeois et al., 2011; Tayeh et al., 2015a
Cameor × Melrose	120	7	8503	736.3§	—	Tayeh et al., 2015a
Kazar × Cameor	84	7	7013	700.7§	—	Tayeh et al., 2015a
Kazar × Melrose	118	7	3917	682.5§	—	Tayeh et al., 2015a

(Continued)

TABLE 1 | Continued

Original cross	# ind.	# LGs	# markers	Map length (cM) [§]	Mapped traits	References
China × Cameor	129	7	7737	833.1 [§]	Flowering time, winter frost damage, branching type, leaf area, leaf chlorophyll content, plant height, yield components, plant biomass, harvest index and seed protein content	Deulvot et al., 2010; Klein et al., 2014; Tayeh et al., 2015a
Cameor × Sommette	144	7	5537	769.1 [§]	—	Tayeh et al., 2015a
Cameor × Cerise	120	7	7206	523.8 [§]	—	Tayeh et al., 2015a
Baccara × PI180693	178	7	4620	705.2 [§]	Aphanomyces root rot resistance, earliness at flowering	Hamon et al., 2011; Duarte et al., 2014; Tayeh et al., 2015a;
Terèse × K586 ^b	139	7	249	1113	Yield components, seed protein content, number, and volume of cotyledon cells, flowering time, plant height, number of basal branches, plant biomass and nitrogen content, nitrogen nutrition index, harvest index	Laucou et al., 1998; Loridon et al., 2005; Aubert et al., 2006; Burstin et al., 2007
Shawnee × Bohatyr	187	8	272	1716	Seed Mineral Content, partial resistance to Fusarium wilt Race 2	Loridon et al., 2005; McPhee et al., 2012
Orb × CDC Striker	90	7	431	—	Cotyledon bleaching resistance, visual seed quality traits	Ubayasena et al., 2010, 2011; Sindhu et al., 2014
Alfetta × CDC Bronco	120	11	223	450	Visual seed quality traits	Ubayasena et al., 2011
Kaspa × Parafield	134	9	868	1916	Salinity tolerance at the seedling stage	Leonforte et al., 2013b
DSP × 90-2131 ^c	111	9	168	1046	Earliness at flowering, plant height, Aphanomyces root rot resistance	Hamon et al., 2013
Kaspa × Yarum	106	13	821	1910	Powdery mildew resistance	Sudheesh et al., 2014
Kaspa × ps1771	106	9	852	1545	Powdery mildew resistance, boron tolerance	Sudheesh et al., 2014
JL15 × J399	77	9	348	1400 [§]	Resistance genes (<i>Pp1</i> and <i>Pp2</i>) to <i>Pseudomonas syringae</i> pv. <i>pisi</i>	Ellis et al., 1992; Hall et al., 1997; Hunter et al., 2001
JL281 × Jl399	53	10	421	2300 [§]	Resistance genes (<i>Pp2</i>) to <i>Pseudomonas syringae</i> pv. <i>pisi</i>	Ellis et al., 1992; Hall et al., 1997; Hunter et al., 2001
JL15 × JL1194	—	10	—	—	—	Ellis et al., 1992; Hall et al., 1997
JL15 × JL61	—	13	—	—	—	Ellis et al., 1992; Hall et al., 1997
JL1201 × JL813	—	3	—	—	—	Ellis et al., 1992; Hall et al., 1997
Carrera × CDC Striker	134	—	245	—	—	Sindhu et al., 2014
1-2347-144 × CDC Meadow ^d	163	7	367	437.2	phytic acid-phosphorus concentration	Sindhu et al., 2014; Shunmugam et al., 2015
Alfetta × P651	144	—	303	—	—	Sindhu et al., 2014
"Afghanistan" (<i>sym2</i>) × A1078-239	19	—	—	—	Tolerance to Fusarium root rot	Weeden and Porter, 2007
CMG × PI 220174	225	—	—	—	Tolerance to Fusarium root rot	Weeden and Porter, 2007
Carman × Reward ^e	71	—	—	—	Resistance to Fusarium root rot	Feng et al., 2011
HUVP 1 × FC 1	136	7	57	634	Resistance to <i>Uromyces fabae</i> (Pers.) de-Barry	Rai et al., 2011
MN313 × OSU1026	45	—	—	—	Tolerance to Aphanomyces euteiches	Weeden et al., 2000

(Continued)

TABLE 1 | Continued

Original cross	# ind.	# LGs	# markers	Map length (cM) [§]	Mapped traits	References
Baccara × K52 ^f	178	—	—	—	Aphanomyces root rot resistance, earliness at flowering	Hamon et al., 2011, 2013
P665 × Messire	362	8	414	1188	Resistance to <i>Mycosphaerella pinodes</i> , <i>Orobanche crenata</i> , <i>Pseudomonas syringae</i> , earliness, root length, aerial biomass, drought tolerance	Valderama et al., 2004; Fondevilla et al., 2008, 2011, 2012; Carrillo et al., 2012; Iglesias-Garcia et al., 2015
MN313 × JI 1794	47	9	242	—	Pod dehiscence	Weeden et al., 2002
JI 1794 × Slow	53	—	—	1289	Tolerance to <i>Fusarium solani</i> f. sp. Pisi; Pod dehiscence, dry seed weight	Timmerman-Vaughan et al., 1996; Weeden et al., 1998, 2002; Hance et al., 2004
Pennant × ATC113	270	9	155	2686	<i>Bruchus pisorum</i> resistance	Byrne et al., 2008; Aryamanesh et al., 2014
(C) CONSENSUS MAPS BASED ON RIL POPULATIONS						
Terese × K586; Champagne × Terese; Shawnee × Bohaty ^c	490	7	462	1430 [§]	—	Loridon et al., 2005
Terese × K586; Champagne × Terese	303	7	363	1458 [§]	—	Aubert et al., 2006
Terese × K586; Terese × Champagne; China × Cameor; Cameor × VavD265; Cameor × Ballet; Ballet × VavD265	1022	7	536	1389	—	Bordat et al., 2011
Baccara × PI 180693; Baccara × 552; Baccara × PI 180693; Baccara × 552; DSP × 90-2131; Puget × 90-2079	356	7	224	1652	—	Hamon et al., 2011
Orb × CDC Striker; Carrera × CDC Striker; 1-2347-144 × CDC Meadow; Alfetta × P651; China × Cameor	594	7	619	1513	—	Hamon et al., 2013
JI296 × DP; Champagne × Terese; China × Cameor; Baccara × PI180693	586	7	939	771.6	—	Sindhu et al., 2014
Kaspa × Yarum; Kaspa × ps1771; Kaspa × Parafield	360	7	2070	1255	—	Duarte et al., 2014
Champagne × Terese; VavD265 × Cameor; Ballet × Cameor; VavD265 × Ballet; Cameor × Meirose; Kazar × Cameor; Kazar × Meirose; China × Cameor; Cameor × Sommette; Cameor × Cerise; Baccara × PI180693; JI296 × DP	1384	7	15079	794.9 [§]	—	Sudheesh et al., 2014 ^g
						Tayeh et al., 2015a

^aNGB589 is an isogenic dwarf derivative of cv. Torsdag that carries a mutation in the gibberellin biosynthesis gene *LE*; line JI1794 is a representative accession of the northern race of *Pisum sativum* var. *humile*.

^bK586 is a ramified mutant obtained from "Torsdag" (Laucou et al., 1998).

^cDSP: Dark Skin Perfection.

^dLine 1-2347-144 was obtained from CDC Bronco through mutagenesis; it retains the majority of the traits of this latter (Varkentin et al., 2012).

^eReciprocal crosses between Cameor and Reward were made and seeds from both reciprocal crosses were used to develop an *F*₈ RIL.

^f552 is a garden pea breeding line derived from the 8th cycle of a recurrent selection program conducted for Aphanomyces root rot resistance (Hamon et al., 2011).

^gA total of 535 SNP markers from this map originates from transcript sequences common with the SNP markers from Duarte et al. (2014); merging the two datasets resulted in a 7-LG 2028-marker 2387-cM consensus map.

[§]Cf. Haldane.

AFLP, amplified fragment length polymorphism; RFLP, random fragment length polymorphism; RAPD, random amplified polymorphic DNA; SSR, simple sequence repeat; STS, sequence tagged site; ISSR, inter simple sequence repeat; CAPS, cleaved amplified polymorphic site; HRM, high-resolution melting; SNP, single nucleotide polymorphism.

TABLE 2 | Recently-developed resources in pea: transcriptome and whole-genome sequencing data, SNP collections, and genotyping platforms.

References	Plant material	Sequencing			Read assembly			Genotyping							
		Genotype(s) ^a	Organ(s)/ Tissue(s)	Contrasted growth condition(s) (if any)	Sequencing Platform	# reads for Assembly	# contigs and singletions	Mean contig length (bp)	Reference	Platform	Marker types	Reference genotype for SNP calling	# robust filtered SNPs	# markers on the genotyping array	
Aubert et al., 2006 Gene-specific	Cameor, Ballet, Champagne, Kazir, Melrose, VavD265, K886, Terese, other sources	Leaf	—	Sanger	—	—	—	—	Deußel et al., 2010	GoldenGate	Genomic, EST or cDNA-derived SNP	—	520	384	325
Franssen et al., 2011	Little Marvel	Cotyledon, seedling, hypocotyl, epicotyl, leaf, flower ^N	With or without 6-h light exposure of the seedlings	454 Roche GS FLX or GS20	2,209,735	MiRA	128,767 contigs and 195,661 singletions ^b	324	Transcriptome Shotgun Assembly Archive of GenBank; accession numbers: J898856-J981123	—	—	—	—	—	—
Kaur et al., 2012	Parafield, Yarum, Kaspa, 96-286	Seed, root, stem, leaf, flower, pod	—	454 Roche GS FLX Titanium	687,200	Next Gene software	13,583 contigs and 57,099 singletions ^c	719	Transcriptome Shotgun Assembly Archive of GenBank; accession numbers: JRE50256-JR96420	Leonforte et al., 2013b	GoldenGate Transcript-based SNP	Consensus reference	956	768	705
Duarie et al., 2014	Lumina, Hardy, Panache, Rocket, Kayanne, Terese, Cherokee, Champagne	Seedlings ^N	—	454 Roche GS FLX Titanium	3,042,418 (267,453-458,682 total reads per cultivar)	MiRA ("est" mode)	68,850 contigs ^d	842	Transcriptome Shotgun Assembly of GenBank under the accession GAM1000000000	Duartre et al., 2014	GoldenGate Transcript-based SNP	Consensus reference	35,455	1920	1620
Sindhu et al., 2014 ^e	CDC Bronco, Alfetta, Cooper, CDC striker, Nitouch, Giro, P651, and P135861 ^f	Seed, seedling, leaf, stem, flower	Normal or dark conditions for Titanium seedlings grown in Petri dishes	454 Roche GS FLX Titanium	4,008,648 (520,797-593,701 total 454 reads per cultivar)	NGen (DNAStar) software	29,725 contigs ^g	—	NCBI-NIH Short Read Archive, BioProject ID: PRJNA237986	Sindhu et al., 2014	GoldenGate Transcript-based SNP	Consensus reference	8822	1536	1491
Ferraro et al., 2014	Courter	Seed coat	—	454 Roche GS FLX Titanium	40,903	MiRA	5,766 contigs and 10,506 singletions	—	—	—	—	—	—	—	—
Alves-Carvalho et al., 2015	Cameor	Seed, root, nodule, stem, apical node, leaf, peduncle, tendril, flower	Hydroponic or aeroponic growth conditions with either high or low-nitrogen supply	Illumina Genome Analyzer II or HiSeq2000	1,018,751, 326	Velvet-Oases and TGICL++	46,099 contigs ^h	1199	http://bios.dillon.inra.fr/FATAI/cgi/pscam.cgi	—	—	—	—	—	—

(Continued)

TABLE 2 | Continued

References	Plant material		Sequencing		Read assembly		References	Platform	Marker types	Reference genotype for SNP calling	# robust filtered SNPs	# markers on the genotyping array	
	Genotype(s) ^a	Organ(s)/ Tissue(s)	Contrasted growth conditions (if any)	Sequencing Platform	# reads for assembly	# contigs and singlettons							
Burstin et al., 2014; Tayeh et al., 2015a Whole-genome	Cameor, Ballet, Cerise, Champagne, China, JI281, Kazar, Melrose, PH180693, Serpette, d'Auvergne, Sommette, TIL336/11, VaiD265, JI1703, JI2202, DG076 ^b	Leaf	–	Illumina HiSeq2000	184,764, 428,1,061, 737,954 total reads per accession ^c	181,125 contigs	756	http://www.ncbi.nlm.nih.gov/bioproject/PR:NA25605/	Tayeh et al., 2015a	Infinium Gene-space assembly-SNP	248,617	15,000	13,156

^aIn all cases, separate sequencing of cDNA libraries from different genotypes were undertaken.^bA second pass assembly is described with 81,449 total unigenes but it lacks about a third of contig annotations obtained for the first pass assembly.^cA total of 45,161 overlapping hits with contigs from Franssen et al. (2011) were identified (10,832 contigs [24%] and 34,329 singlettons [76%]).^d70,337 transcripts (86%) out of 81,449 from Franssen et al. (2011) and 12,776 (95%) out of 13,445 from Kaur et al. (2012) were reported to have a hit against assemblies from Duarte et al. (2014).^eA 3'-transcript profiling was performed: 3'-anchored cDNA libraries were generated and sequenced.^fP651 and P358610 are wild accessions; P651 (*P. fulvum*) and P358610 (*P. sativum abyssinicum*).^gAssembled using CDC Bronco reads only.^hHigh- and low-copy contigs were identified based on comparison of the Unigene set with pea genomic sequences.ⁱJI2202 is a *P. sativum abyssinicum* accession; JI1703, a *P. sativum elatius* accession; and DCG076, a *P. fulvum* accession.^jOnly reads from Cameor ($n = 1,061,737,954$) were used to generate a gene-space assembly of the pea genome.^kNormalization of cDNA libraries.

help to identify candidate genes for the control of seed size (D'Erfurth et al., 2012) and freezing tolerance (Tayeh et al., 2013a,b).

As other legume crops, pea is not easily amenable to genetic transformation (Warkentin et al., 2015): transformation occurs at low rates (Svabova and Griga, 2008) and plant regeneration is difficult. Fortunately, functional validation of candidate genes can benefit from a TILLING population developed from the genotype Cameor (Dalmais et al., 2008) and from the Virus Induced Gene Silencing (VIGS) methodology successfully adapted in pea (Grønlund et al., 2010; Pfleiger et al., 2013).

THE USE OF GENOMIC TOOLS IN MARKER-TRAIT ASSOCIATION STUDIES AND BREEDING PROGRAMS

Genetic maps have proven to be useful to uncover the molecular bases of monogenic characters such as Mendel's characters (see Ellis et al., 2011 for review) and also to decipher the determinism of complex agronomically-important traits (Table 1). QTLs responsible for the genetic control of yield-related traits, seed protein content, aerial and root architecture, and biotic and abiotic stress resistance have been detected under multiple environmental conditions and located on different maps (Table 1). In addition to QTL mapping analyses in bi-parental populations, association analyses have emerged as a complementary approach to dissect quantitative traits in pea by exploiting natural genetic diversity and ancestral recombination events characterizing germplasm collections. Diverse sets of cultivars with distinct geographic origins were used to determine associations between genetic markers and seed mineral nutrient concentration (Kwon et al., 2012; Cheng et al., 2015; Diapari et al., 2015), seed low-carbohydrate content (Cheng et al., 2015), seed lipid content (Ahmad et al., 2015), yield-related traits (Kwon et al., 2012), disease/pest resistance, and morphological traits such as flower color and seed coat color. Genome wide association mapping and subsequent allele mining could build on genomic resources reviewed herein and the phenotyping data available for diverse germplasm collections, as reviewed in Warkentin et al. (2015).

Specific markers linked to major genes were developed for use in breeding, especially for trypsin inhibitors in pea seeds (Page et al., 2002; Duc et al., 2004), flowering (Weller and Ortega, 2015), lodging resistance (Zhang et al., 2006) and resistance to diseases such as powdery mildew (Ghafoor and McPhee, 2012; Reddy et al., 2015), pea enation and seed borne mosaic virus (Frew et al., 2002; Jain et al., 2013), fusarium wilt (McClendon et al., 2002), Ascochyta blight (Jha et al., 2015), and rust (Barilli et al., 2010) (Supplementary Table 1). Some other resistance, flowering or seed composition genes were reviewed by Warkentin et al. (2015). Marker-assisted selection was conducted in early generation (F_2) breeding populations using markers linked in coupling to two major QTLs controlling lodging resistance and was demonstrated more efficient than

phenotypic selection (Zhang et al., 2006). Recently, marker-assisted backcrossing (MABC) was successfully used to introgress one to three of the seven main *Aphanomyces* root rot resistance QTLs (Hamon et al., 2013) into several recipient agronomic lines (Lavaud et al., 2015). Evaluation for resistance of the subsequent 157 BC5/6 Near Isogenic Lines (NILs) validated the effect of the major and some minor QTLs in controlled conditions and showed QTL x genetic background interactions. A MABC strategy was also used to introgress three frost tolerance QTLs among the main four QTLs identified by Lejeune-Hénaut et al. (2008). Field evaluations of 125 QTL-NILs validated the effect of these QTLs in the spring-type genetic background Eden (Hascoët et al., 2014). So far, marker-assisted construction of QTL-NILs has mainly allowed QTL effects to be validated. The rational use of these genetic regions in breeding strategies can now be considered in order to combine favorable alleles at complementary QTLs to improve multiple stress resistance in agronomic material. A list of the markers that should be useful for pea breeding is provided in Supplementary Table 1. In parallel to strategies considering the combination of individual QTLs, genomic selection seems a promising approach in pea, as first suggested by the prediction of the date of beginning of flowering and 1000 seed weight using a subset of 331 SNP markers genotyped in a reference collection of 372 pea accessions (Burstin et al., 2015). Increasing the marker coverage of the genome by using the newly-developed GenoPea 13.2K SNP Array (Tayeh et al., 2015a) further improved prediction accuracies (Tayeh et al., 2015b).

REFERENCES

- Ahmad, S., Kaur, S., Lamb-Palmer, N. D., Lefsrud, M., and Singh, J. (2015). Genetic diversity and population structure of *Pisum sativum* accessions for marker-trait association of lipid content. *Crop J.* 3, 238–245. doi: 10.1016/j.cj.2015.03.005
- Alves-Carvalho, S., Aubert, G., Carrère, S., Cruaud, C., Brochot, A.-L., Jacquin, F., et al. (2015). Full-length de novo assembly of RNA-seq data in pea (*Pisum sativum* L.) provides a gene expression atlas and gives insights into root nodulation in this species. *Plant J.* 84, 1–19. doi: 10.1111/tpj.12967
- Arnoldi, A., Zanoni, C., Lammi, C., and Boschin, G. (2015). The role of grain legumes in the prevention of hypercholesterolemia and hypertension. *CRC Crit. Rev. Plant Sci.* 34, 144–168. doi: 10.1080/07352689.2014.897908
- Aryamanesh, N., Byrne, O., Hardie, D. C., Khan, T., Siddique, K. H. M., and Yan, G. (2012). Large-scale density-based screening for pea weevil resistance in advanced backcross lines derived from cultivated field pea (*Pisum sativum*) and *Pisum fulvum*. *Crop Pasture Sci.* 63, 612–618. doi: 10.1071/CP12225
- Aryamanesh, N., Zeng, Y., Byrne, O., Hardie, D. C., Al-Subhi, A. M., Khan, T., et al. (2014). Identification of genome regions controlling cotyledon, pod wall/seed coat and pod wall resistance to pea weevil through QTL mapping. *Theor. Appl. Genet.* 127, 489–497. doi: 10.1007/s00122-013-2234-2
- Aubert, G., Morin, J., Jacquin, F., Lordin, K., Quillet, M. C., Petit, A., et al. (2006). Functional mapping in pea, as an aid to the candidate gene selection and for investigating synteny with the model legume *Medicago truncatula*. *Theor. Appl. Genet.* 112, 1024–1041. doi: 10.1007/s00122-005-0205-y
- Bagheri, A., Paull, J. G., and Rathjen, A. J. (1994). The response of *Pisum sativum* L. germplasm to high concentrations of soil boron. *Euphytica* 75, 9–17. doi: 10.1007/BF00024526
- Banniza, S., Hashemi, P., Warkentin, T. D., Vandenberg, A., and Davis, A. R. (2005). The relationships among lodging, stem anatomy, degree of lignification, and resistance to mycosphaerella blight in field pea (*Pisum sativum*). *Can. J. Bot.* 83, 954–967. doi: 10.1139/b05-044
- Baranger, A., Aubert, G., Arnaud, G., Lainé, A. L., Deniot, G., Potier, J., et al. (2004). Genetic diversity within *Pisum sativum* using protein- and PCR-based markers. *Theor. Appl. Genet.* 108, 1309–1321. doi: 10.1007/s00122-003-1540-5
- Barilli, E., Satovic, Z., Rubiales, D., and Torres, A. M. (2010). Mapping of quantitative trait loci controlling partial resistance against rust incited by *Uromyces pisi* (Pers.) Wint. in a *Pisum fulvum* L. intraspecific cross. *Euphytica* 175, 151–159. doi: 10.1007/s10681-010-0141-z
- Bastianelli, D., Grosjean, F., Peyronnet, C., Duparque, M., and Regnier, J. M. (1998). Feeding value of pea (*Pisum sativum*, L.) - 1 Chemical composition of different categories of pea. *Anim. Sci.* 67, 609–619. doi: 10.1017/S1357729800033051
- Bastida Garcia, J. L., González-Ronquillo, M., Domínguez Vara, I. A., Romero-Bernal, J., and Castelán Ortega, O. (2011). Effect of field pea (*Pisum sativum* L.) level on intake, digestion, ruminal fermentation and *in vitro* gas production in sheep fed maintenance diets. *Anim. Sci. J.* 82, 654–662. doi: 10.1111/j.1740-0929.2011.00884.x
- Ben-Ze'ev, N., and Zohary, D. (1973). Species relationships in the genus *Pisum* L. *Isr. J. Bot.* 22, 73–91.
- Bogdanova, V. S., Kosterin, O. E., and Yadrikhinskiy, A. K. (2014). Wild peas vary in their cross-compatibility with cultivated pea (*Pisum sativum* subsp. *sativum* L.) depending on alleles of a nuclear-cytoplasmic incompatibility locus. *Theor. Appl. Genet.* 127, 1163–1172. doi: 10.1007/s00122-014-2288-9
- Bohra, A., Pandey, M. K., Jha, U. C., Singh, B., Singh, I. P., Datta, D., et al. (2014). Genomics-assisted breeding in four major pulse crops of developing countries: present status and prospects. *Theor. Appl. Genet.* 127, 1263–1291. doi: 10.1007/s00122-014-2301-3
- Bordat, A., Savois, V., Nicolas, M., Salse, J., Chauveau, A., Bourgeois, M., et al. (2011). Translational genomics in legumes allowed placing *in silico* 5460

PERSPECTIVES

The gradually-developed genomic tools for pea now represent a rich resource for innovative strategies in both basic research and applied breeding. The large set of bi-parental interconnected populations segregating for diverse important agronomic traits, the individual and consensus genetic maps, the dense arrays of genetic markers, the high-throughput SNP genotyping tools, the BAC libraries, the TILLING population, and the whole-genome and transcriptome sequences from a large group of accessions should enhance significant advances in pea breeding in the next few years and foster the use of more diverse genetic resources for pea improvement. The genome sequence when released will further advance the pea genomic breeding revolution.

ACKNOWLEDGMENTS

The present work of NT, GA, IL, MP, JB was done in the frame of the French National Research Agency (ANR) Project Investissements d'Avenir PeaMUST under the grant number ANR-11-BTBR-0002.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2015.01037>

- unigenes on the pea functional map and identified candidate genes in *Pisum sativum* L. G3 1, 93–103. doi: 10.1534/g3.111.000349
- Bourgeois, M., Jacquin, F., Cassecuelle, F., Savoie, V., Belghazi, M., Aubert, G., et al. (2011). A PQL (protein quantity loci) analysis of mature pea seed proteins identifies loci determining seed protein composition. *Proteomics* 11, 1581–1594. doi: 10.1002/pmic.201000687
- Bourion, V., Rizvi, S. M. H., Fournier, S., de Larambergue, H., Galmiche, F., Marget, P., et al. (2010). Genetic dissection of nitrogen nutrition in pea through a QTL approach of root, nodule, and shoot variability. *Theor. Appl. Genet.* 121, 71–86. doi: 10.1007/s00122-010-1292-y
- Brummer, Y., Kaviani, M., and Tosh, S. M. (2015). Structural and functional characteristics of dietary fibre in beans, lentils, peas and chickpeas. *Food Res. Int.* 67, 117–125. doi: 10.1016/j.foodres.2014.11.009
- Burstin, J., Alves-Carvalho, S., Aluome, C., Tayeh, N., Brochot, A. L., Carrère, S., et al. (2014). “Recent pea genomic resources will enhance complementary improvement strategies in this crop,” in *IFLRC VI and ICLGG VII Conference* (Saskatoon, SK).
- Burstin, J., Deniot, G., Potier, J., Weinachter, C., Aubert, G., and Baranger, A. (2001). Microsatellite polymorphism in *Pisum sativum*. *Plant Breed.* 120, 311–317. doi: 10.1046/j.1439-0523.2001.00608.x
- Burstin, J., Gallardo, K., Mir, R. R., Varshney, R. K., and Duc, G. (2011). “Improving protein content and nutrition quality,” in *Biology and Breeding of Food Legumes*, eds A. Pratap and J. Kumar (Wallingford, CT: CAB International), 314–328. doi: 10.1079/9781845937669.0314
- Burstin, J., Marget, P., Huart, M., Moessner, A., Mangin, B., Duchene, C., et al. (2007). Developmental genes have pleiotropic effects on plant morphology and source capacity, eventually impacting on seed protein content and productivity in pea. *Plant Physiol.* 144, 768–781. doi: 10.1104/pp.107.096966
- Burstin, J., Salloignon, P., Chabert-Martinello, M., Magnin-Robert, J. B., Siol, M., Jacquin, F., et al. (2015). Genetic diversity and trait genomic prediction in a pea diversity panel. *BMC Genomics* 16:105. doi: 10.1186/s12864-015-1266-1
- Byrne, O. M., Hardie, D. C., Khan, T. N., Speijers, J., and Yan, G. (2008). Genetic analysis of pod and seed resistance to pea weevil in a *Pisum sativum* x *P. fulvum* interspecific cross. *Aust. J. Agric. Res.* 59, 854–862. doi: 10.1071/AR07353
- Carrillo, E., Rubiales, D., Pérez-De-Luque, A., and Fondevilla, S. (2012). Characterization of mechanisms of resistance against *Didymella pinodes* in *Pisum* spp. *Eur. J. Plant Pathol.* 135, 761–769. doi: 10.1007/s10658-012-0116-0
- Cheng, P., Holdsworth, W., Ma, Y., Coyne, C. J., Mazourek, M., Grusak, M. A., et al. (2015). Association mapping of agronomic and quality traits in USDA pea single-plant collection. *Mol. Breed.* 35, 75. doi: 10.1007/s11032-015-0277-6
- Choi, H.-K., Mun, J.-H., Kim, D.-J., Zhu, H., Baek, J.-M., Mudge, J., et al. (2004). Estimating genome conservation between crop and model legume species. *Proc. Natl. Acad. Sci. U.S.A.* 101, 15289–15294. doi: 10.1073/pnas.0402251101
- Clement, S. L., Hardie, D. C., and Elberson, L. R. (2002). Variation among accessions of *Pisum fulvum* for resistance to pea weevil. *Crop Sci.* 42, 2167–2173. doi: 10.2135/cropsci2002.2167
- Clement, S. L., McPhee, K. E., Elberson, L. R., and Evans, M. A. (2009). Pea weevil, *Bruchus pisorum* L. (Coleoptera: Bruchidae), resistance in *Pisum sativum* x *Pisum fulvum* interspecific crosses. *Plant Breed.* 128, 478–485. doi: 10.1111/j.1439-0523.2008.01603.x
- Clulow, S. A., Matthews, P., and Lewis, B. G. (1991). Genetic analysis of resistance to *Mycosphaerella pinodes* in pea seedlings. *Euphytica* 58, 183–189. doi: 10.1007/BF00022819
- Coyne, C. J., McClendon, M. T., Walling, J. G., Timmerman-Vaughan, G. M., Murray, S., Meksem, K., et al. (2007). Construction and characterization of two bacterial artificial chromosome libraries of pea (*Pisum sativum* L.) for the isolation of economically important genes. *Genome* 50, 871–875. doi: 10.1139/G07-063
- Dahl, W. J., Foster, L. M., and Tyler, R. T. (2012). Review of the health benefits of peas (*Pisum sativum* L.). *Br. J. Nutr.* 108, S3–S10. doi: 10.1017/s0007114512000852
- Dalmais, M., Schmidt, J., Le Signor, C., Moussy, F., Burstin, J., Savoie, V., et al. (2008). UTILLdb, a *Pisum sativum* in silico forward and reverse genetics tool. *Genome Biol.* 9:R43. doi: 10.1186/gb-2008-9-2-r43
- De Caire, J., Coyne, C. J., Brumett, S., and Shultz, J. L. (2012). Additional pea EST-SSR markers for comparative mapping in pea (*Pisum sativum* L.). *Plant Breed.* 131, 222–226. doi: 10.1111/j.1439-0523.2011.01917.x
- D’Erfurth, I., Le Signor, C., Aubert, G., Sanchez, M., Vernoud, V., Darchy, B., et al. (2012). A role for an endosperm-localized subtilase in the control of seed size in legumes. *New Phytol.* 196, 738–751. doi: 10.1111/j.1469-8137.2012.04296.x
- Deulvot, C., Charrel, H., Marty, A., Jacquin, F., Donnadieu, C., Lejeune-Hénaut, I., et al. (2010). Highly-multiplexed SNP genotyping for genetic mapping and germplasm diversity studies in pea. *BMC Genomics* 11:468. doi: 10.1186/1471-2164-11-468
- Diapari, M., Sindhu, A., Warkentin, T. D., Bett, K., and Tar’An, B. (2015). Population structure and marker-trait association studies of iron, zinc and selenium concentrations in seed of field pea (*Pisum sativum* L.). *Mol. Breed.* 35, 30. doi: 10.1007/s11032-015-0252-2
- Dirlewanger, E., Isaac, P. G., Ranade, S., Belajouza, M., Cousin, R., and de Vienne, D. (1994). Restriction fragment length polymorphism analysis of loci associated with disease resistance genes and developmental traits in *Pisum sativum* L. *Theor. Appl. Genet.* 88, 17–27. doi: 10.1007/BF00222388
- Dolezel, J., and Greilhuber, J. (2010). Nuclear genome size: are we getting closer? *Cytometry A* 77A, 635–642. doi: 10.1002/cyto.a.20915
- Dolezel, J., Greilhuber, J., Lucretti, S., Meister, A., Lysak, M. A., Nardi, L., et al. (1998). Plant genome size estimation by flow cytometry: inter-laboratory comparison. *Ann. Bot.* 82, 17–26. doi: 10.1006/anbo.1998.0730
- Dotas, V., Bampidis, V. A., Sinapis, E., Hatzipanagiotou, A., and Papanikolaou, K. (2014). Effect of dietary field pea (*Pisum sativum* L.) supplementation on growth performance, and carcass and meat quality of broiler chickens. *Livest. Sci.* 164, 135–143. doi: 10.1016/j.livsci.2014.03.024
- Duarte, J., Rivière, N., Baranger, A., Aubert, G., Burstin, J., Cornet, L., et al. (2014). Transcriptome sequencing for high throughput SNP development and genetic mapping in Pea. *BMC Genomics* 15:126. doi: 10.1186/1471-2164-15-126
- Duc, G., Blancard, S., Hénault, C., Lecomte, C., Petit, M., Bernicot, M.-H., et al. (2010). Potentiels et leviers pour développer la production et l’utilisation des protéagineux dans le cadre d’une agriculture durable en Bourgogne. *Innov. Agron.* 11, 157–171.
- Duc, G., Marget, P., Page, D., and Domoney, C. (2004). “Facile breeding markers to lower contents of vicine and convicine in faba bean seeds and trypsin inhibitors in pea seeds,” in *Recent Advances of Research in Antinutritional Factors in Legume Seeds and Oilseeds, Proceedings of the Fourth International Workshop on Antinutritional Factors in Legume Seeds and Oilseeds*, eds M. Muzquiz, G. D. Hill, C. Cuadrado, M. M. Pedrosa, and C. Burbano (Toledo), 281–285.
- Dumont, E., Fontaine, V., Vuylsteke, C., Sellier, H., Bodèle, S., Voedts, N., et al. (2009). Association of sugar content QTL and PQL with physiological traits relevant to frost damage resistance in pea under field and controlled conditions. *Theor. Appl. Genet.* 118, 1561–1571. doi: 10.1007/s00122-009-1004-7
- Ellis, T. H. N., Hofer, J. M. I., Timmerman-Vaughan, G. M., Coyne, C. J., and Hellens, R. P. (2011). Mendel, 150 years on. *Trends Plant Sci.* 16, 590–596. doi: 10.1016/j.tplants.2011.06.006
- Ellis, T. H. N., Turner, L., Hellens, R. P., Lee, D., Harker, C. L., Enard, C., et al. (1992). Linkage maps in pea. *Genetics* 130, 649–663.
- Feng, J., Hwang, R., Chang, K. F., Conner, R. L., Hwang, S. F., Strelkov, S. E., et al. (2011). Identification of microsatellite markers linked to quantitative trait loci controlling resistance to Fusarium root rot in field pea. *Can. J. Plant Sci.* 91, 199–204. doi: 10.4141/cjps09176
- Ferraro, K., Jin, A. L., Trinh-Don, N., Reinecke, D. M., Ozga, J. A., and Ro, D.-K. (2014). Characterization of proanthocyanidin metabolism in pea (*Pisum sativum*) seeds. *BMC Plant Biol.* 14:238. doi: 10.1186/s12870-014-0238-y
- Fondevilla, S., Almeida, N. F., Satovic, Z., Rubiales, D., Vaz Patto, M. C., Cubero, J. I., et al. (2011). Identification of common genomic regions controlling resistance to *Mycosphaerella pinodes*, earliness and architectural traits in different pea genetic backgrounds. *Euphytica* 182, 43–52. doi: 10.1007/s10681-011-0460-8
- Fondevilla, S., Martin-Sanz, A., Satovic, Z., Dolores Fernandez-Romero, M., Rubiales, D., and Caminero, C. (2012). Identification of quantitative trait loci involved in resistance to *Pseudomonas syringae* pv. syringae in pea (*Pisum sativum* L.). *Euphytica* 186, 805–812. doi: 10.1007/s10681-011-0592-x
- Fondevilla, S., Satovic, Z., Rubiales, D., Moreno, M. T., and Torres, A. M. (2008). Mapping of quantitative trait loci for resistance to *Mycosphaerella pinodes* in *Pisum sativum* subsp *syriacum*. *Mol. Breed.* 21, 439–454. doi: 10.1007/s11032-007-9144-4

- Ford, R., Le Roux, K., Itman, C., Brouwer, J. B., and Taylor, P. W. J. (2002). Diversity analysis and genotyping in *Pisum* with sequence tagged microsatellite site (STMS) primers. *Euphytica* 124, 397–405. doi: 10.1023/A:1015752907108
- Franssen, S. U., Shrestha, R. P., Bräutigam, A., Bornberg-Bauer, E., and Weber, A. P. M. (2011). Comprehensive transcriptome analysis of the highly complex *Pisum sativum* genome using next generation sequencing. *BMC Genomics* 12:227. doi: 10.1186/1471-2164-12-227
- Frew, T. J., Russell, A. C., and Timmerman-Vaughan, G. M. (2002). Sequence tagged site markers linked to the *sbm1* gene for resistance to pea seedborne mosaic virus in pea. *Plant Breed.* 121, 512–516. doi: 10.1046/j.1439-0523.2002.00761.x
- Ghafoor, A., and McPhee, K. (2012). Marker assisted selection (MAS) for developing powdery mildew resistant pea cultivars. *Euphytica* 186, 593–607. doi: 10.1007/s10681-011-0596-6
- Goldenberg, J. B. (1965). "Afila" a new mutation in pea (*Pisum sativum* L.). *Bole Genetico* 1, 27–31.
- Gong, Y. M., Xu, S. C., Mao, W. H., Hu, Q. Z., Zhang, G. W., Ding, J., et al. (2010). Developing new SSR markers from ESTs of pea (*Pisum sativum* L.). *J. Zhejiang Univ. Sci. B* 11, 702–707. doi: 10.1631/jzus.B100004
- Grønlund, M., Olsen, A., Johansen, E. I., and Jakobsen, I. (2010). Protocol: using virus-induced gene silencing to study the arbuscular mycorrhizal symbiosis in *Pisum sativum*. *Plant Methods* 6:28. doi: 10.1186/1746-4811-6-28
- Hall, K. J., Parker, J. S., Ellis, T. H. N., Turner, L., Knox, M. R., Hofer, J. M. I., et al. (1997). The relationship between genetic and cytogenetic maps of pea. II. Physical maps of linkage mapping populations. *Genome* 40, 755–769. doi: 10.1139/g97-798
- Hamon, C., Baranger, A., Coyne, C. J., McGee, R. J., Le Goff, I., L'Anthoëne, V., et al. (2011). New consistent QTL in pea associated with partial resistance to *Aphanomyces euteiches* in multiple French and American environments. *Theor. Appl. Genet.* 123, 261–281. doi: 10.1007/s00122-011-1582-z
- Hamon, C., Coyne, C. J., McGee, R. J., Lesné, A., Esnault, R., Mangin, P., et al. (2013). QTL meta-analysis provides a comprehensive view of loci controlling partial resistance to *Aphanomyces euteiches* in four sources of resistance in pea. *BMC Plant Biol.* 13:45. doi: 10.1186/1471-2229-13-45
- Hance, S. T., Grey, W., and Weeden, N. F. (2004). Identification of tolerance to *Fusarium solani* in *Pisum sativum* ssp. *elatius*. *Pisum Genet.* 36, 9–13.
- Hanocq, E., Jeuffroy, M. H., Lejeune-Hénaut, I., and Munier-Jolain, N. (2009). Construire des idéotypes pour des systèmes de culture variés en pois d'hiver. *Innov. Agron.* 7, 14–28.
- Harland, S. C. (1948). Inheritance of immunity to mildew in Peruvian forms of *Pisum sativum*. *Heredity* 2, 263–269. doi: 10.1038/hdy.1948.15
- Hascoët, E., Jaminon, O., Devaux, C., Blasius, C., Bahrman, N., Bochard, A.-M., et al. (2014). "Towards fine mapping of frost tolerance QTLs in pea," in *2nd PeaMUST Annual Meeting* (Dijon).
- Hayer, F., Bonnin, E., Carrouée, B., Gaillard, G., Nemecek, T., Schneider, A., et al. (2010). "Designing sustainable crop rotations using Life Cycle assessment of crop combinations," in *9th European IFSO Symposium* (Vienna).
- Hunter, P. J., Ellis, N., and Taylor, J. D. (2001). Association of dominant loci for resistance to *Pseudomonas syringae* pv. *pisii* with linkage groups II, VI and VII of *Pisum sativum*. *Theor. Appl. Genet.* 103, 129–135. doi: 10.1007/s001220100566
- Iglesias-Garcia, R., Prats, E., Fondevilla, S., Satovic, Z., and Rubiales, D. (2015). Quantitative trait loci associated to drought adaptation in pea (*Pisum sativum* L.). *Plant Mol. Biol. Rep.* 1–11. doi: 10.1007/s11105-015-0872-z
- Irzykowska, L., and Wolko, B. (2004). Interval mapping of QTLs controlling yield-related traits and seed protein content in *Pisum sativum*. *J. Appl. Genet.* 45, 297–306.
- Irzykowska, L., Wolko, B., and Swiecicki, W. (2002). Interval mapping of QTLs controlling some morphological traits in pea. *Cell. Mol. Biol. Lett.* 7, 417–422.
- Jain, S., Weeden, N. F., Porter, L. D., Eigenbrode, S. D., and McPhee, K. (2013). Finding linked markers to aid for efficient selection of pea enation mosaic virus resistance in pea. *Crop Sci.* 53, 2392–2399. doi: 10.2135/cropsci2013.04.0211
- Jensen, E. S., Peoples, M. B., Boddey, R. M., Gresshoff, P. M., Hauggaard-Nielsen, H., Alves, B. J. R., et al. (2012). Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. A review. *Agron. Sust. Dev.* 32, 329–364. doi: 10.1007/s13593-011-0056-7
- Jha, A. B., Arganosa, G., Tar'An, B., Diederichsen, A., and Warkentin, T. D. (2013). Characterization of 169 diverse pea germplasm accessions for agronomic performance, *Mycosphaerella* blight resistance and nutritional profile. *Genet. Resour. Crop Evol.* 60, 747–761. doi: 10.1007/s10722-012-9871-1
- Jha, A. B., Tar'An, B., Diapari, M., Sindhu, A., Shunmugam, A., Bett, K., et al. (2015). Allele diversity analysis to identify SNPs associated with ascochyta blight resistance in pea. *Euphytica* 202, 189–197. doi: 10.1007/s10681-014-1254-6
- Jha, A. B., Warkentin, T. D., Gurusamy, V., Tar'An, B., and Banniza, S. (2012). Identification of *Mycosphaerella* blight resistance in wild *Pisum* species for use in pea breeding. *Crop Sci.* 52, 2462–2468. doi: 10.2135/cropsci2012.04.0242
- Kabir, A. H., Paltridge, N. G., Able, A. J., Paull, J. G., and Stangoulis, J. C. R. (2012). Natural variation for Fe-efficiency is associated with upregulation of Strategy I mechanisms and enhanced citrate and ethylene synthesis in *Pisum sativum* L. *Planta* 235, 1409–1419. doi: 10.1007/s00425-011-1583-9
- Kaur, S., Pemberton, L. W., Cogan, N. O. I., Savin, K. W., Leonforte, T., Paull, J., et al. (2012). Transcriptome sequencing of field pea and faba bean for discovery and validation of SSR genetic markers. *BMC Genomics* 13:104. doi: 10.1186/1471-2164-13-104
- Klein, A., Houtin, H., Rond, C., Marget, P., Jacquin, F., Boucherot, K., et al. (2014). QTL analysis of frost damage in pea suggests different mechanisms involved in frost tolerance. *Theor. Appl. Genet.* 127, 1319–1330. doi: 10.1007/s00122-014-2299-6
- Kraft, J. M., Dunne, B., Goulden, D., and Armstrong, S. (1998). A search for resistance in peas to *Mycosphaerella pinodes*. *Plant Dis.* 82, 251–253. doi: 10.1094/PDIS.1998.82.2.251
- Krajewski, P., Bocianowski, J., Gawłowska, M., Kaczmarek, Z., Pniewski, T., Świecicki, W., et al. (2012). QTL for yield components and protein content: a multi-environment study of two pea (*Pisum sativum* L.) populations. *Euphytica* 183, 323–336. doi: 10.1007/s10681-011-0472-4
- Kujala, V. (1953). Felderbse, bei welcher die ganze Blattspreite in Ranken umgewandelt ist. *Arch. Soc. Zool. Bot. Fenn.* 8, 44–45.
- Kwon, S.-J., Brown, A. F., Hu, J., McGee, R., Watt, C., Kisha, T., et al. (2012). Genetic diversity, population structure and genome-wide marker-trait association analysis emphasizing seed nutrients of the USDA pea (*Pisum sativum* L.) core collection. *Genes Genomics* 34, 305–320. doi: 10.1007/s13258-011-0213-z
- Lamprecht, H. (1948). The variation of linkage and the course of crossing-over. *Agri Hort. Genet.* 6, 10–48
- Laucou, V., Haurogné, K., Ellis, N., and Rameau, C. (1998). Genetic mapping in pea. I. RAPD-based genetic linkage map of *Pisum sativum*. *Theor. Appl. Genet.* 97, 905–915. doi: 10.1007/s00120050971
- Laudadio, V., Nahashon, S. N., and Tufarelli, V. (2012). Growth performance and carcass characteristics of guinea fowl broilers fed micronized-dehulled pea (*Pisum sativum* L.) as a substitute for soybean meal. *Poult. Sci.* 91, 2988–2996. doi: 10.3382/ps.2012-02473
- Lavaud, C., Lesné, A., Piriou, C., Le Roy, G., Boutet, G., Moussart, A., et al. (2015). Validation of QTL for resistance to *Aphanomyces euteiches* in different pea genetic backgrounds using near-isogenic lines. *Theor. Appl. Genet.* 128, 2273–2288. doi: 10.1007/s00122-015-2583-0
- Lejeune-Hénaut, I., Hanocq, E., Bethencourt, L., Fontaine, V., Delbreil, B., Morin, J., et al. (2008). The flowering locus *Hr* colocalizes with a major QTL affecting winter frost tolerance in *Pisum sativum* L. *Theor. Appl. Genet.* 116, 1105–1116. doi: 10.1007/s00122-008-0739-x
- Leonforte, A., Forster, J. W., Redden, R. J., Nicolas, M. E., and Salisbury, P. A. (2013a). Sources of high tolerance to salinity in pea (*Pisum sativum* L.). *Euphytica* 189, 203–216. doi: 10.1007/s10681-012-0771-4
- Leonforte, A., Sudheesh, S., Cogan, N. O. I., Salisbury, P. A., Nicolas, M. E., Materne, M., et al. (2013b). SNP marker discovery, linkage map construction and identification of QTLs for enhanced salinity tolerance in field pea (*Pisum sativum* L.). *BMC Plant Biol.* 13:161. doi: 10.1186/1471-2229-13-161
- Liu, X., Glahn, R. P., Arganosa, G. C., and Warkentin, T. D. (2015). Iron bioavailability in low phytate pea. *Crop Sci.* 55, 320–330. doi: 10.2135/cropsci2014.06.0412
- Loridon, K., McPhee, K., Morin, J., Dubreuil, P., Pilet-Nayel, M. L., Aubert, G., et al. (2005). Microsatellite marker polymorphism and mapping in pea (*Pisum*

- sativum* L.). *Theor. Appl. Genet.* 111, 1022–1031. doi: 10.1007/s00122-005-0014-3
- Macas, J., Neumann, P., and Navrátilová, A. (2007). Repetitive DNA in the pea (*Pisum sativum* L.) genome: comprehensive characterization using 454 sequencing and comparison to soybean and *Medicago truncatula*. *BMC Genomics* 8:427. doi: 10.1186/1471-2164-8-427
- Macwilliam, S., Wismer, M., and Kulshreshtha, S. (2014). Life cycle and economic assessment of Western Canadian pulse systems: the inclusion of pulses in crop rotations. *Agric. Syst.* 123, 43–53. doi: 10.1016/j.agsy.2013.08.009
- Madoui, M. A., Labadie, K., Aury, J. M., Coyne, C. J., Warkentin, T., Kreplak, J., et al. (2015). “The international pea genome sequencing project: sequencing and assembly progresses,” in *PLANT and ANIMAL Genome XXIII* (San Diego, CA).
- Marles, M. A. S., Warkentin, T. D., and Bett, K.E. (2013). Genotypic abundance of carotenoids and polyphenolics in the hull of field pea (*Pisum sativum* L.). *J. Sci. Food Agric.* 93, 463–470. doi: 10.1002/jsfa.5782
- McCallum, J. A., Frew, T. J., Gilpin, B. J., and Timmerman-Vaughan, G. M. (1997). A linkage map of the pea (*Pisum sativum* L.) genome containing cloned sequences of known function and expressed sequence tags (ESTs). *Theor. Appl. Genet.* 95, 1289–1299. doi: 10.1007/s001220050695
- McClendon, M. T., Inglis, D. A., McPhee, K. E., and Coyne, C. J. (2002). DNA markers linked to Fusarium wilt race 1 resistance in pea. *J. Am. Soc. Hortic. Sci.* 127, 602–607.
- McGee, R. J., Coyne, C. J., Pilet-Nayel, M.-L., Moussart, A., Tivoli, B., Baranger, A., et al. (2012). Registration of pea germplasm lines partially resistant to aphanomyces root rot for breeding fresh or freezer pea and dry pea types. *J. Plant Regist.* 6, 203–207. doi: 10.3198/jpr2011.03.0139crg
- McPhee, K. E., Inglis, D. A., Gundersen, B., and Coyne, C. J. (2012). Mapping QTL for Fusarium wilt Race 2 partial resistance in pea (*Pisum sativum*). *Plant Breed.* 131, 300–306. doi: 10.1111/j.1439-0523.2011.01938.x
- Mishra, R. K., Gangadhar, B. H., Nookaraju, A., Kumar, S., and Park, S. W. (2012). Development of EST-derived SSR markers in pea (*Pisum sativum*) and their potential utility for genetic mapping and transferability. *Plant Breed.* 131, 118–124. doi: 10.1111/j.1439-0523.2011.01926.x
- Mitchell, D. C., Lawrence, F. R., Hartman, T. J., and Curran, J. M. (2009). Consumption of dry beans, peas, and lentils could improve diet quality in the US population. *J. Am. Diet. Assoc.* 109, 909–913. doi: 10.1016/j.jada.2009.02.029
- Nemecek, T., and Kägi, T. (2007). “Life Cycle Inventories of Swiss and European agricultural production systems,” in *Final Report Ecoinvent V2. 0 No 15a. Agroscope Reckenholz-Tänikon Research Station ART. Zurich; Dübendorf, CH: Swiss Centre for Life Cycle Inventories. SimaPro Ph.D.*, 7.
- Nemecek, T., Von Richthofen, J.-S., Dubois, G., Casta, P., Charles, R., and Pahl, H. (2008). Environmental impacts of introducing grain legumes into European crop rotations. *Eur. J. Agron.* 28, 380–393. doi: 10.1016/j.eja.2007.11.004
- Ochatt, S. J., Benabdelmouna, A., Marget, P., Aubert, G., Moussy, F., Pontecaille, C., et al. (2004). Overcoming hybridization barriers between pea and some of its wild relatives. *Euphytica* 137, 353–359. doi: 10.1023/B:EUPH.0000040476.57938.81
- Page, D., Aubert, G., Duc, G., Welham, T., and Domoney, C. (2002). Combinatorial variation in coding and promoter sequences of genes at the Tri locus in *Pisum sativum* accounts for variation in trypsin inhibitor activity in seeds. *Mol. Genet. Genomics* 267, 359–369. doi: 10.1007/s00438-002-0667-4
- Petkova, V., Nikolova, V., Kalapchieva, S. H., Stoeva, V., Topalova, E., and Angelova, S. (2009). Physiological response and pollen viability of *Pisum sativum* genotypes under high temperature influence. *Iv Balkan Symp. Veg. Potato.* 830, 665–671. doi: 10.17660/actahortic.2009.830.96
- Pflieger, S., Richard, M. M. S., Blanchet, S., Meziadi, C., and Geffroy, V. (2013). VIGS technology: an attractive tool for functional genomics studies in legumes. *Funct. Plant Biol.* 40, 1234–1248. doi: 10.1071/FP13089
- Pilet-Nayel, L., Muehlbauer, F. J., McGee, R. J., Kraft, J. M., Baranger, A., and Coyne, C. J. (2002). Quantitative trait loci for partial resistance to Aphanomyces root rot in pea. *Theor. Appl. Genet.* 106, 28–39.
- Pilet-Nayel, M. L., Muehlbauer, F. J., McGee, R. J., Kraft, J. M., Baranger, A., and Coyne, C. J. (2005). Consistent quantitative trait loci in pea for partial resistance to *Aphanomyces euteiches* isolates from the United States and France. *Phytopathology* 95, 1287–1293. doi: 10.1094/PHYTO-95-1287
- Praca-Fontes, M. M., Carvalho, C. R., and Clarindo, W. R. (2014). Karyotype revised of *Pisum sativum* using chromosomal DNA amount. *Plant Syst. Evol.* 300, 1621–1626. doi: 10.1007/s00606-014-0987-y
- Prioul, S., Frankewitz, A., Deniot, G., Morin, G., and Baranger, A. (2004). Mapping of quantitative trait loci for partial resistance to *Mycosphaerella pinodes* in pea (*Pisum sativum* L.), at the seedling and adult plant stages. *Theor. Appl. Genet.* 108, 1322–1334. doi: 10.1007/s00122-003-1543-2
- Rai, R., Singh, A. K., Singh, B. D., Joshi, A. K., Chand, R., and Srivastava, C. P. (2011). Molecular mapping for resistance to pea rust caused by *Uromyces fabae* (Pers.) de-Bary. *Theor. Appl. Genet.* 123, 803–813. doi: 10.1007/s00122-011-1628-2
- Ray, H., Bett, K., Tar'An, B., Vandenberg, A., Thavarajah, D., and Warkentin, T. (2014). Mineral micronutrient content of cultivars of field pea, chickpea, common bean, and lentil grown in Saskatchewan, Canada. *Crop Sci.* 54, 1698–1708. doi: 10.2135/cropsci2013.08.0568
- Reddy, D. C. L., Preethi, B., Wani, M. A., Aghora, T. S., Aswath, C., and Mohan, N. (2015). Screening for powdery mildew (*Erysiphe pisi* DC) resistance gene-linked SCAR and SSR markers in five breeding lines of *Pisum sativum* L. *J. Hortic. Sci. Biotechnol.* 90, 78–82.
- Rozov, S. M., Kosterin, O., Borisov, A. Y., and Tsyganov, V. (1999). The history of the pea gene map: last revolutions and the new symbiotic genes. *Pisum Genet.* 31, 55–56.
- Sarikamis, G., Yanmaz, R., Ermis, S., Bakir, M., and Yüksel, C. (2010). Genetic characterization of pea (*Pisum sativum*) germplasm from Turkey using morphological and SSR markers. *Genetics Mol. Res.* 9, 591–600. doi: 10.4238/vol9-1gmr762
- Shunmugam, A. S. K., Liu, X., Stonehouse, R., Tar'An, B., Bett, K. E., Sharpe, A. G., et al. (2015). Mapping seed phytic acid concentration and iron bioavailability in a pea recombinant inbred line population. *Crop Sci.* 55, 828–836. doi: 10.2135/cropsci2014.08.0544
- Sindhu, A., Ramsay, L., Sanderson, L. A., Stonehouse, R., Li, R., Condie, J., et al. (2014). Gene-based SNP discovery and genetic mapping in pea. *Theor. Appl. Genet.* 127, 2225–2241. doi: 10.1007/s00122-014-2375-y
- Smýkal, P., Coyne, C. J., Ambrose, M. J., Maxted, N., Schaefer, H., Blair, M. W., et al. (2015). Legume crops phylogeny and genetic diversity for science and breeding. *CRC. Crit. Rev. Plant Sci.* 34, 43–104. doi: 10.1080/07352689.2014.897904
- Smýkal, P., Hybl, M., Corander, J., Jarkovsky, J., Flavell, A. J., and Griga, M. (2008). Genetic diversity and population structure of pea (*Pisum sativum* L.) varieties derived from combined retrotransposon, microsatellite and morphological marker analysis. *Theor. Appl. Genet.* 117, 413–424. doi: 10.1007/s00122-008-0785-4
- Smýkal, P., Jovanović, Ž., Stanisavljević, N., Zlatković, B., Ćupina, B., Đorđević, V., et al. (2014). A comparative study of ancient DNA isolated from charred pea (*Pisum sativum* L.) seeds from an Early Iron Age settlement in southeast Serbia: inference for pea domestication. *Genet. Resour. Crop Evol.* 61, 1533–1544. doi: 10.1007/s10722-014-0128-z
- Stein, H. H., Everts, A. K. R., Sweeter, K. K., Peters, D. N., Maddock, R. J., Wulf, D. M., et al. (2006). The influence of dietary field peas (*Pisum sativum* L.) on pig performance, carcass quality, and the palatability of pork. *J. Anim. Sci.* 84, 3110–3117. doi: 10.2527/jas.2005-744
- Sudheesh, S., Lombardi, M., Leonforte, A., Cogan, N. O. I., Materne, M., Forster, J. W., et al. (2014). Consensus genetic map construction for field pea (*Pisum sativum* L.), trait dissection of biotic and abiotic stress tolerance and development of a diagnostic marker for the er1 powdery mildew resistance gene. *Plant Mol. Biol. Rep.* 33, 1391–1403. doi: 10.1007/s11105-014-0837-7
- Sun, X., Yang, T., Hao, J., Zhang, X., Ford, R., Jiang, J., et al. (2014). SSR genetic linkage map construction of pea (*Pisum sativum* L.) based on Chinese native varieties. *Crop J.* 2, 170–174. doi: 10.1016/j.cj.2014.03.004
- Svabova, L., and Griga, M. (2008). The effect of cocultivation treatments on transformation efficiency in pea (*Pisum sativum* L.). *Plant Celle Tissue Organ Culture* 95, 293–304. doi: 10.1007/s11240-008-9443-4
- Tar'an, B., Warkentin, T., Somers, D. J., Miranda, D., Vandenberg, A., Blade, S., et al. (2003). Quantitative trait loci for lodging resistance, plant height and partial resistance to mycosphaerella blight in field pea (*Pisum sativum* L.). *Theor. Appl. Genet.* 107, 1482–1491. doi: 10.1007/s00122-003-1379-9

- Tar'an, B., Warkentin, T., Somers, D. J., Miranda, D., Vandenberg, A., Blade, S., et al. (2004). Identification of quantitative trait loci for grain yield, seed protein concentration and maturity in field pea (*Pisum sativum* L.). *Euphytica* 136, 297–306. doi: 10.1023/B:EUPH.0000032721.03075.a0
- Tayeh, N., Aloume, C., Falque, M., Jacquin, F., Klein, A., Chauveau, A., et al. (2015a). Development of two major resources for pea genomics: the GenoPea 13.2K SNP Array and a high density, high resolution consensus genetic map. *Plant J.* doi: 10.1111/tpj.13070. [Epub ahead of print].
- Tayeh, N., Bahrman, N., Devaux, R., Bluteau, A., Prosperi, J.-M., Delbreil, B., et al. (2013a). A high-density genetic map of the *Medicago truncatula* major freezing tolerance QTL on chromosome 6 reveals colinearity with a QTL related to freezing damage on *Pisum sativum* linkage group VI. *Mol. Breed.* 32, 279–289. doi: 10.1007/s11032-013-9869-1
- Tayeh, N., Bahrman, N., Sellier, H., Bluteau, A., Blassiau, C., Fourment, J., et al. (2013b). A tandem array of *CBF/DREB1* genes is located in a major freezing tolerance QTL region on *Medicago truncatula* chromosome 6. *BMC Genomics* 14:814. doi: 10.1186/1471-2164-14-814
- Tayeh, N., Klein, A., Le Paslier, M. C., Jacquin, F., Houtin, H., Rond, C., et al. (2015b). Genomic prediction in pea: effect of marker density and training population size and composition on prediction accuracy. *Front. Plant Sci.* 6:941. doi: 10.3389/fpls.2015.00941
- Timmerman-Vaughan, G. M., Frew, T. J., Butler, R., Murray, S., Gilpin, M., Falloon, K., et al. (2004). Validation of quantitative trait loci for Ascochyta blight resistance in pea (*Pisum sativum* L.), using populations from two crosses. *Theor. Appl. Genet.* 109, 1620–1631. doi: 10.1007/s00122-004-1779-5
- Timmerman-Vaughan, G. M., Frew, T. J., Russell, A. C., Khan, T., Butler, R., Gilpin, M., et al. (2002). QTL mapping of partial resistance to field epidemics of ascochyta blight of pea. *Crop Sci.* 42, 2100–2111. doi: 10.2135/cropsci2002.02100
- Timmerman-Vaughan, G. M., McCallum, J. A., Frew, T. J., Weeden, N. F., and Russell, A. C. (1996). Linkage mapping of quantitative trait loci controlling seed weight in pea (*Pisum sativum* L.). *Theor. Appl. Genet.* 93, 431–439. doi: 10.1007/BF00223187
- Timmerman-Vaughan, G. M., Mills, A., Whitfield, C., Frew, T., Butler, R., Murray, S., et al. (2005). Linkage mapping of QTL for seed yield, yield components, and developmental traits in pea. *Crop Sci.* 45, 1336–1344. doi: 10.2135/cropsci2004.0436
- Ubayasena, L., Bett, K., Tar'an, B., Vijayan, P., and Warkentin, T. (2010). Genetic control and QTL analysis of cotyledon bleaching resistance in green field pea (*Pisum sativum* L.). *Genome* 53, 346–359. doi: 10.1139/G10-013
- Ubayasena, L., Bett, K., Tar'an, B., and Warkentin, T. (2011). Genetic control and identification of QTLs associated with visual quality traits of field pea (*Pisum sativum* L.). *Genome* 54, 261–272. doi: 10.1139/g10-117
- Valderrama, M. R., Roman, B., Satovic, Z., Rubiales, D., Cubero, J. I., and Torres, A. M. (2004). Locating quantitative trait loci associated with *Orobanche crenata* resistance in pea. *Weed Res.* 44, 323–328. doi: 10.1111/j.1365-3180.2004.00406.x
- van Leur, J. A. G., Aftab, M., Leonforte, B., Moore, S., and Freeman, A. J. (2007). "Control of Pea seedborne mosaic virus in field pea through resistance breeding," in *Proceedings 16th Biennial Conference of the Australasian Plant Pathology Society* (Adelaide, SA).
- Varshney, R. K., Kudapa, H., Pazhamala, L., Chitikineni, A., Thudi, M., Bohra, A., et al. (2015). Translational genomics in agriculture: some examples in grain legumes. *CRC. Crit. Rev. Plant Sci.* 34, 169–194. doi: 10.1080/07352689.2014.897909
- Wang, T. L., Domoney, C., Hedley, C. L., Casey, R., and Grusak, M. A. (2003). Can we improve the nutritional quality of legume seeds? *Plant Physiol.* 131, 886–891. doi: 10.1104/pp.102.017665
- Warkentin, T. D., Delgerjav, O., Arganosa, G., Rehman, A. U., Bett, K. E., Anbessa, Y., et al. (2012). Development and characterization of low-phytate pea. *Crop Sci.* 52, 74–78. doi: 10.2135/cropsci2011.05.0285
- Warkentin, T. D., Smykal, P., Coyne, C. J., Weeden, N., Domoney, C., Bing, D., et al. (2015). "Pea (*Pisum sativum* L.)" in *Grain Legumes, Series Handbook of Plant Breeding*, ed A. M. De Ron (New York, NY: Springer Science+Business Media), 37–83.
- Weeden, N. F., Brauner, S., and Przyborowski, J. A. (2002). Genetic analysis of pod dehiscence in pea (*Pisum sativum* L.). *Cell. Mol. Biol. Lett.* 7, 657–663.
- Weeden, N. F., Ellis, T. H. N., Timmerman-Vaughan, G. M., Swiecicki, W. K., Rozov, S. M., and Berdnikov, V. A. (1998). A consensus linkage map for *Pisum sativum*. *Pisum Genet.* 30, 1–4.
- Weeden, N. F., McGee, R., Grau, C. R., and Muehlbauer, F. J. (2000). A gene influencing tolerance to common root rot is located on linkage group IV. *Pisum Genet.* 32, 53–55.
- Weeden, N. F., and Porter, L. (2007). The genetic basis of Fusarium root rot tolerance in the 'Afghanistan' pea. *Pisum Genet.* 39, 35–36.
- Weiss, E., and Zohary, D. (2011). The Neolithic Southwest Asian founder crops their biology and archaeobotany. *Curr. Anthropol.* 52, S237–S254. doi: 10.1086/658367
- Wellensiek, S. J. (1925). Genetic monograph on *Pisum*. *Bibliogr. Genet.* 2, 343–476.
- Weller, J. L., Liew, L. C., Hecht, V. F. G., Rajandran, V., Laurie, R. E., Ridge, S., et al. (2012). A conserved molecular basis for photoperiod adaptation in two temperate legumes. *Proc. Natl. Acad. Sci. U.S.A.* 109, 21158–21163. doi: 10.1073/pnas.1207943110
- Weller, J. L., and Ortega, R. (2015). Genetic control of flowering time in legumes. *Front. Plant Sci.* 6:207. doi: 10.3389/fpls.2015.00207
- Willcox, G., Buxo, R., and Herveux, L. (2009). Late Pleistocene and early Holocene climate and the beginnings of cultivation in northern Syria. *Holocene* 19, 151–158. doi: 10.1177/0959683608098961
- Zhang, C., Tar'An, B., Warkentin, T., Tullu, A., Bett, K. E., Vandenberg, B., et al. (2006). Selection for lodging resistance in early generations of field pea by molecular markers. *Crop Sci.* 46, 321–329. doi: 10.2135/cropsci2005.0123
- Zohary, D. (1999). Monophyletic vs. polyphyletic origin of the crops on which agriculture was founded in the Near East. *Genet. Resour. Crop Evol.* 46, 133–142. doi: 10.1023/A:1008692912820
- Zong, X., Redden, R. J., Liu, Q., Wang, S., Guan, J., Liu, J., et al. (2009). Analysis of a diverse global *Pisum* sp collection and comparison to a Chinese local *P. sativum* collection with microsatellite markers. *Theor. Appl. Genet.* 118, 193–204. doi: 10.1007/s00122-008-0887-z

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Tayeh, Aubert, Pilet-Nayel, Lejeune-Hénaut, Warkentin and Burstin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.