



Zinc Oxide Nanoparticles Affect Biomass Accumulation and Photosynthesis in *Arabidopsis*

Xiaoping Wang, Xiyu Yang, Siyu Chen, Qianqian Li, Wei Wang, Chunjiang Hou, Xiao Gao, Li Wang and Shucui Wang*

Key Laboratory of Molecular Epigenetics of Ministry of Education, Northeast Normal University, Changchun, China

OPEN ACCESS

Edited by:

Nelson Marniroli,
University of Parma, Italy

Reviewed by:

Zhenzhu Xu,
Chinese Academy of Sciences, China
Jane Geisler-Lee,
Southern Illinois University
Carbondale, USA

*Correspondence:

Shucui Wang
wangsc550@nenu.edu.cn

Specialty section:

This article was submitted to
Functional Plant Ecology,
a section of the journal
Frontiers in Plant Science

Received: 28 August 2015

Accepted: 21 December 2015

Published: 12 January 2016

Citation:

Wang X, Yang X, Chen S, Li Q,
Wang W, Hou C, Gao X, Wang L
and Wang S (2016) Zinc Oxide
Nanoparticles Affect Biomass
Accumulation and Photosynthesis
in *Arabidopsis*.
Front. Plant Sci. 6:1243.
doi: 10.3389/fpls.2015.01243

Dramatic increase in the use of nanoparticles (NPs) in a variety of applications greatly increased the likelihood of the release of NPs into the environment. Zinc oxide nanoparticles (ZnO NPs) are among the most commonly used NPs, and it has been shown that ZnO NPs were harmful to several different plants. We report here the effects of ZnO NPs exposure on biomass accumulation and photosynthesis in *Arabidopsis*. We found that 200 and 300 mg/L ZnO NPs treatments reduced *Arabidopsis* growth by ~20 and 80%, respectively, in comparison to the control. Pigments measurement showed that Chlorophyll a and b contents were reduced more than 50%, whereas carotenoid contents remain largely unaffected in 300 mg/L ZnO NPs treated *Arabidopsis* plants. Consistent with this, net rate of photosynthesis, leaf stomatal conductance, intercellular CO₂ concentration and transpiration rate were all reduced more than 50% in 300 mg/L ZnO NPs treated plants. Quantitative RT-PCR results showed that expression levels of chlorophyll synthesis genes including *CHLOROPHYLL A OXYGENASE (CAO)*, *CHLOROPHYLL SYNTHASE (CHLG)*, *COPPER RESPONSE DEFECT 1 (CRD1)*, *MAGNESIUM-PROTOPORPHYRIN IX METHYLTRANSFERASE (CHLM)* and *MG-CHELATASE SUBUNIT D (CHLD)*, and photosystem structure gene *PHOTOSYSTEM I SUBUNIT D-2 (PSAD2)*, *PHOTOSYSTEM I SUBUNIT E-2 (PSAE2)*, *PHOTOSYSTEM I SUBUNIT K (PSAK)* and *PHOTOSYSTEM I SUBUNIT K (PSAN)* were reduced about five folds in 300 mg/L ZnO NPs treated plants. On the other hand, elevated expression, though to different degrees, of several carotenoids synthesis genes including *GERANYLGERANYL PYROPHOSPHATE SYNTHASE 6 (GGPS6)*, *PHYTOENE SYNTHASE (PSY)*, *PHYTOENE DESATURASE (PDS)*, and *ZETA-CAROTENE DESATURASE (ZDS)* were observed in ZnO NPs treated plants. Taken together, these results suggest that toxicity effects of ZnO NPs observed in *Arabidopsis* was likely due to the inhibition of the expression of chlorophyll synthesis genes and photosystem structure genes, which results in the inhibition of chlorophylls biosynthesis, leading to the reduce in photosynthesis efficiency in the plants.

Keywords: nanoparticles, ZnO, biomass, chlorophylls, carotenoid, gene expression, *Arabidopsis*

INTRODUCTION

Nanoparticles (NPs), also known as particulate nanomaterials (NMs), are particle materials with at least one dimension in the nanoscale (1–100 nm). Because of their particular properties, such as small size, high surface-to-volume ratio and unique physical and chemical properties, the use of NPs in industries and a wide range of consumer products are increasing greatly (Stampoulis et al., 2009). These positive commercial advances have stimulated a rapidly increasing production of engineered NPs, which made nanotechnologies a rapidly developing field with an expectation that the annual value of nanotechnology-related products is going to reach one trillion dollars in 2015 (Roco, 2005). However, the increasing usage of NPs greatly increased the likelihood of their release into the environment, and has raised concerns about the impacts of NMs on health and the environment (Xia et al., 2009; Shvedova et al., 2010).

Studies in plants have demonstrated that at least some NPs can be uptaken (Etxeberria et al., 2006; Chen et al., 2010; Kurepa et al., 2010; Schwab et al., 2015a), transported (Liu et al., 2009; Cifuentes et al., 2010; Zhao et al., 2012; Wang et al., 2012b), and accumulated in specific subcellular locations such as cell vacuoles, nuclei and plasmodesmata (Kurepa et al., 2010; Schwab et al., 2015b), and NPs can alter plant physiological processes, and influence plant growth and development (Stampoulis et al., 2009; Ma et al., 2010; Burklew et al., 2012; García-Sánchez et al., 2015). For example, ultra small anatase TiO₂ NPs have been shown to be able to enter into plant cells, accumulate in subcellular locations such as cell vacuoles and nuclei of root cells, and cause reorganization and elimination of microtubules, resulting in inhibition of root elongation in *Arabidopsis* (Kurepa et al., 2010; Wang et al., 2011). CuO NPs have been shown to be able to transport in maize via xylem and phloem (Wang et al., 2012b). Whereas silver NPs (Ag NPs) and zinc oxide NPs (ZnO NPs) treatment lead to increase in contents of free radicals, including reactive oxygen and nitrogen species (ROS/RNS) and hydrogen peroxide (H₂O₂) in duckweed (Thwala et al., 2013).

Effects of several NPs, including Ag NPs (Qian et al., 2013; Thwala et al., 2013; Geisler-Lee et al., 2014), aluminum oxide NPs (Al₂O₃ NPs) (Burklew et al., 2012; Riahi-Madvar et al., 2012), silicon dioxide NPs (SiO₂ NPs) (Zhang et al., 2015), and ZnO NPs (Ma et al., 2013) have been studied in several different plant species. These experiments indicated that NPs have negative effects on several different aspects of plant growth and development including seed germination (Ma et al., 2013), root elongation (Wang et al., 2012a), biomass accumulation (Zhao et al., 2013). NPs have also been shown to be able to induce oxidative stress and alter gene expression in plants (Wang et al., 2013a).

ZnO NPs are among the most commonly used NPs in a variety of applications such as personal care, ceramics, paints, pigments, foods, batteries, and semiconductors (Ju-Nam and Lead, 2008), which increased the potential of their direct release to the environment. Thus the ecological risk of ZnO NPs is an important topic. As a matter of fact, the toxicity effects of ZnO NPs have been observed in several different plants species including *Arabidopsis* (Lee et al.,

2010), buckweed (Lee et al., 2013), wheat (Du et al., 2011), dotted duckmeat (Thwala et al., 2013), cucumber (Zhao et al., 2013), rapeseed (Kouhi et al., 2014), alfalfa (Bandyopadhyay et al., 2015), and cowpea (Wang et al., 2013b). Most of the experiments show that the inhibition effects of ZnO NPs on plant growth and development are likely due to the induction of oxidative stress (Hernandez-Viezcas et al., 2011; Thwala et al., 2013; Bandyopadhyay et al., 2015). In consistent with this, transcriptome analysis in *Arabidopsis* have shown that most of the genes induced by ZnO NPs are ontology groups annotated as stress responsive, including both abiotic and biotic stimuli (Landa et al., 2012).

To further investigate the mechanisms of the toxicity effects of ZnO NPs on plant growth and development, we examined the effects of ZnO NPs on biomass accumulation and photosynthesis in *Arabidopsis*. We show that expose to ZnO NPs led to decrease in biomass accumulation in both shoots and roots, and that chlorophylls, but not carotenoid contents were decreased in plants treated ZnO NPs. Consistent with this observation, quantitative RT-PCR results showed that the expression levels of some chlorophyll synthesis genes and photosystem genes examined were decreased in response to ZnO NPs treatment.

MATERIALS AND METHODS

Plant Materials and Growth Conditions

Arabidopsis (*Arabidopsis thaliana*) ecotype Columbia (Col-0) was used for the experiments. Seeds were sown directly into soil pots and kept in a growth room at 22°C with a light density (photosynthetic active radiation) of approximately 120 μmol m⁻²s⁻¹, and a light/dark photoperiod of 16 h/8 h.

ZnO NPs Treatment

Soil pots were prepared by filling the 5.5 cm × 5.5 cm pots with TS-1 white peat bedding substrate (Epagma) moistened thoroughly with suspensions containing 0, 50, 100, 200, 250, and 300 mg/L ZnO NPs with particle size <50 nm (Sigma), respectively. Substrate moistened with supernatants from 300 mg/L ZnO NPs suspensions were used as controls. The supernatants were obtained by centrifuging 300 mg/L ZnO NPs suspensions for 10 min at 4000 rpm, and filtered through 0.22 μm-diameter filters. *Arabidopsis* seeds were then germinated and grown in the soil pots. After germination, extra seedlings were removed to ensure that every pot contains 12 plants. The plants were watered regularly till 4-week-old, then watered once every 2 days for two times with 50 ml suspensions containing ZnO NPs at the same concentrations, or supernatants from 300 mg/L ZnO NPs suspensions as described above.

Measurement of Net Rate of Photosynthesis, Leaf Stomatal Conductance, Intercellular CO₂ Concentration, and Transpiration Rate

Net rate of photosynthesis (P_N), leaf stomatal conductance (g_s), intercellular CO₂ concentration (C_i) and transpiration rate (E) of

the fully expanded fifth rosette leaves of 6-week-old plants were measured by using a portable open-flow, gas-exchange system (LI-6400; LICOR Biosciences, Lincoln, NE, USA) on the morning before the plants were harvested. The ambient CO₂ concentration was 360 ± 10 μmol mol⁻¹, the air temperature was 22°C, and humidity was about 50%. A total of four leaves per pot were measured, and measurements were repeated four times for each leaf.

The net rate of photosynthesis was calculated as CO₂ uptake in micromole per square meter leaf area per second (μmol CO₂ m⁻²s⁻¹), leaf stomatal conductance as water vapor in mole per square meter leaf area per second (mol H₂O m⁻²s⁻¹), intercellular CO₂ concentration as CO₂ in micromole per mole intercellular air (μmol CO₂ mol⁻¹), and the transpiration rate as water loss in millimole per square meter leaf area per second (mmol H₂O m⁻²s⁻¹).

Chlorophylls and Carotenoids Measurement

Chlorophylls and carotenoids contents were measured as described by Lichtenhaler and Wellburn (1983). Briefly, Chlorophylls and Carotenoids were extracted from rosette leaves from 6-week-old plants with 100% alcohol. Absorption of the extracts was measured using a spectrophotometer (721 TYPE, Shanghai analysis instrument co., LTD). Contents of Chlorophyll a (Chl a), Chlorophyll b (Chl b), total chlorophylls (C_T), and Carotenoids (Car) were calculated using the formula described (Lichtenhaler and Wellburn, 1983).

Biomass and Water Content Measurement

Six-week-old plants from each pot were harvested, washed thoroughly with tap water, and then distilled water. After removing excess water with paper towels, the roots and shoots were separated, and fresh weight (FW) was recorded. The samples were then oven-dried at 80°C for 15 min, followed by vacuum-dry at 40°C to a constant mass before dry weight (DW) was recorded. FW and DW per plant were calculated respectively, by divided the FW or DW per pot with 12. Water contents were calculated by using the formula: (FW-DW)/DW.

RNA Isolation and Quantitative RT-PCR (qRT-PCR)

Total RNA were isolated from rosette leaves of 6-week-old *Arabidopsis* plants by using EasyPure™ Plant RNA Kit (Transgene) and following the manufacturer's protocols.

First strand cDNA was synthesized using 2 μg total RNA by Oligo(dT)-primed reverse transcription using the EazyScript First-Strand DNA Synthesis Super Mix (TransGen Biotech) by following the manufacturer's instructions. Quantitative RT-PCR (qRT-PCR) was used to examine the expression of chlorophylls synthesis genes, carotenoids synthesis genes, and photosystem structure genes. *Arabidopsis* gene *ACTIN2* (*ACT2*) were used as an inner control for qRT-PCR. The primers used for qRT-PCR examination of *ACT2*, *MAGNESIUM-PROTOPORPHYRIN IX*

METHYLTRANSFERASE (*CHLM*), *Mg-chelatase subunit D* (*CHLD*), *GERANYLGERANYL PYROPHOSPHATE SYNTHASE 6* (*GGPS6*), *PHYTOENE SYNTHASE* (*PSY*), *PHYTOENE DESATURASE* (*PDS*), *ZETA-CAROTENE DESATURASE* (*ZDS*), *PHOTOSYSTEM I SUBUNIT K* (*PSAK*), and *PHOTOSYSTEM I SUBUNIT K* (*PSAN*) have been described previously (Qin et al., 2007; Stephenson and Terry, 2008; Liu et al., 2015).

The primer pairs used for qRT-PCR examination of *CHLOROPHYLL A OXYGENASE* (*CAO*), *CHLOROPHYLL SYNTHASE* (*CHLG*), *COPPER RESPONSE DEFECT 1* (*CRD1*), *PHOTOSYSTEM I SUBUNIT D-2* (*PSAD2*) and *PHOTOSYSTEM I SUBUNIT E-2* (*PSAE2*) were listed in **Table 1**.

Statistical Analysis

Statistical analysis was performed as described previously (Wang et al., 2015). Briefly, data were analyzed by one-way analysis of variance (ANOVA) using the statistical software SPSS 22.0 (IBM Inc., New York, USA), and compared by student-Newman-Keuls (*q*-test).


RESULTS

ZnO NPs Affect Growth and Biomass Accumulation in *Arabidopsis*

After 1 month of growth, ZnO NPs treated plants showed an obvious decrease in the rosette size, and the decrease in the rosette sizes were positively related with the concentrations of ZnO NPs applied (**Figure 1A**), suggesting that ZnO NPs inhibited *Arabidopsis* growth. Quantitative analysis showed that ZnO NPs at 100 mg/L had little, if any effects on the FW and DW of the plants (**Figures 1B,C**), however, an about 20% of decrease on both fresh and dry weight was observed for 200 mg/L ZnO NPs treated plants, and that for 300 mg/L ZnO NPs treated plants was about 80% (**Figures 1B,C**). Water contents, on the other hand, were reduced only in 300 mg/L ZnO NPs treated plants (**Figure 1D**). Detailed analysis showed that ZnO NPs at high concentration has more severe inhibition effects on root growth than shoot growth (**Figures 1E,F**). As a result, root/shoot ratio reduced about 40% in 300 mg/L ZnO NPs treated plants, whereas ZnO NPs at relative lower concentration has little, if any effects on root/shoot ratio of the plants (**Figure 1G**).

TABLE 1 | Primers used in this study.

| Primers | Sequences |
|----------|-----------------------------|
| CAO-qF | 5'-AGTCCTTCTGCTTTATCTCTC-3' |
| CAO-qR | 5'-TTCTCAACTAATCCACTCTCA-3' |
| CHLG-qF | 5'-GAGATTGTGTGTGCGTGCGG-3' |
| CHLG-qR | 5'-CCAGTGGAGGCCAAGTGACT-3' |
| CRD1-qF | 5'-AAGAGGAACTGGATAGAA-3' |
| CRD1-qR | 5'-AAAGAAGTAACCAAAGGAA-3' |
| PSAD2-qF | 5'-CAAACACACCATCACCAATC-3' |
| PSAD2-qR | 5'-ACCTCGTACCTAAAGCCAAA-3' |
| PSAE-qF | 5'-CACCACCATTGTGTCTTTCT-3' |
| PSAE-qR | 5'-TTGACCTTGGATCCTCTCTT-3' |



Chlorophylls, but not Carotenoids Contents were Affected by ZnO NPs Treatment

Yellow leaf color observed in ZnO NPs treated plants (Figure 1A) indicates that ZnO NPs may affect chlorophylls contents in the plants. To test this, we measured contents of chlorophylls including Chl a and Chl b in the plants. As shown in Figure 2A, ZnO NPs at lower concentrations has little, if any effects on Chl a contents, however, an about 50% reduce in Chl a contents was observed in 300 mg/L ZnO NPs treated plants. On the other hand, even at lower concentrations, i.e., 100 mg/L ZnO NPs treatment decreased Chl b contents in plants, and the degrees of decrease increased following the increase in the ZnO NPs concentrations (Figure 2B). Consistent with this, Chl a/b ratio increased, though not to a high degree in ZnO NPs treated plants (Figure 2C), and total chlorophylls contents also reduced in ZnO NPs treated plants (Figure 2D). On the other hand, ZnO NPs treatment increased carotenoids contents (Figure 2E). However, a positive relation between the increases of carotenoids contents and the concentrations of ZnO NPs was not observed (Figure 2E). As a consequence, carotenoids/chlorophylls ratio

increased in ZnO NPs treated plants, and an about three-fold increase was observed in 300 mg/L ZnO NPs treated plants (Figure 2F).

Leaf Photosynthesis in ZnO NPs Treated Plants was Reduced

Because ZnO NPs treatment resulted in reduction of chlorophylls contents (Figure 2), we suspected that photosynthesis may be affected in ZnO NPs treated plants. To test this, we examined net rate of photosynthesis, leaf stomatal conductance, intercellular CO₂ concentration and transpiration rate of the fully expanded fifth rosette leaves. As shown in Figure 3A, ZnO NPs treatment at all the concentrations tested decreased the net rate of photosynthesis, with an about 60% decrease observed in 300 mg/L ZnO NPs treated plants. An even higher degree of decrease was observed for the leaf stomatal conductance, in the plants treated with 300 mg/L ZnO NPs, the leaf stomatal conductance was only about 30% of that in the control plants treated with water only or with supernatants from 300 mg/L ZnO NPs suspensions (Figure 3B). The intercellular CO₂ concentration was also greatly reduced in ZnO NPs treated




FIGURE 2 | Effects of ZnO NPs on chlorophylls and carotenoids contents in *Arabidopsis* rosette leaves. Chlorophyll a (A), Chlorophyll b (B), Chlorophyll a/b ratio (C), total chlorophylls (D), carotenoids (E) and carotenoids/total chlorophylls ratio (F) in rosette leaves of 6-week-old *Arabidopsis* plants treated with 0, 100, 200, 300 mg/L ZnO NPs, or supernatant from 300 mg/L ZnO NPs suspensions. Chlorophylls and carotenoids were extracted from rosette leaves from 6-week-old *Arabidopsis* plants using alcohol, and measured using a spectrophotometer. Data represent the mean \pm SD of four replicates. Different letters indicate significantly different ($p < 0.05$).

plants, with an about 20% decrease in 200 mg/L ZnO NPs treated plants, and a more than 60% decrease in 300mg/L ZnO NPs treated plants (Figure 3C). A similar trend of decrease on transpiration rate was also observed in plants watered with ZnO NPs, i.e., an about 10, 30, and 60% decrease, respectively in plants treated with 100, 200, and 300 mg/L ZnO NPs (Figure 3D).

Expression Levels of Chlorophyll Synthesis Genes in ZnO NPs Treated Plants were Reduced

Several genes including *CHLOROPHYLL A OXYGENASE* (CAO), *CHLOROPHYLL SYNTHASE* (CHLG), *COPPER RESPONSE DEFECT 1* (CRD1), *MAGNESIUM-PROTOPORPHYRIN*

IX METHYLTRANSFERASE (CHLM), and *MG-CHELATASE SUBUNIT D* (CHLD) have been shown to be involved chlorophyll synthesis (Tanaka et al., 1998; Lange and Ghassemian, 2003; Rzeznicka et al., 2005; Pontier et al., 2007; Kobayashi et al., 2008). Having shown that ZnO NPs treatments reduced chlorophylls contents in *Arabidopsis* (Figure 2), we wanted to further examine if expression of the genes involved in chlorophyll synthesis were affected. Total RNA was isolated from rosette leaves, and quantitative RT-PCR was used to examine the expression of the chlorophyll synthesis genes. As shown in Figure 4, expression of all the genes examined including CAO, CHLG, CRD1, CHLM and CHLD was reduced in the plants treated with ZnO NPs at all concentrations. In 300 mg/L ZnO NPs treated plants, an ~5–7-fold decrease of the genes examined was observed (Figure 4). It should be noted that a slight, i.e., ~1.3-fold,




FIGURE 3 | Effects of ZnO NPs on photosynthesis of *Arabidopsis* plants. (A) Net photosynthetic rate (P_N), (B) stomatal conductance (g_s), (C) intercellular CO_2 concentration (C_i), and (D) transpiration rate (E) in rosette leaves of 6-week-old *Arabidopsis* plants treated with 0, 100, 200, 300 mg/L ZnO NPs, or supernatant from 300 mg/L ZnO NPs suspensions. The fifth rosette leaves were used for the measurement by using a portable open-flow, gas-exchange system. Data represent the mean \pm SD of four replicates. Different letters indicate significantly different ($p < 0.05$).




FIGURE 4 | Effects of ZnO NPs on the expression of chlorophyll synthesis genes in *Arabidopsis*. RNA was isolated from rosette leaves of *Arabidopsis* plants treated with 0, 100, 200, 300 mg/L ZnO NPs, or supernatant from 300 mg/L ZnO NPs suspensions, and qRT-PCR was used to examine the expression of genes involved chlorophyll synthesis. Expression of *ACT2* was used as a reference gene, and expression of corresponding genes in *Arabidopsis* in the absence of ZnO NPs was set as 1. Data represent the mean \pm SD of three replicates. Different letters indicate significantly different ($p < 0.05$).


increase for the expression of *CAO* and *CHLG* was observed in plants treated with supernatant of 300 mg/L ZnO NPs, whereas the expression of *CRD1*, *CHLM* and *CHLD* in plants treated with supernatant of 300 mg/L ZnO NPs remained largely unchanged when compared with that in water watered control plants (Figure 4).

Expression Levels of Carotenoids Synthesis Genes were Increased in ZnO NPs Treated Plants

Because carotenoids contents were increased slightly in ZnO NPs treated plants, we also examined the expression of several genes that have been reported to be involved in carotenoids synthesis, including *GERANYLGERANYL PYROPHOSPHATE SYNTHASE 6 (GGPS6)*, *PHYTOENE SYNTHASE (PSY)*, *PHYTOENE DESATURASE (PDS)*, and *ZETA-CAROTENE DESATURASE (ZDS)* (Lange and Ghassemian, 2003; Dong et al., 2007; Qin et al., 2007; Rodríguez-Villalón et al., 2009). We found that the expression of these genes was increased in response to ZnO NPs treatments, but to different degrees. In 300 mg/L ZnO NPs treated plants, an about 2-, 4-, 2- and 10- fold increase for *GGPS6*, *PSY*, *PDS* and *ZDS*, respectively was observed (Figure 5).

Expression Levels of Photosystem Structure Genes in ZnO NPs Treated Plants were Reduced


In addition to the expression of the chlorophylls and carotenoids synthesis genes, we also examined the expression of photosystem structure genes including *PHOTOSYSTEM I SUBUNIT D-2 (PSAD2)*, *PHOTOSYSTEM I SUBUNIT E-2 (PSAE2)*, *PHOTOSYSTEM I SUBUNIT K (PSAK)*, and *PHOTOSYSTEM I SUBUNIT K (PSAN)* (Varotto et al., 2002; Friso et al., 2004; Ilnatowicz et al., 2004; Peltier et al., 2004). We found that the expression of these genes was also reduced in ZnO NPs treated



plants (Figure 6). As a control, a slight increased expression of all the four genes was observed in plants treated with supernatant of 300 mg/L ZnO NPs (Figure 6).

DISCUSSION

Previous experiments have shown that toxic effects of ZnO NPs on plants are likely caused by the induction of oxidative stress in plants (Hernandez-Viezcas et al., 2011; Thwala et al., 2013; Bandyopadhyay et al., 2015). Consistent with this, most of the genes induced by ZnO NPs in *Arabidopsis* are relative



to stress responses (Landa et al., 2012). We provide evidence here in this report that ZnO NPs are toxic to *Arabidopsis* plants, and their toxic effects may due to the effects of ZnO NPs on the expression of chlorophyll synthesis and photosystem genes.

First, similar to the results obtained in other plants such as cucumber (Zhao et al., 2013), ZnO NPs treated *Arabidopsis* plants have reduced rosette size and biomass accumulation, and the reduction is positively related to the concentrations of ZnO NPs used (Figure 1). Our data also show that the ZnO NPs have similar inhibition effects on biomass accumulation in shoots and roots (Figure 1). Whereas plants treated with supernatant of 300 mg/L ZnO NPs suspensions are similar to control plants in both rosette size and biomass accumulation (Figure 1). These results suggest that ZnO NPs are toxic to *Arabidopsis*, though we could not rule out the possibility that the toxic effects may be partially due to the release of Zn^{2+} by ZnO NPs into the soil (Cornelis et al., 2014), it is unlikely that the toxic effects observed was due to the possible release of Zn^{2+} or other ions by ZnO NPs to the supernatants.

Second, contents of chlorophyll a and chlorophyll b were reduced in rosette leaves of ZnO NPs treated plants, whereas those in plants treated with supernatants of 300 mg/L ZnO NPs suspensions remain largely unaffected (Figure 2). Net rate of photosynthesis, leaf stomatal conductance, intercellular CO_2 concentration and transpiration rate were all reduced in ZnO NPs treated plants (Figure 3). Carotenoids contents, on the other hand, showed slightly increase in ZnO NPs treated plants (Figure 2), suggesting that reduction in photosynthesis is caused by reduced chlorophylls contents. Nevertheless, these results suggest that toxic effects of ZnO NPs on *Arabidopsis* are likely caused by reduced chlorophylls contents in the plants, which in turn limited photosynthesis in the plants, leading to the reduce in biomass accumulation.

Third, consistent with the observation that chlorophylls contents were reduced in ZnO NPs treated plants, expression of chlorophyll synthesis genes was reduced in ZnO NPs treated plants (Figure 4). We also found the expression of photosystem genes was reduced in ZnO NPs treated plants (Figure 6). On the other hand, the expression of some chlorophyll synthesis and photosystem genes was slightly increased in plants treated with supernatant from 300 mg/L ZnO NPs suspensions (Figures 4 and 6), further confirmed that the toxic effects observed in ZnO NPs treated plants were not due to the Zn^{2+} or other ions released by ZnO NPs in to the supernatants. Though carotenoids contents only slightly increased in ZnO NPs treated plants (Figure 2), we found the expression of some carotenoids synthesis genes, especially ZDS, was dramatically increased in ZnO NPs treated plants (Figure 5). Considering that carotenoids is also involved in photosynthesis, and ZnO NPs are toxic to plants, it will be of great interest to find out why ZnO NPs treatment increased the expression of carotenoids synthesis genes in *Arabidopsis*.

It should be noted that when compared with a single plant or plants grown at low density, plants growth at high density showed intraspecific competition (Geisler et al., 2012; Masclaux et al., 2012; Pierik and de Wit, 2014), which led to the upregulation of

photosynthesis genes (Geisler et al., 2012). However, because all the pots in our experiments contained the same number of plants, the difference of the gene expression observed was unlikely due to intraspecific competition.

Nevertheless, our results show that ZnO NPs inhibited plant growth and biomass accumulation in *Arabidopsis*, and that in addition to the induction of oxidative stresses, regulation of chlorophyll synthesis and photosystem genes expression may also contribute to the toxic effects of ZnO NPs in *Arabidopsis*.

REFERENCES

- Bandyopadhyay, S., Plascencia-Villa, G., Mukherjee, A., Rico, C. M., José-Yacamán, M., Peralta-Videa, J. R., et al. (2015). Comparative phytotoxicity of ZnO NPs, bulk ZnO, and ionic zinc onto the alfalfa plants symbiotically associated with *Sinorhizobium meliloti* in soil. *Sci. Total Environ.* 515–516, 60–69. doi: 10.1016/j.scitotenv.2015.02.014
- Burklew, C. E., Ashlock, J., Winfrey, W. B., and Zhang, B. (2012). Effects of aluminum oxide nanoparticles on the growth, development, and microRNA expression of tobacco (*Nicotiana tabacum*). *PLoS ONE* 7:e34783. doi: 10.1371/journal.pone.0034783
- Chen, R., Ratnikova, T. A., Stone, M. B., Lin, S., Lard, M., Huang, G., et al. (2010). Differential uptake of carbon nanoparticles by plant and mammalian cells. *Small* 6, 612–617. doi: 10.1002/smll.200901911
- Cifuentes, Z., Custardoy, L., de la Fuente, J. M., Marquina, C., Ibarra, M. R., Rubiales, D., et al. (2010). Absorption and translocation to the aerial part of magnetic carbon-coated nanoparticles through the root of different crop plants. *J. Nanobiotech* 8:26. doi: 10.1186/1477-3155-8-26
- Cornelis, G., Hund-Rinke, K., Kuhlbusch, T., van den Brink, N., and Nickel, C. (2014). Fate and bioavailability of engineered nanoparticles in soils: a review. *Crit. Rev. Env. Sci. Technol.* 44, 2720–2764. doi: 10.1080/10643389.2013.829767
- Dong, H., Deng, Y., Mu, J., Lu, Q., Wang, Y., Xu, Y., et al. (2007). The *Arabidopsis* spontaneous cell death1 gene, encoding a zeta-carotene desaturase essential for carotenoid biosynthesis, is involved in chloroplast development, photoprotection and retrograde signalling. *Cell Res.* 17, 458–470. doi: 10.1038/cr.2007.37
- Du, W., Sun, Y., Ji, R., Zhu, J., Wu, J., and Guo, H. (2011). TiO₂ and ZnO nanoparticles negatively affect wheat growth and soil enzyme activities in agricultural soil. *J. Environ. Monit.* 13, 822–828. doi: 10.1039/c0em00611d
- Etxeberria, E., Gonzalez, P., Baroja-Fernandez, E., and Romero, J. P. (2006). Fluid phase endocytic uptake of artificial nano-spheres and fluorescent quantum dots by sycamore cultured cells. *Plant Signal. Behav.* 1, 196–200. doi: 10.4161/psb.1.4.3142
- Friso, G., Giacomelli, L., Ytterberg, A. J., Peltier, J. B., Rudella, A., Sun, Q., et al. (2004). In-depth analysis of the thylakoid membrane proteome of *Arabidopsis thaliana* chloroplasts: new proteins, new functions, and a plastid proteome database. *Plant Cell* 16, 478–499. doi: 10.1105/tpc.017814
- García-Sánchez, S., Bernales, I., and Cristobal, S. (2015). Early response to nanoparticles in the *Arabidopsis* transcriptome compromises plant defence and root-hair development through salicylic acid signalling. *BMC Genomics* 16:341. doi: 10.1186/s12864-015-1530-4
- Geisler, M., Gibson, D. J., Lindsey, K. J., Millar, K., and Wood, A. J. (2012). Upregulation of photosynthesis genes, and down-regulation of stress defense genes, is the response of *Arabidopsis thaliana* shoots to intraspecific competition. *Bot. Stud.* 53, 85–96.
- Geisler-Lee, J., Brooks, M., Gerfen, J. R., Wang, Q., Fotis, C., Sparer, A., et al. (2014). Reproductive toxicity and life history study of silver nanoparticle effect, uptake and transport in *Arabidopsis thaliana*. *Nanomaterials* 4, 301–318. doi: 10.3390/nano4020301
- Hernandez-Viezas, J. A., Castillo-Michel, H., Servin, A. D., Peralta-Videa, J. R., and Gardea-Torresdey, J. L. (2011). Spectroscopic verification of zinc absorption and distribution in the desert plant *Prosopis juliflora*-velutina (velvet mesquite) treated with ZnO nanoparticles. *Chem. Eng. J.* 170, 346–352. doi: 10.1016/j.cej.2010.12.021

ACKNOWLEDGMENTS

This work was supported by the International Science & Technology Cooperation Program of China (2014DFA31740), the Key Laboratory of Molecular Epigenetics of MOE (130014542), a startup fund from Northeast Normal University (www.nenu.edu.cn), and the Programme for Introducing Talents to Universities (B07017). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

- Ihnatowicz, A., Pesaresi, P., Varotto, C., Richly, E., Schneider, A., Jahns, P., et al. (2004). Mutants for photosystem I subunit D of *Arabidopsis thaliana*: effects on photosynthesis, photosystem I stability and expression of nuclear genes for chloroplast functions. *Plant J.* 37, 839–852. doi: 10.1111/j.1365-313X.2004.02011.x
- Ju-Nam, Y., and Lead, J. R. (2008). Manufactured nanoparticles: an overview of their chemistry, interactions and potential environmental implications. *Sci. Total Environ.* 400, 396–414. doi: 10.1016/j.scitotenv.2008.06.042
- Kobayashi, K., Mochizuki, N., Yoshimura, N., Motohashi, K., Hisabori, T., and Masuda, T. (2008). Functional analysis of *Arabidopsis thaliana* isoforms of the Mg-chelatase CHL1 subunit. *Photochem. Photobiol. Sci.* 7, 1188–1195. doi: 10.1039/b802604c
- Kouhi, S. M. M., Lahouti, M., Ganjeali, A., and Entezar, M. H. (2014). Comparative phytotoxicity of ZnO nanoparticles, ZnO microparticles, and Zn²⁺ on rapeseed (*Brassica napus* L.): investigating a wide range of concentrations. *Toxicol. Environ. Chem.* 96, 861–868. doi: 10.1080/02772248.2014.994517
- Kurepa, J., Paunesku, T., Vogt, S., Arora, H., Rabatic, B. M., Lu, J., et al. (2010). Uptake and distribution of ultrasmall anatase TiO₂ Alizarin red S nanoconjugates in *Arabidopsis thaliana*. *Nano Lett.* 10, 2296–2302. doi: 10.1021/nl903518f
- Landa, P., Vankova, R., Andrlova, J., Hodek, J., Marsik, P., Storchova, H., et al. (2012). Nanoparticle-specific changes in *Arabidopsis thaliana* gene expression after exposure to ZnO, TiO₂, and fullerene soot. *J. Hazard. Mater.* 241–242, 55–62. doi: 10.1016/j.jhazmat.2012.08.059
- Lange, B. M., and Ghassemian, M. (2003). Genome organization in *Arabidopsis thaliana*: a survey for genes involved in isoprenoid and chlorophyll metabolism. *Plant Mol. Biol.* 51, 925–948. doi: 10.1023/A:1023005504702
- Lee, C. W., Mahendra, S., Zodrow, K., Li, D., Tsai, Y. C., Braam, J., et al. (2010). Developmental phytotoxicity of metal oxide nanoparticles to *Arabidopsis thaliana*. *Environ. Toxicol. Chem.* 29, 669–675. doi: 10.1002/etc.58
- Lee, S., Kim, S., Kim, S., and Lee, I. (2013). Assessment of phytotoxicity of ZnO NPs on a medicinal plant, *Fagopyrum esculentum*. *Environ. Sci. Pollut. Res. Int.* 20, 848–854. doi: 10.1007/s11356-012-1069-8
- Lichtenhaler, H. K., and Wellburn, A. R. (1983). Determination of total carotenoid and chlorophyll a and b of leaf extract in different solvent. *Biochem. Soc. Trans.* 603, 591–592. doi: 10.1042/bst0110591
- Liu, Q., Chen, B., Wang, Q., Shi, X., Xiao, Z., Lin, J., et al. (2009). Carbon nanotubes as molecular transporters for walled plant cells. *Nano Lett.* 9, 1007–1010. doi: 10.1021/nl803083u
- Liu, S., Hu, Q., Luo, S., Li, Q., Yang, X., Wang, X., et al. (2015). Expression of wild-type PtrIAA14.1, a poplar Aux/IAA gene causes morphological changes in *Arabidopsis*. *Front. Plant Sci.* 6:388. doi: 10.3389/fpls.2015.00388
- Ma, H., Williams, P. L., and Diamond, S. A. (2013). Ecotoxicity of manufactured AnO nanoparticles – a review. *Environ. Pollut.* 172, 76–85. doi: 10.1016/j.envpol.2012.08.011
- Ma, X., Geiser-Lee, J., Deng, Y., and Kolmakov, A. (2010). Interactions between engineered nanoparticles (ENPs) and plants: phytotoxicity, uptake and accumulation. *Sci. Total Environ.* 408, 3053–3061. doi: 10.1016/j.scitotenv.2010.03.031
- Masclaux, F., Bruessow, F., Schweizer, F., Gouhier-Darimont, C., Keller, L., and Reymond, P. (2012). Transcriptome analysis of intraspecific competition in *Arabidopsis thaliana* reveals organ-specific signatures related to nutrient acquisition and general stress response pathways. *BMC Plant Biol.* 12:227. doi: 10.1186/1471-2229-12-227

- Peltier, J. B., Ytterberg, A. J., Sun, Q., and van Wijk, K. J. (2004). New functions of the thylakoid membrane proteome of *Arabidopsis thaliana* revealed by a simple, fast, and versatile fractionation strategy. *J. Biol. Chem.* 279, 49367–49383. doi: 10.1074/jbc.M406763200
- Pierik, R., and de Wit, M. (2014). Shade avoidance: phytochrome signalling and other aboveground neighbour detection cues. *J. Exp. Bot.* 65, 2815–2824. doi: 10.1093/jxb/ert389
- Pontier, D., Albrieux, C., Joyard, J., Lagrange, T., and Block, M. A. (2007). Knock-out of the magnesium protoporphyrin IX methyltransferase gene in *Arabidopsis*. Effects on chloroplast development and on chloroplast-to-nucleus signaling. *J. Biol. Chem.* 282, 2297–2304. doi: 10.1074/jbc.M610286200
- Qian, H., Peng, X., Han, X., Ren, J., Sun, L., and Fu, Z. (2013). Comparison of the toxicity of silver nanoparticles and silver ions on the growth of terrestrial plant model *Arabidopsis thaliana*. *J. Environ. Sci.* 25, 1947–1955. doi: 10.1016/S1001-0742(12)60301-5
- Qin, G., Gu, H., Ma, L., Peng, Y., Deng, X. W., Chen, Z., et al. (2007). Disruption of phytoene desaturase gene results in albino and dwarf phenotypes in *Arabidopsis* by impairing chlorophyll, carotenoid, and gibberellin biosynthesis. *Cell Res.* 17, 471–482. doi: 10.1038/cr.2007.40
- Riahi-Madvar, A., Rezaee, F., and Jalali, V. (2012). Effects of alumina nanoparticles on morphological properties and antioxidant system of *Triticum aestivum*. *Ira. J. Plant Physiol.* 3, 595–603.
- Roco, M. C. (2005). Environmentally responsible development of nanotechnology. *Environ. Sci. Technol.* 39, 106A–112A. doi: 10.1021/es053199u
- Rodríguez-Villalón, A., Gas, E., and Rodríguez-Concepción, M. (2009). Colors in the dark: a model for the regulation of carotenoid biosynthesis in etioplasts. *Plant Signal. Behav.* 4, 965–967. doi: 10.4161/psb.4.10.9672
- Rzeznicka, K., Walker, C. J., Westergren, T., Kannangara, C. G., von Wettstein, D., Merchant, S., et al. (2005). Xantha-1 encodes a membrane subunit of the aerobic Mg-protoporphyrin IX monomethyl ester cyclase involved in chlorophyll biosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* 102, 5886–5891. doi: 10.1073/pnas.0501784102
- Schwab, F., Tanner, S., Schulz, R., Rotzetter, A., Stark, W., von Quadt, A., et al. (2015a). Dissolved cerium contributes to uptake of Ce in the presence of differently sized CeO₂-nanoparticles by three crop plants. *Metallomics* 7, 466–477. doi: 10.1039/c4mt00343h
- Schwab, F., Zhai, G., Kern, M., Turner, A., Schnoor, J. L., and Wiesner, M. R. (2015b). Barriers, pathways and processes for uptake, translocation and accumulation of nanomaterials in plants – critical review. *Nanotoxicology* doi: 10.3109/17435390.2015.1048326 [Epub ahead of print].
- Shvedova, A. A., Kagan, V. E., and Fadeel, B. (2010). Close encounters of the small kind: adverse effects of man-made materials interfacing with the nano-cosmos of biological systems. *Annu. Rev. Pharmacol.* 50, 63–88. doi: 10.1146/annurev.pharmtox.010909.105819
- Stampoulis, D., Sinha, S. K., and White, J. C. (2009). Assay-dependent phytotoxicity of nanoparticles to plants. *Environ. Sci. Technol.* 43, 9473–9479. doi: 10.1021/es901695c
- Stephenson, P. G., and Terry, M. J. (2008). Light signalling pathways regulating the Mg-chelatase branchpoint of chlorophyll synthesis during de-etiolation in *Arabidopsis thaliana*. *Photochem. Photobiol. Sci.* 7, 1243–1252. doi: 10.1039/b802596g
- Tanaka, A., Ito, H., Tanaka, R., Tanaka, N. K., Yoshida, K., and Okada, K. (1998). Chlorophyll a oxygenase (CAO) is involved in chlorophyll b formation from chlorophyll a. *Proc. Natl. Acad. Sci. U.S.A.* 95, 12719–12723. doi: 10.1073/pnas.95.21.12719
- Thwala, M., Musee, N., Sikhwivhilu, L., and Wepener, V. (2013). The oxidative toxicity of Ag and ZnO nanoparticles towards the aquatic plant *Spirodela punctata* and the role of testing media parameters. *Environ. Sci. Proc. Impacts* 15, 1830–1843. doi: 10.1039/c3em00235g
- Varotto, C., Pesaresi, P., Jahns, P., Lessnick, A., Tizzano, M., Schiavon, F., et al. (2002). Single and double knockouts of the genes for photosystem I subunits G, K, and H of *Arabidopsis*. Effects on photosystem I composition, photosynthetic electron flow, and state transitions. *Plant Physiol.* 129, 616–624. doi: 10.1104/pp.002089
- Wang, H., Wu, F., Meng, W., White, J. C., Holden, P. A., and Xing, B. (2013a). Engineered nanoparticles may induce genotoxicity. *Environ. Sci. Technol.* 47, 13212–13214. doi: 10.1021/es404527d
- Wang, P., Menzies, N. W., Lombi, E., McKenna, B. A., Johannessen, B., Glover, C. J., et al. (2013b). Fate of ZnO nanoparticles in soils and cowpea (*Vigna unguiculata*). *Environ. Sci. Technol.* 47, 13822–13830. doi: 10.1021/es403466p
- Wang, S., Kurepa, J., and Smalle, J. A. (2011). Ultra-small TiO₂ nanoparticles disrupt microtubular networks in *Arabidopsis thaliana*. *Plant Cell Environ.* 34, 811–820. doi: 10.1111/j.1365-3040.2011.02284.x
- Wang, X., Han, H., Liu, X., Gu, X., Chen, K., and Lu, D. (2012a). Multi-walled carbon nanotubes can enhance root elongation of wheat (*Triticum aestivum*) plants. *J. Nanopart. Res.* 14:841. doi: 10.1007/s11051-012-0841-5
- Wang, Z., Xie, X., Zhao, J., Liu, X., Feng, W., White, J. C., et al. (2012b). Xylem- and phloem-based transport of CuO nanoparticles in maize (*Zea mays* L.). *Environ. Sci. Technol.* 46, 4434–4441. doi: 10.1021/es204212z
- Wang, X., Jiang, P., Ma, Y., Geng, S., Wang, S., and Shi, D. (2015). Physiological strategies of sunflower exposed to salt or alkali stresses: restriction of ion transport in the cotyledon node zone and solute accumulation. *Agron. J.* 107, 2181–2192. doi: 10.2134/agronj15.0012
- Xia, T., Li, N., and Nel, A. E. (2009). Potential health impact of nanoparticles. *Annu. Rev. Publ. Health* 30, 137–150. doi: 10.1146/annurev.publhealth.031308.100155
- Zhang, W., Ebbs, S. D., Musante, C., White, J. C., Gao, C., and Ma, X. (2015). Uptake and accumulation of bulk and nanosized cerium oxide particles and ionic cerium by radish (*Raphanus sativus* L.). *J. Agric. Food Chem.* 63, 382–390. doi: 10.1021/jf505244z
- Zhao, L., Sun, Y., Hernandez-Viezas, J. A., Servin, A. D., Hong, J., Niu, G., et al. (2013). Influence of CeO₂ and ZnO nanoparticles on cucumber physiological markers and bioaccumulation of Ce and Zn: a life cycle study. *J. Agric. Food Chem.* 61, 11945–11951. doi: 10.1021/jf404328e
- Zhao, L. J., Peralta-Videa, J. R., Ren, M. H., Varela-Ramirez, A., Li, C. Q., Hernandez-Viezas, J. A., et al. (2012). Transport of Zn in a sandy loam soil treated with ZnO NPs and uptake by corn plants: electron microprobe and confocal microscopy studies. *Chem. Eng. J.* 184, 1–8. doi: 10.1016/j.cej.2012.01.041

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Wang, Yang, Chen, Li, Wang, Hou, Gao, Wang and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.