



A Sweetpotato Auxin Response Factor Gene (*IbARF5*) Is Involved in Carotenoid Biosynthesis and Salt and Drought Tolerance in Transgenic *Arabidopsis*

Chen Kang¹, Shaozhen He¹, Hong Zhai¹, Ruijie Li¹, Ning Zhao¹ and Qingchang Liu^{1,2*}

¹ Key Laboratory of Sweetpotato Biology and Biotechnology, Ministry of Agriculture, Beijing Key Laboratory of Crop Genetic Improvement, Laboratory of Crop Heterosis and Utilization, Ministry of Education, College of Agronomy and Biotechnology, China Agricultural University, Beijing, China, ² College of Agronomy, Qingdao Agricultural University, Qingdao, China

OPEN ACCESS

Edited by:

Ruth Grene,
Virginia Tech, United States

Reviewed by:

Kazuo Nakashima,
Japan International Research Center
for Agricultural Sciences, Japan
Sang-Soo Kwak,
Korea Research Institute
of Bioscience and Biotechnology
(KRIBB), South Korea

*Correspondence:

Qingchang Liu
liuqc@cau.edu.cn

Specialty section:

This article was submitted to
Plant Abiotic Stress,
a section of the journal
Frontiers in Plant Science

Received: 25 June 2018

Accepted: 17 August 2018

Published: 11 September 2018

Citation:

Kang C, He S, Zhai H, Li R, Zhao N
and Liu Q (2018) A Sweetpotato
Auxin Response Factor Gene
(*IbARF5*) Is Involved in Carotenoid
Biosynthesis and Salt and Drought
Tolerance in Transgenic *Arabidopsis*.
Front. Plant Sci. 9:1307.
doi: 10.3389/fpls.2018.01307

Auxin response factors (ARFs) compose a family of transcription factors and have been found to play major roles in the process of plant growth and development. However, their roles in plant carotenoid biosynthesis and responses to abiotic stresses are rarely known to date. In the present study, we found that the *IbARF5* gene from sweetpotato (*Ipomoea batatas* (L.) Lam.) line HVB-3 increased the contents of carotenoids and enhanced the tolerance to salt and drought in transgenic *Arabidopsis*. The transgenic *Arabidopsis* plants exhibited the increased abscisic acid (ABA) and proline contents and superoxide dismutase (SOD) activity and the decreased H₂O₂ content. Furthermore, it was found that *IbARF5* positively regulated the genes associated with carotenoid and ABA biosynthesis and abiotic stress responses. These results suggest that *IbARF5* is involved in carotenoid biosynthesis and salt and drought tolerance in transgenic *Arabidopsis*. This study provides a novel *ARF* gene for improving carotenoid contents and salt and drought tolerance of sweetpotato and other plants.

Keywords: sweetpotato, *IbARF5*, *Arabidopsis*, carotenoid content, salt and drought tolerance

INTRODUCTION

In nature, more than 750 kinds of carotenoids are characterized structurally, which are widely found in bacteria, fungi, algae, and plants (Hirschberg, 2001; Takaichi, 2011). The biosynthesis pathway of carotenoids has been extensively studied in plants, and nearly all of the key genes have been isolated and characterized (Cunningham and Gantt, 1998; Fraser and Bramley, 2004; Colasuonno et al., 2017; Kang et al., 2018). Abiotic stresses, especially salt and drought, seriously affect the productivity and cultivation expansion of crop plants worldwide, accordingly, to develop their high tolerance to salt and drought is highly desirable (Zhu, 2002; Lindemose et al., 2013; Zhai et al., 2016; Li et al., 2017). As the precursor of abscisic acid (ABA), carotenoids have functional roles in development and environmental adaptation of plants (Schwartz et al., 2003; Nambara and Marion-Poll, 2005; Mehrotra et al., 2014; Li, 2015; Moreno et al., 2016). Thus, increasing the contents of carotenoids helps to enhance the adaptation of plants to harsh environments.

Auxin response factors (ARFs) constitute a family of plant specific transcription factors. A typical ARF protein contains a B3-DNA binding domain in the highly conserved N-terminal

region (Ulmasov et al., 1997; Hagen and Guilfoyle, 2002; Mei et al., 2018). ARFs mediate responses to auxin and have been shown to be implicated in senescence (Ellis et al., 2005), hormone signaling (Li et al., 2006) and developmental programs (Krogan et al., 2012). In rice, *OsARF1* was auxin-regulated and classified as a primary auxin responsive gene (Waller et al., 2002). In *Arabidopsis*, *ARF2* mediated ABA response (Wang et al., 2011); *MP/ARF5* regulated embryo and flower patterning and vascular differentiation (Hardtke and Berleth, 1998; Krogan et al., 2012); *ARF6* and *ARF8* promoted jasmonic acid production and flower maturation (Nagpal et al., 2005); *NPH4/ARF7* and *ARF19* controlled leaf expansion and lateral root growth (Okushima et al., 2005; Wilmoth et al., 2005). In tomato, *SIARF2* regulated lateral root formation and flower senescence (Ren et al., 2017); *ARF4* controlled sugar metabolism (Sagar et al., 2013); *ARF10* increased chlorophyll and sugar accumulation during fruit development (Mei et al., 2018); *ARF5* regulated fruit set and development (Liu et al., 2018). However, the roles of ARFs in plant carotenoid biosynthesis and abiotic stress responses are rarely known to date.

Sweetpotato (*Ipomoea batatas* (L.) Lam.) is an important food crop worldwide, which provides rich carbohydrates and carotenoids for human consumption (Teow et al., 2007; Zhai et al., 2016). This crop can also be used for bioenergy production on marginal lands due to its high adaption to harsh environments (Liu et al., 2014). Sweetpotato breeders are focusing on improving carotenoid contents and abiotic stresses tolerance of this crop.

Kang et al. (2017) summarized the improvement of carotenoids by gene engineering in sweetpotato. Overexpression of the genes related to carotenoid biosynthesis have been shown to increase the contents of carotenoids and enhance the tolerance to abiotic stresses in sweetpotato (Kim et al., 2012, Kim et al., 2013b; Yu et al., 2013; Kim et al., 2014; Li et al., 2017; Kang et al., 2018). To date, ARFs have not been reported in sweetpotato. In this study, we found that the *IbARF5* gene from storage roots of sweetpotato is involved in carotenoid biosynthesis and salt and drought tolerance in transgenic *Arabidopsis*.


MATERIALS AND METHODS



Plant Materials

Sweetpotato line HVB-3 with high carotenoid content was employed to clone the *IbARF5* gene in this study. The expressed sequence tag (EST) for *IbARF5* was obtained from the transcriptome data of HVB-3 developed by Li et al. (2015). *Arabidopsis* wild type (Columbia-0, WT) was used for characterizing the *IbARF5* gene.

Isolation and Sequence Analysis of *IbARF5*

Total RNA was extracted from freshly harvested storage roots of HVB-3 and transcribed into first-strand cDNA according to the method of Kang et al. (2018). The full-length cDNA of





IbARF5 was amplified with specific primers (**Supplementary Table S1**) by rapid amplification of cDNA ends (RACE) method. Genomic DNA isolated from *in vitro*-grown plants of HVB-3 was used to amplify the genomic sequence of *IbARF5*. The *IbARF5* cDNA was analyzed by an online BLAST¹. The open-reading frame (ORF) Finder² was used to predict the ORF of *IbARF5*. The DNAMAN software was applied to align the amino acid sequence of *IbARF5* with those of ARF proteins from different plant species. The MEGA 7.0 software was employed to conduct the phylogenetic analysis with the neighbor-joining (NJ) method. Exon–intron structure was constructed using Splign tool³. The molecular weight and theoretical isoelectric point (*pI*) of *IbARF5* were calculated at http://web.expasy.org/compute_pi/.

Subcellular Localization of *IbARF5*

The *IbARF5* ORF amplified with specific primers (**Supplementary Table S1**) was ligated into pMDC83. pMDC83-*IbARF5-GFP* and pMDC83-*GFP* (as control) were transiently expressed in the onion epidermal cells with a GeneGun (HeliosTM, Biorad, United States). After co-cultivation on Murashige and Skoog (MS) medium (pH 5.8) at 28°C for 24 h, the onion cells were examined under a laser scanning confocal fluorescence microscope (Nikon Inc., Melville, NY, United States).

Transactivation Activity Assay of *IbARF5* in Yeast

Transactivation activity of *IbARF5* in yeast (*Saccharomyces cerevisiae*) was assayed as described by Jiang et al. (2014). The corresponding regions of *IbARF5* were PCR-amplified using specific primers (**Supplementary Table S1**) and integrated into the yeast expression vector pGBKT7 (pBD). Expression vectors pBD-*IbARF5*, pBD-*IbARF5-1*, pBD-*IbARF5-2*, pBD-*IbARF5-3*, pGAL4 (as positive vector), and pBD (as negative vector) were transferred into the yeast strain AH109, respectively. The transactivation activity was determined as described in the yeast protocols handbook (PT3024-1; Clontech, Mountain View, CA, United States).

Expression Analysis of *IbARF5* in Sweetpotato

Total RNA was isolated from storage root, stem, and leaf tissues of the 100-day-old HVB-3 and used to analyze the expression of *IbARF5* by quantitative real-time PCR (qRT-PCR) with its specific primers (**Supplementary Table S1**). *Ibactin* (AY905538) was served as an internal control. Comparative *C_T* method was employed to quantify the gene expression (Schmittgen and Livak, 2008).

After cultured on MS medium for 4 weeks, the HVB-3 plants were treated in liquid MS media containing H₂O (as control), 200 mM NaCl, 20% PEG6000 and 100 μM ABA, respectively, and

sampled at 0, 2, 4, 6, 12, 24, and 48 h after treatment for analyzing the expression of *IbARF5*.

Production of the Transgenic *Arabidopsis* Plants


The overexpression vector pC3301-121-*IbARF5* was constructed through inserting 35S-*IbARF5-NOS* into pCAMBIA3301. The recombinant vector was introduced into the *Agrobacterium tumefaciens* strain GV3101. The dipping flower method was applied to transform *Arabidopsis* and putatively transgenic *Arabidopsis* seeds were sown on MS medium with 12.5 mg L⁻¹ phosphinothricin (PPT) for selecting transgenic plants. Histochemical GUS assay (Jefferson et al., 1987) and PCR analysis were used to identify the transgenic *Arabidopsis* plants. Transgenic *Arabidopsis* was planted in pots with a soil, vermiculite and humus mixture (1:1:1, v/v/v) to obtain T₃ seeds.

Measurement of Carotenoid Contents

Leaves (2-week-old) and seeds of the transgenic *Arabidopsis* plants were applied to extract α-carotene, lutein, β-carotene, β-cryptoxanthin, and zeaxanthin. High performance liquid chromatography (HPLC) system was used to determine their contents (Li et al., 2017).

Assay for Salt and Drought Tolerance

One-week-old *in vitro*-grown seedlings of transgenic *Arabidopsis* and WT were treated on MS media with 200 mM NaCl and 300 mM mannitol, respectively. After 2 weeks, their root length and fresh weight (FW) were investigated. Furthermore, the transgenic and WT seedlings were planted for 2 weeks in pots with a soil, vermiculite and humus mixture (1:1:1, v/v/v) and subsequently irrigated with a 33 mL of 300 mM NaCl solution for each pot once every 2 days for 2 weeks, or stressed by drought for 4 weeks followed by 2 days re-watering. The transgenic plants and WT grown for 6 weeks under normal condition were used as control. The proline and H₂O₂ contents and superoxide dismutase (SOD) activity in the transgenic plants and WT grown in pots for 4 weeks under normal condition, 1 week under 300 mM NaCl stress after 2 weeks of normal treatment,



¹ <https://blast.ncbi.nlm.nih.gov/Blast.cgi>

² <https://www.ncbi.nlm.nih.gov/orffinder/>

³ <https://www.ncbi.nlm.nih.gov/sutils/splign/splign.cgi?textpage=online&level=form>

and 2 weeks under drought stress after 2 weeks of normal treatment, respectively, were determined with Assay Kits (Comin Biotechnology Co., Ltd. Suzhou, China). The ABA content was measured as described by Gao et al. (2011). Twenty-seven plants in three pots with nine plants per pot were treated for each line.

For ABA sensitivity assay, the transgenic and WT seeds were sown on MS media with 0, 0.5, and 1 μM ABA, respectively. After 1 week, their germination and cotyledon opening and greening rates were investigated. Fifty seeds of each line on a plate were analyzed.

Expression Analysis of the Related Genes

Leaves (2-week-old) and seeds of the transgenic *Arabidopsis* plants and WT were applied to analyze the expression of the key genes in carotenoid biosynthesis. The leaves of the transgenic plants and WT potted for 4 weeks under normal condition, 1 week under 300 mM NaCl stress after 2 weeks of normal treatment, and 2 weeks under drought stress after 2 weeks of normal treatment, respectively, were used for analyzing the expression of the genes associated with ABA biosynthesis and abiotic stress responses. The specific primers of *Atactin*




FIGURE 5 | Expression analysis of *IbARF5* in the *in vitro*-grown plants of HVB-3 after different times (h) in response to H₂O, 200 mM NaCl, 20% PEG6000 and 100 μM ABA, respectively. Data are presented as means \pm SE ($n = 3$). * and ** indicate a significant difference from that of WT at $P < 0.05$ and $P < 0.01$, respectively, by Student's *t*-test.

TABLE 1 | Carotenoid contents in leaves of the *IbARF5*-overexpressing *Arabidopsis* plants.

Lines	Carotenoids content ($\mu\text{g g}^{-1}$ FW)					
	α -carotene	Lutein	β -carotene	β -cryptoxanthin	Zeaxanthin	Total
WT	0.149 \pm 0.005	13.275 \pm 0.581	7.601 \pm 0.467	n.d.	0.044 \pm 0.010	21.070 \pm 1.020
L1	0.166 \pm 0.011	17.475 \pm 0.694**	6.751 \pm 0.552	n.d.	0.119 \pm 0.007**	24.511 \pm 1.038**
L4	0.151 \pm 0.004	16.750 \pm 1.221**	6.849 \pm 0.330	n.d.	0.147 \pm 0.016**	23.896 \pm 0.949*
L5	0.145 \pm 0.006	16.775 \pm 0.315**	6.657 \pm 0.755	n.d.	0.147 \pm 0.004**	23.993 \pm 0.854*
L6	0.153 \pm 0.007	17.532 \pm 1.210**	6.717 \pm 0.111	n.d.	0.126 \pm 0.009**	24.529 \pm 1.109**

Leaves from 2-week-old *Arabidopsis* plants were sampled for the quantification of carotenoids. FW, fresh weight; n.d., not detectable. Data are presented as mean \pm SE ($n = 3$). * and ** indicate a significant difference from that of WT at $P < 0.05$ and $P < 0.01$, respectively, by Student's *t*-test.

TABLE 2 | Carotenoid contents in seeds of the *IbARF5*-overexpressing *Arabidopsis* plants.

Lines	Carotenoids content ($\mu\text{g g}^{-1}$ DW)					
	α -carotene	Lutein	β -carotene	β -cryptoxanthin	Zeaxanthin	Total
WT	n.d.	1.470 \pm 0.041	0.178 \pm 0.003	0.020 \pm 0.005	0.089 \pm 0.009	1.755 \pm 0.048
L1	n.d.	1.907 \pm 0.028**	0.270 \pm 0.031*	0.023 \pm 0.006	0.119 \pm 0.010**	2.318 \pm 0.051**
L4	n.d.	2.519 \pm 0.207**	0.262 \pm 0.035*	0.038 \pm 0.006	0.153 \pm 0.008**	2.972 \pm 0.241**
L5	n.d.	2.625 \pm 0.150**	0.309 \pm 0.057**	0.046 \pm 0.002*	0.143 \pm 0.005**	3.123 \pm 0.194**
L6	n.d.	2.303 \pm 0.287**	0.386 \pm 0.059**	0.052 \pm 0.003**	0.144 \pm 0.023**	2.885 \pm 0.313**

Seeds from *Arabidopsis* plants were harvested for the quantification of carotenoids. DW, dry weight; n.d., not detectable. Data are presented as mean \pm SE ($n = 3$). * and ** indicate a significant difference from that of WT at $P < 0.05$ and $P < 0.01$, respectively, by Student's *t*-test.

(NM112764) as internal control and the related genes were listed in **Supplementary Table S1**.

Statistical Analysis

All experiments were conducted with three biological replicates. Data presented as the mean \pm SE were analyzed with Student's *t*-test (two-tailed analysis) at $P < 0.05$ and $P < 0.01$.

RESULTS

Cloning and Sequence Analysis of *IbARF5*

The 3757-bp cDNA of the *IbARF5* gene contained a 2841-bp ORF encoding a 946-aa polypeptide with a molecular weight of 104.84 kDa and a predicted *pI* of 5.17. The *IbARF5* protein shared a high sequence identity with ARF5 proteins in *Nicotiana tabacum* (XP_016465083.1, 74%), *Capsicum annuum* (XP_016568464.1, 72%), *Sesamum indicum* (XP_011083507.1, 72%), *Solanum lycopersicum* (NP_001234545.1, 72%), *Solanum tuberosum* (XP_006342026.1, 72%) and *Vitis vinifera* (XP_003634382.2, 68%). It contained one plant-specific B3-DNA binding domain, one Auxin_resp and one AUX_IAA

(**Supplementary Figure S1**). Phylogenetic analysis showed that *IbARF5* had a close relationship with that of *N. tabacum* (**Figure 1A**). The 4794-bp genomic DNA of *IbARF5* contained 13 exons and 12 introns (**Figure 1B**).

IbARF5 Is Localized to Nuclei

The images from onion epidermal cells indicated that the green fluorescence emitted by *IbARF5*-GFP was exclusively distributed over the nuclei of the cells (**Figure 2**). These results showed that *IbARF5* was localized to nuclei.

IbARF5 Shows Transactivation Activity in Yeast

The yeast two-hybrid system was applied to identify a possible transactivation activity of *IbARF5*. The yeast cells harboring pBD-GAL4, pGBKT7-*IbARF5* and pGBKT7-*IbARF5*-2, respectively, grew well on synthetic dropout (SD) plate without tryptophan and histidine and exhibited β -galactosidase activity, but the cells bearing pBD, pGBKT7-*IbARF5*-1, and pGBKT7-*IbARF5*-3, respectively, failed to grow (**Figure 3A**). These results demonstrated that *IbARF5* might act as a transcription activator, and its transactivation activity was determined by the middle region, *IbARF5*-2 (**Figure 3B**).




FIGURE 6 | Responses of the transgenic *Arabidopsis* seedlings and WT cultured for 2 weeks on MS medium with 200 mM NaCl and 300 mM mannitol, respectively. Data are presented as mean \pm SE ($n = 3$). ** indicates a significant difference from that of WT at $P < 0.01$ by Student's *t*-test.





FIGURE 7 | Responses of the transgenic *Arabidopsis* plants and WT grown in pots under salt and drought stresses. **(A)** Phenotypes of transgenic plants vs. WT grown for 6 weeks under normal condition, 2 weeks under 300 mM NaCl stress after 2 weeks of normal treatment, and 4 weeks under drought stress followed by 2 days re-watering after 2 weeks of normal treatment, respectively. **(B)** ABA, proline and H₂O₂ contents and SOD activity in the transgenic plants and WT grown for 4 weeks under normal condition, 1 week under 300 mM NaCl stress after 2 weeks of normal treatment, and 2 weeks under drought stress after 2 weeks of normal treatment, respectively. Data are presented as mean ± SE (n = 3). ** indicates a significant difference from that of WT at P < 0.01 by Student's t-test.

Expression Patterns of *IbARF5* in Sweetpotato

Quantitative real-time PCR analysis revealed that *IbARF5* exhibited higher expression level in the storage roots of HVB-3 than in its leaves and stems (Figure 4). Its expression in HVB-3 was strongly induced by NaCl, PEG6000 and ABA, and peaked (5.03-fold) at 4 h, (9.68-fold) at 24 h, and (12.18-fold) at 24 h, respectively (Figure 5).

Production of the Transgenic *Arabidopsis* Plants

Putatively transgenic *Arabidopsis* seeds formed the plants on MS medium with 12.5 mg L^{-1} PPT. GUS assay and PCR analysis confirmed that 6 of the randomly sampled 60 plants were transgenic plants, named L1, L2, . . . , L6, respectively, from which T₃ were generated. *IbARF5* showed high expression levels in the transgenic *Arabidopsis* plants, especially L1, L4, L5, and L6 (Supplementary Figure S2).

IbARF5 Increases Carotenoid Contents


The lutein and zeaxanthin contents were significantly increased, but α -carotene and β -carotene contents were not changed

and β -cryptoxanthin was not detected in leaves of L1, L4, L5, and L6 (Table 1). In seeds of these four transgenic plants, the lutein, β -carotene, and zeaxanthin contents were significantly increased, but α -carotene was not detected and β -cryptoxanthin content was significantly increased only in L5 and L6 (Table 2). The total carotenoid contents in leaves and seeds were increased by 1.13–1.16 folds and 1.32–1.78 folds, respectively (Tables 1 and 2).

IbARF5 Enhances Salt and Drought Tolerance

Four transgenic *Arabidopsis* plants, L1, L4, L5, and L6, and WT seedlings showed no significant differences in rooting and FW on MS medium without stresses (Figure 6). However, the transgenic plants exhibited good rooting and increased FW in contrast to WT on MS media with 200 mM NaCl and 300 mM mannitol, respectively (Figure 6).

The transgenic plants and WT grown in pots showed similar growth trends under normal conditions (Figure 7A). Under NaCl and drought stresses, the transgenic plants showed good growth, while WT almost died (Figure 7A). Furthermore, it was found that the ABA and proline contents were increased, SOD activity



was enhanced and H_2O_2 content was decreased in the transgenic plants (Figure 7B).

No obvious differences in seed germination were observed between the transgenic plants and WT under normal condition (Figure 8). Under the treatment with different ABA concentrations, the germination of the transgenic seeds was more sensitive to ABA-elicited inhibition than that of WT though both germination rate and cotyledon opening and greening rate of the transgenic and WT seeds declined (Figure 8). These results demonstrated that *IbARF5* might participate in the ABA signaling pathway.

IbARF5 Up-Regulates the Genes Involved in Carotenoid and ABA Biosynthesis and Abiotic Stress Responses


The genes encoding the key enzymes in carotenoid biosynthesis, geranylgeranyl pyrophosphate (GGPS), ζ -carotene desaturase (ZDS), phytoene synthase (PSY), ϵ -carotene hydroxylase (ϵ -CHY), β -lycopene cyclase (β -LCY) and β -carotene hydroxylase (β -CHY) except for phytoene desaturase (PDS)

and ϵ -lycopene cyclase (ϵ -LCY) were systematically up-regulated in leaves of the transgenic *Arabidopsis* plants (Figure 9). *GGPS*, ϵ -*CHY*, β -*LCY*, and β -*CHY* exhibited the increased expression levels, but *ZDS*, *PSY*, and ϵ -*LCY* showed no changes in expression level and *PDS* was down-regulated in the transgenic seeds (Figure 9). Under NaCl and drought stresses, the genes encoding the key enzymes in ABA biosynthesis, zeaxanthin epoxidase (*ZEP*), 9-cis-epoxycarotenoid dioxygenase (*NCED*), and xanthoxin dehydrogenase (*ABA2*) were up-regulated, and abiotic stress-responsive genes encoding pyrroline-5-carboxylate synthase (*P5CS*), *SOD*, ascorbate peroxidase (*APX*), and dehydroascorbate reductase (*DHAR*) were also found to be up-regulated (Figure 10).

DISCUSSION


IbARF5 Increases Carotenoid Contents and Salt and Drought Tolerance

In plants, ARFs encode important transcription factors which regulate the expression of genes in response to auxin (Guilfoyle and Hagen, 2007). Several ARF transcription factor genes have



been cloned from *Arabidopsis*, rice and tomato, and were found to play crucial roles in plant growth and developmental processes (Harper et al., 2000; Waller et al., 2002; Ellis et al., 2005; Okushima et al., 2005; Wilmoth et al., 2005; Wang et al., 2011; Ren et al., 2017; Liu et al., 2018). However, there is no report on the ARF transcription factors in improving carotenoid contents and abiotic stress tolerance in plants. Previous studies demonstrated that *AtARF5* affected lateral organ development,

primary root initiation, flower primordium initiation, and cotyledon development in *Arabidopsis* (Krogan et al., 2012) and *SlARF5* controlled fruit set and development in tomato (Liu et al., 2018). In the present study, the *IbARF5* gene was isolated from sweetpotato line HVB-3 with high carotenoid content. We found that its overexpression significantly increased the content of carotenoids and enhanced the tolerance to salt and drought in transgenic *Arabidopsis* (Tables 1, 2 and Figures 6, 7).



IbARF5 Up-Regulates the Genes Involved in Carotenoid Biosynthesis

It has been shown that carotenoid biosynthesis is mainly regulated at the transcript level of genes encoding the biosynthetic enzymes (Römer and Fraser, 2005; Sandmann et al., 2006; Li et al., 2017; Kang et al., 2018). In this study, we found that the key genes in carotenoid biosynthesis, *GGPS*, *ZDS*, *PSY*, ϵ -*CHY*, β -*LCY*, and β -*CHY* in leaves and *GGPS*, ϵ -*CHY*, β -*LCY*, and β -*CHY* in seeds of transgenic *Arabidopsis* were significantly up-regulated (Figure 9), which corresponded with the increase of carotenoid contents in transgenic *Arabidopsis* (Tables 1 and 2). These findings suggest that *IbARF5* positively controls the expression of carotenoid biosynthetic genes, which resulted in the increased carotenoid contents in transgenic *Arabidopsis*. Overexpression of the *Orange* gene (*IbOr*) from sweetpotato increased carotenoid accumulation and abiotic stress tolerance in transgenic sweetpotato, potato, and alfalfa (Kim et al., 2013a; Goo et al., 2015; Wang et al., 2015). Furthermore, it was proved that similar to *AtOr* of *Arabidopsis*, *IbOr* directly interacted with *PSY* and increased carotenoid accumulation (Park et al., 2016; Kim et al., 2018). Therefore, the precise underlying mechanisms of *IbARF5* in plant carotenoid accumulation need to be further investigated. In addition, we are developing the *IbARF5*-overexpressing sweetpotato plants for further analyzing its roles in carotenoid accumulation of the storage roots.

IbARF5 Up-Regulates the Genes Involved in ABA Biosynthesis

Carotenoids, especially β -branch carotenoids, serve as precursors for ABA biosynthesis and play a crucial role in plant tolerance and adaptation to abiotic stresses (Demmig-Adams and Adams, 2002; Xiong and Zhu, 2003; Sah et al., 2016). ABA regulates the expression of ABA-dependent stress-responsive genes and the increased level of ABA has been found to enhance the tolerance to salt and drought (Tuteja, 2007; Vishwakarma et al., 2017). It was reported that overexpression of *IbMIPS1*, *IbZDS*, and *IbLCYB2* increased the level of ABA, which led to the enhanced tolerance to salt and drought in sweetpotato (Zhai et al., 2016; Li et al., 2017; Kang et al., 2018). In this study, the *IbARF5*-overexpressing *Arabidopsis* seeds showed the increased sensitivity to ABA in germination (Figure 8). The ABA biosynthetic genes *IbZEP*, *IbNCED*, and *IbABA2* were up-regulated and ABA level was also significantly increased in transgenic *Arabidopsis* (Figures 7B and 10). These results suggest that overexpression of *IbARF5* confers salt and drought tolerance by up-regulating the ABA biosynthetic genes and increasing ABA level in transgenic *Arabidopsis*.

IbARF5 Up-Regulates Abiotic Stress-Responsive Genes and Changes Abiotic Stress-Associated Components

It is reported that the high level of ABA increases the transcript level of *P5CS*, which leads to more accumulation of proline under abiotic stresses (Sripinyowanich et al., 2013). Proline plays a pivotal role in maintaining osmotic balance, protecting integrity membrane and increasing reactive oxygen species (ROS)

scavenging capacity, and its elevated level enhances salt and drought tolerance in plants (Yoshida et al., 1997; Maggio et al., 2002; Neisiani et al., 2009; Gill and Tuteja, 2010; Kang et al., 2018). SOD as the first line of defense against ROS is induced by abiotic stresses to promote ROS scavenging (Wang et al., 2009). In the present study, *P5CS*, *SOD*, *APX*, and *DHAR* were up-regulated, proline level and SOD activity were increased and H_2O_2 content was decreased in transgenic *Arabidopsis* under salt and drought stresses (Figure 7B). Therefore, it is thought that the enhanced tolerance to salt and drought is due to up-regulation of abiotic stress-responsive genes and change of abiotic stress-associated components in transgenic *Arabidopsis*.

CONCLUSION

This study reveals, for the first time, that the *IbARF5* gene from sweetpotato is involved in carotenoid biosynthesis and salt and drought tolerance of plants. Its overexpression increased the contents of carotenoids and conferred the tolerance to salt and drought by up-regulating the key genes involved in carotenoid and ABA biosynthesis and abiotic stress responses in transgenic *Arabidopsis*. This study provides a novel *ARF* gene for improving carotenoid contents and salt and drought tolerance of sweetpotato and other plants.

AUTHOR CONTRIBUTIONS

QL and CK conceived and designed the experiments. CK and RL performed the experiments. CK and SH analyzed the data. QL, HZ, and NZ contributed reagents, materials, and analysis tools. QL and CK wrote the paper. All authors read and approved the final manuscript.

FUNDING

This work was supported by China Agriculture Research System (CARS-10, Sweetpotato) and National Natural Science Foundation of China (31461143017).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2018.01307/full#supplementary-material>

FIGURE S1 | Sequence alignment of *IbARF5* with its homologs from other plants. Characteristic regions of ARF5 are indicated above the *IbARF5* sequence. —, Plant-specific B3-DNA binding domain; —, Auxin_resp; —, AUX_IAA.

FIGURE S2 | Expression analysis of *IbARF5* in the transgenic *Arabidopsis* plants. The *Arabidopsis actin* gene was used as an internal control. Data are presented as means \pm SE ($n = 3$). ** indicates a significant difference from that of WT at $P < 0.01$ by Student's *t*-test.

TABLE S1 | Primers used in this study.

REFERENCES

- Colasuonno, P., Lozito, M. L., Marcotuli, I., Nigro, D., Giancaspro, A., Mangini, G., et al. (2017). The carotenoid biosynthetic and catabolic genes in wheat and their association with yellow pigments. *BMC Genomics* 18:122. doi: 10.1186/s12864-016-3395-6
- Cunningham, F. X., and Gantt, E. (1998). Genes and enzymes of carotenoid biosynthesis in plants. *Annu. Rev. Plant Physiol. Mol. Biol.* 49, 557–583. doi: 10.1146/annurev.arplant.49.1.557
- Demmigadams, B., and Adams, W. W. (2002). Antioxidants in photosynthesis and human nutrition. *Science* 298, 2149–2153. doi: 10.1126/science.1078002
- Ellis, C. M., Nagpal, P., Young, J. C., Hagen, G., Guilfoyle, T. J., and Reed, J. W. (2005). Auxin response factor1 and auxin response factor2 regulate senescence and floral organ abscission in *Arabidopsis thaliana*. *Development* 132, 4563–4574. doi: 10.1242/dev.02012
- Fraser, P. D., and Bramley, P. M. (2004). The biosynthesis and nutritional uses of carotenoids. *Prog. Lipid Res.* 43, 228–265. doi: 10.1016/j.plipres.2003.10.002
- Gao, S., Yuan, L., Zhai, H., Liu, C. L., He, S. Z., and Liu, Q. C. (2011). Transgenic sweetpotato plants expressing an LOS5 gene are tolerant to salt stress. *Plant Cell Tissue Organ Cult.* 107, 205–213. doi: 10.1007/s11240-011-9971-1
- Gill, S. S., and Tuteja, N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol. Biochem.* 48, 909–930. doi: 10.1016/j.plaphy.2010.08.016
- Goo, Y., Han, E., Jeong, J. C., Kwak, S., Yu, J., Kim, Y., et al. (2015). Overexpression of the sweet potato IbOr gene results in the increased accumulation of carotenoid and confers tolerance to environmental stresses in transgenic potato. *C. R. Biol.* 338, 12–20. doi: 10.1016/j.crv.2014.10.006
- Guilfoyle, T. J., and Hagen, G. (2007). Auxin response factors. *Curr. Opin. Plant Biol.* 10, 453–460.
- Hagen, G., and Guilfoyle, T. (2002). Auxin-responsive gene expression: genes, promoters and regulatory factors. *Plant Mol. Biol.* 49, 373–385.
- Hardtke, C. S., and Berleth, T. (1998). The *Arabidopsis* gene MONOPTEROS encodes a transcription factor mediating embryo axis formation and vascular development. *EMBO J.* 17, 1405–1411. doi: 10.1093/emboj/17.5.1405
- Harper, R. M., Stowe-Evans, E. L., Luesse, D. R., Muto, H., Tatematsu, K., Watahiki, M. K., et al. (2000). The NPH4 locus encodes the auxin response factor ARF7, a conditional regulator of differential growth in aerial *Arabidopsis* tissue. *Plant Cell* 12, 757–770. doi: 10.1105/tpc.12.5.757
- Hirschberg, J. (2001). Carotenoid biosynthesis in flowering plants. *Curr. Opin. Plant Biol.* 4, 210–218. doi: 10.1016/S1369-5266(00)00163-1
- Jefferson, R. A., Kavanagh, T. A., and Bevan, M. W. (1987). GUS fusions: β -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* 6, 3901–3907.
- Jiang, X., Zhang, C., Lü, P., Jiang, G., Liu, X., Dai, F., et al. (2014). RhNAC3, a stress-associated NAC transcription factor, has a role in dehydration tolerance through regulating osmotic stress-related genes in rose petals. *Plant Biotechnol. J.* 12, 38–48. doi: 10.1111/pbi.12114
- Kang, C., Zhai, H., Xue, L. Y., Zhao, N., He, S. Z., and Liu, Q. C. (2018). A lycopene β -cyclase gene, IbLCYB2, enhances carotenoid contents and abiotic stress tolerance in transgenic sweetpotato. *Plant Sci.* 272, 243–254. doi: 10.1016/j.plantsci.2018.05.005
- Kang, L., Park, S., Ji, C. Y., Kim, H. S., Lee, H., and Kwak, S. (2017). Metabolic engineering of carotenoids in transgenic sweetpotato. *Breed. Sci.* 67, 27–34. doi: 10.1270/jsbbs.16118
- Kim, S. H., Ahn, Y. O., Ahn, M., Jeong, J. C., Lee, H., and Kwak, S. (2013a). Cloning and characterization of an Orange gene that increases carotenoid accumulation and salt stress tolerance in transgenic sweetpotato cultures. *Plant Physiol. Biochem.* 70, 445–454. doi: 10.1016/j.plaphy.2013.06.011
- Kim, S. H., Kim, Y., Ahn, Y. O., Ahn, M., Jeong, J. C., Lee, H., et al. (2013b). Downregulation of the lycopene ϵ -cyclase gene increases carotenoid synthesis via the β -branch-specific pathway and enhances salt-stress tolerance in sweetpotato transgenic calli. *Physiol. Plant.* 147, 432–442. doi: 10.1111/j.1399-3054.2012.01688.x
- Kim, S. H., Ahn, Y. O., Ahn, M., Lee, H., and Kwak, S. (2012). Down-regulation of β -carotene hydroxylase increases β -carotene and total carotenoids enhancing salt stress tolerance in transgenic cultured cells of sweetpotato. *Phytochemistry* 74, 69–78. doi: 10.1016/j.phytochem.2011.11.003
- Kim, S. H., Jeong, J. C., Park, S., Bae, J., Ahn, M., Lee, H., et al. (2014). Down-regulation of sweetpotato lycopene β -cyclase gene enhances tolerance to abiotic stress in transgenic calli. *Mol. Biol. Rep.* 41, 8137–8148. doi: 10.1007/s11033-014-3714-4
- Kim, S. H., Ji, Y. C., Lee, C., Kim, S., Park, S., and Kwak, S. (2018). Orange: a target gene for regulating carotenoid homeostasis and increasing plant tolerance to environmental stress in marginal lands. *J. Exp. Bot.* 69, 3393–3400. doi: 10.1093/jxb/ery023
- Krogan, N. T., Ckurshumova, W., Marcos, D., Caragea, A. E., and Berleth, T. (2012). Deletion of MP/ARF5 domains III and IV reveals a requirement for Aux/IAA regulation in *Arabidopsis* leaf vascular patterning. *New Phytol.* 194, 391–401. doi: 10.1111/j.1469-8137.2012.04064.x
- Li, J., Dai, X., and Zhao, Y. (2006). A role for auxin response factor 19 in auxin and ethylene signaling in *Arabidopsis*. *Plant Physiol.* 140, 899–908. doi: 10.1104/pp.105.070987
- Li, R. J., Kang, C., Song, X. J., Yu, L., Liu, D. G., He, S. Z., et al. (2017). A ζ -carotene desaturase gene, IbZDS, increases β -carotene and lutein contents and enhances salt tolerance in transgenic sweetpotato. *Plant Sci.* 262, 39–51. doi: 10.1016/j.plantsci.2017.05.014
- Li, R. J., Zhai, H., Kang, C., Liu, D. G., He, S. Z., and Liu, Q. C. (2015). De novo transcriptome sequencing of the orange-fleshed sweet potato and analysis of differentially expressed genes related to carotenoid biosynthesis. *Int. J. Genomics* 2015:843802. doi: 10.1155/2015/843802
- Li, T. (2015). Recent advances in understanding carotenoid-derived signaling molecules in regulating plant growth and development. *Front. Plant Sci.* 6:790. doi: 10.3389/fpls.2015.00790
- Lindemose, S., O'Shea, C., Jensen, M., and Skriver, K. (2013). Structure, function and networks of transcription factors involved in abiotic stress responses. *Int. J. Mol. Sci.* 14, 5842–5878. doi: 10.3390/ijms14035842
- Liu, D. G., He, S. Z., Zhai, H., Wang, L. J., Zhao, Y., Wang, B., et al. (2014). Overexpression of IbP5CR enhances salt tolerance in transgenic sweetpotato. *Plant Cell Tissue Organ Cult.* 117, 1–16. doi: 10.1007/s11240-013-0415-y
- Liu, S. Y., Zhang, Y. W., Feng, Q. S., Qin, L., Pan, C. T., Lamin-Samu, A. T., et al. (2018). Tomato auxin response factor 5 regulates fruit set and development via the mediation of auxin and gibberellin signaling. *Sci. Rep.* 8:2971. doi: 10.1038/s41598-018-21315-y
- Maggio, A., Miyazaki, S., Veronese, P., Fujita, T., and Ibeas, J. I. (2002). Does proline accumulation play an active role in stress-induced growth reduction? *Plant J.* 31, 699–712. doi: 10.1046/j.1365-313X.2002.01389.x
- Mehrotra, R., Bhalothia, P., Bansal, P., Basantani, M. K., Bharti, V., and Mehrotra, S. (2014). Abscisic acid and abiotic stress tolerance-different tiers of regulation. *J. Plant Physiol.* 171, 486–496. doi: 10.1016/j.jplph.2013.12.007
- Mei, L. Y., Yuan, Y. J., Wu, M. B., Gong, Z. H., et al. (2018). SlARF10, an auxin response factor, is required for chlorophyll and sugar accumulation during tomato fruit development. *bioRxiv* [Preprint]. doi: 10.1101/253237
- Moreno, J. C., Cerda, A., Simpson, K., Lopez-Diaz, I., Carrera, E., Handford, M., et al. (2016). Increased *Nicotiana tabacum* fitness through positive regulation of carotenoid, gibberellin and chlorophyll pathways promoted by *Daucus carota* lycopene β -cyclase (Dlcyb1) expression. *J. Exp. Bot.* 67, 2325–2338. doi: 10.1093/jxb/erw037
- Nagpal, P., Ellis, C. M., Weber, H., Ploense, S. E., Barkawi, L. S., Guilfoyle, T. J., et al. (2005). Auxin response factors ARF6 and ARF8 promote jasmonic acid production and flower maturation. *Development* 132, 4107–4118. doi: 10.1242/dev.01955
- Nambara, E., and Marion-Poll, A. (2005). Abscisic acid biosynthesis and catabolism. *Annu. Rev. Plant Biol.* 56, 165–185. doi: 10.1146/annurev.arplant.56.032604.144046
- Neisiani, F. F., Sanavy, S. A. M. M., Ghanati, F., and Dolatabadian, A. (2009). Effect of foliar application of pyridoxine on antioxidant enzyme activity, proline accumulation and lipid peroxidation of Maize (*Zea mays* L.) under water deficit. *Not. Bot. Horti Agrobot. Cluj Napoca* 37, 116–121.
- Okushima, Y., Overvoorde, P. J., Arima, K., Alonso, J. M., Chan, A., Chang, C., et al. (2005). Functional genomic analysis of the AUXIN RESPONSE FACTOR gene family members in *Arabidopsis thaliana*: unique and overlapping functions of ARF7 and ARF19. *Plant Cell* 17, 444–463. doi: 10.1105/tpc.104.028316

- Park, S., Kim, H. S., Jung, Y. J., Kim, S. H., Ji, C. Y., Wang, Z., et al. (2016). Orange protein has a role in phytoene synthase stabilization in sweetpotato. *Sci. Rep.* 6:33563. doi: 10.1038/srep33563
- Ren, Z., Liu, R., Gu, W., and Dong, X. (2017). The *Solanum lycopersicum* auxin response factor SLARF2 participates in regulating lateral root formation and flower organ senescence. *Plant Sci.* 256, 103–111. doi: 10.1016/j.plantsci.2016.12.008
- Römer, S., and Fraser, P. D. (2005). Recent advances in carotenoid biosynthesis, regulation and manipulation. *Planta* 221, 305–308. doi: 10.1007/s00425-005-1533-5
- Sagar, M., Chervin, C., Mila, I., Hao, Y., Roustan, J. P., Benichou, M., et al. (2013). SLARF4, an auxin response factor involved in the control of sugar metabolism during tomato fruit development. *Plant Physiol.* 161, 1362–1374. doi: 10.1104/pp.113.213843
- Sah, S. K., Reddy, K. R., and Li, J. X. (2016). Abscisic acid and abiotic stress tolerance in crop plants. *Front. Plant Sci.* 7:571. doi: 10.3389/fpls.2016.00571
- Sandmann, G., Mer, S. R., and Fraser, P. D. (2006). Understanding carotenoid metabolism as a necessity for genetic engineering of crop plants. *Metab. Eng.* 8, 291–302. doi: 10.1016/j.ymben.2006.01.005
- Schmittgen, T. D., and Livak, K. J. (2008). Analyzing real-time PCR data by the comparative CT method. *Nat. Protoc.* 3, 1101–1108. doi: 10.1038/nprot.2008.73
- Schwartz, S. H., Qin, X., and Zeevaert, J. A. (2003). Elucidation of the indirect pathway of abscisic acid biosynthesis by mutants, genes, and enzymes. *Plant Physiol.* 131, 1591–1601. doi: 10.1104/pp.102.017921
- Sripinyowanich, S., Klomsakul, P., Boonburapong, B., Bangyeekhun, T., Asami, T., Gu, H., et al. (2013). Exogenous ABA induces salt tolerance in indica rice (*Oryza sativa* L.): the role of OsP5CS1 and OsP5CR gene expression during salt stress. *Environ. Exp. Bot.* 86, 94–105. doi: 10.1016/j.envexpbot.2010.01.009
- Takaichi, S. (2011). Carotenoids in algae distributions biosyntheses and functions. *Mar. Drugs* 9, 1101–1118. doi: 10.3390/md9061101
- Teow, C. C., Truong, V., McFeeters, R. F., Thompson, R. L., Pecota, K. V., and Yencho, G. C. (2007). Antioxidant activities, phenolic and β -carotene contents of sweet potato genotypes with varying flesh colours. *Food Chem.* 103, 829–838. doi: 10.1016/j.foodchem.2006.09.033
- Tuteja, N. (2007). Abscisic acid and abiotic stress signaling. *Plant Signal. Behav.* 2, 135–138.
- Ulmasov, T., Hagen, G., and Guilfoyle, T. J. (1997). ARF1, a transcription factor that binds to auxin response elements. *Science* 276, 1865–1868. doi: 10.1126/science.276.5320.1865
- Vishwakarma, K., Upadhyay, N., Kumar, N., Yadav, G., Singh, J., Mishra, R. K., et al. (2017). Abscisic acid signaling and abiotic stress tolerance in plants: a review on current knowledge and future prospects. *Front. Plant Sci.* 8:161. doi: 10.3389/fpls.2017.00161
- Waller, F., Furuya, M., and Nick, P. (2002). OsARF1, an auxin response factor from rice, is auxin-regulated and classifies as a primary auxin responsive gene. *Plant Mol. Biol.* 50, 415–425. doi: 10.1023/A:1019818110761
- Wang, L., Hua, D., He, J., Duan, Y., Chen, Z., Hong, X., et al. (2011). Auxin Response Factor 2 (ARF2) and its regulated homeodomain gene HB33 mediate abscisic acid response in *Arabidopsis*. *PLoS Genet.* 7:e1002172. doi: 10.1371/journal.pgen.1002172
- Wang, W. B., Kim, Y. H., Lee, H. S., Kim, K. Y., and Deng, X. (2009). Analysis of antioxidant enzyme activity during germination of alfalfa under salt and drought stresses. *Plant Physiol. Biochem.* 47, 570–577. doi: 10.1016/j.plaphy.2009.02.009
- Wang, Z., Ke, Q., Kim, M. D., Kim, S. H., Ji, C. Y., Jeong, J. C., et al. (2015). Transgenic alfalfa plants expressing the sweetpotato Orange gene exhibit enhanced abiotic stress tolerance. *PLoS One* 10:e0126050. doi: 10.1371/journal.pone.0126050
- Wilmoth, J. C., Wang, S., Tiwari, S. B., Joshi, A. D., Hagen, G., Guilfoyle, T. J., et al. (2005). NPH4/ARF7 and ARF19 promote leaf expansion and auxin-induced lateral root formation. *Plant J.* 43, 118–130. doi: 10.1111/j.1365-313X.2005.02432.x
- Xiong, L. M., and Zhu, J. K. (2003). Regulation of abscisic acid biosynthesis. *Plant Physiol.* 133, 29–36. doi: 10.1104/pp.103.025395
- Yoshihara, Y., Kiyosue, T., Nakashima, K., Yamaguchi-Shinozaki, K., and Shinozaki, K. (1997). Regulation of levels of proline as an osmolyte in plants under water stress. *Plant Cell Physiol.* 38, 1095–1102. doi: 10.1093/oxfordjournals.pcp.a029093
- Yu, L., Zhai, H., Chen, W., He, S. Z., and Liu, Q. C. (2013). Cloning and functional analysis of lycopene ϵ -cyclase (IbLCYE) gene from sweetpotato, *Ipomoea batatas* (L.) lam. *J. Integr. Agr.* 12, 773–780. doi: 10.1016/S2095-3119(13)60299-3
- Zhai, H., Wang, F. B., Si, Z. Z., Huo, J. X., Xing, L., An, Y. Y., et al. (2016). A myo-inositol-1-phosphate synthase gene, IbMIPS1, enhances salt and drought tolerance and stem nematode resistance in transgenic sweet potato. *Plant Biotechnol. J.* 14, 592–602. doi: 10.1111/pbi.12402
- Zhu, J. K. (2002). Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.* 53, 247–273. doi: 10.1146/annurev.arplant.53.091401.143329

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Kang, He, Zhai, Li, Zhao and Liu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.