



Fast-Growing SMBHs in Fast-Growing Galaxies, at High Redshifts: The Role of Major Mergers As Revealed by ALMA

Benny Trakhtenbrot^{1*}, Paulina Lira², Hagai Netzer³, Claudia Cicone^{1,4}, Roberto Maiolino^{5,6} and Ohad Shemmer⁷

¹ Department of Physics, ETH Zurich, Zurich, Switzerland, ² Departamento de Astronomía, Universidad de Chile, Santiago, Chile, ³ Raymond and Beverly Sackler Faculty of Exact Sciences, School of Physics and Astronomy and the Wise Observatory, Tel-Aviv University, Tel-Aviv, Israel, ⁴ INAF-Osservatorio Astronomico di Brera, Milan, Italy, ⁵ Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom, ⁶ Kavli Institute for Cosmology, University of Cambridge, Cambridge, United Kingdom, ⁷ Department of Physics, University of North Texas, Denton, TX, United States

OPEN ACCESS

Edited by:

Paola Marziani,
Osservatorio Astronomico di Padova
(INAF), Italy

Reviewed by:

Francoise Combes,
Observatoire de Paris, France
Daniela Bettoni,
Osservatorio Astronomico di Padova
(INAF), Italy

*Correspondence:

Benny Trakhtenbrot
benny.trakhtenbrot@phys.ethz.ch

Specialty section:

This article was submitted to
Milky Way and Galaxies,
a section of the journal
Frontiers in Astronomy and Space
Sciences

Received: 30 September 2017

Accepted: 06 November 2017

Published: 30 November 2017

Citation:

Trakhtenbrot B, Lira P, Netzer H, Cicone C, Maiolino R and Shemmer O (2017) Fast-Growing SMBHs in Fast-Growing Galaxies, at High Redshifts: The Role of Major Mergers As Revealed by ALMA. *Front. Astron. Space Sci.* 4:49. doi: 10.3389/fspas.2017.00049

We present a long-term, multi-wavelength project to understand the epoch of fastest growth of the most massive black holes by using a sample of 40 luminous quasars at $z \simeq 4.8$. These quasars have rather uniform properties, with typical accretion rates and black hole masses of $L/L_{\text{Edd}} \simeq 0.7$ and $M_{\text{BH}} \simeq 10^9 M_{\odot}$. The sample consists of “FIR-bright” sources with a previous *Herschel*/SPIRE detection, suggesting $\text{SFR} > 1,000 M_{\odot} \text{ yr}^{-1}$, as well as of “FIR-faint” sources for which *Herschel* stacking analysis implies a typical SFR of $\sim 400 M_{\odot} \text{ yr}^{-1}$. Six of the quasars have been observed by ALMA in [C II] $\lambda 157.74 \mu\text{m}$ line emission and adjacent rest-frame $150 \mu\text{m}$ continuum, to study the dusty cold ISM. ALMA detected companion, spectroscopically confirmed sub-mm galaxies (SMGs) for three sources—one FIR-bright and two FIR-faint. The companions are separated by $\sim 14\text{--}45$ kpc from the quasar hosts, and we interpret them as major galaxy interactions. Our ALMA data therefore clearly support the idea that major mergers may be important drivers for rapid, early SMBH growth. However, the fact that not all high-SFR quasar hosts are accompanied by interacting SMGs, and their ordered gas kinematics observed by ALMA, suggest that other processes may be fueling these systems. Our analysis thus demonstrates the diversity of host galaxy properties and gas accretion mechanisms associated with early and rapid SMBH growth.

Keywords: sub-millimeter galaxies, galaxy mergers, quasars: supermassive black holes, Quasars: host galaxies, high-redshift galaxies

1. INTRODUCTION

The highest-redshift quasars, observed at $z \sim 5\text{--}7$ suggest that supermassive black holes (SMBHs) with $M_{\text{BH}} \simeq 10^9 M_{\odot}$ existed about 1 Gyr after the big bang, which challenges our understanding of BH formation and early growth, and how these processes relate to the galaxies that host the earliest SMBHs.

In order to account for the observed high BH masses of the earliest quasars, many models have promoted the possibility of high-mass BH seed formation, in dense stellar populations in

proto-galaxies and/or through the direct collapse of gaseous halos (see, e.g., Natarajan, 2011; Volonteri, 2012, for reviews). Regardless of the seed mass, the subsequent BH growth must proceed at high accretion rates and high duty cycles. The former is indeed directly observed, as the accretion rate of high- z quasars approaches $L/L_{\text{Edd}} \simeq 1$ (e.g., Kurk et al., 2007; Willott et al., 2010; De Rosa et al., 2011; Trakhtenbrot et al., 2011). The latter requirement is found to be somewhat more challenging. One way to efficiently fuel SMBH accretion is through major mergers of gas rich galaxies (Sanders et al., 1988; Hopkins et al., 2006). Such mergers would be more common in dense large-scale environments. Moreover, several simulations have suggested that over-dense large-scale environments would expedite the growth of the most massive early BHs, as large amounts of inter galactic gas could stream onto the SMBHs host galaxies (Dekel et al., 2009; Di Matteo et al., 2012; Dubois et al., 2012; Costa et al., 2014).

Regardless of the exact mechanism driving the nearly continuous SMBH fueling, the low angular momentum gas is expected to trigger intense star formation (SF) throughout the host, and any interacting galaxy. Several observations of high-redshift quasars (including our own; see §2) have indeed identified intense SF, with growth rates exceeding $\text{SFR} \sim 1,000 M_{\odot} \text{ yr}^{-1}$ (e.g., Mor et al., 2012; Netzer et al., 2014, 2016). Although these high SFRs are suggestive of merger activity, the low spatial resolution of the far-IR (FIR) data prohibited any detailed investigation of this possibility. Other dedicated searches for close companions have identified some examples of major mergers (Wagg et al., 2012), but most searches did not yield convincing evidence for merger activity (e.g., Willott et al., 2005). Similarly, wide-field imaging campaigns aimed at determining whether high- z quasars are found in over-dense large-scale environments yielded ambiguous results (Willott et al., 2005; Kim et al., 2009; Husband et al., 2013; Bañados et al., 2013; Simpson et al., 2014).

Here we describe a pilot study with ALMA that aims to identify major galaxy-galaxy interactions among a sample of six fast-growing SMBHs at $z \simeq 4.8$. The full presentation of this study was recently published in *The Astrophysical Journal* (Trakhtenbrot et al., 2017, T17 hereafter), and here we only provide a brief summary of the sample, the ALMA observations, and our main results. The interested reader is encouraged to refer to T17 for any additional details. Throughout this work, we assume a cosmological model with $\Omega_{\Lambda} = 0.7$, $\Omega_{\text{M}} = 0.3$, and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

2. SAMPLE AND ALMA OBSERVATIONS

Our sample of six quasars is drawn from a larger sample of 40 sources at $z \simeq 4.8$, for which reliable estimates of M_{BH} , L/L_{Edd} , and integrated host SFRs are available through our long-term, multi-wavelength observational effort, conducted using the VLT, Gemini, *Spitzer*, and *Herschel* facilities. The $z \simeq 4.8$ quasars typically have $M_{\text{BH}} \simeq 10^9 M_{\odot}$ and $L/L_{\text{Edd}} \simeq 0.7$, and the sample covers a rather limited range in these two quantities (see Trakhtenbrot et al., 2011, T11 hereafter). The host galaxies, on the other hand, exhibit a wide range in

SFRs. While $\sim 75\%$ of the systems have $\text{SFR} \sim 400 M_{\odot} \text{ yr}^{-1}$, as determined from *Herschel* stacking analysis (“FIR-faint” systems), the outstanding 25% are individually detected and have $\text{SFR} \sim 1,000\text{--}4,000 M_{\odot} \text{ yr}^{-1}$ (“FIR-bright” systems; see Netzer et al., 2014, 2016). The *Herschel* data available prior to the ALMA campaign is therefore suggestive of a scenario where major mergers may be in play in at least in a fraction of these systems. The six quasars selected for our pilot ALMA study are equally split between “FIR-bright” and “FIR-faint” subsets, in an attempt to address this possibility.

The ALMA band-7 observations were designed to detect and resolve, at kpc scales, the emission from the prominent [C II] $\lambda 157.74 \mu\text{m}$ line and the adjacent continuum. While the continuum emission probes the spatial distribution of cold dusty ISM in the quasar hosts, the [C II] emission line—which is an efficient ISM coolant—probes their kinematics and can be used to spectroscopically confirm the nature of any companion galaxies (e.g., Maiolino et al., 2009; Wagg et al., 2012; Wang et al., 2013; Neri et al., 2014). We used the extended C34-4 configuration of ALMA, providing a resolution of $\sim 0.3''$ at 330 GHz. This corresponds to about 2 kpc at $z \simeq 4.8$. The ALMA field of view covers distances of $\sim 6.8''$, or almost 50 kpc, from the quasar locations. The chosen spectral setup provided four windows, each covering 1,875 MHz ($\sim 1,650 \text{ km s}^{-1}$), at a resolution of $\sim 30 \text{ km s}^{-1}$. On-source integrations lasted between 11 and 54 min, with longer integrations for the “FIR-faint” sources. The resulting limiting flux densities were $F_{\nu} \sim (4.2\text{--}9.2) \times 10^{-2} \text{ mJy/beam}$ (rms). At the redshifts of the quasars, and under reasonable assumptions regarding the possible shapes of their FIR SEDs, this corresponds to lower limits of roughly $4\text{--}11 M_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ (at the 3σ level).


3. RESULTS

The host galaxies of all six quasars are robustly detected, and (marginally) resolved, in both continuum and [C II] emission. As an example, we show in **Figure 1** the continuum and [C II] emission maps of one of the “FIR-faint” sources in our sample, SDSS J092303.53+024739.5 ($z_{\text{QSO}} = 4.6589$; J0923 hereafter).

In what follows, we highlight our main findings from the analysis of these data. We demonstrate these findings using different diagrams for the aforementioned source J0923. We note that many of the choices we made through the analysis of the ALMA data were motivated by recent sub-mm studies of $z \gtrsim 5$ quasars (Wang et al., 2013; Willott et al., 2015; Venemans et al., 2016). The reader is referred to T17 for a detailed discussion of our analysis and assumptions.

3.1. Quasar Hosts

We measure a wide range in (spatially-integrated) 345 GHz continuum flux densities, between $F_{\nu} \simeq 1.6\text{--}18.5 \text{ mJy}$. This wide range in continuum levels is reminiscent of that of the FIR luminosities and SFRs measured from the *Herschel*/SPIRE data (which covered rest-frame wavelengths of $\sim 45\text{--}90 \mu\text{m}$). Indeed, we find that the new ALMA continuum measurements are generally in very good agreement with the *Herschel* measurements, under reasonable assumptions regarding the




shape of the FIR SED, namely a gray-body with dust temperature $T_d = 47 \text{ K}$ and $\beta = 1.6$. Some sources require somewhat warmer dust temperatures (up to $T_d \simeq 60 \text{ K}$). Moreover, most sources are consistent with the FIR SED templates of Chary and Elbaz (2001). Importantly, we note that the ALMA continuum measurements for the FIR-faint sources, are consistent with the extrapolation of the *stacking* measurements of the *Herschel* data, thus reassuring that our interpretation of the *Herschel* results was robust. **Figure 2** demonstrates these findings for J0923.

By combining the new ALMA continuum measurements and the assumed FIR SEDs, we estimate (spatially integrated) total FIR luminosities of $L(8\text{--}1,000 \mu\text{m}) \simeq (1.9\text{--}35.5) \times 10^{12} L_\odot$. These luminosities translate to host SFRs in the range $\text{SFR} \sim 190\text{--}3,500 M_\odot \text{ yr}^{-1}$. This is, again, consistent with our *Herschel*-based findings, but now robustly resolving the hosts, which is crucial in several cases (see §3.2 below).


The spatially resolved [C II] line emission maps allow us to study the kinematics of the hosts, and estimate their dynamical masses. Most sources (at least four out of six) show [C II] velocity gradients that are consistent with rotation, as shown in the left panel of **Figure 3** for J0923. We therefore assume a simple model of an inclined rotating disk for the [C II]-emitting ISM in the hosts. Following common practices with similar data, we can then deduce *dynamical* host masses, by combining the size of the [C II]-emitting region ($D_{[\text{C II}]}$) with the typical velocity of the gas (FWHM [C II]), corrected for the inclination of the disk (i):

$$M_{\text{dyn}} = 9.8 \times 10^8 \left(\frac{D_{[\text{C II}]}}{\text{kpc}} \right) \left[\frac{\text{FWHM} [\text{C II}]}{100 \text{ km s}^{-1}} \right]^2 \sin^{-2}(i) M_\odot. \quad (1)$$

The inclination of each system is estimated from the spatial shape (morphology) of the [C II] emitting region, available from our resolved ALMA data (i.e., the major-to-minor axis ratio).



The resulting dynamical masses cover a rather limited range, $M_{\text{dyn}} \simeq (3.7\text{--}7.4) \times 10^{10} M_\odot$. By assuming that the dynamical masses are dominated by the stellar components, and considering the wide range in SFRs, this means that the lower-SFR (FIR-faint) hosts are consistent with the so-called “main sequence” of SF galaxies (e.g., Speagle et al., 2014; Steinhardt et al., 2014, and references therein), while the high-SFR (FIR-bright)



hosts would lie above it. Moreover, given the narrow range in M_{BH} and L/L_{Edd} of the quasars themselves, it appears that these host properties are not directly linked to the SMBH properties.

3.2. Companion Galaxies

Our most intriguing finding is related to the detection of several gas rich companions, which are likely interacting with the quasar hosts.

We robustly detect companion galaxies for three of the six quasar hosts, in both continuum and [C II] emission. These companions are separated by $\sim 14\text{--}45$ kpc and $|\Delta v| < 450 \text{ km s}^{-1}$ from the quasar hosts, thus being truly physically related to the quasars systems. An additional continuum source that lacks [C II] emission is detected ~ 25 kpc away from one of the FIR-bright systems, which also has a more distant spectroscopically-confirmed companion. **Figure 4** demonstrates the spectral proximity of the companion of J0923 to the quasar host.

Following the same procedures as those used for the quasar hosts, we find that the companion galaxies have continuum fluxes that translate to SFRs of $\simeq 100\text{--}200 M_{\odot} \text{ yr}^{-1}$, and dynamical masses of $M_{\text{dyn}} \simeq (2.1\text{--}10.7) \times 10^{10} M_{\odot}$. Compared to the respective quasar host masses, the companions have mass ratios $|q| \lesssim 2:1$, suggestive of *major* galaxy interactions. Moreover, the companion galaxies are consistent with being on the main sequence of SF galaxies.

4. DISCUSSION AND CONCLUSION

The most intriguing finding of our ALMA study is the identification of spectroscopically-confirmed companion galaxies for three out of the six quasar hosts in our sample. Considering the small field of view (FoV) of our ALMA data ($\sim 13.5''$ or ~ 100 kpc in diameter), the number of sub-mm bright

galaxies we find is much higher than what is found in “blind” surveys. For example, surveys of rest-frame UV selected SF galaxies predict roughly 0.01 galaxies with $\text{SFR} \simeq 100 M_{\odot} \text{ yr}^{-1}$ in a single ALMA FoV (e.g., Bouwens et al., 2015; Stark, 2016). Even more complete surveys of [C II]-emitting galaxies at $z \gtrsim 5$ predict of about 0.05 galaxies per each of our ALMA pointings (e.g., Aravena et al., 2016). We therefore conclude that fast-growing $z \sim 5$ SMBHs reside in over-dense environments in the early universe, and that their fast accumulation of mass may be related to enhanced major-merger activity. Further support for this scenario was recently presented in a large ALMA study of $z \sim 6$ quasars, using identical methods to those we used in our study (Decarli et al., 2017).

The naive expectation from the previously available *Herschel* data would be that the high-SFR (FIR-bright) systems would be associated with major mergers, while the lower-SFR (FIR-faint) systems would show no signs of interaction. Our ALMA data show a very different picture. Two of the three companions are found near FIR-faint systems, and only one is associated with a FIR-bright system. Conversely, two of the three FIR-bright systems in our sample are not associated with companion, interacting galaxies. Although in principle these quasar hosts may be in an advanced merger stage (which would remain unresolved in our data), the signatures of rotationally-dominated gas structures would not support this scenario. This is exemplified in the system J1341, which has $\text{SFR} \simeq 3,000 M_{\odot} \text{ yr}^{-1}$, and shows signatures of rotation-dominated gas and no companion galaxies (see **Figure 3**, right). The two lower-SFR systems with companion galaxies are expected to experience a later increase in SFR. This means that the low SFRs we deduced for the “FIR-faint” T11 $z \simeq 4.8$ systems cannot be simply due to the onset of “AGN feedback” in the final stages of an episode of SMBH and host growth. However, a larger sample is needed to clarify which of all these processes dominates the growth of the general $z \sim 5$ SMBH population.




FIGURE 4 | Spatially-integrated spectra of the [C II] emission line in the quasar host (**left**) and the interacting companion (**right**) of J0923. The relative velocities (top x-axis) are calculated relative to the rest-frame UV Mg II λ 2798 broad emission line of the quasar. These data confirm the physical association of the companion source to the quasar host.

The companion galaxies detected with ALMA were *not* seen in our *Spitzer* data. Given their SFRs and (dynamical) masses, and what is known about the population of rest-frame UV selected SF galaxies at $z \simeq 4.8$ (e.g., Steinhardt et al., 2014; Stark, 2016), we conclude that this is due to significant dust obscuration. This may explain the fact that many previous studies were unable to identify companions and/or over-dense environments for $z \gtrsim 5$ quasars. High resolution, spectroscopic sub-mm observations are therefore crucial for the study of mergers and environments among the highest-redshift quasars.

We are currently leading an ALMA cycle-4 program that would provide similar data for a dozen additional $z \simeq 4.8$ quasars from the T11 sample, bringing the total number of such quasars with resolved host ISM kinematics, and close companion mapping, to 18. Analysis of the ALMA data for the 12 additional sources is ongoing. Moreover, we were recently awarded *HST*/WFC3/IR time to map the *stellar* component in the host galaxies, and in the close companions of the six quasars described here. The *HST* data will also probe the larger-scale environments of the quasars, out to ~ 400 kpc, allowing us to detect any additional (unobscured) companions that may be present in this field.

AUTHOR CONTRIBUTIONS

BT led the interpretation of the ALMA data, the preparation of the paper describing our results (T17), and presented the results as a contributed talk in the “Quasars at All Cosmic Epochs” meeting. PL was the PI of the ALMA proposal and led the data analysis. All authors participated in different aspects of

the analysis, interpretation, and preparation of this study for publication.

FUNDING

HN acknowledges support by the Israel Science Foundation grant 284/13. CC gratefully acknowledges support from the Swiss National Science Foundation Professorship grant PP00P2_138979/1. CC also acknowledges funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 664931. RM acknowledges support by the Science and Technology Facilities Council (STFC) and the ERC Advanced Grant 695671 “QUENCH.”

ACKNOWLEDGMENTS

We are grateful for the organizers of the “Quasars at All Cosmic Epochs” meeting for the opportunity to present our results in a stimulating and friendly environment. We thank K. Schawinski, L. Mayer, R. Teyssier, P. Capelo, M. Dotti, and D. Fiacconi for useful discussions during the preparation of the paper describing our results (T17). The work described here made use of the ALMA data set ADS/JAO.ALMA#2013.1.01153.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), NSC and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ.

REFERENCES

- Aravena, M., Decarli, R., Walter, F., Bouwens, R. J., Oesch, P. A., Carilli, C. L., et al. (2016). ALMA spectroscopic survey in the hubble ultra deep field: Search for [CII] line and dust emission in $6 < z < 8$ galaxies. *Astrophys. J.* 833:71. doi: 10.3847/1538-4357/833/1/71
- Bañados, E., Venemans, B. P., Walter, F., Kurk, J., Overzier, R., and Ouchi, M. (2013). The galaxy environment of a QSO at $z \sim 5.7$. *Astrophys. J.* 77:178. doi: 10.1088/0004-637X/773/2/178
- Bouwens, R. J., Illingworth, G. D., Oesch, P. A., Trenti, M., Labbé, I., Bradley, L., et al. (2015). UV luminosity functions at redshifts $z \sim 4$ to $z \sim 10$: 10,000 galaxies from HST legacy fields. *Astrophys. J.* 803:34. doi: 10.1088/0004-637X/803/1/34
- Chary, R.-R., and Elbaz, D. (2001). Interpreting the cosmic infrared background: constraints on the evolution of the dustenshrouded star formation rate. *Astrophys. J.* 556, 562–581. doi: 10.1086/321609
- Costa, T., Sijacki, D., Trenti, M., and Haehnelt, M. G. (2014). The environment of bright QSOs at $z \sim 6$: star-forming galaxies and X-ray emission. *Monthly Not. R. Astron. Soc.* 439, 2146–2174. doi: 10.1093/mnras/stu101
- De Rosa, G., Decarli, R., Walter, F., Fan, X., Jiang, L., Kurk, J. D., et al. (2011). Evidence for non-evolving Fe II/Mg II ratios in rapidly accreting $z \sim 6$ QSOs. *Astrophys. J.* 739:56. doi: 10.1088/0004-637X/739/2/56
- Decarli, R., Walter, F., Venemans, B. P., Bañados, E., Bertoldi, F., Carilli, C. L., et al. (2017). Rapidly star-forming galaxies adjacent to quasars at redshifts exceeding 6. *Nature* 545:457. doi: 10.1038/nature22358
- Dekel, A., Birnboim, Y., Engel, G., Freundlich, J., Goerdt, T., Mumcuoglu, M., et al. (2009). Cold streams in early massive hot haloes as the main mode of galaxy formation. *Nature* 457, 451–454. doi: 10.1038/nature07648
- Di Matteo, T., Khandai, N., DeGraf, C., Feng, Y., Croft, R., Lopez, J., et al. (2012). Cold flows and the first quasars. *Astrophys. J.* 745:L29. doi: 10.1088/2041-8205/745/2/L29
- Dubois, Y., Pichon, C., Haehnelt, M., Kimm, T., Slyz, A., Devriendt, J., et al. (2012). Feeding compact bulges and supermassive black holes with low angular momentum cosmic gas at high redshift. *Monthly Not. R. Astron. Soc.* 423, 3616–3630. doi: 10.1111/j.1365-2966.2012.21160.x
- Hopkins, P. F., Hernquist, L., Cox, T. J., Di Matteo, T., Robertson B. E., and Springel, V. (2006). A unified, merger driven model of the origin of starbursts, quasars, the cosmic X ray background, supermassive black holes, and galaxy spheroids. *Astrophys. J. Suppl. Series* 163, 1–49. doi: 10.1086/499298
- Husband, K., Bremer, M. N., Stanway, E. R., Davies, L. J. M., Lehnert, M. D., and Douglas, L. S. (2013). Are $z \sim 5$ QSOs found in the most massive high redshift over-densities? *Monthly Not. R. Astron. Soc.* 432, 2869–2877. doi: 10.1093/mnras/stt642
- Kim, S., Stiavelli, M., Trenti, M., Pavlovsky, C. M., Djorgovski, S. G., Scarlata, C., et al. (2009). The environments of high-redshift quasi-stellar objects. *Astrophys. J.* 695, 809–817. doi: 10.1088/0004-637X/695/2/809
- Kurk, J. D., Walter, F., Fan, X., Jiang, L., Riechers, D. A., Rix, H., et al. (2007). Black hole masses and enrichment of $z \sim 6$ SDSS quasars. *Astrophys. J.* 669, 32–44. doi: 10.1086/521596
- Maiolino, R., Caselli, P., Nagao, T., Walmsley, M., De Breuck, C., and Meneghetti, M. (2009). Strong [CII] emission at high redshift. *Astron. Astrophys.* 500, L1–L4. doi: 10.1051/0004-6361/200912265
- Mor, R., Netzer, H., Trakhtenbrot, B., Shemmer, O., and Lira, P. (2012). Extreme star formation in the host galaxies of the fastest growing supermassive black holes at $z = 4.8$. *Astrophys. J.* 749:L25. doi: 10.1088/2041-8205/749/2/L25
- Natarajan, P. (2011). The formation and evolution of massive black hole seeds in the early Universe. *Bull. Astron. Soc. India* 39:145. doi: 10.1142/9789814374774_0013
- Neri, R., Downes, D., Cox, P., and Walter, F. (2014). High-resolution C+ imaging of HDF850.1 reveals a merging galaxy at $z = 5.185$. *Astron. Astrophys.* 562:A35. doi: 10.1051/0004-6361/201322528
- Netzer, H., Lani, C., Nordon, R., Trakhtenbrot, B., Lira, P., and Shemmer, O. (2016). Star formation black hole growth and dusty tori in the most luminous AGNs at $z = 2-3.5$. *Astrophys. J.* 819:123. doi: 10.3847/0004-637X/819/2/123
- Netzer, H., Mor, R., Trakhtenbrot, B., Shemmer, O., and Lira, P. (2014). Star formation and black hole growth at $z \simeq 4.8$. *Astrophys. J.* 791:34. doi: 10.1088/0004-637X/791/1/34
- Sanders, D. B., Soifer, B. T., Elias, J. H., Madore, B. F., Matthews, K., Neugebauer, G., et al. (1988). Ultraluminous infrared galaxies and the origin of quasars. *Astrophys. J.* 325:74. doi: 10.1086/165983
- Simpson, C., Mortlock, D. J., Warren, S., Cantalupo, S., Hewett, P. C., McLure, R. J., et al. (2014). No excess of bright galaxies around the redshift 7.1 quasar ULAS J1120+0641. *Monthly Not. R. Astron. Soc.* 442, 3454–3461. doi: 10.1093/mnras/stu1116
- Speagle, J. S., Steinhardt, C. L., Capak, P. L., and Silverman, J. D. (2014). A highly consistent framework for the evolution of the star-forming “mail sequence” from $\sim 0-6$. *Astrophys. J. Suppl. Series* 214:15. doi: 10.1088/0067-0049/214/2/15
- Stark, D. P. (2016). Galaxies in the first billion years after the big bang. *Annu. Rev. Astron. Astrophys.* 54, 761–803. doi: 10.1146/annurev-astro-081915-023417
- Steinhardt, C. L., Speagle, J. S., Capak, P. L., Silverman, J. D., Carollo, C. M., Dunlop, J. S., et al. (2014). Star formation at $4 < z < 6$ from the *Spitzer* large area survey with hyper-suprime-Cam (SPLASH). *Astrophys. J.* 791:L25. doi: 10.1088/2041-8205/791/2/L25
- Trakhtenbrot, B., Lira, P., Netzer, H., Cicone, C., Maiolino, R., and Shemmer, O. (2017). ALMA observations show major mergers among the host Galaxies of fast-growing, high-redshift, supermassive black holes. *Astrophys. J.* 836:8. doi: 10.3847/1538-4357/836/1/8
- Trakhtenbrot, B., Netzer, H., Lira, P., and Shemmer, O. (2011). Black hole mass and growth rate at $z \simeq 4.8$: a short episode of fast growth followed by short duty cycle activity. *Astrophys. J.* 730:7. doi: 10.1088/0004-637X/730/1/7
- Venemans, B. P., Walter, F., Zschaechner, L., Decarli, R., Rosa, G. D., Findlay, J. R., et al. (2016). Bright [C II] and dust emission in three $z < 6.6$ quasar host Galaxies observed by ALMA. *Astrophys. J.* 816:37. doi: 10.3847/0004-637X/816/1/37
- Volonteri, M. (2012). The formation and evolution of massive black holes. *Science* 337, 544–547. doi: 10.1126/science.1220843
- Wagg, J., Wiklind, T., Carilli, C. L., Espada, D., Peck, A., Riechers, D., et al. (2012). [C II] Line emission in massive star-forming galaxies at $z = 4.7$. *Astrophys. J.* 752:L30. doi: 10.1088/2041-8205/752/2/L30
- Wang, R., Wagg, J., Carilli, C. L., Walter, F., Lentati, L., Fan, X., et al. (2013). Star formation and gas kinematics of quasar host Galaxies at $z \sim 6$: new insights from ALMA. *Astrophys. J.* 773:24. doi: 10.1088/2041-8205/774/2/L24
- Willott, C. J., Albert, L., Arzoumanian, D., Bergeron, J., Crampton, D., Delorme, P., et al. (2010). Eddington-limited accretion and the black hole mass function at redshift 6. *Astron. J.* 140, 546–560. doi: 10.1088/0004-6256/140/2/546
- Willott, C. J., Bergeron, J., and Omont, A. (2015). Star formation rate and dynamical mass of 10^8 solar mass black hole host galaxies at redshift 6. *Astrophys. J.* 801:123. doi: 10.1088/0004-637X/801/2/123
- Willott, C. J., Percival, W. J., McLure, R. J., Crampton, D., Hutchings, J. B., Jarvis, M. J., et al. (2005). Imaging of SDSS $z > 6$ quasar fields: gravitational lensing, companion galaxies, and the host dark matter halos. *Astrophys. J.* 626, 657–665. doi: 10.1086/430168

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer DB and handling Editor declared their shared affiliation.

Copyright © 2017 Trakhtenbrot, Lira, Netzer, Cicone, Maiolino and Shemmer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.