



Erratum: Submillimeter and Far-Infrared Polarimetric Observations of Magnetic Fields in Star-Forming Regions

Frontiers Production Office*

Frontiers Media SA, Lausanne, Switzerland

Keywords: molecular clouds, far-infrared (FIR), magnetic fields, star formation, submillimeter astronomy, polarimetry

OPEN ACCESS

Approved by:

Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*Correspondence:

Frontiers Production Office
production.office@frontiersin.org

Specialty section:

This article was submitted to
Stellar and Solar Physics,
a section of the journal
Frontiers in Astronomy and Space
Sciences

Received: 14 August 2019

Accepted: 15 August 2019

Published: 19 August 2019

Citation:

Frontiers Production Office (2019)
Erratum: Submillimeter and
Far-Infrared Polarimetric Observations
of Magnetic Fields in Star-Forming
Regions.
Front. Astron. Space Sci. 6:58.
doi: 10.3389/fspas.2019.00058

An Erratum on

Submillimeter and Far-Infrared Polarimetric Observations of Magnetic Fields in Star-Forming Regions

by Pattle, K., and Fissel, L. (2019). *Front. Astron. Space Sci.* 6:15. doi: 10.3389/fspas.2019.00015

Due to a production error, a copyrighted figure was used as part of **Figure 4** in the original article. This figure and the corresponding part label have been removed from the article and the updated **Figure 4** can be seen below.

The publisher apologizes for this error and states that this does not change the scientific conclusions of the article in any way. The original version of this article was updated on July 4th, 2019.

REFERENCES

- Fissel, L. M., Ade, P. A. R., Angilè, F. E., Ashton, P., Benton, S. J., Devlin, M. J., et al. (2016). Balloon-borne submillimeter polarimetry of the Vela C molecular cloud: systematic dependence of polarization fraction on column density and local polarization-angle dispersion. *Astrophys. J.* 824:134. doi: 10.3847/0004-637X/824/2/134
- Soler, J. D., Ade, P. A. R., Angilè, F. E., Ashton, P., Benton, S. J., Devlin, M. J., et al. (2017). The relation between the column density structures and the magnetic field orientation in the Vela C molecular complex. *Astron. Astrophys.* 603:A64. doi: 10.1051/0004-6361/201730608

Copyright © 2019 Frontiers Production Office. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 4 | Polarization observations of the Vela C cloud from the BLASTPol telescope. **(A)** Magnetic field orientation (texture) inferred from the BLASTPol 500 μm data overlaid on a *Herschel*-derived column density (N_{H}) map—Soler et al. (2017), reproduced with permission © ESO. Four sub-region are labeled, while the shaded pink region indicates where the dust is heated by a compact H_{II} region. **(B)** Map of polarized intensity with contours of total intensity (green) overlaid (from Figure 3 of Fissel et al., 2016© AAS. Reproduced with permission).