



Selective Serotonin Reuptake Inhibitor-Treatment Does Not Show Beneficial Effects on Cognition or Amyloid Burden in Cognitively Impaired and Cognitively Normal Subjects

OPEN ACCESS

Edited by:

Kristina Endres,
Johannes Gutenberg University
Mainz, Germany

Reviewed by:

Georg S. Kranz,
The Hong Kong Polytechnic
University, Hong Kong SAR, China
Andreas Fellgiebel,
Johannes Gutenberg University
Mainz, Germany

*Correspondence:

Caroline Bouter
caroline.bouter@
med.uni-goettingen.de

†Data used in the preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in the analysis or writing of this report.

A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Specialty section:

This article was submitted to Alzheimer's Disease and Related Dementias, a section of the journal Frontiers in Aging Neuroscience

Received: 24 February 2022

Accepted: 24 May 2022

Published: 23 June 2022

Yvonne Bouter¹ and Caroline Bouter^{2*} for the Alzheimer's Disease Neuroimaging Initiative†

¹ Department of Psychiatry and Psychotherapy, University Medical Center Göttingen (UMG), Georg-August-University, Göttingen, Germany, ² Department of Nuclear Medicine, University Medical Center Göttingen (UMG), Georg-August-University, Göttingen, Germany

Preclinical studies indicate that selective serotonin reuptake inhibitors (SSRI) have beneficial effects on Alzheimer-related pathologies. Therefore, the aim of this study was to evaluate the influence of SSRI-treatment on amyloid burden in ¹⁸F-Florbetapir-positron emission tomography (PET) and on cognition in cognitively normal and cognitively impaired subjects. We included $n = 755$ cognitively impaired and $n = 394$ cognitively normal participants from the Alzheimer's Disease Neuroimaging Initiative (ADNI) that underwent at least one ¹⁸F-Florbetapir-PET. Standardized uptake ratios (SUVR) and the Alzheimer Disease Assessment Scale-cognitive subscale (ADAS) scores as well as follow-up results were compared between subgroups with a history of SSRI-treatment (SSRI+) and without SSRI-treatment (SSRI-) as well as in subgroups of SSRI+/Depression+ and SSRI+/Depression- and SSRI-/Depression+ and SSRI-/Depression-. ¹⁸F-Florbetapir-PET did not show significant differences of SUVR between the SSRI+ and SSRI- groups in both, cognitively impaired and cognitively normal participants. There were no differences in subgroups of SSRI+/Depression+ and SSRI+/Depression- and SSRI-/Depression+ and SSRI-/Depression-. However, SUVR showed a dose-dependent inverse correlation to the duration of medication in cognitively normal and in cognitively impaired patients. SSRI-treatment did not show an effect on ADAS scores. Furthermore, there was no effect on follow-up SUVR or on follow-up ADAS scores. Overall, SSRI-treatment did not show beneficial effects on amyloid load nor on cognition.

Keywords: positron emission tomography, selective serotonin reuptake inhibitors, amyloid-PET imaging, florbetaben, Alzheimer's disease

INTRODUCTION

Alzheimer's disease (AD), the most common form of dementia, is a progressive neurodegenerative disease characterized by memory loss and decline of cognitive function. Despite extensive treatment efforts, AD remains incurable and novel therapies to prevent, slow down or delay the onset of the disease are urgently needed. Without effective therapies, the number of patients with dementia worldwide is estimated to reach more than 130 million by 2050 (Cummings et al., 2016).

Given the need for disease-modifying therapies for AD, drug repurposing may be a promising approach. Preclinical studies indicate that antidepressants, particularly selective serotonin reuptake inhibitors (SSRI), have beneficial effects on AD-related biomarkers including amyloid plaques, one of the major pathological hallmarks in AD (Cirrito et al., 2011, 2020; Sheline et al., 2014). Neuritic plaques consist of aggregated A β peptides that are formed within neurons by sequentially cleavage of the amyloid precursor protein (APP). Dysregulation of A β production and A β clearance leads to accumulation of hydrophobic A β forms and the formation of extracellular plaques (Chen et al., 2017; Guo et al., 2020). While the pathway of APP processing is well-characterized, mechanisms of its regulation are not yet fully understood. APP processing can be modulated by several signaling pathways including NMDA, acetylcholine and serotonin signaling systems. Serotonin receptors (5-HT α) might affect APP processing and A β levels. *In vitro* studies could show that activation of 5-HT α , 5-HT β , and 5-HT γ increases non-amyloidogenic APP processing (Nitsch et al., 1996; Robert et al., 2001; Pakaski et al., 2005; Shen et al., 2011). In addition, treatment with SSRI reduced A β levels and amyloid plaque burden in different mouse models of AD (Tucker et al., 2005; Nelson et al., 2007; Cirrito et al., 2011, 2020). Furthermore, beneficial effects of antidepressant medication on cognition could be shown in depressed patients, while there was no effect on non-depressed individuals (Prado et al., 2018).

However, whether chronic SSRI-treatment influences amyloid burden and cognition in humans remains unclear. A few studies suggest a possible positive benefit of SSRI treatment on the risk of developing AD, whereas the effect on amyloid burden remains controversial (Kessing et al., 2009; Cirrito et al., 2011; Bartels et al., 2018, 2020).

The aim of this study is the evaluation of the influence of SSRI-treatment on amyloid burden in ^{18}F -Florbetapir positron emission tomography (PET) in cognitively normal and cognitively impaired individuals.

MATERIALS AND METHODS


Data used in this study were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database.¹ The ADNI was launched in 2003 as a public-private partnership, led by Principal Investigator Michael W. Weiner, MD. The primary

¹<http://adni.loni.usc.edu>

goal of ADNI has been to test whether serial magnetic resonance imaging (MRI), positron emission tomography (PET), other biological markers, and clinical and neuropsychological assessment can be combined to measure the progression of mild cognitive impairment (MCI) and early Alzheimer's disease (AD). ADNI is a multicentered project that aims to improve clinical research on AD unifying data on demographics, clinical and cognitive assessments as well as on genetic, biochemical and imaging biomarkers. Standardized protocols and unlimited data access allow analysis of an enormous amount of data from more than 1800 patients that have been included to the ADNI database to date. Groups of AD patients, patients with mild cognitive impairment (MCI), and cognitively normal elderly controls were included in four phases starting in 2004.

Study Sample

In this study, ADNI data from all four phases, ADNI-1, ADNI-GO, ADNI-2, and ADNI-3, were downloaded from the ADNI database (see text footnote 1) on May 7, 2021. Patients aged 57–93 with at least one available ^{18}F -Florbetapir-PET were downloaded ($n = 1296$; **Figure 1**). Patient characteristics, medical history, medication, Mini Mental State Examination (MMSE), Clinical Dementia Rating score (CDR), the Alzheimer's Disease Assessment Scale–Cognitive Subscale (ADAS), depression-history, education years, ApoE4 status and PET results were assessed. Patients with insufficient or inconsistent data on cognition or prior medication were excluded ($n = 22$). $N = 20$ patients were excluded as SSRI-treatment was paused before the baseline PET and $n = 105$ patients without significant objective amnesic dysfunction but with subjective memory concerns were excluded.



According to the ADNI protocol, participants were categorized as cognitively normal and cognitively impaired with subgroups of MCI and Alzheimer's dementia (AD). AD patients showed a significant subjective and objective memory loss (based on scores on the WMS-R Logical Memory II subscale) affecting activities of daily living with a MMSE below 26 and CDR of 0.5 or 1 plus. MCI patients were defined as patients showing significant memory issues with a MMSE from 24 to 30 and CDR of 0.5 plus an abnormal score on the WMS-R Logical Memory II subscale. Cognitively normal participants showed no signs of dementia with normal cognition (MMSE from 24 to 30 and a CDR of 0 plus a WMS-R Logical Memory II subscale score above education-adjusted cutoffs).

Detailed information about the ADNI inclusion and exclusion criteria can be found online.²

For our analysis, all groups and subgroups were divided into history of SSRI treatment (SSRI+) and without SSRI treatment (SSRI-).

A total of $n = 164$ patients were currently treated with an SSRI at the time of the PET (Citalopram: $n = 69$; Escitalopram $n = 16$; Fluoxetine $n = 24$; Paroxetine $n = 7$; Sertraline $n = 48$; **Table 1**).

Alzheimer's Disease Neuroimaging Initiative ¹⁸F-Florbetapir-PET/CT

¹⁸F-Florbetapir-PET/CT was performed according to a standardized imaging protocol with 4×5 min frames acquired 50–70 min post-injection of $370 (\pm 10\%)$ MBq ¹⁸F-Florbetapir. All PET images were reviewed for quality control by the ADNI PET QC team and transmitted in DICOM format to the Laboratory of Neuroimaging (LONI) for storage. In order to uniformize data from different origins, PET image data were pre-processed with motion correction, time frame averaging, reorientation in a standardized $160 \times 160 \times 96$ matrix with a voxel size of 1.5 mm and smoothing with a scanner-specific filter function determined from Hoffman phantom scans during the certification process (more details on adni.loni.usc.edu).

PET images were further processed with FreeSurfer v7.1.1 for an MRI-based definition of multiple cortical regions as well as reference regions for normalization.

Amyloid load was quantitatively analyzed using standardized uptake ratios (SUVr) of composite regions normalized by cerebellar uptake. In order to analyze possible region-specific

differences, SUVr of frontal, parietal, temporal and cingulate cortex regions were also obtained. As described before, intensity normalization of SUVr was performed using a FreeSurfer-defined whole cerebellum region for cross-sectional analysis and a FreeSurfer-defined composite reference region for longitudinal studies.

Statistical Analysis

Statistical analysis was performed using SPSS Statistics version 27 (IBM, Armonk, NY, United States) and GraphPad Prism version 9 (GraphPad Software, San Diego, CA, United States).

Differences between groups were tested using non-parametric tests (as data did not pass normality tests, e.g., Shapiro–Wilk test $p = 0.0006$), therefore Mann–Whitney U test or Kruskal–Wallis test followed by *post hoc* multiple comparison were used as indicated. Univariate analysis of covariance (ANCOVA) was used for covariate adjustment as indicated. Chi-square test or Fisher's exact test were used for categorical variables. Relationships between two variables were assessed using Spearman correlation and simple linear regression.

For follow-up PET and ADAS data changes were calculated as absolute and relative change between baseline and follow-up (Δ SUVr/ Δ ADAS; $\Delta\%$ SUVr/ $\Delta\%$ ADAS).

Significance levels are given as follows: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

RESULTS

Baseline Characteristics

A total of $n = 755$ cognitively impaired (AD and MCI patients) and $n = 394$ cognitively normal participants from ADNI phases ADNI-1 ($n = 82$), ADNI-GO ($n = 181$), ADNI-2 ($n = 682$), and ADNI-3 ($n = 204$) that underwent at least one ¹⁸F-Florbetapir-PET were included in this study. Baseline characteristics for all patients categorized into diagnostic groups are shown in **Table 2**. Patients were categorized as AD ($n = 160$), MCI ($n = 595$) and cognitively normal ($n = 394$) according to clinical symptoms and results of neuropsychological assessments following the ADNI protocol (**Figure 1**).

The group of AD patients was slightly older compared to the MCI group (Kruskal–Wallis test with Dunn's multiple comparison test, $p = 0.0093$) while there were no differences between all other groups (Kruskal–Wallis test, $p > 0.07$). Groups of cognitively impaired patients showed a higher number of male patients compared to cognitively normal controls (Chi-square test, $p = 0.0002$).

MMSE, CDR and ADAS showed significant differences between AD and MCI as well as between AD or MCI compared to cognitively normal patients after adjusting for age and gender as covariates (ANCOVA, $p < 0.001$, **Table 2**).

There were no differences between age, gender, MMSE scores, ApoE4-alleles or diagnosis between cognitively impaired SSRI+ and SSRI- patients ($p > 0.11$; **Table 2**). There were no differences between age, MMSE scores or ApoE4-alleles between cognitively normal SSRI+ and SSRI- patients ($p > 0.19$; **Table 2**). Slight differences were observed in gender distribution with

²<https://adni.loni.usc.edu/study-design/>

TABLE 1 | Selective serotonin reuptake inhibitor (SSRI) treatment.

	SSRI+		Duration of medication	Dose
	<i>n</i>	%	Months (SD)	mg (SD)
All	164	100	56.27 (64.92)	
Citalopram	69	42	43.75 (50.61)	19.64 (10.79)
Escitalopram	16	10	47.27 (47.41)	14.06 (6.88)
Fluoxetine	24	15	113.5 (94.96)	28.33 (14.94)
Paroxetine	7	4	39.0 (28.65)	25.71 (12.72)
Sertraline	48	29	54.23 (63.08)	72.86 (43.71)

TABLE 2 | Patient characteristics.

Characteristics	Total		SSRI+		SSRI-		
1. Cognitively impaired patients							
	Total N = 755		SSRI+ N = 131		SSRI- N = 624		
	Mean	SD	Mean	SD	Mean	SD	p
Age	73.59	7.9	73.04	7.6	73.7	8.0	0.4095
MMSE	25.96	3.9	25.52	4.01	26.1	3.8	0.2549
ADAS	19.66	11.2	22.43	12.1	19.0	10.8	0.0126
Education (years)	16	2.7	15.73	2.6	16.06	2.7	0.185
	N	%	N	%	N	%	
<i>Diagnosis</i>							
AD	160	21	30	21	130	21	>0.9999
MCI	595	79	101	79	494	79	
<i>Gender</i>							
Female	326	44	65	49	261	42	0.1146
Male	429	56	66	51	363	58	
<i>ApoE4 alleles</i>							
0	378	50	56	43	322	52	0.2385
1	287	38	57	43	230	37	
2	90	12	18	14	72	12	
<i>SSRIs</i>							
All			131	100			
Citalopram			52	40			
Escitalopram			9	7			
Fluoxetine			22	16			
Paroxetine			5	4			
Sertraline			39	32			
<i>History of depression</i>							
Depression	249	33	94	69	155		<0.001
No depression	506	67	37	31	469		
2. Objectively cognitively normal patients							
	Total N = 394		SSRI+ N = 33		SSRI- N = 361		
	Mean	SD	Mean	SD	Mean	SD	p
Age	74.44	7.13	72.08	5.8	74.2	6.9	0.1930
MMSE	28.99	1.21	28.68	1.7	29.0	1.2	0.3556
ADAS	9.3	5.0	11.5	7.8	9.1	4.5	0.0383
Education (years)	16.54	2.56	15.64	2.25	16.66	2.57	0.0138
	N	%	N	%	N	%	
<i>Gender</i>							
Female	215		26	79	189	52	0.0035
Male	179		7	21	172	48	
<i>ApoE4 alleles</i>							
0	282		24	73	258	72	0.5615
1	100		9	27	91	25	
2	12		-	0	12	3	
<i>SSRI</i>							
All			33	100			
Citalopram			17	52			
Escitalopram			7	21			
Fluoxetine			2	6			
Paroxetine			2	6			
Sertraline			5	15			
<i>History of depression</i>							
Depression	51	13	19	58	32	9	<0.001
No depression	343	87	14	42	329	91	

AD, Alzheimer's dementia; MCI, mild cognitive impairment. Gray shades highlight patients with SSRI medication.

proportionately more women in the SSRI+ group (Chi-square test; $p = 0.002$; **Table 2**) which was considered irrelevant for further analysis. Amyloid load did not differ between female and male patients (Mann–Whitney U test, $p = 0.1308$).

$N = 300$ patients had a history of depression. The amount of patients with a history of depression was significantly higher in the SSRI+ group compared to the SSRI– group (Chi-square test; $p < 0.001$; **Table 2**).

Positron Emission Tomography Results

A baseline PET was available in all patients ($n = 1149$). SUVRs were significantly higher in the AD subgroup compared to the MCI subgroup as well as in AD or MCI compared to cognitively normal controls after adjusting for age and gender as covariates (**Figure 2A**, ANCOVA, $p < 0.001$).

Selective Serotonin Reuptake Inhibitors Treatment

There were no significant differences of SUVR between the SSRI+ and SSRI– patients regardless of the diagnosis (**Figure 2A**, Mann–Whitney U test; AD: $p = 0.667$; MCI: $p = 0.169$; cognitively normal controls: $p = 0.188$).

SUVR showed a significant negative correlation with the duration of SSRI-treatment in cognitively normal SSRI+ as well as in cognitively impaired SSRI+ patients (**Figures 2B,C**, Spearman correlation, cognitive normal: $p = 0.0038$; $r = -0.4965$; cognitively impaired: $p = 0.025$; $r = -0.1958$).

SUVRs did not differ between groups treated with citalopram, escitalopram, fluoxetine, paroxetine or sertraline, neither in SSRI+ CI patients (Kruskal–Wallis test; $p = 0.3276$) nor in SSRI+ cognitively normal controls (Kruskal–Wallis test; $p = 0.8296$).

There was no dose-dependency of SUVR in any of the groups treated with citalopram, escitalopram, fluoxetine, paroxetine or sertraline, neither in SSRI+ cognitively impaired patients (Kruskal–Wallis test; $p > 0.14$) nor in SSRI+ cognitive normal patients (Kruskal–Wallis test; $p > 0.3$).

Regional Differences

Regional differences of amyloid load were analyzed in frontal, cingulate, parietal and temporal cortex. There were no significant differences between regional SUVR of SSRI+ and SSRI– regardless of the diagnosis (Kruskal–Wallis test, $p > 0.5$ in all groups).

History of Depression

^{18}F -Florbetapir uptake did not show significant differences between SSRI+ and SSRI– patients with and without history of depression regardless of the diagnosis (**Figure 3**; Kruskal–Wallis test; AD: $p = 0.8304$; MCI: $p = 0.5308$; cognitive normal controls: $p = 0.1937$).

Follow-Up ^{18}F -Florbetapir-PET

At least one follow-up ^{18}F -Florbetapir-PET was available in $n = 650$ patients. Mean follow-up period was 26.49 months for one follow-up PET. A second or third follow-up was only available in the MCI and cognitively normal group with a mean

period of 57.34 months for a second follow-up PET scan ($n = 361$) and 74.86 months for a third follow-up PET scan ($n = 184$).

In order to analyze changes of amyloid deposition, differences of SUVR between the baseline PET and available follow-up PETs were calculated (ΔSUVR). Longitudinal changes of SUVR did not show any significant differences between SSRI+ and SSRI– AD-patients after one follow-up (**Table 3** and **Figure 4A**; Mann–Whitney U test; $p = 0.4269$).

There were no significant differences between SSRI+ and SSRI– MCI-patients after the first, second or third follow-up (**Table 3** and **Figure 4A**; Mann–Whitney U test; first follow-up: $p = 0.9501$; second follow up: $p = 0.3405$; third follow-up: $p = 0.2387$). Furthermore, we did not detect significant differences between ΔSUVR in the cognitively normal group either (**Table 3** and **Figure 4A**; Mann–Whitney U test; first follow-up: $p = 0.6367$; second follow up: $p = 0.8637$; third follow-up: $p = 0.921$).

Cognition

In order to analyze differences in cognition as well as longitudinal changes of cognition between SSRI+ and SSRI– groups the Alzheimer Disease Assessment Scale-cognitive subscale (ADAS) scores were used. Baseline ADAS as well as follow-up scores were collected as close to the date of the corresponding baseline and follow-up PET scan as possible.

Baseline ADAS was available in all patients ($n = 1149$). ADAS scores correlated with amyloid burden in PET (**Figure 5A**, Spearman $r = 0.4396$; $p < 0.001$). ADAS was significantly higher in AD patients compared to MCI patients and in MCI and AD patients compared to cognitively normal controls after adjusting for age, gender and education as covariates (**Figure 5B**; ANCOVA; $p < 0.001$). There were no significant differences between SSRI+ and SSRI– subgroups after adjusting for age, gender and education as covariates (**Figure 5B**, ANCOVA, AD: $p = 0.238$; MCI: $p = 0.153$; NMC: $p = 0.063$).

At least one follow-up ADAS score was available in $n = 646$ patients with a mean follow-up of 25.5 months. A second and third follow-up ADAS score was available in the MCI and cognitively normal group with a mean period of 60.67 months for a second follow-up ($n = 356$) and 73.93 months for a third follow-up ($n = 179$). In order to analyze longitudinal changes of cognition, differences of ADAS between the baseline score and available follow-up scores were calculated (ΔADAS ; **Table 3**).

There were no significant differences of the ADAS score between SSRI+ and SSRI– AD-patients after one follow-up (**Table 3** and **Figure 4B**; Mann–Whitney U test; $p = 0.2593$). Furthermore, we did not detect significant differences between ΔADAS in the MCI-group nor in the cognitively normal group after the first, second or third follow-up (**Table 3** and **Figure 4B**; Mann–Whitney U test; MCI: first follow-up: $p = 0.5651$; second follow up: $p = 0.5142$; third follow-up: $p = 0.2465$; cognitively normal group: first follow-up: $p = 0.2084$; second follow up: $p = 0.595$; third follow-up: $p = 0.359$).

ApoE4

ApoE4 carriers showed significantly higher SUVR compared to non-carriers in cognitively impaired patients as well as in cognitively normal subjects (Mann–Whitney U test; $p < 0.001$;





TABLE 3 | Follow up SUVR and ADAS.

	Delta SUVR			Delta ADAS		
	FU1	FU2	FU3	FU1	FU2	FU3
SSRI+ AD	+0.039 (4%)	na	na	+9.5 (29%)	na	na
SSRI- AD	+0.018 (1.8%)	na	na	+4.4 (14%)	na	na
SSRI+ MCI	+0.014 (1.6%)	+0.029 (3.3%)	+0.045 (4.9%)	+2.6 (14%)	+4.3 (22%)	+6.9 (36%)
SSRI- MCI	+0.014 (1.6%)	+0.021 (2.5%)	+0.018 (2.1%)	+1.6 (10%)	+3.7 (23%)	+3.1 (19%)
SSRI+ CN	+0.007 (1%)	+0.023 (3%)	+0.03 (3.8%)	+1.4 (11%)	+1.7 (13%)	+6.7 (46%)
SSRI- CN	+0.013 (1.7%)	+0.019 (2.5%)	+0.02 (2.6%)	-0.1 (1.1%)	+0.9 (10%)	+1.5 (17%)

AD, Alzheimer's dementia; MCI, mild cognitive impairment; FU, follow up; CN, cognitively normal controls. Gray shades highlight patients with SSRI medication.

Figure 6). However, there were no differences in the SUVR between SSRI+ and SSRI- ApoE4 carriers regardless of the diagnosis (Mann-Whitney U test; AD: $p = 0.7909$; MCI: $p = 0.3644$; cognitively normal: $p = 0.9065$; **Figure 6**).

DISCUSSION

Amyloid plaques are a major pathological hallmark of AD and the modulation of plaques, APP processing, and related signaling pathways has been a research focus for novel AD-therapies for many years. The role of different neurotransmitter systems in the development of AD is still unclear. Recent evidence suggests that the serotonergic system, that has mainly been studied in mood disorders so far, might play a role in the pathogenesis of AD (Cirrito et al., 2011; Ramos-Rodriguez et al., 2013).

Mid-life or late-life depression is considered a risk factor for the development of AD. However, whether depression is a prodromal symptom or a true etiologic risk factor for AD remains controversial. Both, AD and depression share similar neuropathological changes as neuroinflammation, elevated oxidative stress markers and synaptic dysfunctions and an increased cortical amyloid burden is also seen in patients with a lifetime history of depression (Chung et al., 2015; Herbert and Lucassen, 2016; Mahgoub and Alexopoulos, 2016).

Therefore, the modulation of serotonergic systems with SSRIs might be a potential treatment option for AD patients with or without a history of depression.

Several preclinical studies demonstrated beneficial effects of SSRI-treatment on AD-pathologies in different mouse models of AD suggesting a modulation of APP processing by SSRIs. Cirrito et al. (2011) first showed a serotonin-dependent reduction of A β levels in the brain interstitial fluid (ISF) in APP/PS1 mice after short-term treatment with citalopram or fluoxetine. Furthermore, reduced cortical and hippocampal plaque burden as well as decreased A β levels in ISF and cerebrospinal fluid (CSF) were also detected after chronic citalopram administration (Cirrito et al., 2011). Beneficial effects on plaque load and A β -levels following citalopram treatment could also be confirmed by a couple of other studies (Sheline et al., 2014; Wei et al., 2017; Zhang et al., 2018). Sheline and colleagues demonstrated a reduced formation of plaques as well as decreased growth of preexisting plaques in APP/PS1 mice after 28 days of citalopram treatment (Sheline et al., 2014). Similar to the results obtained



after citalopram treatment, several studies demonstrated that fluoxetine treatment decreases amyloid plaques and soluble A β levels in the brain in different AD mouse models (Cirrito et al., 2011; Wang et al., 2014, 2016; Qiao et al., 2016; Jin et al., 2017; Ma et al., 2017; Sun et al., 2017; Zhou et al., 2019). Furthermore, there is limited evidence that paroxetine has also beneficial effects on the AD pathology as Nelson et al. (2007) showed reduced accumulation of tau-protein and cortical A β 1-40 levels after chronic paroxetine treatment in 3xTg mice and Olesen et al. (2017) detected a reduction of plaque load in the hippocampus of APP/PS1-mice (Nelson et al., 2007; Olesen et al., 2016, 2017). In addition, escitalopram was able to reduce tau-hyperphosphorylation in cultures of A β 1-42 treated hippocampal neurons of fetal brains obtained from rats (Wang et al., 2016) and in a recent study, Cirrito and colleagues showed a significantly reduced plaque load in APP/PS1 mice after 28 days of escitalopram treatment (Cirrito et al., 2020).

While these preclinical results hint to a beneficial effect of SSRIs on AD-pathologies, studies on their impact in humans are very limited. Bartels et al. (2018) showed that long-term use of SSRIs delayed the conversion from MCI to AD in patients with a previous depression for 3 years (Bartels et al., 2018). Similarly, Burke et al. (2018) showed that the risk of cognitively normal patients with a prior history of depression developing AD was neutralized by SSRI treatment compared to untreated patients (Burke et al., 2018). A retrospective study of Down syndrome patients with a previous history of depression showed that SSRI use for more than 90 days significantly delayed the onset of dementia (Tsiouris et al., 2014).

Sheline et al. (2014) demonstrated a significant reduction of A β production and CSF A β concentration after a single high dose application of citalopram in healthy volunteers (Sheline et al., 2014). However, it should be noted that the study was performed on young participants, presumably before plaque formation while our cohort was much older. However, in a recent study, the same group showed that short-term longitudinal treatment of escitalopram decreases CSF A β 42 levels in cognitively normal older adults (Sheline et al., 2020).

In the current study, we could not detect any differences of amyloid burden or cognition between patients with SSRI treatment compared to untreated patients regardless of a previous depression.


So far, only two studies have evaluated the effects of SSRI-treatment on amyloid burden using amyloid-PET showing



controversial findings. Brendel et al. (2018) showed no significant reduction of amyloid deposition in ¹⁸F-Florbetapir-PET between SSRI-treated and untreated MCI and AD patients with a prior history of depression (Brendel et al., 2018). Results are in line with our findings, as we did not detect significant differences in amyloid burden between SSRI-treated and untreated patients in a comparable patient cohort. In addition, we did not find any significant differences in MCI and AD groups without prior depression nor in cognitively normal patient groups with or without depression. Consistent with our findings, in a previous

study by Bartels et al. (2018) CSF Aβ levels were unaffected by SSRI treatment in MCI patients with a history of depression.

In contrast to those findings and to our findings, Cirrito et al. (2011) showed a significant lower cortical binding of the ¹¹C-labeled amyloid tracer Pittsburgh Compound-B (¹¹C-PiB) in cognitively normal patients that have received fluoxetine, citalopram or escitalopram treatment (mean duration 35 months) compared to patients with no history of SSRI use. The Thioflavin-S derivative ¹¹C-PiB was the first amyloid tracer applied in human PET studies with high sensitivity and specificity for



the detection of amyloid plaques. Limitations of the ^{11}C -labeled tracer due to its availability and short half-life were overcome with the development of ^{18}F -labeled amyloid tracers, including ^{18}F -Florbetapir. ^{18}F -labeled tracers show comparable binding properties and clinical performance to ^{11}C -PiB and therefore, differences cannot be mainly explained by the use of different amyloid tracers (Johnson et al., 2013; Barthel and Sabri, 2017; Villemagne et al., 2017). In the study by Cirrito et al. (2011), all participants that took SSRIs had a history of depression while our cohort also included patients that used SSRI due to other reasons than depression which might be the main reason of discrepant results. However, after separating participants with and without history of depression, there were no differences between SSRI+ and SSRI– subjects in our cohort.

We could show that the duration of SSRI treatment correlated inversely with amyloid burden in both, cognitively normal and cognitively impaired patients, consistent with the findings of Cirrito et al. (2011). The time of onset and duration of exposure to SSRIs seems to be a crucial factor as SSRI-treatment might have preventive properties reducing $\text{A}\beta$ accumulation. Hypothesizing that SSRIs influence $\text{A}\beta$ metabolism promoting the non-amylogenic APP processing pathway leading to a lower amount of soluble neurotoxic $\text{A}\beta$ forms and lower amyloid plaque burden, only long-term treatment might contain AD pathologies in a protective way. Furthermore, the development of AD pathologies takes many years and only short-term treatment might not be sufficient in order to reduce $\text{A}\beta$ accumulation. This theory is supported by the results of a large Danish population-based study in which patients who received only one prescription of SSRI antidepressants had an increased rate of dementia compared to subjects unexposed to antidepressants, while continued long-term antidepressant

treatment (six to nine prescriptions) was associated with a reduced rate of dementia (Kessing et al., 2009). Additional factors that might be modulated by long term SSRI-treatment include depressive symptoms, stress and neuroinflammation with a possible beneficial effect on cognition.

Preclinical studies indicate that the decrease of $\text{A}\beta$ levels after SSRI treatment is dose-dependent (Cirrito et al., 2011; Sheline et al., 2014). A dose-dependent decrease of $\text{A}\beta$ levels was first demonstrated by Cirrito et al. (2011) as they showed a reduction of ISF $\text{A}\beta$ -levels by 12–16% using 5 mg/kg of citalopram while 10 mg/kg reduced $\text{A}\beta$ -levels by 24%. However, in our cohort, we could not detect a dose-effect on the cerebral amyloid load in ^{18}F -Florbetapir-PET. One possible explanation of these divergences might be an inter-individual variation in the efficacy of SSRI-treatment. Thus, the dose of SSRI treatment may vary greatly between individual patients and bias a grouped analysis.

It has previously been shown that the ApoE $\epsilon 4$ allele influences the response to different antidepressants in geriatric depression. Murphy et al. (2003) showed that patients carrying the $\epsilon 4$ allele showed a rapid onset of the noradrenergic and specific serotonergic antidepressant mirtazapine action, whereas paroxetine-treated patients with the $\epsilon 4$ allele were slow to respond (Murphy et al., 2003). However, ApoE genotype had no effect on the treatment outcome with SSRIs in the current study. Although, ApoE4 carriers showed significantly higher SUVR compared to non-carriers in cognitively impaired patients as well as in cognitively normal subjects.

Limitations of this study include the retrospective setting. Especially inconsistencies in SSRI-treatment protocols (doses, type of SSRI, duration of treatment) have to be considered as an important limitation. Furthermore, more detailed information on the number and extent of depressive episodes were not available in the dataset. In addition, inconsistencies between the groups regarding the history of depression might have led to a possible bias. Another limitation is the limited number of patients in some of the studied sub-groups.

Overall, the effect of SSRI-treatment on AD-pathologies in humans remains controversial. While preclinical studies and first clinical data showed promising results, a beneficial effect neither on amyloid load nor cognition could be confirmed so far.

CONCLUSION

The effect of SSRI-treatment on AD-pathologies remains unclear. However, preclinical data and the negative correlation between the time of SSRI treatment and amyloid load continue to suggest a possible beneficial effect of SSRI-treatment on the pathogenesis of AD. A controlled randomized prospective study on the effect of SSRIs on AD-pathologies is necessary in order to overcome limitations of previous studies evaluating treatment effects in a dose- and time-controlled manner. Furthermore, more information of the mechanism of action of SSRI-treatment are needed, including secondary and tertiary preventive properties, e.g., possible modulation of depressive symptoms and stress of SSRIs in the development of AD.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

All ADNI participants provided informed written consent, which was approved by each site's Institutional Review Board. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

CB and YB designed the project, analyzed the data, and wrote the manuscript. Both authors contributed to revising the manuscript and approved the final version.

ACKNOWLEDGMENTS

We acknowledge support by the Open Access Publication Funds of the Georg-August-University, Göttingen University. Data collection and sharing for this project was funded by the Alzheimer's Disease Neuroimaging Initiative (ADNI; National Institutes of Health Grant U01 AG024904) and DOD

ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd. and its affiliated company Genentech, Inc.; Fujirebio US; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development, LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

REFERENCES

- Bartels, C., Belz, M., Vogelgsang, J., Hessmann, P., Bohlken, J., Wiltfang, J., et al. (2020). To Be Continued? Long-Term Treatment Effects of Antidepressant Drug Classes and Individual Antidepressants on the Risk of Developing Dementia: a German Case-Control Study. *J. Clin. Psychiatry* 81:19m13205. doi: 10.4088/JCP.19m13205
- Bartels, C., Wagner, M., Wolfgruber, S., Ehrenreich, H., Schneider, A., and Alzheimer's Disease Neuroimaging, I. (2018). Impact of SSRI Therapy on Risk of Conversion From Mild Cognitive Impairment to Alzheimer's Dementia in Individuals With Previous Depression. *Am. J. Psychiatry* 175, 232–241. doi: 10.1176/appi.ajp.2017.17040404
- Barthel, H., and Sabri, O. (2017). Clinical Use and Utility of Amyloid Imaging. *J. Nucl. Med.* 58, 1711–1717.
- Brendel, M., Jaworska, A., Overhoff, F., Blume, T., Probst, F., Gildehaus, F. J., et al. (2018). Efficacy of chronic BACE1 inhibition in PS2APP mice depends on the regional Abeta deposition rate and plaque burden at treatment initiation. *Theranostics* 8, 4957–4968. doi: 10.7150/thno.27868
- Burke, S. L., Maramaldi, P., Cadet, T., and Kukull, W. (2018). Decreasing hazards of Alzheimer's disease with the use of antidepressants: mitigating the risk of depression and apolipoprotein E. *Int. J. Geriatr. Psychiatry* 33, 200–211. doi: 10.1002/gps.4709
- Chen, G. F., Xu, T. H., Yan, Y., Zhou, Y. R., Jiang, Y., Melcher, K., et al. (2017). Amyloid beta: structure, biology and structure-based therapeutic development. *Acta Pharmacol. Sin.* 38, 1205–1235. doi: 10.1038/aps.2017.28
- Chung, J. K., Plitman, E., Nakajima, S., Chow, T. W., Chakravarty, M. M., Caravaggio, F., et al. (2015). Lifetime History of Depression Predicts Increased Amyloid-beta Accumulation in Patients with Mild Cognitive Impairment. *J. Alzheimers Dis.* 45, 907–919.
- Cirrito, J. R., Disabato, B. M., Restivo, J. L., Verges, D. K., Goebel, W. D., Sathyan, A., et al. (2011). Serotonin signaling is associated with lower amyloid-beta levels and plaques in transgenic mice and humans. *Proc. Natl. Acad. Sci. U. S. A.* 108, 14968–14973. doi: 10.1073/pnas.1107411108
- Cirrito, J. R., Wallace, C. E., Yan, P., Davis, T. A., Gardiner, W. D., and Doherty, B. M. (2020). Effect of escitalopram on Abeta levels and plaque load in an Alzheimer mouse model. *Neurology* 95, e2666–e2674. doi: 10.1212/WNL.0000000000010733
- Cummings, J., Aisen, P. S., Dubois, B., Frolich, L., Jack, C. R. Jr., and Jones, R. W. (2016). Drug development in Alzheimer's disease: the path to 2025. *Alzheimers Res. Ther.* 8:39. doi: 10.1186/s13195-016-0207-9
- Guo, T., Zhang, D., Zeng, Y., Huang, T. Y., Xu, H., and Zhao, Y. (2020). Molecular and cellular mechanisms underlying the pathogenesis of Alzheimer's disease. *Mol. Neurodegener.* 15:40. doi: 10.1186/s13024-020-00391-7
- Herbert, J., and Lucassen, P. J. (2016). Depression as a risk factor for Alzheimer's disease: genes, steroids, cytokines and neurogenesis - What do we need to know? *Front. Neuroendocrinol.* 41, 153–171. doi: 10.1016/j.yfrne.2015.12.001
- Jin, L., Gao, L. F., Sun, D. S., Wu, H., Wang, Q., Ke, D., et al. (2017). Long-term Ameliorative Effects of the Antidepressant Fluoxetine Exposure on Cognitive Deficits in 3 x TgAD Mice. *Mol. Neurobiol.* 54, 4160–4171. doi: 10.1007/s12035-016-9952-9
- Johnson, K. A., Sperling, R. A., Gidicsin, C. M., Carmasin, J. S., Maye, J. E., Coleman, R. E., et al. (2013). Florbetapir (F18-AV-45) PET to assess amyloid burden in Alzheimer's disease dementia, mild cognitive impairment, and normal aging. *Alzheimers Dement.* 9, S72–S83. doi: 10.1016/j.jalz.2012.10.007
- Kessing, L. V., Sondergard, L., Forman, J. L., and Andersen, P. K. (2009). Antidepressants and dementia. *J. Affect. Disord.* 117, 24–29.
- Ma, J., Gao, Y., Jiang, L., Chao, F. L., Huang, W., Zhou, C. N., et al. (2017). Fluoxetine attenuates the impairment of spatial learning ability and prevents neuron loss in middle-aged APP^{sw}/PSEN1^{de9} double transgenic Alzheimer's disease mice. *Oncotarget* 8, 27676–27692. doi: 10.18632/oncotarget.15398
- Mahgoub, N., and Alexopoulos, G. S. (2016). Amyloid Hypothesis: is There a Role for Anti-amyloid Treatment in Late-Life Depression? *Am. J. Geriatr. Psychiatry* 24, 239–247. doi: 10.1016/j.jagp.2015.12.003

- Murphy, G. M., Kremer, C., Rodrigues, H., Schatzberg, A. F., and Mitrazapine Versus Paroxetine Study, G. (2003). The apolipoprotein E epsilon4 allele and antidepressant efficacy in cognitively intact elderly depressed patients. *Biol. Psychiatry* 54, 665–673. doi: 10.1016/s0006-3223(03)00174-4
- Nelson, R. L., Guo, Z., Halagappa, V. M., Pearson, M., Gray, A. J., and Matsuoka, Y. (2007). Prophylactic treatment with paroxetine ameliorates behavioral deficits and retards the development of amyloid and tau pathologies in 3xTgAD mice. *Exp. Neurol.* 205, 166–176. doi: 10.1016/j.expneurol.2007.01.037
- Nitsch, R. M., Deng, M., Growdon, J. H., and Wurtman, R. J. (1996). Serotonin 5-HT_{2a} and 5-HT_{2c} receptors stimulate amyloid precursor protein ectodomain secretion. *J. Biol. Chem.* 271, 4188–4194. doi: 10.1074/jbc.271.8.4188
- Olesen, L. O., Bouzinova, E. V., Severino, M., Sivasaravanaparan, M., Hasselstrom, J. B., Finsen, B., et al. (2016). Behavioural Phenotyping of APPswe/PS1deltaE9 Mice: age-Related Changes and Effect of Long-Term Paroxetine Treatment. *PLoS One* 11:e0165144. doi: 10.1371/journal.pone.0165144
- Olesen, L. O., Sivasaravanaparan, M., Severino, M., Babcock, A. A., Bouzinova, E. V., and West, M. J. (2017). Neuron and neuroblast numbers and cytogenesis in the dentate gyrus of aged APPswe/PS1dE9 transgenic mice: effect of long-term treatment with paroxetine. *Neurobiol. Dis.* 104, 50–60. doi: 10.1016/j.nbd.2017.04.021
- Pakaski, M., Bjelik, A., Huguycz, M., Kasa, P., Janka, Z., and Kalman, J. (2005). Imipramine and citalopram facilitate amyloid precursor protein secretion in vitro. *Neurochem. Int.* 47, 190–195. doi: 10.1016/j.neuint.2005.03.004
- Prado, C. E., Watt, S., and Crowe, S. F. (2018). A meta-analysis of the effects of antidepressants on cognitive functioning in depressed and non-depressed samples. *Neuropsychol. Rev.* 28, 32–72. doi: 10.1007/s11065-018-9369-5
- Qiao, J., Wang, J., Wang, H., Zhang, Y., Zhu, S., and Adilijiang, A. (2016). Regulation of astrocyte pathology by fluoxetine prevents the deterioration of Alzheimer phenotypes in an APP/PS1 mouse model. *Glia* 64, 240–254. doi: 10.1002/glia.22926
- Ramos-Rodriguez, J. J., Molina-Gil, S., Rey-Brea, R., Berrocoso, E., and Garcia-Alloza, M. (2013). Specific serotonergic denervation affects tau pathology and cognition without altering senile plaques deposition in APP/PS1 mice. *PLoS One* 8:e79947. doi: 10.1371/journal.pone.0079947
- Robert, S. J., Zugaza, J. L., Fischmeister, R., Gardier, A. M., and Lezoualc'h, F. (2001). The human serotonin 5-HT₄ receptor regulates secretion of non-amyloidogenic precursor protein. *J. Biol. Chem.* 276, 44881–44888. doi: 10.1074/jbc.M109008200
- Sheline, Y. I., Snider, B. J., Beer, J. C., Seok, D., Fagan, A. M., and Suckow, R. F. (2020). Effect of escitalopram dose and treatment duration on CSF Abeta levels in healthy older adults: a controlled clinical trial. *Neurology* 95, e2658–e2665. doi: 10.1212/WNL.00000000000010725
- Sheline, Y. I., West, T., Yarasheski, K., Swarm, R., Jasielc, M. S., and Fisher, J. R. (2014). An antidepressant decreases CSF Abeta production in healthy individuals and in transgenic AD mice. *Sci. Transl. Med.* 6:236re234.
- Shen, F., Smith, J. A., Chang, R., Bourdet, D. L., Tsuruda, P. R., Obedencio, G. P., et al. (2011). 5-HT₄ receptor agonist mediated enhancement of cognitive function in vivo and amyloid precursor protein processing in vitro: a pharmacodynamic and pharmacokinetic assessment. *Neuropharmacology* 61, 69–79. doi: 10.1016/j.neuropharm.2011.02.026
- Sun, D. S., Gao, L. F., Jin, L., Wu, H., Wang, Q., Zhou, Y., et al. (2017). Fluoxetine administration during adolescence attenuates cognitive and synaptic deficits in adult 3xTgAD mice. *Neuropharmacology* 126, 200–212. doi: 10.1016/j.neuropharm.2017.08.037
- Tsiouris, J. A., Patti, P. J., and Flory, M. J. (2014). Effects of antidepressants on longevity and dementia onset among adults with Down syndrome: a retrospective study. *J. Clin. Psychiatry* 75, 731–737. doi: 10.4088/JCP.13m08562
- Tucker, S., Ahl, M., Bush, A., Westaway, D., Huang, X., and Rogers, J. T. (2005). Pilot study of the reducing effect on amyloidosis in vivo by three FDA pre-approved drugs via the Alzheimer's APP 5' untranslated region. *Curr. Alzheimer. Res.* 2, 249–254. doi: 10.2174/1567205053585855
- Villemagne, V. L., Dore, V., Bourgeat, P., Burnham, S. C., Laws, S., Salvado, O., et al. (2017). Abeta-amyloid and Tau Imaging in Dementia. *Semin. Nucl. Med.* 47, 75–88. doi: 10.1053/j.semnuclmed.2016.09.006
- Wang, J., Zhang, Y., Xu, H., Zhu, S., Wang, H., He, J., et al. (2014). Fluoxetine improves behavioral performance by suppressing the production of soluble beta-amyloid in APP/PS1 mice. *Curr. Alzheimer. Res.* 11, 672–680. doi: 10.2174/1567205011666140812114715
- Wang, Y. J., Ren, Q. G., Gong, W. G., Wu, D., Tang, X., Li, X. L., et al. (2016). Escitalopram attenuates beta-amyloid-induced tau hyperphosphorylation in primary hippocampal neurons through the 5-HT_{1A} receptor mediated Akt/GSK-3beta pathway. *Oncotarget* 7, 13328–13339. doi: 10.18632/oncotarget.7798
- Wei, Z., Junhong, G., Xiaoyuan, N., Jie, W., Zhaojun, W., Meina, W., et al. (2017). Citalopram Ameliorates Impairments in Spatial Memory and Synaptic Plasticity in Female 3xTgAD Mice. *Biomed. Res. Int.* 2017:1238687. doi: 10.1155/2017/1238687
- Zhang, Q., Yang, C., Liu, T., Liu, L., Li, F., Cai, Y., et al. (2018). Citalopram restores short-term memory deficit and non-cognitive behaviors in APP/PS1 mice while halting the advance of Alzheimer's disease-like pathology. *Neuropharmacology* 131, 475–486. doi: 10.1016/j.neuropharm.2017.12.021
- Zhou, C. N., Chao, F. L., Zhang, Y., Jiang, L., Zhang, L., Fan, J. H., et al. (2019). Fluoxetine delays the cognitive function decline and synaptic changes in a transgenic mouse model of early Alzheimer's disease. *J. Comp. Neurol.* 527, 1378–1387. doi: 10.1002/cne.24616

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Citation: Bouter Y and Bouter C (2022) Selective Serotonin Reuptake Inhibitor-Treatment Does Not Show Beneficial Effects on Cognition or Amyloid Burden in Cognitively Impaired and Cognitively Normal Subjects. *Front. Aging Neurosci.* 14:883256. doi: 10.3389/fnagi.2022.883256

Copyright © 2022 Bouter and Bouter. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.