

OPEN ACCESS

EDITED AND REVIEWED BY Allison B. Reiss, New York University, United States

*CORRESPONDENCE
Natalya V. Ponomareva

☑ ponomareva@neurology.ru

RECEIVED 27 August 2024 ACCEPTED 30 August 2024 PUBLISHED 19 September 2024

CITATION

Ponomareva NV, Klyushnikov SA, Abramycheva N, Konovalov RN, Krotenkova M, Kolesnikova E, Malina D, Urazgildeeva G, Kanavets E, Mitrofanov A, Fokin V, Rogaev E and Illarioshkin SN (2024) Corrigendum: Neurophysiological hallmarks of Huntington's disease progression: an EEG and fMRI connectivity study. Front. Aging Neurosci. 16:1487201. doi: 10.3389/fnagi.2024.1487201

COPYRIGHT

© 2024 Ponomareva, Klyushnikov, Abramycheva, Konovalov, Krotenkova, Kolesnikova, Malina, Urazgildeeva, Kanavets, Mitrofanov, Fokin, Rogaev and Illarioshkin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: Neurophysiological hallmarks of Huntington's disease progression: an EEG and fMRI connectivity study

Natalya V. Ponomareva^{1,2*}, Sergey A. Klyushnikov¹, Natalia Abramycheva¹, Rodion N. Konovalov¹, Marina Krotenkova¹, Ekaterina Kolesnikova¹, Daria Malina¹, Gusel Urazgildeeva¹, Elena Kanavets¹, Andrey Mitrofanov³, Vitaly Fokin¹, Evgeny Rogaev^{2,4} and Sergey N. Illarioshkin¹

¹Research Center of Neurology, Moscow, Russia, ²Center for Genetics and Life Science, Sirius University of Science and Technology, Sochi, Russia, ³Research Center of Mental Health, Moscow, Russia, ⁴Department of Psychiatry, Umass Chan Medical School, Shrewsbury, MA, United States

KEYWORDS

EEG, functional MRI, Huntington's disease, preclinical stage, HTT gene, CAG repeats, cognitive functions

A Corrigendum on

Neurophysiological hallmarks of Huntington's disease progression: an EEG and fMRI connectivity study

by Ponomareva, N. V., Klyushnikov, S. A., Abramycheva, N., Konovalov, R. N., Krotenkova, M., Kolesnikova, E., Malina, D., Urazgildeeva, G., Kanavets, E., Mitrofanov, A., Fokin, V., Rogaev, E., and Illarioshkin, S. N. (2023). *Front. Aging Neurosci.* 15:1270226. doi: 10.3389/fnaqi.2023.1270226

In the published article, there was an error in the caption for Figure 3 and Figure 4, where these captions were erroneously switched. The corrected captions for Figure 3 and Figure 4 appear below:

Figure 3. The patterns of functional links corresponding to the differences in fMRI resting-state functional connectivity (rsFC) among the individuals with HTT mutations, including those with PreHD and EMHD, and the control group of healthy individuals (p-FDR < 0.05). Abbreviations are the same as in Table 3.

Figure 4. Patterns of circuits in which fMRI resting-state functional connectivity (rsFC) is associated with EEG alpha relative power among the Huntington's disease (HD) mutation carriers and healthy individuals. Results of regression analysis. Abbreviations are the same as in Table 4.

The authors apologize for these errors and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.