



Sensitive and Rapid Phenotyping of Microbes With Soluble Methane Monooxygenase Using a Droplet-Based Assay

Hyewon Lee¹, Ji In Baek^{1,2}, Su Jin Kim¹, Kil Koang Kwon¹, Eugene Rha¹, Soo-Jin Yeom^{1,3}, Haseong Kim^{1,4}, Dae-Hee Lee^{1,4}, Dong-Myung Kim^{1,2} and Seung-Goo Lee^{1,4*}

¹ Synthetic Biology and Bioengineering Research Center, Korea Research Institute of Bioscience and Biotechnology, Daejeon, South Korea, ² Department of Chemical Engineering and Applied Chemistry, Chungnam National University, Daejeon, South Korea, ³ School of Biological Sciences and Technology, Chonnam National University, Gwangju, South Korea, ⁴ Department of Biosystems and Bioengineering, KRIBB School of Biotechnology, University of Science and Technology, Daejeon, South Korea

OPEN ACCESS

Edited by:

Shuguang Xie,
Peking University, China

Reviewed by:

Jianhong Xu,
Tsinghua University, China
Eugene Zhang,
Huazhong University of Science
and Technology, China

*Correspondence:

Seung-Goo Lee
sglee@kribb.re.kr

Specialty section:

This article was submitted to
Bioprocess Engineering,
a section of the journal
Frontiers in Bioengineering and
Biotechnology

Received: 26 November 2019

Accepted: 31 March 2020

Published: 24 April 2020

Citation:

Lee H, Baek JI, Kim SJ, Kwon KK, Rha E, Yeom S-J, Kim H, Lee D-H, Kim D-M and Lee S-G (2020) Sensitive and Rapid Phenotyping of Microbes With Soluble Methane Monooxygenase Using a Droplet-Based Assay. *Front. Bioeng. Biotechnol.* 8:358. doi: 10.3389/fbioe.2020.00358

Methanotrophs with soluble methane monooxygenase (sMMO) show high potential for various ecological and biotechnological applications. Here, we developed a high throughput method to identify sMMO-producing microbes by integrating droplet microfluidics and a genetic circuit-based biosensor system. sMMO-producers and sensor cells were encapsulated in monodispersed droplets with benzene as the substrate and incubated for 5 h. The sensor cells were analyzed as the reporter for phenol-sensitive transcription activation of fluorescence. Various combinations of methanotrophs and biosensor cells were investigated to optimize the performance of our droplet-integrated transcriptional factor biosensor system. As a result, the conditions to ensure sMMO activity to convert the starting material, benzene, into phenol, were determined. The biosensor signals were sensitive and quantitative under optimal conditions, showing that phenol is metabolically stable within both cell species and accumulates in picoliter-sized droplets, and the biosensor cells are healthy enough to respond quantitatively to the phenol produced. These results show that our system would be useful for rapid evaluation of phenotypes of methanotrophs showing sMMO activity, while minimizing the necessity of time-consuming cultivation and enzyme preparation, which are required for conventional analysis of sMMO activity.

Keywords: synthetic biology, transcriptional factor-based biosensors, microfluidics, large-scale phenotyping, cell-cell communication

INTRODUCTION

Methane-oxidizing bacteria, called methanotrophs, have been extensively investigated in biotechnological applications such as methane mitigation, environmental remediation, and biochemical production (Semrau, 2011; Shindell et al., 2012; Strong et al., 2015). Methanotrophs are the only catalysts that can convert methane in a single step under mild conditions via methane monooxygenase (MMO) (Conrado and Gonzalez, 2014). MMOs are classified into two categories: membrane-bound particulate MMO (pMMO) and cytoplasmic soluble MMO (sMMO) (Sirajuddin and Rosenzweig, 2015). While pMMO is naturally predominant and has a relatively high affinity for

methane, sMMO has a higher specific activity and turnover frequency (Lawton and Rosenzweig, 2016; Ross and Rosenzweig, 2017). In addition, sMMO can use various carbon substrates including aromatics (Grosse et al., 1999). Despite its high applicability, sMMO engineering is difficult because of its unsuccessful expression in traditional laboratory strains such as *Escherichia coli* (West et al., 1992). Typically, to improve performance, methanotrophs are genetically engineered using synthetic biology approaches (Lloyd et al., 1999; Borodina et al., 2007; Crombie and Murrell, 2011; Smith and Murrell, 2011; Puri et al., 2015; Tapscott et al., 2019). In addition, a novel methanotroph strain with the desired phenotype of interest can be isolated from environmental samples.

For biocatalyst development in synthetic biology, phenotyping microbial cells with high enzyme activity often remains as the bottleneck. Although conventional methods including high-performance liquid chromatography (HPLC), gas chromatography, and mass spectrometry have been used to detect the sMMO activity (Shah et al., 1995; Grosse et al., 1999), these approaches show low sensitivity, and are time-consuming and labor-intensive (Lim et al., 2018). Phenotyping methods using naphthalene (Graham et al., 1992) or coumarin (Miller et al., 2002) shorten the detection time by facilitating parallel processing of multiple samples ($\sim 10^3$); however, their moderate throughput renders them impractical for analyzing large-scale ($\sim 10^6$) libraries. Therefore, highly sensitive and rapid systems are needed for large-scale quantitative phenotyping of methanotrophs with sMMO activity.

Recent advances in synthetic biology have enabled the establishment of genetically engineered biosensors for high-throughput screening of biocatalysts by converting enzyme activity into a fluorescence signal upon transcriptional activation (van der Helm et al., 2018). Few biosensors have been developed for engineering methanotroph-derived enzymes (Rohlfhill et al., 2017; Selvamani et al., 2017; Lee et al., 2019). However, the target ligands were cellular metabolites such as methanol and formaldehyde, which are rapidly assimilated by methanotrophs with limited extracellular transport. Thus, applying these biosensors for phenotyping native methanotrophs is inappropriate.

This limitation can be overcome by the use of phenol-mediated genetically encoded biosensor named GESS, which can quantitatively measure sMMO activity with benzene substrate. We previously developed a GESS biosensor that uses a dimethylphenol regulator activated by phenolic compounds (Choi et al., 2014; Kim et al., 2016a, 2018; Yeom et al., 2018). GESS is not limited to detecting intracellular enzymes and has been extended to microbe phenotyping based on cell-to-cell communication (Kim et al., 2016b). In the microbe-phenotyping-GESS (MP-GESS), sensor cells surrounding the novel microbe with target enzyme activity express fluorescent reporters through extracellular transcription regulation. This versatile MP-GESS is simple and efficient; however, this culture-based assay requires long sample preparation time to obtain microbial colonies, especially for slow-growing bacteria. Also, its qualitative and labor-intensive prototyping process requires improvement.

To facilitate large-scale quantitative analysis, droplet-based microfluidics have been used for high-throughput analysis (Thorsen et al., 2001; Meyer et al., 2015; Ngara and Zhang, 2018). Microfluidic systems generate water-in-oil droplets at rates of 1000s of samples per second (Guo et al., 2012). All compartments are miniaturized bioreactors with precisely controlled reaction volumes, cell numbers, reagent concentrations, and incubation times (Joensson and Andersson Svahn, 2012). Therefore, droplet-based microfluidics has been successfully used as a screening platform with broad applications, such as for the directed evolution of enzymes, identification of novel metagenomic enzymes, and production of extracellular chemicals (Agresti et al., 2010; Hosokawa et al., 2015; Kaushik et al., 2017; Siedler et al., 2017; Giuffrida et al., 2018; de Almeida et al., 2019). It is possible to extend this technology to methanotroph phenotyping, which has not been performed before. We expect that the GESS biosensor and droplet microfluidics (GESSlet) system developed here can be used as a high-throughput screening platform for methanotroph with sMMO activity with further modifications.

Moreover, GESSlet can be extended to many applications involving cell-to-cell communication-based methanotroph phenotyping. Recently, the synthetic community of methanotrophs and non-methanotrophs has gained considerable attention as a promising platform of methane metabolism (Chistoserdova, 2018; Chistoserdova and Kalyuzhnaya, 2018). Methanotrophs can support non-methanotrophs by potentially releasing methane-derived carbons (Kalyuzhnaya et al., 2013; Tavormina et al., 2017; Yu and Chistoserdova, 2017); therefore, characterizing methanotrophs based on their communication with other microbial community members is crucial.

MATERIALS AND METHODS

Chemicals

Chemicals and cloning reagents (restriction endonucleases, T4 DNA ligase, and DNA polymerase) were purchased from Sigma-Aldrich (St. Louis, MO, United States) and New England Biolabs (Ipswich, MA, United States), respectively. Plasmid DNA isolation and DNA extraction were conducted using plasmid preparation kits (Promega, Madison, WI, United States). Oligonucleotides were commercially synthesized and sequenced by Genotech (Daejeon, South Korea).

GESS Preparation

As a genetic circuit system, the plasmid pGESSv4 was previously constructed with the *dmpR* transcriptional activator along with its own promoter, P_X , and EGFP with the P_{dmpR} promoter from *Pseudomonas putida* KCTC 1452 (Choi et al., 2014). To enhance the system sensitivity, the newly constructed pGESS_{sfGFP} plasmid has a constitutive P_{HCE} promoter instead of P_X to achieve high *dmpR* expression and the strong reporter gene *sf-gfp* along with the P_{dmpR} promoter. For optimization, three *E. coli* strains (DH5 α , C2566, and MG1655) were used as host bacteria for pGESS_{sfGFP}. For phenotyping of wild-type methanotrophs with high reliability, the GESS_{sfGFP} genetic construct was integrated into the *E. coli* DH5 α chromosome as described previously

(Kim et al., 2018; Yeom et al., 2018). The *E. coli* sensor cells were cultured in Luria-Bertani medium at 37°C and stored at –80°C in 15% glycerol until further analysis.

To use the GESS microbial sensor, the frozen cell stock was thawed on ice and plated to grow single colonies, which were inoculated in 5 mL of Luria-Bertani medium at 37°C for 16 h. After culturing the cells, the optical density of the bacterial suspension was measured at 600 nm with a spectrophotometer. The cells were harvested by centrifugation at 1000 × *g* for 20 min and then resuspended in M9 medium (47.8 mM Na₂HPO₄, 22 mM KH₂PO₄, 18.7 mM NH₄Cl, 8.6 mM NaCl, 2 mM MgSO₄, 0.1 mM CaCl₂, 0.1% acetate, and 0.01% thiamine) with 10–20 mM benzene.

Methanotroph Culture

Five methanotrophs strains were obtained: *Methylococcus capsulatus* Bath from American Type Culture Collection (ATCC 33009, Manassas, VA, United States), *Methylosinus sporium* 5 and *Methylocella silvestris* BL2 from Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ 17706 and 15510, Braunschweig, Germany), *Methylosinus trichosporium* OB3b from Korean Collection for Type Cultures (KCTC 12760, Jeollabuk-do, South Korea), and *Methylomonas* sp. DH-1 from Eun Yeol Lee (Kyung Hee University, Yongin-si, South Korea). All methanotrophs were cultured in nitrate mineral salts (NMSs) medium, pH 6.8 or pH 5.8 (*M. silvestris* BL2), as described previously (Whittenbury et al., 1970). Frozen stocks of methanotroph in 10% (v/v) dimethyl sulfoxide were thawed, washed, and inoculated into 3 mL NMS media in glass bottles. The bottles were sealed with rubber stoppers and aluminum caps. The headspace was filled with a methane-air gas mixture (30:70), and the cultures were incubated at 30 or 42°C with agitation (200 rpm). After 24 h, the methanotroph culture was transferred to 200-mL sealed serum bottles containing 30 mL of NMS medium. Under a methane-air gas mixture, the cultures were incubated at 30 or 42°C with agitation for 2 days. Cells were harvested by centrifugation at 1000 × *g* for 10 min and resuspended at the desired concentration in M9 medium.

Operation of the Microfluidic Device

For high-throughput co-encapsulation of methanotrophs and GESS, we used the μEncapsulator 1–2 Reagent Droplet Chip (50 μm, fluorophilic, Dolomite Microfluidics, Royston, United Kingdom). Monodisperse droplets with diameters of 40–50 μm were produced at the flow focusing junction, which has two aqueous inlets, two oil inlets, and an outlet channel. This chip interface interacts directly with the Sample Reservoir Chip (Dolomite Microfluidics), where the methanotroph sample and GESS/benzene solution were loaded through separate channels. Both channels were connected to a single pressure pump through PTFE tubing (Kinesis, Berlin Township, NJ, United States) with a Y-connector. The tubing sections were filled with FC-40 oil (Sigma-Aldrich), which served as the displacement fluid. By precisely controlling the pressure pump, a pressure of 300 mbar was used for aqueous sample delivery. An oil phase consisting of FC-40 with 2% Pico-Surf (Dolomite Microfluidics) was delivered through a single port of the microfluidic chip using a pressure

pump at 500 mbar, and the flow was split equally into the two channels. Pico-Surf, a fluorinated surfactant, was added to stabilize the droplets.

Fluorescence Plate Reader Analysis

To validate the GESS biosensors, GESS cells reacted with or without *M. trichosporium* cells in M9 media containing 1–20 mM benzene at 37°C were analyzed with the Victor X multilabel plate reader (PerkinElmer, Waltham, MA, United States) or Infinite 200 PRO multi-functional microplate reader (Tecan, Männedorf, Switzerland) using black-walled 96-well polystyrene plates at excitation and emission wavelengths of 485 and 535 nm, respectively. To determine the specific fluorescence, the optical density was determined by measuring the absorbance at 600 nm in a Victor X multilabel plate reader or Infinite 200 PRO multi-functional microplate reader.

HPLC Analysis

To verify sMMO activity for the conversion of benzene to phenol, *M. trichosporium* cells were treated with benzene in NMS medium, and phenol formed in the reaction mixture was analyzed by HPLC (1260 infinity, Agilent Technologies, Santa Clara, CA, United States) fitted with an Agilent Zorbax Eclipse XDB-C18 column. The mobile phase comprising a mixture of acetonitrile and water (50:50) was delivered at a flow rate of 1 mL/min at 23°C. The UV absorbance of the samples was measured at 270 nm.

Microscopic Observation

Droplets containing GESS and methanotrophs were incubated at 37°C for 5 h. Droplets were imaged using an Axiovert A1 inverted microscope (Carl Zeiss, Oberkochen, Germany) equipped with a GFP filter (excitation at 455–495 nm, emission at 505–555 nm). Images were analyzed using ImageJ software (National Institutes of Health, Bethesda, MD, United States) (Schneider et al., 2012). Further, methanotrophic bacteria in droplets were imaged and analyzed at 20 or 63x magnification in bright-field mode with the Axiovert A1 inverted microscope.

Flow Cytometry Analysis

After incubation at 37°C for 5 h, the droplets were broken by adding 200 μL of 1H, 1H, 2H, 2H-perfluoro-1-octanol. The sample tube was agitated and centrifuged for 30 s at 1000 × *g*. The supernatant was transferred to 1 mL of phosphate-buffered saline. Fluorescence intensity was measured with a FACSAria™ Flow Cytometer Sorter (BD Biosciences, Franklin Lakes, NJ, United States) with a blue laser source (488 nm). Data were acquired using BD CellQuest Pro (version 4.0.2, BD Biosciences) and analyzed using FlowJo software (Tree Star, Ashland, OR, United States). Debris were excluded via forward and side scattering measurements. The target intensity was calculated by averaging the total fluorescence of GESS cells that had been reacted with methanotrophs and benzene. The fluorescence of GESS cells without methanotroph was averaged to calculate the background signal. The net signal

was determined by subtracting the background signal from the target signal.


RESULTS AND DISCUSSION


Assay Development

To quantify sMMO activity in methanotrophs based on cell-to-cell communication, we applied our transcription factor-based microbial sensor, GESS. A previous study reported that the GESS sensor enabled identification of phenol-generating enzymes in a high-throughput manner. sMMO converts the substrate benzene into phenol, which diffuses into GESS sensor cells to induce the expression of a fluorescent reporter through extracellular transcription regulation (Figure 1). In an initial trial with *M. trichosporium* OB3b, the fluorescence signal intensity increased over time (Figure 2A). Also, liquid chromatography analysis indicated that phenol produced by *M. trichosporium* increased over time with either 1 or 10 mM benzene as the substrate. Moreover, *M. trichosporium* did not degrade phenol for 24 h (see Supplementary Figure 1). This indicates that phenol was generated by sMMO-catalyzed oxidation of benzene and not degraded by the methanotroph or GESS metabolism, unlike

methanol and formaldehyde. Indeed, *in vivo* MMO activity is typically measured by oxidizing substrates other than methane, such as propylene, to prevent further metabolism of the product. Thus, by using phenol-mediated GESS, the *in vivo* activity of sMMO in microbes can be measured. Notably, negative control strains such as *E. coli* DH5 and *Mesorhizobium loti* showed negligible signals with 1 mM benzene (Supplementary Figure 2) due to an absence of monooxygenase activity.

In this assay, for sensitive measurement of sMMO, we engineered the GESS plasmid by replacing the reporter gene with *sf-gfp* and promoter for a transcription factor under a strong constitutive promoter P_{HCE} . Furthermore, we empirically selected the host cell for the GESS biosensor for robust detection. The host cell greatly influences the assay duration, sensitivity, and specificity as a device containing genetic circuit-based sensing molecules (Wang et al., 2013). Both the *E. coli* B and K strains have been widely used to develop genetically encoded biosensors and produce recombinant proteins. As shown in Figure 2B, GESS with the B strain such as C2566 showed a lower fold-change (38-fold) because of its high background signal compared to K-12 strains such as DH5 α and MG1655. The DH5 α strain showed the highest fold-change (76-fold). Additionally, to reliably detect sMMO in methanotrophs, we integrated the





GESS genetic construct into the DH5 α chromosome to minimize cellular heterogeneity due to variations in the copy numbers of the sensing circuit gene.

Phenol-mediated GESS enables specific detection of sMMO in methanotrophs. Of the two MMOs in methanotrophs, sMMO can oxidize not only methane but also various substrates including aromatic compounds. To evaluate the specificity of GESS, we tuned sMMO and pMMO expression using copper at different concentrations, which reciprocally regulates the expression of sMMO and pMMO in methanotrophs (Murrell et al., 2000). As shown in **Supplementary Figure 3a**, SDS-PAGE shows protein bands corresponding to sMMO and pMMO. *M. trichosporium* cells produced sMMO in the absence of copper. When the copper concentration was high (10 and 20 μ M of copper), copper enhanced pMMO expression and formation of

the intracytoplasmic membrane in *M. trichosporium* as observed in previous studies (Collins et al., 1991; Nielsen et al., 1996; Semrau et al., 2010; Dassama et al., 2017) (**Supplementary Figure 3b**). When *M. trichosporium* cells were cultured at various copper concentrations, we introduced GESS sensors to the reaction tube and analyzed the fluorescence signal intensity after incubation. Notably, positive (10 μ M phenol) and negative (none – no substrate) control signals from different copper concentrations exhibited negligible differences. As shown in **Figure 2C**, our assay scheme yielded a measurable signal only in the absence of copper when 1 mM benzene was used as the substrate. However, *Methylocella* strains are known to constitutively express sMMO even when cultured with copper. The GESS biosensors were thus applied to measure sMMO activity in *Methylocella silvestris* BL2 cultured with and

without 10 μM copper. With 1 mM benzene, both showed significant signals (**Supplementary Figure 4**), demonstrating the high specificity of GESS for detecting methanotrophs with sMMO activity.


Moreover, we predicted that 1% of benzene was converted to phenol because the fluorescence intensities were similar on using 1 mM benzene without copper and 10 μM phenol (**Figure 2C**). In this context, 0.1 mM benzene without copper may be converted to 1 μM phenol, resulting in a minimally distinguishable signal. We further optimized the benzene concentration to maximize GESS biosensor sensitivity for sMMO detection. The signals from the negative controls (no *M. trichosporium* cells) with 1–20 mM benzene displayed negligible differences in intensity (**Supplementary Figure 5**). For *M. trichosporium* cells cultured without copper, the fluorescent signal increased when the benzene concentration was increased from 1 to 10 mM; however, the signal decreased drastically when 20 mM benzene was used, possibly due to the chemical toxicity (**Figure 2D**). Thus, we used 10 mM benzene for all subsequent analyses.

Integration of GESS-Based sMMO Detection With Droplet Fluidics

Using the GESS-based assay for analyzing methanotrophs with sMMO presents several practical challenges. Using agar plates, a previous study reported that GESS biosensors enabled easy and simple identification of active clones with desired enzyme activity (Kim et al., 2016b); however, colony-based detection methods showed relatively poor dynamic ranges (Leemhuis et al., 2009). In addition, this method relies on a time-consuming cell culture step that requires days to weeks for phenotyping of slow-growing bacteria. To improve the assay quality and throughput of GESS-based sMMO detection, we used a droplet-based microfluidic technique for isolating methanotrophic cells in picoliter-sized compartments with GESS biosensors (**Figure 1A**). In the present GESSlet (GESS biosensor and droplet microfluidics), two different aqueous and oil phases

were delivered into a glass microfluidic device (Dolomite) using pressure pumps. A few 1000 45- μm water-in-oil droplets were produced in 1 s with minimum variations (coefficient of variation of 0.59–3.22%, **Figure 1B**). Enzymatic turnover of benzene in a confined compartment increased the phenol product concentrations, thus improving the assay sensitivity. Additionally, we optimized the OD of GESS biosensors, which determines the number of sensor cells encapsulated. As the GESS OD decreased, the phenol-induced target signal was decreased significantly whereas the change in the background signal was negligible. Thus, as shown in **Supplementary Figure 6**, the optimal OD of GESS biosensors is 1–2. We determined that OD \sim 1.0 (\sim 50 cells encapsulated per droplet) maximizes fold change and minimizes droplet-to-droplet variability for reliable measurement. Moreover, this assay platform requires only small sample volumes ($<100 \mu\text{L}$) for 1000s of measurements, which is beneficial for methanotroph phenotyping by overcoming the challenges in culturing.

Thereafter, we evaluated the compatibility of GESS with the droplet system. GESS biosensors and *M. trichosporium* cells mixed at a 1:10 ratio were isolated in droplets with 10 mM benzene. After incubation for 1–5 h at 37°C, co-cultured cells (10% GESS and 90% *M. trichosporium*) were recovered from the droplets and their fluorescence values were measured by flow cytometry. As illustrated in **Figure 3A**, the proportion of fluorescently labeled cells gradually increased with incubation time. Particularly, we observed that nearly 10% of all cells showed fluorescence after 5 h, indicating that all GESS cells responded. Moreover, time-lapse imaging of GESS biosensors in the droplet revealed that fluorescence indeed increased within every droplet as the incubation time was increased (**Figure 3B**). The reaction time was set to 5 h to maximize sensitivity, while minimizing the assay duration. If necessary, this assay duration can be further reduced and still yield reasonable signals after 1 h of incubation. Based on these results, the droplet system was successfully integrated with the GESS biosensor for efficient measurement of sMMO with minimal sample preparation.





Detection Performance

For phenotyping analysis of microbes with desired enzyme activity, sensitivity is most relevant in the context of the total number of cells required for quantification. To investigate the sensitivity of the GESSlet system, we first calibrated the number of cells per droplet. Cell encapsulation in droplets is a random process that relies on the cell density in the suspension. We calculated the cell number in each droplet from the cell density measured in bulk solution and droplet volume measured using microscopic imaging analysis. A previous study predicted that a 40-pL droplet contains four methanotrophic cells on average, as 0.1 OD of inoculum corresponds to $\sim 9.28 \times 10^7$ cells/mL (Myung et al., 2016). We experimentally verified the number of cells encapsulated in a single droplet through microscopic observation. GESS cells showing fluorescence following a pre-reaction with phenol were encapsulated at different concentrations. At each concentration (input OD = 0.02, 0.05, and 0.1), the average number of cells in each droplet was 0.8, 1.8, and 4.1 cells/droplet, respectively (Supplementary Figure 7), which agrees with the theoretical calculation (0.9, 2.2, and 4.4 cells/droplet). In addition, a Poisson distribution of cell numbers was observed to be consistent with the random distribution of cells within the droplets.

We next examined the sensitivity of the GESSlet system by plotting a calibration curve. In each droplet containing GESS sensors with 10 mM benzene, we incubated an average of 0–436 methanotrophic cells for 5 h at 37°C. After chemically breaking the droplets, the fluorescent signal of individual cells was quantified by flow cytometry. As shown in Figure 4A, GESS revealed a methanotroph-dose-dependent response; the fluorescence signal increased with an increasing number of methanotroph cells. From the calibration curve shown in Figure 4B, the assay revealed that the methanotroph detection limit was 4 cells per droplet on average for identifying sMMO activity. Considering the variation in encapsulated cells, sMMO activity can be measured from at least 10 methanotrophic cells,

which is substantially lower than that using a conventional assay platform with a microplate reader (Supplementary Figure 8). This indicates that the GESS integrated with droplet fluidics is highly sensitive.

This sensitivity can be further enhanced using a modified GESS with higher sensitivity (Choi et al., 2014) or incorporating the amplification of target methanotrophs in a droplet. Water-in-oil emulsions provided favorable conditions for methanotroph



culturing with enhanced methane delivery without agitation, thus successfully achieving methanotroph growth (Myung et al., 2016). Moreover, our GESSlet system requires only a few rounds of generations derived from a single cell. Thus, we expect that our versatile system may be extended to single-cell analysis and high-throughput screening for methanotrophs with sMMO activity with substantial further development.

Functional Phenotyping of Methanotrophs With sMMO Activity

To verify the applicability of the GESSlet, we performed functional phenotyping for five different methanotrophic bacteria based on their sMMO activities. Each methanotroph strain loaded into the microfluidic chip was successfully encapsulated in a water-in-oil droplet with GESS sensors and 10 mM benzene. Following off-chip incubation, the droplets were broken into a continuous aqueous phase, and the fluorescence intensity was measured. As expected, the GESS biosensors showed prominent signals from the three methanotrophs (*Methylococcus capsulatus* (Bath) and *M. trichosporium*, *Methylosinus sporium* 5, and *M. silvestris* BL2) with different sMMO activities, whereas GESS with *Methylomonas* sp. DH-1 displayed minimum background fluorescence (Figure 5). In addition, *M. capsulatus* cells cultured with 10 μ M copper generated a negligible fluorescence signal from the GESS sensors because of the absence of sMMO activity. On the contrary, *M. silvestris* BL2 yielded a fluorescence signal both when cultured with and without copper because of its copper-independent expression of sMMO. Moreover, sMMO of *M. trichosporium* cells showed higher specific activity than that of *M. capsulatus*, which agrees with a previous report (Shaofeng et al., 2007). Therefore, the GESSlet successfully quantified different levels of sMMO activity from various methanotrophs, demonstrating its potential applications for methanotroph screening.

To apply this system in methanotroph screening in the future, the procedure of phenotyping and sorting of microbes of interest needs to be further improved. A gelling agent could be incorporated into our flexible assay system for imaging and sorting of GESS-methanotroph compartments by flow cytometry in a high-throughput manner (Duarte et al., 2017). Addition of gelling reagents would make the GESSlet a powerful system for screening desired methanotrophs for developing industrially applicable enzymes and engineering microbial cell factories based on cell-to-cell communication. During an initial trial of screening natural samples, we isolated *Hydrogenophaga* strains, which are reported to have a symbiotic relationship with methanotrophs. Previous studies have reported that *Hydrogenophaga* strains are co-cultured during a methanotroph isolation process. This indicates the possible applicability of our GESSlet system.

In summary, we developed a technique for cell-to-cell communication-based phenotyping of methanotrophs with sMMO activity using a genetically designed a whole-cell biosensor named GESS and a droplet microfluidic system. sMMO has attracted attention from applied microbiologists and biochemical engineers because of its broad substrate specificity.

Our GESS biosensor has many advantages for identifying methanotrophs with high sMMO activity, such as its ability to observe *in vivo* sMMO activity without metabolic inhibition and its high specificity. Additionally, this assay demonstrates high throughput and highly sensitive measurement, which is beneficial in quantitative large-scale analysis. For application to the screening of methanotrophs with sMMO activity, it is possible that gelling agents could be incorporated into the GESSlet system for efficiently and rapidly sorting compartments of GESS-methanotrophs of interest. This would be beneficial for the screening of large-scale libraries of methanotrophs, which would typically be very time-consuming (weeks or possibly months) using conventional methods. Moreover, the choice of partner microbe in our system is highly flexible. Thus, this versatile system can be easily extended to many other studies such as the characterization of the co-metabolism of methanotroph–non-methanotroph communities.

DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

AUTHOR CONTRIBUTIONS

HL designed and performed the experiments, analyzed the data, and drafted the manuscript. JB, SK, and KK performed the experiments. ER, S-JY, HK, D-HL, and D-MK conceived the research and assisted in data interpretation. S-GL conceived the initial project, assisted in research design, and analyzed the data, and revised the manuscript. All authors reviewed the manuscript.

FUNDING

This work was supported by the C1 Gas Refinery Program funded by the Ministry of Science and ICT (2018M3D3A1A01055732), National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (2018R1C1B6004086 and 2018R1A2B3004755), and Korea Research Institute of Bioscience and Biotechnology Research Initiative Program.

ACKNOWLEDGMENTS

We are grateful to Eun Yeol Lee (Kyung Hee University) for providing *Methylomonas* sp. DH-1, and Sung-Keun Rhee (Chungbuk National University) for constructive feedback and discussion.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fbioe.2020.00358/full#supplementary-material>

REFERENCES

- Agresti, J. J., Antipov, E., Abate, A. R., Ahn, K., Rowat, A. C., Baret, J. C., et al. (2010). Ultrahigh-throughput screening in drop-based microfluidics for directed evolution. *Proc. Natl. Acad. Sci. U.S.A.* 107, 4004–4009. doi: 10.1073/pnas.0910781107
- Borodina, E., Nichol, T., Dumont, M. G., Smith, T. J., and Murrell, J. C. (2007). Mutagenesis of the “leucine gate” to explore the basis of catalytic versatility in soluble methane monooxygenase. *Appl. Environ. Microbiol.* 73, 6460–6467. doi: 10.1128/aem.00823-07
- Chistoserdova, L. (2018). Applications of methylotrophs: can single carbon be harnessed for biotechnology? *Curr. Opin. Biotechnol.* 50, 189–194. doi: 10.1016/j.copbio.2018.01.012
- Chistoserdova, L., and Kalyuzhnaya, M. G. (2018). Current trends in methylotrophy. *Trends Microbiol.* 26, 703–714. doi: 10.1016/j.tim.2018.01.011
- Choi, S. L., Rha, E., Lee, S. J., Kim, H., Kwon, K., Jeong, Y. S., et al. (2014). Toward a generalized and high-throughput enzyme screening system based on artificial genetic circuits. *ACS Synth. Biol.* 3, 163–171. doi: 10.1021/sb400112u
- Collins, M. L., Buchholz, L. A., and Remsen, C. C. (1991). Effect of copper on *Methylomonas albus* BG8. *Appl. Environ. Microbiol.* 57, 1261–1264. doi: 10.1128/aem.57.4.1261-1264.1991
- Conrado, R. J., and Gonzalez, R. (2014). Chemistry. Envisioning the bioconversion of methane to liquid fuels. *Science* 343, 621–623. doi: 10.1126/science.1246929
- Crombie, A., and Murrell, J. C. (2011). Development of a system for genetic manipulation of the facultative methanotroph *Methylocella silvestris* BL2. *Methods Enzymol.* 495, 119–133. doi: 10.1016/B978-0-12-386905-0.00008-5
- Dassama, L. M., Kenney, G. E., and Rosenzweig, A. C. (2017). Methanobactins: from genome to function. *Metalomics* 9, 7–20. doi: 10.1039/c6mt00208k
- de Almeida, J. M., Alnoch, R. C., De Souza, E. M., Mitchell, D. A., and Krieger, N. (2019). Metagenomics: is it a powerful tool to obtain lipases for application in biocatalysis? *Biochim. Biophys. Acta Proteins Proteom.* 1868:140320. doi: 10.1016/j.bbapap.2019.140320
- Duarte, J. M., Barbier, I., and Schaerli, Y. (2017). Bacterial microcolonies in gel beads for high-throughput screening of libraries in synthetic biology. *ACS Synth. Biol.* 6, 1988–1995. doi: 10.1021/acssynbio.7b00111
- Giuffrida, M. C., Cigliana, G., and Spoto, G. (2018). Ultrasensitive detection of lysozyme in droplet-based microfluidic devices. *Biosens Bioelectron.* 104, 8–14. doi: 10.1016/j.bios.2017.12.042
- Graham, D. W., Korich, D. G., Leblanc, R. P., Sinclair, N. A., and Arnold, R. G. (1992). Applications of a colorimetric plate assay for soluble methane monooxygenase activity. *Appl. Environ. Microbiol.* 58, 2231–2236. doi: 10.1128/aem.58.7.2231-2236.1992
- Grosse, S., Laramée, L., Wendlandt, K. D., McDonald, I. R., Miguez, C. B., and Kleber, H. P. (1999). Purification and characterization of the soluble methane monooxygenase of the type II methanotrophic *Bacterium Methylocystis* sp. strain W1 14. *Appl. Environ. Microbiol.* 65, 3929–3935. doi: 10.1128/aem.65.9.3929-3935.1999
- Guo, M. T., Rotem, A., Heyman, J. A., and Weitz, D. A. (2012). Droplet microfluidics for high-throughput biological assays. *Lab Chip* 12, 2146–2155. doi: 10.1039/c2lc21147e
- Hosokawa, M., Hoshino, Y., Nishikawa, Y., Hirose, T., Yoon, D. H., Mori, T., et al. (2015). Droplet-based microfluidics for high-throughput screening of a metagenomic library for isolation of microbial enzymes. *Biosens Bioelectron.* 67, 379–385. doi: 10.1016/j.bios.2014.08.059
- Joensson, H. N., and Andersson Svahn, H. (2012). Droplet microfluidics—a tool for single-cell analysis. *Angew Chem. Int. Ed. Engl.* 51, 12176–12192. doi: 10.1002/anie.201200460
- Kalyuzhnaya, M. G., Yang, S., Rozova, O. N., Smalley, N. E., Clubb, J., Lamb, A., et al. (2013). Highly efficient methane biocatalysis revealed in a methanotrophic bacterium. *Nat. Commun.* 4:2785. doi: 10.1038/ncomms3785
- Kaushik, A. M., Hsieh, K., Chen, L., Shin, D. J., Liao, J. C., and Wang, T. H. (2017). Accelerating bacterial growth detection and antimicrobial susceptibility assessment in integrated picoliter droplet platform. *Biosens Bioelectron.* 97, 260–266. doi: 10.1016/j.bios.2017.06.006
- Kim, S. K., Kim, S. H., Subhadra, B., Woo, S. G., Rha, E., Kim, S. W., et al. (2018). A genetically encoded biosensor for monitoring isoprene production in engineered *Escherichia coli*. *ACS Synth. Biol.* 7, 2379–2390. doi: 10.1021/acssynbio.8b00164
- Kim, H., Kwon, K. K., Seong, W., and Lee, S. G. (2016a). Multi-enzyme screening using a high-throughput genetic enzyme screening system. *J. Vis. Exp.* e54059, 1–7. doi: 10.3791/54059
- Kim, H., Rha, E., Seong, W., Yeom, S. J., Lee, D. H., and Lee, S. G. (2016b). A cell-cell communication-based screening system for novel microbes with target enzyme activities. *ACS Synth. Biol.* 5, 1231–1238. doi: 10.1021/acssynbio.5b00287
- Lawton, T. J., and Rosenzweig, A. C. (2016). Methane-oxidizing enzymes: an upstream problem in biological gas-to-liquids conversion. *J. Am. Chem. Soc.* 138, 9327–9340. doi: 10.1021/jacs.6b04568
- Lee, J. Y., Sung, B. H., Oh, S. H., Kwon, K. K., Lee, H., Kim, H., et al. (2019). C1 compound biosensors: design, functional study, and applications. *Int. J. Mol. Sci.* 20:2253. doi: 10.3390/ijms20092253
- Leemhuis, H., Kelly, R. M., and Dijkhuizen, L. (2009). Directed evolution of enzymes: library screening strategies. *IUBMB Life* 61, 222–228. doi: 10.1002/iub.165
- Lim, H. G., Jang, S., Jang, S., Seo, S. W., and Jung, G. Y. (2018). Design and optimization of genetically encoded biosensors for high-throughput screening of chemicals. *Curr. Opin. Biotechnol.* 54, 18–25. doi: 10.1016/j.copbio.2018.01.011
- Lloyd, J. S., Finch, R., Dalton, H., and Murrell, J. C. (1999). Homologous expression of soluble methane monooxygenase genes in *Methylosinus trichosporium* OB3b. *Microbiology* 145(Pt 2), 461–470. doi: 10.1099/13500872-145-2-461
- Meyer, A., Pellaux, R., Potot, S., Becker, K., Hohmann, H. P., Panke, S., et al. (2015). Optimization of a whole-cell biocatalyst by employing genetically encoded product sensors inside nanolitre reactors. *Nat. Chem.* 7, 673–678. doi: 10.1038/nchem.2301
- Miller, A. R., Keener, W. K., Watwood, M. E., and Roberto, F. F. (2002). A rapid fluorescence-based assay for detecting soluble methane monooxygenase. *Appl. Microbiol. Biotechnol.* 58, 183–188. doi: 10.1007/s00253-001-0885-4
- Murrell, J. C., McDonald, I. R., and Gilbert, B. (2000). Regulation of expression of methane monooxygenases by copper ions. *Trends Microbiol.* 8, 221–225. doi: 10.1016/s0966-842x(00)01739-x
- Myung, J., Kim, M., Pan, M., Criddle, C. S., and Tang, S. K. (2016). Low energy emulsion-based fermentation enabling accelerated methane mass transfer and growth of poly(3-hydroxybutyrate)-accumulating methanotrophs. *Bioresour. Technol.* 207, 302–307. doi: 10.1016/j.biortech.2016.02.029
- Ngara, T. R., and Zhang, H. (2018). Recent advances in function-based metagenomic screening. *Genomics Proteomics Bioinformatics.* 6, 405–415. doi: 10.1016/j.gpb.2018.01.002
- Nielsen, A. K., Gerdes, K., Degn, H., and Murrell, J. C. (1996). Regulation of bacterial methane oxidation: transcription of the soluble methane monooxygenase operon of *Methylococcus capsulatus* (Bath) is repressed by copper ions. *Microbiology* 142(Pt 5), 1289–1296. doi: 10.1099/13500872-142-5-1289
- Puri, A. W., Owen, S., Chu, F., Chavkin, T., Beck, D. A., Kalyuzhnaya, M. G., et al. (2015). Genetic tools for the industrially promising methanotroph *Methylomicrobium buryatense*. *Appl. Environ. Microbiol.* 81, 1775–1781. doi: 10.1128/AEM.03795-14
- Rohllhill, J., Sandoval, N. R., and Papoutsakis, E. T. (2017). Sort-Seq approach to engineering a formaldehyde-inducible promoter for dynamically regulated *Escherichia coli* growth on methanol. *ACS Synth. Biol.* 6, 1584–1595. doi: 10.1021/acssynbio.7b00114
- Ross, M. O., and Rosenzweig, A. C. (2017). A tale of two methane monooxygenases. *J. Biol. Inorg. Chem.* 22, 307–319. doi: 10.1007/s00775-016-1419-y
- Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012). NIH image to imageJ: 25 years of image analysis. *Nat. Methods* 9, 671–675. doi: 10.1038/nmeth.2089
- Selvamani, V., Ganesh, I., Maruthamuthu, M. K., Eom, G. T., and Hong, S. H. (2017). Engineering chimeric two-component system into *Escherichia coli* from *Paracoccus denitrificans* to sense methanol. *Biotechnol. Bioproc. Eng.* 22, 225–230. doi: 10.4014/jmb.1611.11070
- Semrau, J. D. (2011). Bioremediation via methanotrophy: overview of recent findings and suggestions for future research. *Front. Microbiol.* 2:209. doi: 10.3389/fmicb.2011.00209
- Semrau, J. D., Dispirito, A. A., and Yoon, S. (2010). Methanotrophs and copper. *FEMS Microbiol. Rev.* 34, 496–531. doi: 10.1111/j.1574-6976.2010.00212.x

- Shah, N. N., Hanna, M. L., Jackson, K. J., and Taylor, R. T. (1995). Batch cultivation of *Methylosinus trichosporium* OB3B: IV. Production of hydrogen-driven soluble or particulate methane monooxygenase activity. *Biotechnol. Bioeng.* 45, 229–238. doi: 10.1002/bit.260450307
- Shaofeng, H., Shuben, L., Jiayin, X., Jianzhong, N., Chungu, X., Haidong, T., et al. (2007). Purification and biochemical characterization of soluble methane monooxygenase hydroxylase from *Methylosinus trichosporium* IMV 3011. *Biosci. Biotechnol. Biochem.* 71, 122–129.
- Shindell, D., Kuylenstierna, J. C., Vignati, E., Van Dingenen, R., Amann, M., Klimont, Z., et al. (2012). Simultaneously mitigating near-term climate change and improving human health and food security. *Science* 335, 183–189. doi: 10.1126/science.1210026
- Siedler, S., Khatri, N. K., Zsohar, A., Kjaerbolling, I., Vogt, M., Hammar, P., et al. (2017). Development of a bacterial biosensor for rapid screening of yeast p-coumaric acid production. *ACS Synth. Biol.* 6, 1860–1869. doi: 10.1021/acssynbio.7b00009
- Sirajuddin, S., and Rosenzweig, A. C. (2015). Enzymatic oxidation of methane. *Biochemistry* 54, 2283–2294. doi: 10.1021/acs.biochem.5b00198
- Smith, T. J., and Murrell, J. C. (2011). Mutagenesis of soluble methane monooxygenase. *Methods Enzymol.* 495, 135–147. doi: 10.1016/B978-0-12-386905-0.00009-7
- Strong, P. J., Xie, S., and Clarke, W. P. (2015). Methane as a resource: can the methanotrophs add value? *Environ. Sci. Technol.* 49, 4001–4018. doi: 10.1021/es504242n
- Tapscott, T., Guarnieri, M. T., and Henard, C. A. (2019). Development of a CRISPR/Cas9 system for *Methylococcus capsulatus* *in vivo* gene editing. *Appl. Environ. Microbiol.* 85:AEM.00340-19. doi: 10.1128/AEM.00340-19
- Tavormina, P. L., Kellermann, M. Y., Antony, C. P., Tocheva, E. I., Dalleska, N. F., Jensen, A. J., et al. (2017). Starvation and recovery in the deep-sea methanotroph *Methyloprofundus sedimenti*. *Mol. Microbiol.* 103, 242–252. doi: 10.1111/mmi.13553
- Thorsen, T., Roberts, R. W., Arnold, F. H., and Quake, S. R. (2001). Dynamic pattern formation in a vesicle-generating microfluidic device. *Phys. Rev. Lett.* 86, 4163–4166. doi: 10.1103/physrevlett.86.4163
- van der Helm, E., Genee, H. J., and Sommer, M. O. A. (2018). The evolving interface between synthetic biology and functional metagenomics. *Nat. Chem. Biol.* 14, 752–759. doi: 10.1038/s41589-018-0100-x
- Wang, Y. H., Wei, K. Y., and Smolke, C. D. (2013). Synthetic biology: advancing the design of diverse genetic systems. *Annu. Rev. Chem. Biomol. Eng.* 4, 69–102. doi: 10.1146/annurev-chembioeng-061312-103351
- West, C. A., Salmond, G. P., Dalton, H., and Murrell, J. C. (1992). Functional expression in *Escherichia coli* of proteins B and C from soluble methane monooxygenase of *Methylococcus capsulatus* (Bath). *J. Gen. Microbiol.* 138, 1301–1307. doi: 10.1099/00221287-138-7-1301
- Whittenbury, R., Phillips, K. C., and Wilkinson, J. F. (1970). Enrichment, isolation and some properties of methane-utilizing bacteria. *J. Gen. Microbiol.* 61, 205–218. doi: 10.1099/00221287-61-2-205
- Yeom, S. J., Kim, M., Kwon, K. K., Fu, Y., Rha, E., Park, S. H., et al. (2018). A synthetic microbial biosensor for high-throughput screening of lactam biocatalysts. *Nat. Commun.* 9, 5053. doi: 10.1038/s41467-018-07488-0
- Yu, Z., and Chistoserdova, L. (2017). Communal metabolism of methane and the rare Earth element switch. *J. Bacteriol.* 199:e0328-17. doi: 10.1128/JB.00328-17

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Lee, Baek, Kim, Kwon, Rha, Yeom, Kim, Lee, Kim and Lee. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.