



Engineering *Yarrowia lipolytica* to Produce Itaconic Acid From Waste Cooking Oil

Lanxin Rong¹, Lin Miao¹, Shuhui Wang¹, Yaping Wang¹, Shiqi Liu¹, Zhihui Lu¹, Baixiang Zhao¹, Cuiying Zhang¹, Dongguang Xiao¹, Krithi Pushpanathan², Adison Wong^{2*} and Aiqun Yu^{1*}

¹State Key Laboratory of Food Nutrition and Safety, Key Laboratory of Industrial Fermentation Microbiology of the Ministry of Education, Tianjin Key Laboratory of Industrial Microbiology, College of Biotechnology, Tianjin University of Science and Technology, Tianjin, China, ²Food, Chemical and Biotechnology Cluster, Singapore Institute of Technology, Dover, Singapore

OPEN ACCESS

Edited by:

Hua Ling,
National University of Singapore,
Singapore

Reviewed by:

Yuanda Song,
Shandong University of Technology,
China
Xiao-Jun Ji,
Nanjing Tech University, China

*Correspondence:

Adison Wong
adison.wong@
singaporetech.edu.sg
Aiqun Yu
yuaqun@tust.edu.cn

Specialty section:

This article was submitted to
Synthetic Biology,
a section of the journal
Frontiers in Bioengineering and
Biotechnology

Received: 03 March 2022

Accepted: 04 April 2022

Published: 25 April 2022

Citation:

Rong L, Miao L, Wang S, Wang Y,
Liu S, Lu Z, Zhao B, Zhang C, Xiao D,
Pushpanathan K, Wong A and Yu A
(2022) Engineering *Yarrowia lipolytica*
to Produce Itaconic Acid From Waste
Cooking Oil.
Front. Bioeng. Biotechnol. 10:888869.
doi: 10.3389/fbioe.2022.888869

Itaconic acid (IA) is a high-value organic acid with a plethora of industrial applications. In this study, we seek to develop a microbial cell factory that could utilize waste cooking oil (WCO) as raw material for circular and cost-effective production of the abovementioned biochemical. Specifically, we expressed cis-aconitic acid decarboxylase (CAD) gene from *Aspergillus terreus* in either the cytosol or peroxisome of *Yarrowia lipolytica* and assayed for production of IA on WCO. To further improve production yield, the 10 genes involved in the production pathway of acetyl-CoA, an intermediate metabolite necessary for the synthesis of cis-aconitic acid, were individually overexpressed and investigated for their impact on IA production. To minimize off-target flux channeling, we had also knocked out genes related to competing pathways in the peroxisome. Impressively, IA titer up to 54.55 g/L was achieved in our engineered *Y. lipolytica* in a 5 L bioreactor using WCO as the sole carbon source.

Keywords: itaconic acid, *Y. lipolytica*, waste cooking oil, peroxisome, subcellular engineering

INTRODUCTION

Carboxylic acids are important building blocks in the chemical industry. Among them, itaconic acid (IA) is favorably listed by the US Department of Energy as one of top 12 biochemical to be produced from renewable resources (Werpy and Petersen, 2004), with a forecasted market potential of \$260 million in 2025 (Sriariyanun, 2019). IA is an unsaturated dicarboxylic acid that is characteristically stable in acidic, neutral and moderately alkaline conditions. Due to its advantageous properties, IA is often used as a co-monomer in the manufacture of synthetic fibers, coatings, adhesives, thickeners and binders (Willke and Vorlop, 2001; Zhao et al., 2018), and as substitutes for petrochemical-based acrylic or methacrylic acids (Nuss and Gardner, 2013). Traditionally, to meet the growing demand for IA, industries resort to fossil resources through petrochemical refinery processes to produce IA at scale. However, these methods often suffer from low efficiency and generate large amount of waste in the process, such as spent heavy metal catalysts and organic solvents (Krull et al., 2017). Furthermore, fossil resources are finite and will eventually be depleted. For these reasons, bio-based production of IA using microbial cell factories are increasingly being pursued.

Filamentous fungi such as *Aspergillus terreus* (Kuenz et al., 2012), *Ustilago maydis* (Geiser et al., 2016) and *Ustilago cynodontis* (Hosseinpour Tehrani et al., 2019b) have been demonstrated to naturally produce IA at high titers. In one example, the fermentation of *A. terreus* at industrial scale is able to generate a titer of 160 g/L IA (Krull et al., 2017), a value that is close to the theoretical yield. In

another example, up to 220 g/L IA was achieved by fermentation of *U. maydis* (Hosseinpour Tehrani et al., 2019a). Despite having high production titers, current bioprocesses involving filamentous fungi are not without challenges. Critically, the highly branched mycelial filaments of filamentous fungi give rise to high broth viscosity during fermentation, leading to poor aeration and mixing in stirred-tank bioreactors (Kubicek et al., 2011; Porro and Branduardi, 2017). Increasing impeller speed, on the other hand, is not an option due to the shear-sensitive nature of filamentous fungi. Moreover, fermentation of most filamentous fungi requires the addition of alkali to maintain a neutral pH condition which is a cause of concern as this increases the probability of bacterial contamination during cultivation (Cui et al., 2017; Li et al., 2021). To circumvent issues associated with filamentous fungi bioprocessing, scientists have applied systems metabolic engineering principles to enable heterologous production of IA in several strains of bacteria and yeasts (Table 1).

The industrial microbe *Yarrowia lipolytica* is an unconventional oleaginous yeast that is also classified by the US Food and Drug Administration as 'generally regarded as safe' (GRAS) (Zhao et al., 2021b). *Y. lipolytica* possesses unique physiological and metabolic features compared to the most widely used chassis strains *Escherichia coli* and *Saccharomyces cerevisiae*, which enhance its merits as a microbial cell factory (Liu et al., 2015). Firstly, *Y. lipolytica* has good tolerance for external environment stresses, such as low temperatures, high salt concentrations and acidic pH (Gonçalves et al., 2014). Secondly, the oleaginous yeast is able to utilize a myriad of carbon substrates for growth, including waste cooking oil (WCO) (Zinjarde, 2014; Pang et al., 2019; Li et al., 2022). This permits the valorization of waste streams and reduces the overall cost of production. Thirdly, *Y. lipolytica* is richly endowed with multiple pathways for the generation and accumulation of

intracellular acetyl-CoA, which are important intermediaries of IA biosynthesis (Zhou et al., 2012; Ng et al., 2020). Finally, the yeast exhibits high tolerance for IA, thus allowing for accumulation of IA within (Zhao et al., 2019).

In our previous studies, we successfully engineered *Y. lipolytica* to produce limonene and bisabolene, where WCO was employed as the sole carbon source (Pang et al., 2019; Zhao et al., 2021b; Li et al., 2022). Motivated by earlier successes, we herein investigated the feasibility of producing IA from engineered *Y. lipolytica* on WCO (Figure 1). We expressed cis-aconitic acid decarboxylase (CAD) gene from *A. terreus* in either the cytosol or peroxisome of *Y. lipolytica* and assayed for production of IA in the extracellular supernatant. To further improve the final yield, the 10 genes involved in the production pathway of acetyl-CoA, an intermediate metabolite necessary for the synthesis of cis-aconitic acid, were each singly overexpressed. To minimize off-target flux channeling, we had also knocked out genes related to competing pathways in the peroxisome. Finally, IA titer up to 54.55 g/L was obtained in the engineered *Y. lipolytica* with a yield of 0.3 g/g WCO and a maximum productivity of 0.6 g/L/h without pH control in the 5 L bioreactor. At the time of writing, this is the highest titer of IA obtained with an engineered yeast cell factory.

MATERIALS AND METHODS

Strains, Plasmids, Primers, and Cultivation Media

The *E. coli* strain DH5 α was used as the host in this study for the cloning and plasmid construction. *E. coli* strains were routinely cultured at 37°C in Luria-Bertani (LB) media (1% tryptone, 0.5% yeast extract, and 1% sodium chloride contained) or on LB agar plates supplemented with 100 μ g/ml of ampicillin. *Y. lipolytica*

TABLE 1 | Representative examples of IA production in engineered microbial hosts.

Parental strain	Engineering strategy	Fermentation condition	Carbon source	Titer	References
<i>E. coli</i>	CAD \uparrow , CS \uparrow , ICD \downarrow , ICL Δ , PTA Δ , PYK Δ , SUCS Δ	Fed-batch bioreactor	Glucose and glutamic acid	32.00 g/L	Harder et al. (2016)
	CAD \uparrow , ACO \uparrow , ICD Δ	Fed-batch bioreactor	LB + Glucose	4.34 g/L	Okamoto et al. (2014)
<i>S. cerevisiae</i>	CAD \uparrow , CS \uparrow , ACO \uparrow , PTA Δ , LDH Δ	Bioreactor	LB + Glucose	0.69 g/L	Vuoristo et al. (2015)
	CAD \uparrow , ADE3 Δ , BNA2 Δ , TES1 Δ	Large-scale bioreactor	Glucose	0.17 g/L	Blazek et al. (2014)
<i>Halomonas bluephagenesis</i>	CAD \uparrow , ACO \uparrow , ICD \downarrow	Batch shake flask	Citrate	63.60 g/L	Zhang et al. (2021a)
<i>Corynebacterium glutamicum</i>	CAD \uparrow , MALE \uparrow , ICD \downarrow	Shake flask	Glucose	7.80 g/L	Otten et al. (2015)
<i>Pichia kudriavzevii</i> <i>Y. lipolytica</i>	CAD \uparrow , MTT \uparrow , ICD Δ	Fed-batch bioreactor	Glucose	1.23 g/L	Sun et al. (2020)
	CAD \uparrow , ACO \uparrow , AMPD \downarrow	Bioreactor	Glucose	4.60 g/L	Blazek et al. (2015)
	CAD \uparrow , MTT \uparrow	Fed-batch bioreactor	Glucose	22.02 g/L	Zhao et al. (2019)
	CAD-ePTS1 \uparrow , POT1 \uparrow , ICL Δ	Bioreactor	Waste cooking oil	54.55 g/L	This study

\uparrow , Gene overexpression; \downarrow , gene knockdown; Δ , gene knockout; CAD, cis-aconitic acid decarboxylase gene; CS, citrate synthase gene; ICD, isocitrate dehydrogenase gene; ICL, isocitrate lyase gene; PTA, phosphate acetyltransferase gene; PYK, pyruvate kinase gene; SUCS, succinyl-CoA synthetase gene; ACO, aconitase gene; LDH, lactate dehydrogenase gene; ADE3, cytoplasmic trifunctional C1-tetrahydrofolate (THF) synthase gene; BNA2, a putative tryptophan 2,3-dioxygenase or indoleamine 2,3-dioxygenase gene; TES1, peroxisomal acyl-CoA thioesterase gene; MALE, maltose-binding protein gene; MTT, mitochondrial tricarboxylate transporter gene; AMPD, adenosine monophosphate deaminase gene; POT1, peroxisomal thiolase gene.




FIGURE 1 | Simplified schematic of IA biosynthetic pathway in *Y. lipolytica*. Engineered *Y. lipolytica* uptakes and converts extracellular carbon sources such as glucose and waste cooking oil into IA products. Genes and metabolites of the native TCA and glyoxylate cycle pathway are identified in black, while heterologously introduced genes are shown in green and the endogenous genes used in this paper are shown in red. LIP2, lipases; POX1-6, six different acyl-CoA oxidases; MFE1, multifunctional enzyme; POT1, peroxisomal thiolase; PEX10, a proteins required for peroxisome assembly; CAT, carnitine acetyltransferases; ICL, isocitrate lyase; CAD, iso-aconitic acid decarboxylase; ACO, aconitase; MLS, malate synthase; MDH, malate dehydrogenase; CIT, citrate synthase.

Po1g *KU70* Δ was used as the base strain in this study, which has been generated from the parental strain Po1g (a commonly used host strain for protein expression). This strain was used as it is known that the rate of precise homologous recombination (HR) increased substantially for deletion of the *KU70* gene in Po1g (Yu et al., 2016). Routine cultivation of *Y. lipolytica* strains was carried out at 30°C in YPD medium (1% yeast extract, 2% peptone and 2% dextrose contained) while the yeast synthetic complete medium (YNB) (0.67% yeast nitrogen base without amino acids, 2% glucose, 1.5% bacto agar) lacking the appropriate nutrients was used for the screening of transformants. The fermentation experiment used YPO medium containing WCO (1% yeast extract, 2% peptone, 1.18% WCO and 0.2% tween-80 contained), and the initial pH of cultivation media was 5.73. Among them, the amount of WCO added is calculated based on the same C atoms as glucose in YPD medium. The strains and plasmids used in this study are listed in **Supplementary Table S1**. The PCR primers used in this study were synthesized by Genewiz (Jiangsu, China) and are listed in **Supplementary Table S2**.

Plasmid Construction

The *Y. lipolytica* expression vector pYLEX1 used in this study possesses the strong promoter hp4d, and its detailed information was provided in Li et al. (2021). Using primers CAD1-F/R and CAD2-F/R that were synthesized according to the existing sequence (GenBank ID: AB326105.1) in NCBI GenBank, two fragments of the CAD gene without introns were amplified from the *A. terreus* HAT418 genome and cloned into pYLEX1 to yield pYLEX1-CAD through adapted homologous recombination. The construction process of plasmid pYLEX1-CAD is depicted in **Supplementary Figure S1**. The sequences of the oligonucleotides used to amplify all the genes are listed in **Supplementary Table S2** in the Additional file. Subsequently, the expression cassettes of other gene candidates were cloned into pYLEX1-CAD individually (**Supplementary Figure S2**). All recombinant plasmids were constructed using the One Step Cloning Kit from Vazyme Biotech Co., Ltd. (Nanjing, China). Transformants were plated on LB-ampicillin agar plates and incubated overnight at 37°C. Single colonies were inoculated into LB-ampicillin and cultured overnight

at 37°C with shaking at 225 rpm. Plasmids were isolated, and the genes were verified by DNA sequencing.

Following that, all plasmids were linearized using the *Spe* I enzyme and then transformed into the *Y. lipolytica* Po1g *KU70Δ* competent cells using lithium acetate/single-stranded vector DNA/polyethylene glycol method. The linearized plasmids introduced were integrated at the pBR322 locus of the strain Po1g *KU70Δ*. After 2 to 3 days of culture, the positive *Y. lipolytica* transformants were selected on YNB-LEU plates and subsequently confirmed by genomic DNA PCR analysis (Yu et al., 2016). Accordingly, in this study, the engineered *Y. lipolytica* Po1g *KU70Δ* strain was used as the host for all genetic modifications with gene knockouts and chromosomal expression constructs introduced *via* engineered pYLEX1 plasmids.

Yeast Cultivation

Seed inoculum of *Y. lipolytica* were first cultured in a 20 ml tube with 5 ml YPD medium and incubated for 24 h in a shaking incubator set at 30°C and 220 rpm. Next, a 250 ml flask was filled with 50 ml YPO medium and inoculated at the seeding density of OD₆₀₀ 0.1. The inoculated finished shake flasks were grown in a shaking incubator set at 30°C and 220 rpm. Fermented yeast cultures were collected on the fourth day and analyzed by GC-MS to determine and identify the IA content.

Gene Knockout

The *ICLΔ* strain was generated by knocking out the ORF region gene of *ICL* *via* the homologous recombination (HR) mechanism, which replaced *ICL* with the hygromycin B resistance marker gene (*HPH*) amplified from pSH69-Hph using the primer pairs ICL-Hph-F/R. To this end, two targeting arms (upstream and downstream flanking sequences of *ICL*), each approximately 1,000 bp in length, were amplified using PCR from the genomic DNA of Po1g-2G and ligated to the 5' and 3' ends of the *HPH* gene, respectively. After transformation of the *ICL* disruption cassette into *Y. lipolytica* cells, a gene replacement event occurs *via* double-crossover homologous recombination within the two flanking homology arms at the targeted locus. Transformants were grown in the YPDH solid medium (30°C, under dark conditions) supplemented with hygromycin and chosen randomly. The correct *ICLΔ* strain was confirmed by PCR with ICL-Hph-knock-F and ICL-Hph-knock-R primers. The construction of the *CATΔ* strain was carried out using a similar procedure.

Visualizing Fluorescence Distribution by Laser Scanning Confocal Microscopy

To test the peroxisomal targeting ability of enhanced peroxisome targeting signal ePTS1, yeast cells expressing *hrGFPO-ePTS1* were cultured in 50 ml YPD medium for 24 h. For simultaneous visualization of hrGFPO and Nile red, precultures incubated in 50 ml YPD were stained by adding Nile red solution (1 mg/ml) in acetone to the cell suspension (0.1 v/v) and incubated for 60 min in the dark at room temperature. The stained cells were washed with normal saline and resuspended in potassium phosphate buffer (pH 7.4) before being transferred onto glass slides to visualize hrGFPO at 488 nm and Nile red at 561 nm with an Olympus FV1000 confocal laser scanning microscope.

Esterification of the Fermented Supernatant

2 ml of the fermented supernatant was added to 1.5 ml of 10% HCl-CH₃OH solution, which was esterified at 62°C for 3 h. Then, 2 ml of n-hexane was added and the resultant mixture was violently shaken for 1 min to dissolve the dimethyl itaconate. After centrifugation (6,000 rpm, 5 min), the upper organic phase was transferred into another clean bottle for detection.

GC-MS Analysis

0.6 μl of the upper organic phase from the above Section was analyzed by GC-MS using an Agilent 7890A GC with a 5975C MSD equipped with an HP-5MS column (30 m × 0.25 mm × 0.25 μm, Agilent, Santa Clara, CA, United States). The GC oven temperature was initially held at 60°C for 2 min, and then ramped up to 250°C at a rate of 10°C/min and held for 9 min. The split ratio was 10:1. Helium was used as the carrier gas, with an inlet pressure of 13.8 psi. The injector was maintained at 250°C and the ion source temperature was set to 220°C. The final data analysis was performed using the Enhanced Data Analysis software (Agilent, Santa Clara, CA, United States) to obtain the standard curve of dimethyl itaconate, and the area obtained after the sample is analyzed and detected by the instrument is brought into the formula of the standard curve to obtain the output of dimethyl itaconate. The titer of IA is obtained by converting with the esterification rate obtained in the above section.

Statistical Analysis

Differences in titers between the control strain and other strains were evaluated using SPSS 22.0 software for Windows (SPSS, Chicago, IL, United States). One-way ANOVA analyses were carried out with a confidence interval of 95% and statistical significance between the groups and the relevant control was considered if *p*-value < 0.05.

Bioreactor Fermentations


Bioreactor fermentation was batched processed using an optimal medium formulation containing 59 g/L WCO, 16 g/L yeast extract, 8 g/L peptone and 10 g/L tween-80. The strain was first seeded in 50 ml YPD medium in 250 ml shake flasks, cultured at 30°C and 220 rpm for 16 h. Following that, the bioreactor containing 3 L of YPO medium were inoculated with the seed cultures at an OD₆₀₀ of 1.

Fermentation without any pH control was carried out in a 5 L stirred fermenter (Shanghai Baoxing Bioengineering Equipment Co., Ltd., Shanghai, China) at 30°C and 1 vvm. The bioreactor pressure was maintained at 0.06 MPa. The impeller stirring speed was 400 rpm.

RESULTS AND DISCUSSION

Heterologous Expression of *A. terreus* Cis-Aconitic Acid Decarboxylase in *Y. lipolytica*

In *A. terreus*, IA is generated from the decarboxylation of the TCA intermediate cis-aconitic acid by the CAD enzyme (Bonnarme et al., 1995; Tevz et al., 2010). To test if *A. terreus*'s CAD gene can be expressed successfully in *Y. lipolytica* without codon optimization,



we first cloned the associated gene from *A. terreus* HAT418 strain into *Y. lipolytica* strain Po1g *KU70Δ*, with the gene's intron spliced out. In the gene sequencing analysis that followed, we discovered that the actual PCR-amplified gene sequence was different from the genome sequence shown in NCBI database. Our sequence data for *A. terreus* HAT418 *CAD* gene was submitted to GenBank under the accession number MT862134.1. Overexpression of the *CAD* gene in *Y. lipolytica* Po1g *KU70Δ* resulted in the creation of strain Po1g-CAD. We subjected both the engineered strain with cytosolic *CAD* and control strain without *CAD* to shake flask fermentation and assayed for IA continuously over a period of 6 days. We confirmed that IA was produced only in the engineered *Y. lipolytica* but not in its wild type. IA levels were first detected in the supernatant on day 2 and they increased gradually with time until a maximum yield of 33.12 mg/L was obtained on day 4 (Figure 2). This is contrasted with the Po1g *KU70Δ* original strain where no IA production detected, thereby confirming that the heterologous expression of the *A. terreus* *CAD* is necessary for IA production in *Y. lipolytica*.

Peroxisomal Targeting of Heterologous *Cis*-Aconitic Acid Decarboxylase Gene Improved Itaconic Acid Production

β -oxidation of long chain fatty acids in eukaryotes are known to occur mainly in the peroxisomes (Wache et al., 2001; Hanko et al., 2018). In *Y. lipolytica*, this process produces acetyl-CoA which then enters the glyoxylate cycle for synthesis of the IA precursor, *cis*-aconitic acid (Dominguez et al., 2010; Koivistoinen et al., 2013; Xu et al., 2017). Several studies have shown that subcellular localization of specific enzymes or metabolic pathways not only increase product conversion efficiency, but is also able to suppress the undesirable effects of competitive metabolic inhibition (Zhu et al., 2018; Yang



et al., 2019; Zhu et al., 2021). As such, this approach of subcellular compartmentalization is adopted in our study and complemented with the use of WCO as the substrate to enable sustainable, efficient and low-cost production of IA. To this end, IA production from the glyoxylate cycle in *Y. lipolytica* was ensured by targeting the involved heterologous enzymes to the peroxisomal matrix through the addition of enhanced peroxisomal targeting signal (ePTS1) after its gene sequence. The ePTS1 applied in this instance has been shown to be localized in *S. cerevisiae* (DeLoache et al., 2016).

Two separate dyes, Nile red and green fluorescence, were employed for staining of the yeast cells to validate the peroxisomal targeting ability of ePTS1. In an earlier study, it was shown that hrGFPO, encoding the green fluorescence protein, was most strongly expressed in Po1g *KU70Δ* (Zhao et al., 2021a). The plasmid with sequence ePTS1 added after the hrGFPO protein sequence was retransformed into yeast, resulting in strain Po1g-hrGFPO-ePTS1 (Figure 3A). Nile red fluorescence, on the other hand, was used to stain the peroxisomes of the yeast cells. To determine if ePTS1 could be successfully localized to peroxisomes in *Y. lipolytica*, Laser Scanning Confocal Microscopy (LSCM) was performed to observe the location of the two different fluorescence in yeast cells. As shown in Figure 3B, a green fluorescent protein with localization signal ePTS1, which exhibits green light under microscope irradiation, was expressed in the engineered yeast. Yeast cells after Nile red staining also show localized red fluorescence under the microscope. Combining these two images, we observed that the green and red shades overlap almost completely and produce a bright yellow light. Therefore, it can be confirmed that ePTS1 plays a role in determining the location of the peroxisome could be used as a peroxisomal targeting sequence for *Y. lipolytica*.

Subsequently, the plasmid pYLEX1-CAD-ePTS1 constructed through the ligation of ePTS1 downstream of the *CAD* gene was integrated into the *Y. lipolytica* Po1g *KU70Δ* chromosomes of the strain. The resulting engineered strain was cultured in the YPO medium and the 6 days course of IA production titers and biomass were shown in Figure 4. The titers of IA increased continuously from the beginning of cultivation up to day 4 with the highest titer having reached 1.58 g/L. Following this, the titers of IA gradually stabilized, likely owing to WCO depletion. Notably, we also compared the use of WCO and glucose in this subcellular compartmentalized approach to generate IA under the same conditions. The use of WCO had resulted in almost 100-folds increase in IA titer as compared to glucose (13.68 mg/L of IA) as the carbon source, hence implying that WCO was superior to glucose for IA production in these conditions. We also observed that the overproduction of IA has a positive effect on the cell growth. Together, our results demonstrate that the expression and localization of *CAD* in the peroxisomes of *Y. lipolytica* can lead to substantial increase in IA production.

Overexpression of Endogenous Genes Involved in the Acetyl-CoA Production Pathway of *Y. lipolytica*

To further enhance IA production in *Y. lipolytica*, we attempted to study the pathway genes involved in the conversion of oils to fatty



acids and the utilization of fatty acids to raise the flux of precursor acetyl-CoA. The β -oxidation of fatty acids is a four-reaction cycle comprising of oxidation, hydration, dehydrogenation, and thiolysis, which results in one molecule of acetyl-CoA released in the peroxisome (Braga and Belo, 2016). In *Y. lipolytica*, the first step of fatty acid β -oxidation can be catalyzed by six different acyl-CoA oxidases (POX1-6) (Beopoulos et al., 2008). The second and third steps of β -oxidation are catalyzed by a multifunctional enzyme (MFE1) (Black et al., 2000; Dulermo et al., 2013), and the final step is catalyzed by peroxisomal thiolase (POT1) (Wang et al., 2020). As such, the genes involved in the β -oxidation pathway were

overexpressed in an attempt to increase the flux towards IA. Ten genes, consisting of *LIP2* (encoding lipases, Zhang et al., 2021b), *POX1-6* (Ledesma-Amaro and Nicaud, 2016), *MFE1* (Haddouche et al., 2010), *POT1* (Smith et al., 2000), and *PEX10* (encoding a proteins required for peroxisome assembly, Zhang et al., 2021b), were overexpressed individually and investigated for their effects on IA overproduction to determine the genes critical for IA biosynthesis in the acetyl-CoA production pathway. To this end, ten strains were constructed on the basis of the strain expressing *CAD-ePTS1* gene, including the ten endogenous genes in the acetyl-CoA production pathway of *Y. lipolytica*; all genes were integrated into the chromosomes of *Y. lipolytica* Po1g *KU70Δ*. These ten engineered strains were then cultured in YPO medium for 6 days in shake flasks. The IA titers of the strains showed that the overexpression of the individual corresponding genes could improve IA production compared to the control strain expressing only the respective *CAD-ePTS1* gene. Among them, the *POT1*-overexpressed strain (hereafter named Po1g-2G), achieved the highest titers of 2.42 g/L for IA after 4 days of cultivation (Figure 5). The results indicated that overexpression of this key enzymes can effectively promote the fatty acid degradation process and release the most acetyl-CoA molecules for IA biosynthesis. This observation is consistent with several other studies in which *POT1* has already been demonstrated to be the key rate-limiting enzyme in the β -oxidation pathway (Ma et al., 2020; Zhang et al., 2021b). Therefore, the engineered strain Po1g-2G was used for subsequent engineering efforts to boost IA production.

Effects of Deletion of the Carnitine Acetyltransferases and Isocitrate Lyase Genes on Itaconic Acid Production in *Y. lipolytica*

The yield of IA can be further improved by reducing the loss of the precursor acetyl-CoA and preventing the synthesis of cis-aconitic acid from the glyoxylate cycle into downstream products such as succinic




FIGURE 5 | Effects of overexpressing genes involved in the acetyl-CoA production pathway on IA production. The genes involved in the acetyl-CoA production pathway, consisting of *LIP2*, *POX1-6*, *MFE1*, *POT1* and *PEX10*, were overexpressed individually. Titers of IA produced by the strains were quantified after 6 days of cultivation in shake flasks with YPO medium. All values presented are the mean of three biological replicates \pm standard deviation. * $p < 0.05$, significantly different from control by ANOVA.

acid. The carnitine acetyltransferases (CAT) is responsible for transporting acetyl-CoA between different organelles, which can reversibly link the acetyl units to the carrier molecule carnitine (Strijbis et al., 2008; Strijbis et al., 2010). Meanwhile, the isocitrate lyase (ICL) manages the conversion of isocitrate into succinic acid and glyoxylic acid (Koivistoinen et al., 2013). To verify if either of these enzymes assume a major role in IA production, the corresponding genes singly were deleted from Po1g-2G, resulting in the creation of strains Po1g-2G-CAT Δ and Po1g-2G-ICL Δ . After cultivating the resulting strains in shake flasks in YPO medium, it was found that higher IA production reaching 3.33 g/L was observed in ICL knockout strain as compared to CAT knockout strain with 2.8 g/L titers. This suggests that blocking the downstream pathway improves IA production while blocking the efflux effect of the acetyl coenzyme in the peroxisome is not as advantageous. Therefore, the Po1g-2G-ICL Δ strain was selected as the final optimized strain.


Itaconic Acid Production by the Engineered *Y. lipolytica* in Bioreactor

One of the most crucial issues in platform chemicals production is in achieving a high product titers consistently (Gao et al., 2016; Li et al., 2020). To investigate the performance of IA-producing *Y. lipolytica* at

conditions that are more relevant for large-scale application, a 5 L bioreactor was employed. Unlike the procedure conducted in the shaking flask fermentation method, here, the composition of the growth medium was altered and the approach of adding sufficient WCO substrate at once was adopted to avoid the problems caused by fed batch fermentation. In the phase of active cell growth between 24 and 96 h, the Po1g-2G-ICL Δ strain intensively produced IA. During this period, the average specific rate of IA synthesis was 0.8 g/L/h, and the maximum specific rate of 2.3 g/L/h was observed between the 76–96-h intervals (Figure 6). Hence, the maximum titer of IA was 54.55 g/L after the 96-h reaction in the fermenter. At the time of writing, this is the highest IA production achieved by a yeast host reported worldwide. As such, *Y. lipolytica* would be a promising industrial host for IA production from renewable feedstock. Our study also demonstrated that the circular bioeconomy concept can be an effective model for scale-up production of valuable biochemical, in particular with the valorization of WCO as raw material.

CONCLUSION

With increasing global interest in environmental protection and sustainable development, the use of low-cost waste to produce valuable platform chemicals in the industrial scale is gaining



attention. In the few studies conducted to date, IA production in engineered strains of *Y. lipolytica* was predominantly using glucose as the primary carbon source (Blazek et al., 2015; Zhao et al., 2019). Even so, production titers had remained suboptimal (Table 1). This had limited the feasibility of large-scale industrial adoption. Here, we employed the cheap raw material WCO to increase acetyl-CoA availability for conversion into IA in the peroxisome of *Y. lipolytica*. By applying both systems metabolic engineering and bioprocessing optimization strategies in unison, we achieved IA titers of 3.33 g/L in shake flasks and up to 54.55 g/L in stirred-tank bioreactor on WCO as the carbon source without the need for pH control. This amounted to more than 34-folds as compared to the initial titers of 1.58 g/L IA before the optimization of strain and fermentation conditions. In this study, as the supernatant may contain WCO that was not consumed completely, IA cannot be detected directly by HPLC. We used esterification of the supernatant to detect the yield of dimethyl itaconate. While this method, in principle, can be used to determine the theoretical final yield of IA from the esterification rate, it is not the best approach to quantify the exact yield of IA. The development of a more robust and higher throughput method of analysis should be considered in future studies. Furthermore, the yield of organic acids produced by *Y. lipolytica* is primarily affected by the genetic mechanism and various environmental factors, such as the carbon source, nitrogen source, temperature, pH, iron concentration, and dissolved oxygen levels. As such, since bioreactor fermentation with WCO as the sole carbon source is still relatively understudied, further optimization of the fermentation

REFERENCES

Beopoulos, A., Mrozova, Z., Thevenieau, F., Le Dall, M.-T., Hapala, I., Papanikolaou, S., et al. (2008). Control of Lipid Accumulation in the Yeast

conditions could improve IA yields. Nonetheless, the present work on the production of IA by WCO still provides valuable insights that will facilitate further efforts in the biosynthesis of this compound. The results obtained suggest that the oleaginous yeast *Y. lipolytica* is an attractive platform as it provides a viable and scalable pathway to the overproduction of IA and most notably, one that is sustained by waste conversion. However, given the extensive knowledge on IA gene regulation and fermentation conditions, it is believed that higher productivities of IA can be achieved through further engineering of this yeast strain and the optimization of fermentation conditions in subsequent studies.

DATA AVAILABILITY STATEMENT

The datasets presented in this study can be found in online repositories. The names of the repository/repository and accession number(s) can be found in the article/Supplementary Material.

AUTHOR CONTRIBUTIONS

DX, CZ, AW, and, AY conceived and designed the study. LR, LM, SW, YW, SL, ZL, and BZ performed the experiments. LR and KP analyzed data and wrote the manuscript. DX, CZ, AW, and AY critically revised the manuscript. All authors have read and approved the final manuscript.

FUNDING

This work was funded by the Natural Science Foundation of Tianjin, China (17JCYBJC40800), the Research Foundation of Tianjin Municipal Education Commission, China (2017ZD03), the Innovative Research Team of Tianjin Municipal Education Commission, China (TD13-5013), Tianjin Municipal Science and Technology Project (18PTSYJC00140, 19PTSYJC00060), Startup Fund for "Haihe Young Scholars" of Tianjin University of Science and Technology, the Thousand Young Talents Program of Tianjin, China. AW was supported by the Ministry of Education, Singapore (R-MOE-A401-F028) and Lee Foundation (T-LEE-T201-A001).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fbioe.2022.888869/full#supplementary-material>

Yarrowia lipolytica. *Appl. Environ. Microbiol.* 74 (24), 7779–7789. doi:10.1128/AEM.01412-08

Black, P. N., Færgeman, N. J., and DiRusso, C. C. (2000). Long-chain Acyl-CoA-dependent Regulation of Gene Expression in Bacteria, Yeast and Mammals. *J. Nutr.* 130 (2S Suppl. 1), 305S–309S. doi:10.1093/jn/130.2.305S

- Blazek, J., Miller, J., Pan, A., Gengler, J., Holden, C., Jamoussi, M., et al. (2014). Metabolic Engineering of *Saccharomyces cerevisiae* for Itaconic Acid Production. *Appl. Microbiol. Biotechnol.* 98, 8155–8164. doi:10.1007/s00253-014-5895-0
- Blazek, J., Hill, A., Jamoussi, M., Pan, A., Miller, J., and Alper, H. S. (2015). Metabolic Engineering of *Yarrowia lipolytica* for Itaconic Acid Production. *Metab. Eng.* 32, 66–73. doi:10.1016/j.ymben.2015.09.005
- Bonnarme, P., Gillet, B., Sepulchre, A. M., Role, C., Beloeil, J. C., and Ducrocq, C. (1995). Itaconate Biosynthesis in *Aspergillus terreus*. *J. Bacteriol.* 177 (12), 3573–3578. doi:10.1128/jb.177.12.3573-3578.10.1128/jb.177.12.3573-3578.1995
- Braga, A., and Belo, I. (2016). Biotechnological Production of γ -decalactone, a Peach like Aroma, by *Yarrowia lipolytica*. *World J. Microbiol. Biotechnol.* 32 (10), 169. doi:10.1007/s11274-016-2116-2
- Cui, Z., Gao, C., Li, J., Hou, J., Lin, C. S. K., and Qi, Q. (2017). Engineering of Unconventional Yeast *Yarrowia lipolytica* for Efficient Succinic Acid Production from Glycerol at Low pH. *Metab. Eng.* 42, 126–133. doi:10.1016/j.ymben.2017.06.007
- DeLoache, W. C., Russ, Z. N., and Dueber, J. E. (2016). Towards Repurposing the Yeast Peroxisome for Compartmentalizing Heterologous Metabolic Pathways. *Nat. Commun.* 7, 11152. doi:10.1038/ncomms11152
- Dominguez, A., Deive, F. J., Angeles Sanromán, M., and Longo, M. A. (2010). Biodegradation and Utilization of Waste Cooking Oil by *Yarrowia lipolytica* CECT 1240. *Eur. J. Lipid Sci. Technol.* 112, 1200–1208. doi:10.1002/ejlt.201000049
- Dulermo, T., Tréton, B., Beopoulos, A., Kabran Gnankon, A. P., Haddouche, R., and Nicaud, J.-M. (2013). Characterization of the Two Intracellular Lipases of *Y. lipolytica* Encoded by TGL3 and TGL4 Genes: New Insights into the Role of Intracellular Lipases and Lipid Body Organisation. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* 1831 (9), 1486–1495. doi:10.1016/j.bbalip.2013.07.001
- Gao, C., Yang, X., Wang, H., Rivero, C. P., Li, C., Cui, Z., et al. (2016). Robust Succinic Acid Production from Crude Glycerol Using Engineered *Yarrowia lipolytica*. *Biotechnol. Biofuels* 9 (1), xz2179. doi:10.1186/s13068-016-0597-8
- Geiser, E., Przybilla, S. K., Friedrich, A., Buckel, W., Wierckx, N., Blank, L. M., et al. (2016). *Ustilago maydis* Produces Itaconic Acid via the Unusual Intermediate *Trans-aconitate*. *Microb. Biotechnol.* 9 (1), 116–126. doi:10.1111/1751-7915.12329
- Gonçalves, F. A. G., Colen, G., and Takahashi, J. A. (2014). *Yarrowia lipolytica* and its Multiple Applications in the Biotechnological Industry. *Scientific World J.* 2014, 1–14. doi:10.1155/2014/476207
- Haddouche, R., Delessert, S., Sabirova, J., Neuvéglise, C., Poirier, Y., and Nicaud, J.-M. (2010). Roles of Multiple Acyl-CoA Oxidases in the Routing of Carbon Flow towards β -oxidation and Polyhydroxyalkanoate Biosynthesis in *Yarrowia lipolytica*. *FEMS. Yeast. Res.* 10 (7), 917–927. doi:10.1111/j.1567-1364.2010.00670.x
- Hanko, E. K. R., Denby, C. M., Sánchez i Nogué, V., Lin, W., Ramirez, K. J., Singer, C. A., et al. (2018). Engineering β -oxidation in *Yarrowia lipolytica* for Methyl Ketone Production. *Metab. Eng.* 48, 52–62. doi:10.1016/j.ymben.2018.05.018
- Harder, B.-J., Bettenbrock, K., and Klant, S. (2016). Model-based Metabolic Engineering Enables High Yield Itaconic Acid Production by *Escherichia coli*. *Metab. Eng.* 38, 29–37. doi:10.1016/j.ymben.2016.05.008
- Hosseinpour Tehrani, H., Becker, J., Bator, I., Saur, K., Meyer, S., Rodrigues Lóia, A. C., et al. (2019a). Integrated Strain- and Process Design Enable Production of 220 g L⁻¹ Itaconic Acid with *Ustilago maydis*. *Biotechnol. Biofuels* 12, 263. doi:10.1186/s13068-019-1605-6
- Hosseinpour Tehrani, H., Tharmasothirajan, A., Track, E., Blank, L. M., and Wierckx, N. (2019b). Engineering the Morphology and Metabolism of pH Tolerant *Ustilago cynodontis* for Efficient Itaconic Acid Production. *Metab. Eng.* 54, 293–300. doi:10.1016/j.ymben.2019.05.004
- Koivisto, O. M., Kuivaniemi, J., Barth, D., Turkia, H., Pitkänen, J.-P., Penttilä, M., et al. (2013). Glycolic Acid Production in the Engineered Yeasts *Saccharomyces cerevisiae* and *Kluyveromyces lactis*. *Microb. Cell Fact.* 12, 82. doi:10.1186/1475-2859-12-82
- Krull, S., Hevekerl, A., Kuenz, A., and Prüße, U. (2017). Process Development of Itaconic Acid Production by a Natural Wild Type Strain of *Aspergillus terreus* to Reach Industrially Relevant Final Titrers. *Appl. Microbiol. Biotechnol.* 101 (10), 4063–4072. doi:10.1007/s00253-017-8192-x
- Kubicek, C. P., Punt, P., and Visser, J. (2011). “Production of Organic Acids by Filamentous Fungi,” in *Industrial Applications*. Editor M. Hofrichter (Berlin, Heidelberg: Springer), 215–234. doi:10.1007/978-3-642-11458-8_10
- Kuenz, A., Gallenmüller, Y., Wille, T., and Vorlop, K.-D. (2012). Microbial Production of Itaconic Acid: Developing a Stable Platform for High Product Concentrations. *Appl. Microbiol. Biotechnol.* 96 (5), 1209–1216. doi:10.1007/s00253-012-4221-y
- Ledesma-Amaro, R., and Nicaud, J.-M. (2016). *Yarrowia lipolytica* as a Biotechnological Chassis to Produce Usual and Unusual Fatty Acids. *Prog. Lipid Res.* 61, 40–50. doi:10.1016/j.plipres.2015.12.001
- Li, J., Rong, L., Zhao, Y., Li, S., Zhang, C., Xiao, D., et al. (2020). Next-generation Metabolic Engineering of Non-conventional Microbial Cell Factories for Carboxylic Acid Platform Chemicals. *Biotechnol. Adv.* 43, 107605. doi:10.1016/j.biotechadv.2020.107605
- Li, J., Zhu, K., Miao, L., Rong, L., Zhao, Y., Li, S., et al. (2021). Simultaneous Improvement of Limonene Production and Tolerance in *Yarrowia lipolytica* through Tolerance Engineering and Evolutionary Engineering. *ACS Synth. Biol.* 10, 884–896. doi:10.1021/acssynbio.1c00052
- Li, S., Rong, L., Wang, S., Liu, S., Lu, Z., Miao, L., et al. (2022). Enhanced Limonene Production by Metabolically Engineered *Yarrowia lipolytica* from Cheap Carbon Sources. *Chem. Eng. Sci.* 249, 117342. doi:10.1016/j.ces.2021.117342
- Liu, H.-H., Ji, X.-J., and Huang, H. (2015). Biotechnological Applications of *Yarrowia lipolytica*: Past, Present and Future. *Biotechnol. Adv.* 33 (8), 1522–1546. doi:10.1016/j.biotechadv.2015.07.010
- Ma, Y. R., Li, W. J., Mai, J., Wang, J. P., Wei, Y. J., Ledesma-Amaro, R., et al. (2020). Engineering *Yarrowia lipolytica* for Sustainable Production of the Chamomile Sesquiterpene (–)- α -Bisabolol. *Green. Chem.* 23, 780–787. doi:10.1039/d0gc03180a
- Ng, T.-K., Yu, A.-Q., Ling, H., Pratomo Juwono, N. K., Choi, W. J., Leong, S. S. J., et al. (2020). Engineering *Yarrowia lipolytica* towards Food Waste Bioremediation: Production of Fatty Acid Ethyl Esters from Vegetable Cooking Oil. *J. Biosci. Bioeng.* 129 (1), 31–40. doi:10.1016/j.jbiosc.2019.06.009
- Nuss, P., and Gardner, K. H. (2013). Attributional Life Cycle Assessment (ALCA) of Polyitaconic Acid Production from Northeast US Softwood Biomass. *Int. J. Life Cycle Assess.* 18, 603–612. doi:10.1007/s11367-012-0511-y
- Okamoto, S., Chin, T., Hiratsuka, K., Aso, Y., Tanaka, Y., Takahashi, T., et al. (2014). Production of Itaconic Acid Using Metabolically Engineered *Escherichia coli*. *J. Gen. Appl. Microbiol.* 60 (5), 191–197. doi:10.2323/jgam.60.191
- Otten, A., Brocker, M., and Bott, M. (2015). Metabolic Engineering of *Corynebacterium glutamicum* for the Production of Itaconate. *Metab. Eng.* 30, 156–165. doi:10.1016/j.ymben.2015.06.003
- Pang, Y., Zhao, Y., Li, S., Zhao, Y., Li, J., Hu, Z., et al. (2019). Engineering the Oleaginous Yeast *Yarrowia lipolytica* to Produce Limonene from Waste Cooking Oil. *Biotechnol. Biofuels.* 12, 241. doi:10.1186/s13068-019-1580-y
- Porro, D., and Branduardi, P. (2017). “Production of Organic Acids by Yeasts and Filamentous Fungi,” in *Biotechnology of Yeasts and Filamentous Fungi*. Editor A. Sibirny (Cham: Springer), 205–223. doi:10.1007/978-3-319-58829-2_7
- Smith, J. J., Brown, T. W., Eitzen, G. A., and Rachubinski, R. A. (2000). Regulation of Peroxisome Size and Number by Fatty Acid β -Oxidation in the Yeast *Yarrowia lipolytica*. *J. Biol. Chem.* 275 (26), 20168–20178. doi:10.1074/jbc.M909285199
- Sriariyanun, M., Heitz, J. H., Yasurin, P., Asavasanti, S., and Tantayotai, P. (2019). Itaconic Acid: a Promising and Sustainable Platform Chemical? *Appl. Sci. Eng. Prog.* 12 (1), 14410–14416. doi:10.14416/j.asep.2019.05.002
- Strijbis, K., van Roermund, C. W. T., Visser, W. F., Mol, E. C., van den Burg, J., MacCallum, D. M., et al. (2008). Carnitine-dependent Transport of Acetyl Coenzyme A in *Candida albicans* is Essential for Growth on Nonfermentable Carbon Sources and Contributes to Biofilm Formation. *Eukaryot. Cel.* 7 (4), 610–618. doi:10.1128/EC.00017-08
- Strijbis, K., van Roermund, C. W., van den Burg, J., van den Berg, M., Hardy, G. P. M., Wanders, R. J., et al. (2010). Contributions of Carnitine Acetyltransferases to Intracellular Acetyl Unit Transport in *Candida albicans*. *J. Biol. Chem.* 285 (32), 24335–24346. doi:10.1074/jbc.M109.094250
- Sun, W., Vila-Santa, A., Liu, N., Prozorov, T., Xie, D., Faria, N. T., et al. (2020). Metabolic Engineering of an Acid-Tolerant Yeast Strain *Pichia kudriavzevii* for Itaconic Acid Production. *Metab. Eng. Commun.* 10, e00124. doi:10.1016/j.mec.2020.e00124
- Tevz, G., Benčina, M., and Legiša, M. (2010). Enhancing Itaconic Acid Production by *Aspergillus terreus*. *Appl. Microbiol. Biotechnol.* 87 (5), 1657–1664. doi:10.1007/s00253-010-2642-z

- Vuoristo, K. S., Mars, A. E., Sangra, J. V., Springer, J., Eggink, G., Sanders, J. P. M., et al. (2015). Metabolic Engineering of Itaconate Production in *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 99 (1), 221–228. doi:10.1007/s00253-014-6092-x
- Wache, Y., Aguedo, M., Choquet, A., Gatfield, I. L., Nicaud, J.-M., and Belin, J.-M. (2001). Role of β -Oxidation Enzymes in γ -Decalactone Production by the Yeast *Yarrowia lipolytica*. *Appl. Environ. Microbiol.* 67 (12), 5700–5704. doi:10.1128/AEM.67.12.5700-5704.2001
- Wang, J., Ledesma-Amaro, R., Wei, Y., Ji, B., and Ji, X.-J. (2020). Metabolic Engineering for Increased Lipid Accumulation in *Yarrowia lipolytica* - A Review. *Bioresour. Techn.* 313, 123707. doi:10.1016/j.biortech.2020.123707
- Werpy, T., and Petersen, G. (2004). *Top Value Added Chemicals from Biomass: Volume I Results of Screening for Potential Candidates from Sugars and Synthesis Gas*. Golden, CO, US: National Renewable Energy Lab.
- Willke, T., and Vorlop, K.-D. (2001). Biotechnological Production of Itaconic Acid. *Appl. Microbiol. Biotechnol.* 56, 289–295. doi:10.1007/s002530100685
- Xu, P., Qiao, K., and Stephanopoulos, G. (2017). Engineering Oxidative Stress Defense Pathways to Build a Robust Lipid Production Platform in *Yarrowia lipolytica*. *Biotechnol. Bioeng.* 114 (7), 1521–1530. doi:10.1002/bit.26285
- Yang, K., Qiao, Y., Li, F., Xu, Y., Yan, Y., Madzak, C., et al. (2019). Subcellular Engineering of Lipase Dependent Pathways Directed towards Lipid Related Organelles for Highly Effectively Compartmentalized Biosynthesis of Triacylglycerol Derived Products in *Yarrowia lipolytica*. *Metab. Eng.* 55, 231–238. doi:10.1016/j.ymben.2019.08.001
- Yu, A.-Q., Pratomo, N., Ng, T.-K., Ling, H., Cho, H.-S., Leong, S. S. J., et al. (2016). Genetic Engineering of an Unconventional Yeast for Renewable Biofuel and Biochemical Production. *JoVE* 115, 54371. doi:10.3791/54371
- Zhang, J., Jin, B., Hong, K., Lv, Y., Wang, Z., and Chen, T. (2021a). Cell Catalysis of Citrate to Itaconate by Engineered *Halomonas bluephagenesis*. *ACS Synth. Biol.* 10, 3017–3027. doi:10.1021/acssynbio.1c00320
- Zhang, Q., Yu, S., Lyu, Y., Zeng, W., and Zhou, J. (2021b). Systematically Engineered Fatty Acid Catabolite Pathway for the Production of (2S)-Naringenin in *Saccharomyces cerevisiae*. *ACS Synth. Biol.* 10 (5), 1166–1175. doi:10.1021/acssynbio.1c00002
- Zhao, M., Lu, X., Zong, H., Li, J., and Zhuge, B. (2018). Itaconic Acid Production in Microorganisms. *Biotechnol. Lett.* 40 (3), 455–464. doi:10.1007/s10529-017-2500-5
- Zhao, C., Cui, Z., Zhao, X., Zhang, J., Zhang, L., Tian, Y., et al. (2019). Enhanced Itaconic Acid Production in *Yarrowia lipolytica* via Heterologous Expression of a Mitochondrial Transporter MTT. *Appl. Microbiol. Biotechnol.* 103 (5), 2181–2192. doi:10.1007/s00253-019-09627-z
- Zhao, Y., Liu, S., Lu, Z., Zhao, B., Wang, S., Zhang, C., et al. (2021a). Hybrid Promoter Engineering Strategies in *Yarrowia lipolytica*: Isoamyl Alcohol Production as a Test Study. *Biotechnol. Biofuels.* 14 (1), 149. doi:10.1186/s13068-021-02002-z
- Zhao, Y., Zhu, K., Li, J., Zhao, Y., Li, S., Zhang, C., et al. (2021b). High-efficiency Production of Bisabolene from Waste Cooking Oil by Metabolically Engineered *Yarrowia lipolytica*. *Microb. Biotechnol.* 14 (6), 2497–2513. doi:10.1111/1751-7915.13768
- Zhou, J., Yin, X., Madzak, C., Du, G., and Chen, J. (2012). Enhanced α -ketoglutarate Production in *Yarrowia lipolytica* WSH-Z06 by Alteration of the Acetyl-CoA Metabolism. *J. Biotechnol.* 161 (3), 257–264. doi:10.1016/j.jbiotec.2012.05.025
- Zhu, J., Schwartz, C., and Wheeldon, I. (2018). Controlled Intracellular Trafficking Alleviates an Expression Bottleneck in *S. cerevisiae* Ester Biosynthesis. *Metab. Eng. Commun.* 8, e00085. doi:10.1016/j.mec.2018.e00085
- Zhu, K., Kong, J., Zhao, B., Rong, L., Liu, S., Lu, Z., et al. (2021). Metabolic Engineering of Microbes for Monoterpenoid Production. *Biotechnol. Adv.* 53, 107837. doi:10.1016/j.biotechadv.2021.107837
- Zinjarde, S. S. (2014). Food-related Applications of *Yarrowia lipolytica*. *Food Chem.* 152, 1–10. doi:10.1016/j.foodchem.2013.11.117

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Rong, Miao, Wang, Wang, Liu, Lu, Zhao, Zhang, Xiao, Pushpanathan, Wong and Yu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.