



OPEN ACCESS

EDITED BY

Artur Ribeiro,
University of Minho, Portugal

REVIEWED BY

Jorge Padrão,
University of Minho, Portugal
Paul P. De Waal,
Dsm-firmenich, Netherlands

*CORRESPONDENCE

Dave Siak-Wei Ow,
✉ dave_ow@a-star.edu.sg
✉ dave_ow@bti.a-star.edu.sg
Fong Tian Wong,
✉ wongft@imcb.a-star.edu.sg

RECEIVED 14 January 2025

ACCEPTED 11 July 2025

PUBLISHED 24 July 2025

CITATION

Ho PL, Chua YF, Quek JP, Ng SK, Wong FT and
Ow DS-W (2025) Functional expression and
secretion of basic fibroblast growth factor in
Lactococcus lactis.
Front. Bioeng. Biotechnol. 13:1560426.
doi: 10.3389/fbioe.2025.1560426

COPYRIGHT

© 2025 Ho, Chua, Quek, Ng, Wong and Ow.
This is an open-access article distributed under
the terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Functional expression and secretion of basic fibroblast growth factor in *Lactococcus lactis*

Pooi Leng Ho¹, Yu Feng Chua¹, Jun Ping Quek¹, Say Kong Ng¹,
Fong Tian Wong^{2*} and Dave Siak-Wei Ow^{1*}

¹Bioprocessing Technology Institute (BTI), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore, ²Institute of Molecular and Cell Biology (IMCB), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore

Introduction: Cultivated meat, produced by in vitro cell culture in bioreactors, offers a sustainable alternative to traditional meat sources. A significant challenge in its production is the high cost of mitogenic growth factors, which are essential supplements in serum-free media for cultivating meat cells. One strategy to reduce cost involves minimizing purification cost by using a food-grade host to secrete growth factors. In this study, we investigate the production of recombinant FGF2 (Fibroblast Growth Factor 2) through secretion in *Lactococcus lactis*, a Generally Recognized As Safe (GRAS) organism.

Method: To enhance the secretion in *L. lactis*, we employed the USP45 secretory peptide and secretion propeptide (PP1) in the design of our recombinant FGF2-G3. Optimization was performed on various culture parameters that influence protein expression, including media formulation, nisin concentration, induction timing, temperature, and culture duration. Secreted FGF2-G3 produced under optimized conditions was purified and tested for bioactivity on *Anguilla japonica* pre-adipocytic cells, Aj1C-2x.

Results and Discussion: We have generated a recombinant *L. lactis* strain and an optimal expression strategy to enable the production of secreted bioactive growth factors. Our results demonstrate that this system can produce FGF2 which were able to promote the proliferation of fish *Anguilla japonica* pre-adipocytic cells. Despite minimal purification beyond affinity purification and buffer exchange, we were able to obtain comparable specific activity to commercial FGF2. The final yields can be derived at 1.97 mg/L and through simple protein purification and buffer exchange. Finally, this study highlights the potential use of *L. lactis* secretion as an endotoxin-free alternative, compared to *E. coli*, for production of growth factors for use in cultivated meat production.

KEYWORDS

cultivated meat, serum-free media, fibroblast growth factor 2, FGF2, *Lactococcus lactis*, recombinant protein expression, precision fermentation

Introduction

Cultivated meat, also known as cultured meat, has gained significant interest in recent years due to global movement towards achieving sustainability development goals. It provides a prospective alternative meat source to support demand from increasing population, reduce environmental impact from animal agriculture and avoids animal-borne diseases (Campbell et al., 2017; Godfray et al., 2018; Lynch and Pierrehumbert, 2019; Stephens et al., 2018). Cultivated meat is produced through *in vitro* culturing of animal cells in cell culture media. However, foetal bovine serum (FBS), a typical ingredient found in culture media, is derived from extracting blood serum of bovine fetuses from animal slaughter houses (Jochems et al., 2002; Kadim et al., 2015; Lee et al., 2022; Post et al., 2020; Reiss et al., 2021). This conflicts with the concept of producing meat via an animal-free approach. Thus, serum-free media formulations that are capable of sustaining cell culture were developed (Badenes et al., 2016; Das et al., 2009; Messmer et al., 2022; Skrivergaard et al., 2023; Stout et al., 2022). Subsequently, supplementation of growth factors, such as fibroblast growth factors or insulin-like growth factors, into serum-free media formulations are essential to mimic proliferative and developmental effects of FBS (Park et al., 2013; Santos et al., 2023; Venkatesan et al., 2022; Yu et al., 2023).

One such growth factor of interest is the basic fibroblast growth factor, also known as fibroblast growth factor 2 (FGF2), which is a member of the cytokine family. It acts by binding to cell surface receptors (FGFR), activating mitogenic pathways such as PI3k/Akt pathway, MAPK/ERK pathway and JNK pathway. Activation of these pathways regulates cellular responses such as growth, proliferation, migration, maintenance and differentiation (Ahmad et al., 2023; Bikfalvi et al., 1997; Yun et al., 2010). Production of recombinant growth factors, including FGF2, for supplementation into serum-free media are most frequently done in prokaryotic expression system using *Escherichia coli*. However, the recombinant proteins are produced intracellularly and has high tendency for inclusion bodies formation, which subsequently involve expensive and tedious downstream protein refolding and purification processes. Furthermore, host-cell derived impurities, in particular endotoxins, poses health risks to humans (Baneyx and Mujacic, 2004; Kaur et al., 2018; Petsch and Anspach, 2000; Sahdev et al., 2008; Thomas and Baneyx, 1996). This calls for an alternative endotoxin free expression system that is also economical for production of recombinant proteins.

Lactococcus lactis, a Gram-positive lactic acid bacterium that is widely used in food and therapeutic applications (Bahey-El-Din et al., 2010; Kumari et al., 2011; Song et al., 2017), presents a good alternative host for recombinant protein expression. The key feature of using *Lactococcus lactis* expression system is its ability to secrete recombinant proteins into culture medium, minimising the need for cell lysis, complex protein purification and refolding. Moreover, *L. lactis* does not produce lipopolysaccharides and has few extracellular

proteases, that causes endotoxin toxicity and proteolytic degradation respectively (Frelet-Barrand, 2022; Garcia-Fruitos, 2012; Morello et al., 2008). As *L. lactis* are microaerophilic, they only require a simple static fermentation process without aeration, this makes possible for a simple and direct scale-up to industrial scale. Among the various *L. lactis* expression systems developed, the most widely used is the nisin-controlled gene expression (NICE) system, consisting of a nisRK regulatory gene integrated into bacterial host chromosome and an expression vector with nisA promoter to tightly regulate gene expression (de Ruyter et al., 1996; Mierau and Kleerebezem, 2005; Mierau et al., 2005a; Zhou et al., 2006). *L. lactis* expression system has been applied for production of several growth factor proteins (Cao et al., 2020; Gao et al., 2012; Huynh and Li, 2015; Zhou et al., 2021). In our lab, we have recently reported on valorisation of mammalian spent culture media waste to support intracellular FGF2 production in bioreactors (Rizal et al., 2024). However, comprehensive research regarding the production and secretion of functional FGF2 from *L. lactis* is not available. Hence, we set forth herein to investigate the possibility of employing *L. lactis* NICE expression system to produce and secrete biologically active FGF2. To enhance the secretion efficiency, we fused USP45 secretory peptide and secretion propeptide 1 (PP1) (Lim et al., 2017) to a thermostable FGF2 variant, FGF2-G3 (Dvorak et al., 2018). Together with optimisation of media formulation and culture conditions, we were able to obtain ~2 mg/L of secreted FGF2-G3. Furthermore, FGF2-G3 purified from the medium was able to stimulate proliferation of the Japanese eel *Anguilla japonica* pre-adipocytic cells, comparable to commercial FGF2. Together, these results signal the potential application of *L. lactis* protein secretion system as an alternative strategy for recombinant FGF2 and potentially other growth factor production to circumvent issues faced with *E. coli* for cultured meat development.

Materials and methods

Bacterial strain, plasmid and cloning of FGF2-G3 gene

L. lactis NZ9000 and pNZ8148 plasmid (BoCa Scientific, United States) were used for cloning and expression studies. Sequence of the thermostable human FGF2-G3 was obtained from Dvorak et al. (2018). To enhance expression and secretion into medium, USP45 secretion peptide (Accession ABY84357) and propeptide 1 (PP1) (Lim et al., 2017) were fused at the N-terminus of FGF2-G3 sequence. For ease of purification with Ni-NTA affinity chromatography and Western blot detection, we have also included His₆ sequence at the N-terminus. The nucleotide sequences corresponding to the amino acids were codon optimized and synthesized by IDT (Singapore) for expression in *L. lactis*. The coding sequence was cloned into the multiple cloning site (MCS) of pNZ8148 vector using NeBuilder HiFi Assembly (New England Biolabs, United States) (Fusion protein sequence available in Supplementary Figure S1), transformed into *L. lactis* NZ9000 and plated onto M17 agar plate containing 0.5% (w/v) glucose and 10 µg/mL chloramphenicol to screen for positive recombinant clones. The positive recombinant clones were further sequenced to ensure no mutations prior to protein expression with *L. lactis* NZ9000.

Abbreviations: EGF, Epidermal Growth Factor; FBS, Foetal Bovine Serum; FGF2, Fibroblast Growth Factor 2; GRAS, Generally Recognized As Safe; IGF, Insulin-like Growth Factor; NICE, Nisin Controlled Gene Expression System; TGF-β1, Transforming Growth Factor β1.

Culture optimization for FGF2-G3 expression and secretion in *L. lactis*

Productivity of FGF2-G3 production in *L. lactis* was assessed with varying M17 media and glucose concentration. They were expressed in either M17 (supplemented with 0.5% (w/v) glucose), 2xM17 (supplemented with 0.5% (w/v) glucose) or 2xM17 (supplemented with 2% (w/v) glucose). All cultures were also supplemented with 10 µg/mL chloramphenicol for selection and maintenance of cells containing pNZ8148-FGF2-G3 plasmids. To increase production and secretion level, FGF2-G3 expression was further optimized with different nisin inducer concentrations (10, 25, 50 ng/mL), induction time points (OD_{600nm} 0.5, 1.0, 2.0), incubation temperatures (20, 25, 30, 35°C) and expression duration (4, 20 h post-induction). These parameters were investigated individually in 10 mL culture volume. The cultures were harvested by centrifugation at 3,845 g, 4°C for 20 min with Sorvall ST 40 centrifuge (Thermo Fisher Scientific, United States). The media fraction was aspirated and concentrated with a 10 kDa MWCO centrifugal filter (Merck Millipore, Ireland) to obtain secreted protein fraction, while the cell pellet was resuspended in PBS, incubated at 37°C for 1 h with 1 mg/mL lysozyme and 0.05 U/µL mutanolysin and then lysed via sonication. The cell lysate was centrifuged, supernatant collected as intracellular soluble fraction, and the lysed cell pellet resuspended in 6M urea to obtain intracellular insoluble fraction. All 3 fractions (secreted, soluble, insoluble) were analysed on SDS-PAGE to determine productivity level.

Optimized production of secreted FGF2-G3

A larger production volume of FGF2-G3 was performed under optimized conditions. 250 mL of 2xM17 (supplemented with 2% (w/v) glucose and chloramphenicol 10 µg/mL) was inoculated with pre-culture and incubated at 30°C until it reaches OD_{600nm} 1.0. Nisin was then added to final concentration of 25 ng/mL to induce expression of FGF2-G3, and induction was done at 35°C for 20 h. The culture was centrifuged and the media fraction containing secreted FGF2-G3 was collected and purified via affinity chromatography.

Purification of FGF2-G3

Media fraction containing secreted FGF2-G3 was concentrated, and buffer exchanged to buffer A (50 mM NaH₂PO₄, 300 mM NaCl, 10 mM Imidazole, pH 8) using crossflow filtration system. The concentrated sample was loaded onto Ni-NTA Agarose (Qiagen, United States) column and incubated for 60 min. The column was first washed with 10 mM imidazole, next with 20 mM imidazole, and finally eluted with 250 mM imidazole. The eluted fraction was buffer exchanged to PBS for protein evaluation with SDS-PAGE and bioactivity assay. The purified FGF2-G3 was quantified using Bradford method, with BSA as standard.

Western blot and SDS-PAGE

An equal amount of protein from each fraction were run on NuPage 4%–12% Bis-Tris SDS-PAGE gel (Thermo Fisher Scientific,



United States) and transferred onto nitrocellulose membrane using a Trans-Blot SD Semi-Dry Electrophoretic Transfer Cell (Bio-Rad, United States). The membrane was subsequently probed with His-Tag Antibody HRP Conjugate (Merck Millipore, United States) and detected with Clarity Western ECL Blotting Substrate (Bio-Rad, United States). Expression level of FGF2-G3 was determined by densitometric analysis of digital images using ImageJ software (National Institute of Health, United States).

To analyse purity of the large-scale purified media fraction eluted from Ni-NTA agarose column, 2 µg of protein was run on NuPage 4%–12% Bis-Tris SDS-PAGE gel (Thermo Fisher Scientific, United States) and stained with InstantBlue Coomassie Protein stain (Abcam, UK).

Bioactivity assay of FGF2-G3

Biological activity of the purified FGF2-G3 was assessed using *Anguilla japonica* (Japanese eel) pre-adipocytic cells, Aj1C-2x (Sugii et al., 2011). The cells were cultured in Dulbecco's modified Eagle's medium (DMEM)/F12 media (Thermo Fisher Scientific, United States) supplemented with reduced fetal bovine serum (2.5%) and 10 ng/mL of FGF2 at 27°C with 5% CO₂ in a humidified incubator. Cell density and viability were determined using Vi-CELL XR Cell Viability Analyzer (Beckman Coulter, United States), according to manufacturer's instructions. Each well contains fresh DMEM/F12 medium supplemented with 2.5% FBS and varying concentrations of purified recombinant FGF2-G3 or positive control. Commercial heat stable FGF2 (PHG0368, Thermo Fisher Scientific) was used as positive control. Aj1C-2x cells were seeded into 96-well plates at seeding density of 2 × 10⁴ cells/well. After culturing for 3 days, cell viability was determined with CyQuant XTT cell viability assay (X12223, Thermo Fischer Scientific), according to manufacturer's instructions. The absorbance reading of the media control was subtracted from the absorbance reading of each sample to determine the specific absorbance reading for each sample. Subsequently, the specific absorbance reading of each sample were normalized against the specific absorbance reading of the cells cultivated in basal media with 2.5% FBS (0 ng/mL FGF2) for Aj1C-2x cells to calculate relative absorbance fold changes. Cellular health was examined through inverted microscope under ×4 magnification (Nikon eclipse Ti with NIS-Elements AR 4.30.02 software, Nikon).

Biological activity of the purified FGF2-G3 was also further evaluated in a 6-well plate format. Aj1C-2x cells were seeded at a seeding density of 3 × 10⁵ cells/well in a 6-well plate using DMEM/F12 medium (Thermo Fisher Scientific, United States) supplemented with 2.5% FBS and varying concentrations of purified recombinant FGF2-G3 or commercial heat-stable FGF2 (PHG0368, Thermo Fisher Scientific). After 4 days of culture, cellular health was examined through inverted microscope (Nikon eclipse Ti with NIS-Elements AR 4.30.02 software, Nikon). Subsequently, cells were dissociated using TrypLE™ Express Enzyme (Thermo Fisher Scientific, United States) and viable cell density were measured using a Vi-CELL XR Cell Viability Analyzer (Beckman Coulter, United States), following the manufacturer's protocol.



Results and discussion

An enhancer propeptide to enhance secretion

To determine whether FGF-2 can be expressed and secreted out of *L. lactis*, we constructed an expression plasmid containing FGF2-G3, a modified and stable version that has nine amino acid mutations (Dvorak et al., 2018), with N-terminal fused to USP45 signal peptide. Our initial attempts to express the fusion protein resulted in low intracellular soluble yield and no protein was secreted. In previous study conducted by Lim et al. (2017), it was reported that addition of a short secretion propeptide 1 (PP1) to a USP45 fusion protein could significantly enhance protein secretion efficiency. Hence, we added PP1 between USP45 and FGF2-G3 as shown in Figure 1A.

L. lactis cells containing the two different recombinant plasmids (with and without PP1) were cultured in M17 media, supplemented with 0.5% glucose, and expression was induced at OD_{600nm} 0.5 with nisin at final concentration of 10 ng/mL. FGF2-G3 was allowed to be expressed for 4 h at 30°C. The expression and secretion were assessed by Western blot. Appearance of bands corresponding to two different fusion FGF2-G3 constructs (with and without PP1) were observed. It was noted that addition of PP1 not only facilitates secretion of the fusion protein, but also increases the soluble yield (Figure 1B).

Optimisation of cultivation parameters

Next, optimization of culture and expression conditions were examined to increase protein secretion yield. Optimization of culture conditions began with expressing FGF2-G3 in varying M17 and glucose concentrations (M17 + 0.5% (w/v) glucose, 2xM17 + 0.5% (w/v) glucose, 2xM17 + 2% (w/v) glucose). 2xM17 + 2% (w/v) glucose medium proved to be the best for secretion of FGF2-G3 (Figure 2A).

Doubling the amount of M17 increased the soluble protein fractions. Looking at the growth curve of cells grown in different medias (Figure 2B), we hypothesize that doubling M17 not only provides additional nitrogen-based nutrients to support cell metabolism for a higher cell density culture, but also increases the buffering capacity against lactic acid produced during fermentation, due to increased amount of Disodium-β-glycerophosphate, a buffering agent found in M17 medium composition (Hayek et al., 2019; Terzaghi and Sandine, 1975; Zhang et al., 2009). These then worked in concert with increased glucose concentration, permitting *L. lactis* to extend its growth phase for higher FGF2-G3 production and, in particular, secretion titer. Further culture optimizations were performed using 2xM17 + 2% (w/v) glucose medium.

FGF2-G3 production also improved when the culture was induced at a higher cell density of OD_{600nm} 1.0 (Figure 2C). This is anticipated since higher cell density would also mean more plasmids available for induction, which in-turn raises protein expression titer. With the increase in cell density, more nisin may be needed for complete induction and thus we proceeded to determine new optimal nisin concentration. Culture induction was


performed when cells reached OD_{600nm} 1.0 with nisin at final concentrations of 10, 25 or 50 ng/mL. The expression level of FGF2-G3 increased when nisin concentration was up from 10 to 25 ng/mL (Figure 2C), indicating correlation between cell density and nisin concentration needed for maximal induction. There was no further increase in expression level with 50 ng/mL nisin. Induction was also tested at OD_{600nm} 2.0, but it did not lead to higher FGF2-G3 yield.

As the FGF2-G3 used in this study has been engineered for stability, we predicted that translation and folding within *L. lactis* is not limiting, rather the rate of translation/transcription can be further improved via fermentation optimisation. Temperature and expression duration are two important post-induction conditions for optimization as they balance between bacterial growth, functional protein yield and protein degradation. It has been suggested that lower temperature improves proper protein folding and solubility (Mierau et al., 2005; Sahdev et al., 2008; Yu et al., 2021), and prolonged expression should also be avoided as it leads to higher tendency for protein degradation caused by protein instability, lactate accumulation that disrupts energy metabolism for protein expression and/or triggering of cell stress response (Zhou et al., 2006). Interestingly, results for the present study showed an increase in secreted protein yield along with increasing temperature and expression duration, with 35°C and 20 h being the optimal temperature and harvest time-point (Figure 2D).

The highest expression and secretion of FGF2-G3 was achieved when cells were cultured in 2xM17 medium supplemented with 2% (w/v) glucose and induced at OD_{600nm} 1.0 with 25 ng/mL nisin for 20 h at 35°C. The culture was scaled up from 10 mL to 250 mL for subsequent purification and bioactivity testing. The secreted fraction of the overexpressed FGF2-G3 was purified using immobilized metal affinity chromatography, which utilized Ni-NTA resin matrix, and the identity and purity were determined with Western blot and Coomassie staining (Figures 3A,B). The purified yield achieved herein is about 1.97 mg for 1 L of fermentation media, which is comparable to previous study by Rizal et al. (2024) where 2.6 mg/L of intracellular FGF2-G3 was produced from bioreactor fermentation. Based on an estimated requirement of 50 ng/mL recombinant FGF2-G3 in cultivated meat culture, secreted growth factors produced from 1 L fermentation provides enough growth factors for 39.4 L of cultivated meat culture. Secretion titers could be further raised with i) plasmid modifications, such as replacing promoters, ii) increasing membrane porosity by addition of chemicals, such as peptides and detergents, into culture media and iii) using bioreactor fermentation with controlled environment.

Biological activity assessment of the purified FGF2

To assess the biological activity of the purified FGF2-G3, growth stimulation on *Anguilla japonica* (Japanese eel) pre-adipocytic cells, Aj1C-2x, was measured using XTT assay and compared against a commercial heat stable FGF2 (positive control). As shown in Figure 3C, an increase in metabolic activity was detected when cells were cultured with the purified FGF2-G3, indicating its ability to promote cell proliferation and exhibited comparable bioactivity



profile to the commercial FGF2. This is further supported by an increase in cell density observed through microscopy and cell count using a cell viability analyzer (Figures 3D,E). The positive result suggests that our secreted FGF2-G3 can be used to stimulate fish stem cells for cultivated fish meat and cultivating adipocytes (fats) for enhancement of meat texture and flavour. Future bioactivity testing can be performed on mammalian cell lines, such as bovine, porcine and chicken muscle cells, to widen its application range in cultivated meat.

Conclusion

In summary, this study demonstrated that functional FGF2 can be expressed and secreted using the *L. lactis* expression system. We employed a multi-modal optimization strategy that included a secretion-enhancing propeptide and further cultivation optimizations. Specifically, we utilized a nutrient-rich 2xM17 medium containing 2% (w/v) glucose. The ability of the *L. lactis*-produced FGF2-G3 to stimulate growth of Japanese eel cells suggests that *L. lactis* could be used as a safer

alternative for production of growth factors for cultured meat development. The secretion of recombinant proteins into culture medium by *L. lactis* simplifies the production process and presents opportunities to lower production cost for cultured meat. Future efforts could be directed at expressing other growth factors, such as EGF, IGF and TGF β 1, in *L. lactis* and scaling-up in bioreactors for precision fermentation of recombinant growth factors.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary Material](#), further inquiries can be directed to the corresponding authors.

Ethics statement

Ethical approval was not required for the studies on animals in accordance with the local legislation and institutional requirements because only commercially available established cell lines were used.

Author contributions

PH: Conceptualization, Data curation, Formal Analysis, Investigation, Methodology, Validation, Visualization, Writing – Original draft, Project administration. YC: Formal Analysis, Investigation, Methodology, Validation, Writing – Review and editing. JQ: Formal Analysis, Validation, Writing – Review and editing. SN: Writing – Review and editing. FW: Writing – Review and editing. DO: Conceptualization, Funding acquisition, Project administration, Supervision, Writing – Review and editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. This research was supported by the Agency for Science, Technology and Research

(A*STAR), Singapore, and was funded by Singapore Food Story (SFS) R&D Programme (H20H8a0003, W23W2D0009 and NRF-SFSRND2FF-0003).

Acknowledgments

We would like to thank Shigeki Sugii and Lamony Chew for providing us with the *Anguilla japonica* pre-adipocytic cell lines and Zarra Iwana Huang for her assistance in culturing work.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fbioe.2025.1560426/full#supplementary-material>

References

- Ahmad, S. S., Chun, H. J., Ahmad, K., Shaikh, S., Lim, J. H., Ali, S., et al. (2023). The roles of growth factors and hormones in the regulation of muscle satellite cells for cultured meat production. *J. Anim. Sci. Technol.* 65 (1), 16–31. doi:10.5187/jast.2022.e114
- Badenes, S. M., Fernandes, T. G., Cordeiro, C. S., Boucher, S., Kuninger, D., Vemuri, M. C., et al. (2016). Defined essential 8 medium and vitronectin efficiently support scalable xeno-free expansion of human induced pluripotent stem cells in stirred microcarrier culture systems. *PLoS One* 11 (3), e0151264. doi:10.1371/journal.pone.0151264
- Bahey-El-Din, M., Gahan, C. G., and Griffin, B. T. (2010). Lactococcus lactis as a cell factory for delivery of therapeutic proteins. *Curr. Gene Ther.* 10 (1), 34–45. doi:10.2174/156652310790945557
- Baneyx, F., and Mujacic, M. (2004). Recombinant protein folding and misfolding in *Escherichia coli*. *Nat. Biotechnol.* 22 (11), 1399–1408. doi:10.1038/nbt1029
- Bikfalvi, A., Klein, S., Pintucci, G., and Rifkin, D. B. (1997). Biological roles of fibroblast growth factor-2. *Endocr. Rev.* 18 (1), 26–45. doi:10.1210/edrv.18.1.0292
- Campbell, B. M., Beare, D. J., Bennett, E. M., Hall-Spencer, J. M., Ingram, J. S. I., Jaramillo, F., et al. (2017). Agriculture production as a major driver of the Earth system exceeding planetary boundaries. *Ecol. Soc.* 22 (4), art8. doi:10.5751/es-09595-220408
- Cao, W. Y., Dong, M., Hu, Z. Y., Wu, J., Li, Y. C., and Xu, H. D. (2020). Recombinant Lactococcus lactis NZ3900 expressing bioactive human FGF21 reduced body weight of Db/Db mice through the activity of brown adipose tissue. *Benef. Microbes* 11 (1), 67–78. doi:10.3920/BM2019.0093
- Das, M., Rumsey, J. W., Bhargava, N., Gregory, C., Reidel, L., Kang, J. F., et al. (2009). Developing a novel serum-free cell culture model of skeletal muscle differentiation by systematically studying the role of different growth factors in myotube formation. *Vitro Cell Dev. Biol. Anim.* 45 (7), 378–387. doi:10.1007/s11626-009-9192-7
- de Ruyter, P. G., Kuipers, O. P., and de Vos, W. M. (1996). Controlled gene expression systems for Lactococcus lactis with the food-grade inducer nisin. *Appl. Environ. Microbiol.* 62 (10), 3662–3667. doi:10.1128/aem.62.10.3662-3667.1996
- Dvorak, P., Bednar, D., Vanacek, P., Balek, L., Eiselleova, L., Stepankova, V., et al. (2018). Computer-assisted engineering of hyperstable fibroblast growth factor 2. *Biotechnol. Bioeng.* 115 (4), 850–862. doi:10.1002/bit.26531
- Frelet-Barrand, A. (2022). Lactococcus lactis, an attractive cell factory for the expression of functional membrane proteins. *Biomolecules* 12 (2), 180. doi:10.3390/biom12020180

- Gao, G., Qiao, J. J., Yang, C. H., Jiang, D. Z., Li, R. Q., Su, J. J., et al. (2012). Functional expression of mouse insulin-like growth factor-I with food-grade vector in *Lactococcus lactis* NZ9000. *Lett. Appl. Microbiol.* 54 (5), 404–409. doi:10.1111/j.1472-765X.2012.03222.x
- Garcia-Fruitos, E. (2012). Lactic acid bacteria: a promising alternative for recombinant protein production. *Microb. Cell Fact.* 11, 157. doi:10.1186/1475-2859-11-157
- Godfray, H. C. J., Aveyard, P., Garnett, T., Hall, J. W., Key, T. J., Lorimer, J., et al. (2018). Meat consumption, health, and the environment. *Science* 361 (6399), eaam5324. doi:10.1126/science.aam5324
- Hayek, S. A., Gyawali, R., Aljaloud, S. O., Krastanov, A., and Ibrahim, S. A. (2019). Cultivation media for lactic acid bacteria used in dairy products. *J. Dairy Res.* 86 (4), 490–502. doi:10.1017/S002202991900075X
- Huynh, E., and Li, J. (2015). Generation of *Lactococcus lactis* capable of coexpressing epidermal growth factor and trefoil factor to enhance *in vitro* wound healing. *Appl. Microbiol. Biotechnol.* 99 (11), 4667–4677. doi:10.1007/s00253-015-6542-0
- Jochems, C. E., van der Valk, J. B., Stafleu, F. R., and Baumans, V. (2002). The use of fetal bovine serum: ethical or scientific problem? *Altern. Lab. Anim.* 30 (2), 219–227. doi:10.1177/026119290203000208
- Kadim, I. T., Mahgoub, O., Baqir, S., Faye, B., and Purchas, R. (2015). Cultured meat from muscle stem cells: a review of challenges and prospects. *J. Integr. Agric.* 14 (2), 222–233. doi:10.1016/S2095-3119(14)60881-9
- Kaur, J., Kumar, A., and Kaur, J. (2018). Strategies for optimization of heterologous protein expression in *E. coli*: roadblocks and reinforcements. *Int. J. Biol. Macromol.* 106, 803–822. doi:10.1016/j.ijbiomac.2017.08.080
- Kumari, A., Catanzaro, R., and Marotta, F. (2011). Clinical importance of lactic acid bacteria: a short review. *Acta Biomed.* 82 (3), 177–180. Available online at: <https://www.ncbi.nlm.nih.gov/pubmed/22783712>.
- Lee, D. Y., Lee, S. Y., Yun, S. H., Jeong, J. W., Kim, J. H., Kim, H. W., et al. (2022). Review of the current research on fetal bovine serum and the development of cultured meat. *Food Sci. Anim. Resour.* 42 (5), 775–799. doi:10.5851/kosfa.2022.e46
- Lim, P. Y., Tan, L. L., Ow, D. S., and Wong, F. T. (2017). A propeptide toolbox for secretion optimization of flavobacterium meningosepticum endopeptidase in *Lactococcus lactis*. *Microb. Cell Fact.* 16 (1), 221. doi:10.1186/s12934-017-0836-0
- Lynch, J., and Pierrehumbert, R. (2019). Climate impacts of cultured meat and beef cattle. *Front. Sustain. Food Syst.* 3, 5. doi:10.3389/fsufs.2019.00005
- Messmer, T., Klevernic, I., Furquim, C., Ovchinnikova, E., Dogan, A., Cruz, H., et al. (2022). A serum-free media formulation for cultured meat production supports bovine satellite cell differentiation in the absence of serum starvation. *Nat. Food* 3 (1), 74–85. doi:10.1038/s43016-021-00419-1
- Mierau, I., and Kleerebezem, M. (2005). 10 years of the nisin-controlled gene expression system (NICE) in *Lactococcus lactis*. *Appl. Microbiol. Biotechnol.* 68 (6), 705–717. doi:10.1007/s00253-005-0107-6
- Mierau, I., Leij, P., van Swam, I., Blommestein, B., Floris, E., Mond, J., et al. (2005a). Industrial-scale production and purification of a heterologous protein in *Lactococcus lactis* using the nisin-controlled gene expression system NICE: the case of lysostaphin. *Microb. Cell Fact.* 4, 15. doi:10.1186/1475-2859-4-15
- Mierau, I., Olieman, K., Mond, J., and Smid, E. J. (2005b). Optimization of the *Lactococcus lactis* nisin-controlled gene expression system NICE for industrial applications. *Microb. Cell Fact.* 4, 16. doi:10.1186/1475-2859-4-16
- Morello, E., Bermudez-Humaran, L. G., Llull, D., Sole, V., Miraglio, N., Langella, P., et al. (2008). *Lactococcus lactis*, an efficient cell factory for recombinant protein production and secretion. *J. Mol. Microbiol. Biotechnol.* 14 (1-3), 48–58. doi:10.1159/000106082
- Park, Y. H., Gong, S. P., Kim, H. Y., Kim, G. A., Choi, J. H., Ahn, J. Y., et al. (2013). Development of a serum-free defined system employing growth factors for preantral follicle culture. *Mol. Reprod. Dev.* 80 (9), 725–733. doi:10.1002/mrdr.22204
- Petsch, D., and Anspach, F. B. (2000). Endotoxin removal from protein solutions. *J. Biotechnol.* 76 (2-3), 97–119. doi:10.1016/S0168-1656(99)00185-6
- Post, M. J., Levenberg, S., Kaplan, D. L., Genovese, N., Fu, J., Bryant, C. J., et al. (2020). Scientific, sustainability and regulatory challenges of cultured meat. *Nat. Food* 1 (7), 403–415. doi:10.1038/s43016-020-0112-z
- Reiss, J., Robertson, S., and Suzuki, M. (2021). Cell sources for cultivated meat: applications and considerations throughout the production workflow. *Int. J. Mol. Sci.* 22 (14), 7513. doi:10.3390/ijms22147513
- Rizal, J., Mainali, P., Quek, J. P., Lee Ling, T., Bi, J., Chan, A. J., et al. (2024). Valorisation of spent cultivated meat media for recombinant FGF2 production in *GRAS Lactococcus lactis*. *bioRxiv* 2024. doi:10.1101/2024.08.01.606190
- Sahdev, S., Khattar, S. K., and Saini, K. S. (2008). Production of active eukaryotic proteins through bacterial expression systems: a review of the existing biotechnology strategies. *Mol. Cell Biochem.* 307 (1-2), 249–264. doi:10.1007/s11010-007-9603-6
- Santos, A. C. A., Camarena, D. E. M., Roncoli Reigado, G., Chambergo, F. S., Nunes, V. A., Trindade, M. A., et al. (2023). Tissue engineering challenges for cultivated meat to meet the real demand of a global market. *Int. J. Mol. Sci.* 24 (7), 6033. doi:10.3390/ijms24076033
- Skrivergaard, S., Young, J. F., Sahebkhatri, N., Semper, C., Venkatesan, M., Savchenko, A., et al. (2023). A simple and robust serum-free media for the proliferation of muscle cells. *Food Res. Int.* 172, 113194. doi:10.1016/j.foodres.2023.113194
- Song, A. A., In, L. L. A., Lim, S. H. E., and Rahim, R. A. (2017). A review on *Lactococcus lactis*: from food to factory. *Microb. Cell Fact.* 16 (1), 55. doi:10.1186/s12934-017-0669-x
- Stephens, N., Di Silvio, L., Dunsford, I., Ellis, M., Glencross, A., and Sexton, A. (2018). Bringing cultured meat to market: technical, socio-political, and regulatory challenges in cellular agriculture. *Trends Food Sci. and Technol.* 78, 155–166. doi:10.1016/j.tifs.2018.04.010
- Stout, A. J., Mirlani, A. B., Rittenberg, M. L., Shub, M., White, E. C., Yuen, J. S. K., Jr., et al. (2022). Simple and effective serum-free medium for sustained expansion of bovine satellite cells for cell cultured meat. *Commun. Biol.* 5 (1), 466. doi:10.1038/s42003-022-03423-8
- Sugii, S., Kida, Y., Berggren, W. T., and Evans, R. M. (2011). Feeder-dependent and feeder-independent iPSC cell derivation from human and mouse adipose stem cells. *Nat. Protoc.* 6 (3), 346–358. doi:10.1038/nprot.2010.199
- Terzaghi, B. E., and Sandine, W. E. (1975). Improved medium for lactic streptococci and their bacteriophages. *Appl. Microbiol.* 29 (6), 807–813. doi:10.1128/am.29.6.807-813.1975
- Thomas, J. G., and Baneyx, F. (1996). Protein misfolding and inclusion body formation in recombinant *Escherichia coli* cells overexpressing Heat-shock proteins. *J. Biol. Chem.* 271 (19), 11141–11147. doi:10.1074/jbc.271.19.11141
- Venkatesan, M., Semper, C., Skrivergaard, S., Di Leo, R., Mesa, N., Rasmussen, M. K., et al. (2022). Recombinant production of growth factors for application in cell culture. *iScience* 25 (10), 105054. doi:10.1016/j.isci.2022.105054
- Yu, B., Sun, W., Huang, Z., Sun, G., Li, L., Gu, J., et al. (2021). Large-scale preparation of highly stable recombinant human acidic fibroblast growth factor in *Escherichia coli* BL21(DE3) plysS strain. *Front. Bioeng. Biotechnol.* 9, 641505. doi:10.3389/fbioe.2021.641505
- Yu, I. S., Choi, J., Kim, M. K., and Kim, M. J. (2023). The comparison of commercial serum-free media for hanwoo satellite cell proliferation and the role of fibroblast growth factor 2. *Food Sci. Anim. Resour.* 43 (6), 1017–1030. doi:10.5851/kosfa.2023.e68
- Yun, Y. R., Won, J. E., Jeon, E., Lee, S., Kang, W., Jo, H., et al. (2010). Fibroblast growth factors: biology, function, and application for tissue regeneration. *J. Tissue Eng.* 2010, 218142. doi:10.4061/2010/218142
- Zhang, G., Mills, D. A., and Block, D. E. (2009). Development of chemically defined media supporting high-cell-density growth of lactococci, enterococci, and streptococci. *Appl. Environ. Microbiol.* 75 (4), 1080–1087. doi:10.1128/AEM.01416-08
- Zhou, X. X., Li, W. F., Ma, G. X., and Pan, Y. J. (2006). The nisin-controlled gene expression system: construction, application and improvements. *Biotechnol. Adv.* 24 (3), 285–295. doi:10.1016/j.biotechadv.2005.11.001
- Zhou, Y., Chen, P., Shi, S., Li, X., Shi, D., Zhou, Z., et al. (2021). Expression of gallus epidermal growth factor (gEGF) with food-grade *Lactococcus lactis* expression system and its biological effects on broiler chickens. *Biomolecules* 11 (1), 103. doi:10.3390/biom11010103